

# *Trypanosoma cruzi* in dogs: electrocardiographic and echocardiographic evaluation, in Malinalco, State of Mexico

Sandra Díaz González-Vieyra<sup>1</sup>

Ninfa Ramírez-Durán<sup>2</sup>

Ángel H Sandoval-Trujillo<sup>3</sup>

Juan C Vázquez-Chagoyán<sup>1</sup>

Humberto G Monroy-Salazar<sup>1</sup>

Alberto Barbabosa-Pliego<sup>1</sup>

<sup>1</sup>Research Center of Advanced Studies in Animal Health, Veterinary Husbandry School, <sup>2</sup>Medical and Ambiental Microbiology, Research Center of Advanced Studies in Health Science, School of Medicine, Autonomous University of the State of Mexico, Toluca, Mexico;

<sup>3</sup>Department of Biological Systems, Metropolitan Autonomous University, Xochimilco, Mexico City, Mexico

**Abstract:** Chagas disease caused by *Trypanosoma cruzi* is an important public health problem in Latin America. Dogs are considered a risk factor for human Chagas disease, a sentinel for *T. cruzi* infection in endemic regions and an animal model to study pathological aspects of the disease. The potential use of dogs as indicators of human cardiac pathogenicity of local *T. cruzi* strains has been studied insufficiently. We studied electrocardiographic (EKG) and echocardiographic (ECG) alteration frequencies observed in an open population of dogs in Malinalco, Mexico, and determined if such frequencies were statistically associated with *T. cruzi* infection in dogs. Animals (n = 139) were clinically examined and owners were asked to answer a questionnaire about dogs' living conditions. Two commercial serological tests (IHA, ELISA) were conducted to detect anti-*T. cruzi* serum antibodies. Significant differences between seropositive and seronegative animals in cardiomyopathic frequencies were detected through EKG and ECG ( $P < 0.05$ ). Thirty dogs (21.58%) were serologically positive to anti-*T. cruzi* antibodies (to ELISA and IHA assays), of which nine (30%) had EKG and/or ECG alterations. From the remaining 104 (78.42%) seronegative animals, five (4.5%) had EKG and/or ECG abnormalities. Our data support the hypothesis that most EKG and ECG alterations found in dogs from Malinalco could be associated with *T. cruzi* infection. Considering the dog as a sentinel and as an animal model for Chagas disease in humans, our findings suggest that the *T. cruzi* strains circulating in Malinalco have the potential to produce cardiomyopathies in infected humans.

**Keywords:** chagas disease, *Trypanosoma cruzi*, cardiomyopathy, electrocardiography, echocardiography, Malinalco, México

## Introduction

Chagas disease is caused by *Trypanosoma cruzi* and is transmitted by a hematophagous insect vector ("kissing bug") of the Reduviidae family. Approximately 10 million people are infected with *T. cruzi* in 19 countries in Latin America and ~50,000,000 people live at risk of infection.<sup>1</sup> In Mexico, 1.6–5.8 million people may be infected with *T. cruzi*.<sup>2–5</sup> Our group has been studying Chagas disease in the State of Mexico, located under the Tropic of Cancer, where most critical transmission areas in Mexico are located.<sup>4,5</sup> In a previous study we reported *T. cruzi* prevalences of 7.1% and 21% for humans and dogs, respectively,<sup>6</sup> and more recently we have described an epidemiologic study using dogs and triatomines to assess parasite circulation in the Tejupilco municipality.<sup>7</sup> Malinalco is a town located in the south-center region of the State of Mexico, from which no previous reports on *T. cruzi* circulation have been published. However, it is a neighboring area of Tejupilco municipality and shares geographic characteristics with this region. It is also a neighbor of the State of Morelos, where *T. cruzi* circulation has been

Correspondence: Alberto Barbabosa-Pliego  
Centro de Investigación de Estudios Avanzados en Salud Animal, Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma del Estado de México  
Tel +52 722 2965555  
Fax +52 722 2968980  
Email abarbabosa@yahoo.com.mx

previously reported,<sup>6–10</sup> and from Zumpahuacan, from which a pathogenic *T. cruzi* strain has been previously reported and characterized.<sup>11</sup> In the latter study we used dogs as a model to compare the pathogenicity of a regional *T. cruzi* strain vs a reference strain (Sylvio X-10), and described clinical (electrocardiographic and echocardiographic) and pathological (macroscopic and microscopic) cardiac alterations caused by these strains and found that, although the two strains were pathogenic, they had differences in virulence, as reported for other strains.<sup>12</sup> Therefore, epidemiologic studies of Chagas disease in a specific geographical area should consider the pathogenicity of the regional circulating *T. cruzi* strains. Dogs are considered an excellent animal model to study Chagas disease since it mimics the clinical and pathological signs of the disease in humans.<sup>13,14</sup> Accordingly, the objectives of the present study were: first, to use the house-owned dogs from Malinalco to evaluate the feasibility to use them as sentinels to determine prevalence of *T. cruzi* infection in humans in the city and, second, to evaluate the pathogenicity of the circulating strains with an electrocardiographical and echocardiographical epidemiologic study.

## Animals, materials, and methods

### Study area

Malinalco (Figure 1) is located in the south-eastern area of the State of Mexico (between 19°01'58" to 18°45'18" N and 99°35'24" to 99°25'34" W) with an average altitude of 1750 m. It has seasonal climate variations (dry season November–May and rainy season June–October) with an average annual temperature of 20°C. According to the 2005


National Census Program,<sup>15</sup> Malinalco has a population of 22,970 and the main economic activities are agriculture, livestock production, and tourism. According to the 2008 State of Mexico Rabies Vaccination Program, Malinalco has a total population of 2160 house dogs.

### Animals

House-owned dogs ( $n = 139$ ) from Malinalco were studied to assess the prevalence of *T. cruzi* infection in these animals and to study the impact of the infection on dogs' heart conditions. The sample size was calculated with free software "Sample Size Calculator Software",<sup>16</sup> with the following parameters: 95% confidence interval, 5% error, a universe of 2160 house dogs, and 10% estimated prevalence of *T. cruzi* infection. All dogs were evaluated serologically for anti-*T. cruzi* antibodies and electrocardiographically and echocardiographically for heart health. Dogs were included in the study after personalized owners-informed invitation and consent to participate in the study. Owners were asked to answer a brief questionnaire to gather information regarding dogs' breed, weight, sex, age, health status and habits to access the house, existence of triatomines (kissing bug) within or around the house, and pesticide spraying frequency, if any. Additionally, all dogs had to go through a veterinarian physical examination to evaluate their general health status.

### Sample collection

Dogs were handled in the presence of their owners to keep stress and risk to animals and handlers at a minimum. Blood samples (7 mL) were obtained by puncture of the cephalic



**Figure 1** Location of Malinalco, State of Mexico.

vein and collected into tubes (Vacutainer®, Becton-Dickinson, Mexico, DF, Mexico), sera was obtained by centrifugation at 2000 rpm for 20 minutes, and stored at  $-20^{\circ}\text{C}$  until use. Sera samples were analyzed for anti-*T. cruzi* antibodies by IHA (Polychaco, Laboratorio-Lemos SRL, Buenos Aires, Argentina) (with 98% sensitivity and 99% specificity according to the manufacturer's specifications) and ELISA (Laboratorio-Lemos SRL, Buenos Aires, Argentina) Chagas diagnostic kits (with 100% sensitivity and 100% specificity, according to the manufacturer's specifications). All assays were performed according to manufacturer's instructions, except that the HRP-labeled, anti-human IgG antibody in the ELISA kit was replaced with HRP-labeled anti-dog IgG (Koma Biotech, Seoul, Korea) as a secondary antibody.<sup>11</sup> The cut-off value for IHA was set at  $\geq 1:16$  serum dilution and for ELISA at  $0.129 \text{ OD}_{450 \text{ nm}} + 0.1$  (or  $+2\text{-SD}$ ). Samples were considered positive when reactive for both assays.

## Electrocardiography (EKG)

Changes in cardiac rhythm and electrical conduction were monitored in all dogs with an EK-8 electrocardiographic machine (Burdick Stylus, Milton, WI), set at 120V, 60 Hz, 20 amps, and 25 watts, and six leads were considered at 25 mm/second at 1 mV, standardized to 1 cm (Figure 2).

## Echocardiography (ECG)

An ultrasonograph machine (SSD-500, 5 MHz probe; Aloka, Wallingford, CT), with two-dimensional and M-mode


functions was used to analyze ventricular function through short-axis images and right parasternal views (Figure 3).

Statistical analysis was conducted through contingency tables and using EPIDAT (v 3.1, 2006, OPS/OMS). Where the immunologic outcome (seropositive or seronegative) was contrasted vs the following variables: EKG (cardiomyopathy or normal), ECG (cardiomyopathy or normal), age ( $<1$ , 1–2.9, 3–6.9 or  $>7$  years old), housing (indoor or outdoor), sex (male or female), and weight ( $<10$  or  $>10$  kg). Statistical analysis included odd ratios (OR), prevalence ratio (PR) and Chi square ( $\chi^2$ ), with 95% confidence intervals and  $P < 0.05$ .


## Results

Dogs' physical examination showed that all animals were apparently healthy. However, the study found 21.58% (30/139) dogs seropositive to anti-*T. cruzi* IgG antibodies through IHA and ELISA serology tests.

Differences in the frequency of electrocardiographic and echocardiographic abnormalities were found ( $P < 0.05$ ) between seropositive and seronegative animals, where 9/30 (30%) and 5/109 (4.5%) dogs presented electrocardiographic abnormalities (Table 1; Figure 2), and 10/30 (33.3%) and 17/109 (15.5%) dogs resulted with echocardiographic abnormalities for seropositive and seronegative animals, respectively (Table 2; Figure 2). The main abnormalities found in seropositive dogs with echocardiography were: right ventricle dilation in 13.3% (4/30), increased motility and or asyn-



**Figure 2** Electrocardiographic analysis in DII at 25 mm, of seropositive dogs to *Trypanosoma cruzi*, in Malinalco village, State of Mexico. (A) Sinus pacemaker; (B) aberrant conduction complex QRS; (C) sinus arrhythmia, normal in dogs; (D) wave R, electric alternating; (E) wave S deep; (F) normal.



**Figure 3** BM-mode echocardiography of seropositive dogs to *Trypanosoma cruzi*, in the Malinalco village, State of Mexico. **(A)** Increased thickness of the interventricular septum and left ventricular wall; **(B)** asynchronous interventricular septum motility; **(C)** normal.

chony of the interventricular septum in 13.3% (4/30), left ventricle dilated in 13.3% (4/30), and left ventricular wall thickened in 26.6% (8/30). Echocardiographic abnormalities from seronegative dogs were similar to those found in seropositive dogs but at a lower frequency (Table 2; Figure 3).

Factors such as age, housing conditions, weight, and sex were examined for association with seropositivity (Tables 3–6). No statistical differences could be found for these variables, however some tendencies could be observed in the age of the animals and outdoor housing conditions, which were associated with larger prevalences, suggesting that these variables could be considered as possible risk factors in future studies which should be performed at a larger scale. Overall, a larger proportion of female animals were sampled (86:53 female/male), and a larger proportion

(25.5%) of this group was seropositive when compared to males (15.0%). However, no statistical difference was found. The analysis of the influence of dogs’ breed on seropositivity prevalence was not possible because the sample size was not large enough due to the diversity of breeds found in the study.

Discussion

The use of dogs as sentinels to assess human risk of infection was first proposed by Gürtler and colleagues.<sup>13</sup> More recently, our group demonstrated that *T. cruzi* is found in humans, dogs, and triatomines and found a direct correlation of anti-*T. cruzi* sero-prevalence between humans (7%) and dogs (21%) in the southern region of the State of Mexico.<sup>6–8</sup> We have also previously demonstrated that regionally circulating *T. cruzi* strains are pathogenic for dogs, either after natural or experimental infections.<sup>11</sup>

**Table 1** Electrocardiographic alterations found in *Trypanosoma cruzi* seropositive and seronegative dogs from Malinalco village, State of Mexico

Electrocardiography	Dogs			
	Seropositive (n = 30)		Seronegative (n = 109)	
	n	%	n	%
Abnormal	9	30 <sup>a</sup>	5	4.5 <sup>b</sup>
Normal	21	70	104	95.5
	30	100%	109	100%
Atrium-ventricular block (1° grade)	1	3.3	0	0
Supra-ventricular arrhythmia (wandering sinus pacemaker)	3	10	2	1.8
Ventricular arrhythmia (ectopic focus/electric alternating)	2	6.6	1	0.9
Right bundle branch block	3	10	1	0.9
T wave (high)	0	0	1	0.9
R wave (small)	8	26.6	3	2.7

**Notes:**  $\chi^2$  value = 26.836, with 1 df; OR = 15.600; PR = 9.342.  $P < 0.05$ .  $\chi^2$  shows the difference between groups, this difference can be interpreted as an association between seropositivity and risk of developing a cardiomyopathy since OR and PR support analysis risk.

**Table 2** Echocardiographic alterations found in *Trypanosoma cruzi* seropositive and seronegative dogs from Malinalco village, State of Mexico

Echocardiography	Dogs			
	Seropositive (n = 30)		Seronegative (n = 109)	
	n	%	n	%
Abnormal	10	33.3	17	15.5
Normal	20	66.7	92	84.5
	30	100%	109	100%
RV (size increment)	4	13.3		
IS (size increment/asynchronous motility)	4	13.3		
IS (size increment)			4	3.6
LV (size increment)	4	13.3	6	5.5
LVW (size increment)	8	26.6	14	12.8

**Notes:**  $\chi^2$  value = 4.728, with 1 df; OR = 2.705; PR = 2.137.  $P < 0.05$ .  $\chi^2$  shows the difference between groups, this difference can be interpreted as an association between seropositivity and risk of developing a cardiomyopathy since OR and PR support analysis risk.

**Abbreviations:** RV, right ventricle; IS, interventricular septum; LV, left ventricle; LVW, left ventricular wall.



**Table 3** Age associated with seropositivity to *T. cruzi* in dogs from Malinalco village, State of Mexico

Age (years)	Seropositive		Seronegative		Total
	n	%	n	%	
<1	0	0.0	21	100	21
1.1–3	8	16.0	42	84.0	50
3.1–7	10	28.5	25	71.5	35
7.1–10	12	36.36	21	63.63	33
Total	30	21.6	109	78.4	139

**Notes:**  $\chi^2$  value = 11.571, with 1 df; PR for level 1 = 0.0, PR for level 2 = 0.166, PR for level 3 = 0.285, PR for level 4 = 0.363.  $P < 0.05$ . Even though  $\chi^2$  shows differences between groups, this difference can only be interpreted as a tendency, since PR does not support analysis risk.

Here we report a 21.58% seroprevalence anti-*T. cruzi* in dogs found in Malinalco (a tourist town in a municipality of the southern region of the State of Mexico, with no previous seroprevalence reports). These findings agree with our previous reports in the southern region of the State of Mexico, and considering climatic, geographic, and ecological similarities of the regions studied, this outcome is not surprising. Here we also report the association of seroprevalence of IgG anti-*T. cruzi* antibodies found in dogs from Malinalco and the occurrence of electrocardiographic and echocardiographic abnormalities in this species. We also analyzed some variables that have been associated with the seroprevalence of dogs to *T. cruzi*, as risk factors; such as age, weight, sex, and housing.

Through EKG and ECG we found that *T. cruzi* seropositive animals develop cardiac abnormalities with higher frequencies than seronegative dogs. This finding was expected, since it has been previously demonstrated by us and others that this parasite induces cardiac abnormalities in dogs, and that some *T. cruzi* strains found in the south of the State of Mexico are cardiopathogenic for dogs.<sup>7,11</sup> It is of general knowledge that cardiomyopathies can be found through EKG and ECG among apparently healthy, uninfected, open populations of dogs. The type and frequencies of these pathologies may vary according to the

**Table 4** Housing associated to seropositivity to *Trypanosoma cruzi* in dogs from Malinalco village, State of Mexico

Dog housing	Seropositive		Seronegative		Total
	n	%	n	%	
Indoors	2	6.5	29	93.5	31
Outdoors	28	25.92	80	74.07	108
Total	30	21.6	109	78.4	139

**Notes:**  $\chi^2$  value = 5.39, with 1 df; OR = 0.19; PR = 0.25.  $P < 0.05$ . Even though  $\chi^2$  shows the difference between groups, this difference can only be interpreted as a tendency, since OR and PR do not support analysis risk.

**Table 5** Body weight associated to seropositivity to *Trypanosoma cruzi* in dogs from Malinalco village, State of Mexico

Weight (kg)	Seropositive		Seronegative		Total (%)
	n	%	n	%	
≤10.0	2	9.5	19	90.5	21
>10	28	24.6	90	75.4	118
Total	30	21.6	109	78.4	139

**Notes:**  $\chi^2$  value = 2.125, with 1 df; OR = 0.338; PR = 0.382.  $P < 0.05$ . No difference was found between groups.

breed and age.<sup>17,18</sup> Here we report electrocardiographic and echocardiographic alterations in an open population of dogs and analyzed their relationship with the seroprevalence of IgG antibodies against *T. cruzi*. The prevalence of cardiac abnormalities among house dogs living in Malinalco was 10.07%. However, as predicted, cardiomyopathic frequencies were different ( $P < 0.05$ ) among anti-*T. cruzi* seropositive (30%) and seronegative (4.5%) animals (Tables 1 and 2). These findings support our previous report in which we concluded that *T. cruzi* strains circulating in the southern region of the State of Mexico are pathogenic to dogs.<sup>11</sup>

Dogs have been considered sentinels for *T. cruzi* circulation in endemic areas,<sup>12</sup> and it has also been proposed as a model for Chagas disease since infected dogs mimic some of the signs and pathologies seen in humans,<sup>12–14</sup> therefore we considered it interesting to evaluate the dog as a sentinel, so that in future it could be used as a tool to estimate human risk of developing cardiomyopathies derived from infection with local *T. cruzi* strains in different endemic regions. For instance, in a previous study, we reported a correlation of 1:3 between human (7.1%) and dog (21%) anti-*T. cruzi* seroprevalences in the southern region of the State of Mexico.<sup>6</sup> Therefore, assuming that the same correlation exists in cardiac abnormalities due to *T. cruzi* infection, we could predict that from an estimated population of 2160 house dogs and 22,970 people living in Malinalco,<sup>15</sup> there should be around 444 and 1608 individuals infected, and

**Table 6** Sex associated to seropositivity to *Trypanosoma cruzi* in dogs from Malinalco village, State of Mexico

Sex	Seropositive		Seronegative		Total
	n	%	n	%	
Male	8	15.0	45	85.0	53
Female	22	25.5	64	74.5	86
Total	30	21.6	109	78.4	139

**Notes:**  $\chi^2$  value = 2.125, with 1 df; OR = 0.517; PR = 0.590.  $P < 0.05$ . No difference was found between groups.

113 and 410, individuals with cardiomyopathies derived from *T. cruzi* infection for dogs and humans, respectively. This estimate is merely speculative and further research is necessary to obtain conclusive evidence, however this approach could be helpful as an indirect method to estimate epidemiologic information necessary for the design of official public health programs.

When we analyzed some variables that could be participating as risk factors of infection, we were able to find some tendencies of association but not statistically significant correlations. Previous studies have reported age as an accumulative risk factor of infection.<sup>19–21</sup> In this study we observed a strong tendency of age to be associated to seroprevalence, however no statistical significance was found ( $P > 0.05$ ) (Table 3). Future studies should include a larger sample size, to better estimate age as a risk factor. Another factor that has been reported as a risk factor is dog housing (indoor–outdoor). In this study we were not able to demonstrate that housing (indoor vs outdoor), is a risk factor in this endemic region. But, we observed that most owned dogs involved in the present study (77.69%), are regularly kept outside the house or are able to go in and out at their own will; all of these dogs were considered outdoor animals, and the animals that were kept mostly inside the house were included in the indoor group; the prevalences were 6.5% and 25.92% for indoor and outdoor groups, respectively (Table 4). A tendency to have a lower seroprevalence was observed in dogs that were maintained exclusively indoors when compared to outdoor dogs. Most likely a larger sample size is necessary to demonstrate the differences associated with the housing risk factor. On the other hand, other variables analyzed, such as dog weight and sex, do not seem to play important roles in the likelihood of infection (Tables 5 and 6).

We conclude that a relatively high anti-*T. cruzi* antibody seroprevalence in house-owned dogs from Malinalco village is related to elevated frequencies of electrocardiographic and echocardiographic abnormalities. Further studies should be conducted to fully evaluate dogs as sentinels of infection-induced, cardiac abnormalities in humans.

Furthermore, our observations emphasize the need of state health agencies to conduct more aggressive, epidemiologic surveillance programs and implement vector control strategies in the State of Mexico, since the seroprevalence found in dogs suggests that there is a latent risk of human infection in Malinalco.

## Acknowledgment

This work was supported by grants of (FE039/2008) SIyEA of Universidad Autónoma del Estado de México.

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Schofield CJ, Jeannine J, Salvatella R. The future of Chagas disease control. *Trends Parasitol.* 2006;22:583–588.
- Velasco-Castrejón O, Guzmán-Bracho C. [Importance of Chagas disease in Mexico]. *Rev Latinoam Microbiol.* 1986;28:275–283. Spanish.
- Velazco-Castrejón O. [Chagas disease seroepidemiology in Mexico]. *Mexico's Public Health Journal.* 1992;34:186–196. Spanish.
- Guzmán-Bracho C. Epidemiology of Chagas disease in Mexico: an update. *Trends Parasitol.* 2001;17:372–376.
- Cruz-Reyes A, Pickering-López JM. Chagas disease in México: an analysis of geographical distribution during the past 76 years – a review. *Mem Inst Oswaldo Cruz.* 2006;101:345–354.
- Estrada-Franco JG, Vandanajay B, Díaz-Albiter H, et al. Human *Trypanosoma cruzi* infection and seropositivity in dogs, México. *Emerg Infect Dis.* 2006;12:624–630.
- Barbabosa-Pliego A, Campos P, Olivares D, et al. Prevalence of *Trypanosoma cruzi* in dogs (*Canis familiaris*) and triatomines during 2008 in a sanitary region of the State of Mexico, Mexico. *Vector Borne Zoonotic Dis.* 2011;11:151–156.
- Medina-Torres I, Vázquez-Chagoyán JC, Rodríguez-Vivas RI, Montes de Oca-Jiménez R. Risk factors associated with triatomines and its infection with *Trypanosoma cruzi* in rural communities from the southern region of the State of Mexico, Mexico. *Am J Trop Med Hyg.* 2010;82:49–54.
- Cortés Jiménez M, Noguera Torres B, Alejandro Aguilar R, Isita Torneli L, Ramírez Moreno E. Frequency of triatomines infected with *Trypanosoma cruzi* collected in Cuernavaca city, Morelos, México. *Rev Latinoam Microbiol.* 1996;38:115–119.
- Bautista NL, García de la Torre GS, de Haro-Arteaga I, Salazar-Shettino PM. Importance of *Triatoma pallidipennis* (Hemiptera: Reduviidae) as a vector of *Trypanosoma cruzi* (Kinetoplastida: Trypanosomatidae) in the state of Morelos, Mexico, and possible ecotypes. *J Med Entomol.* 1999;36:233–235.
- Barbabosa-Pliego A, Díaz-Albiter HM, Ochoa-García L, et al. *Trypanosoma cruzi* circulating in the southern region of the State of Mexico (*Zumpahuacan*) are pathogenic: a dog model. *Am J Trop Med Hyg.* 2009;81:390–395.
- Andrade ZA. The canine model of Chagas disease. *Mem Inst Oswaldo Cruz.* 1984;79:77–83.
- Gürtler RE, Solard ND, Lauricela MA, Haedo M. Dynamic of transmission of *Trypanosoma cruzi* in a rural area of Argentina. III Persistence of *T. cruzi* parasitemia among canine reservoirs in a two-years follow-up. *Rev Inst Med Trop Sao Paulo.* 1986;28:213–219.
- Caliari MV, Machado R, Lana M, et al. Quantitative analysis of cardiac lesion in chronic canine Chagasic cardiomyopathy. *Rev Inst Med Trop Sao Paulo.* 2002;44:273–278.
- INEGI. 2005. National Institute of Statistics, Geography and Informatics (INEGI). Available at: <http://www.inegi.org.mx>. Accessed August 11, 2011.
- Roasft. Roasoft. Sample size calculator. Roasoft, Inc. 2004. Available at: <http://www.roasoft.com/samplesize.html>. Accessed January 28, 2008.
- Dukes-McEwan J, Borgarelli M, Tidholm A, Vollmar AC, Häggström J. Proposed guidelines for the diagnosis of canine idiopathic dilated cardiomyopathy. *J Vet Cardiol.* 2003;5:7–19.
- Oyama MA, Sisson DD, Solter PF. Prospective screening for occult cardiomyopathy in dogs by measurement of plasma atrial natriuretic peptide, B-type natriuretic peptide, and cardiac troponin-I concentrations. *Am J Vet Res.* 2007;68:42–47.

19. Gürtler RE, Cecere MC, Lauricella MA, et al. Incidence of *Trypanosoma cruzi* infection among children following domestic reinfestation after insecticide spraying in rural northwestern Argentina. *Am J Trop Med Hyg.* 2005;73:95–103.
20. Diosque P, Padilla AM, Cimino RO, et al. Chagas disease in rural areas of Chaco Province, Argentina: epidemiologic survey in humans, reservoirs, and vectors. *Am J Trop Med Hyg.* 2004;71:590–593.
21. Pascon da EJP, Pereira NG, Sousa GM, Junior DP, Camacho AA. Clinical characterization of chagasic cardiomyopathy in dogs. *Pesq Vet Bras.* 2010;30:115–120.

#### Research and Reports in Tropical Medicine

### Publish your work in this journal

Research and Reports in Tropical Medicine is an international, peer-reviewed, open access journal publishing original research, case reports, editorials, reviews and commentaries on all areas of tropical medicine, including: Diseases and medicine in tropical regions; Entomology; Epidemiology; Health economics issues; Infectious disease; Laboratory

Submit your manuscript here: <http://www.dovepress.com/research-and-reports-in-tropical-medicine-journal>

science and new technology in tropical medicine; Parasitology; Public health medicine/health care policy in tropical regions; and Microbiology. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress