

# Effects of *CYP2D6* and *CYP3A5* polymorphisms on tamoxifen and its metabolites in Thai breast cancer patients

Wanaporn  
Charoenchokthavee<sup>1</sup>  
Nutthada Aareepium<sup>2</sup>  
Duangchit Panomvana<sup>2</sup>  
Virote Sriuranpong<sup>2</sup>

<sup>1</sup>Department of Pharmacy Practice, Faculty of Pharmaceutical Science, Chulalongkorn University, <sup>2</sup>Medical Oncology Unit, Department of Medicine, Faculty of Medicine, Chulalongkorn University, Pathumwan, Bangkok, Thailand

**Purpose:** This study aimed to determine the effects of *CYP2D6* and *CYP3A5* polymorphisms on the levels of tamoxifen (TAM) and its metabolites in the plasma of breast cancer patients. The protocol was designed to test the associations between *CYP2D6*, *CYP3A5* genotypes and phenotypes (extensive metabolizer [EM], intermediate metabolizer [IM] and poor metabolizer [PM]) and TAM, *N*-desmethyl tamoxifen (NDMT), endoxifen (END) and 4-hydroxytamoxifen (4OHT) concentrations.

**Patients and methods:** One hundred and thirty-four Thai breast cancer patients from the Thai Tamoxifen Project undergoing TAM treatment who met the inclusion/exclusion criteria were recruited. Plasma samples were assessed for the concentrations of TAM and its metabolites using high-performance liquid chromatography. The data are presented as actual values and metabolic ratios (MR). The hypotheses were tested using Kruskal–Wallis or Mann–Whitney *U* test, including the simple main effects analysis.

**Results:** The patients had stage 0–IV breast cancer. The mean age and body mass index were 51.6±11.6 years and 24.0±4.3, respectively. Also, 53.0% of them were premenopausal, 10.4% were perimenopausal and 36.6% were postmenopausal, while 23.1% were *CYP2D6*-EM/*CYP3A5*-EM and 20.9% carried only *CYP2D6* and *CYP3A5* incomplete alleles. The median concentrations of TAM, NDMT, END and 4OHT were 374.7 (interquartile range [IQR] 230.2) ng/mL, 1,064.9 (IQR 599.6) ng/mL, 54.5 (IQR 52.5) ng/mL and 5.0 (IQR 3.1) ng/mL, respectively. MR (TAM-NDMT) and MR (NDMT-END) were statistically different ( $p=0.013$  and  $p=0.014$ , respectively), while MR (4OHT-END) was not statistically different within the *CYP2D6* phenotype ( $p=0.594$ ). MR (TAM-4OHT) was not statistically different within the *CYP2D6* phenotype ( $p=0.079$ ), but it was potentially different from *CYP3A5*-PM ( $p=0.056$ ). None of the MR was statistically different within the *CYP3A5* phenotype.

**Conclusion:** *CYP2D6* polymorphisms appear to affect END concentration through an NDMT subpathway and potentially affect 4OHT concentrations through a 4OHT subpathway in *CYP3A5*-PM group.

**Keywords:** endoxifen, cytochrome P450, single nucleotide polymorphisms, pharmacogenetics, pharmacogenomics, human

## Introduction

Tamoxifen (TAM) has been widely used for breast cancer prevention and treatment.<sup>1</sup> Interindividual variability because of the effects of single-nucleotide polymorphisms in genes encoding cytochrome P450 (CYP450) enzymes involved in Phase I of the TAM metabolic pathway has been shown in many studies.<sup>2–5</sup> The CYP450 enzymes (CYP2D6, CYP2C9, CYP2C19, CYP3A4 and CYP3A5) are essential for the transformation of TAM from a prodrug form to its active metabolites (endoxifen [END] and 4-hydroxytamoxifen [4OHT]).<sup>1,6,7</sup> The END and 4OHT

Correspondence: Wanaporn Charoenchokthavee  
Pharmaceutical Care Unit, Department of Pharmacy, Vajira Hospital, Faculty of Medicine Vajira Hospital, Navamindradhiraj University, 681 Samsen Road, Vajira, Dusit, Bangkok 10300, Thailand  
Email wanaporn@edu.vajira.ac.th

active metabolites demonstrate a 30- to 100-fold higher suppression of cell proliferation and nearly 100-fold higher affinity for the estrogen receptor, compared with TAM. However, the 4OHT levels are found to be five to ten times lower than those of END in the plasma.<sup>6</sup> END can be formed through two subpathways: the main pathway involves *N*-desmethyl tamoxifen (NDMT subpathway; 90%) and the other pathway is through 4OHT (4OHT subpathway; 10%).<sup>1,2,8–10</sup> Previous studies have investigated the associations between TAM and its metabolites, including studying the relevant CYP450 enzyme activity in the TAM metabolic subpathways,<sup>9–11</sup> but conclusive results have not been obtained because of the heterogeneity of the studies.<sup>12,13</sup> The prevalence and type of incomplete functional allele (null allele and reduced allele) that is involved in CYP450 enzyme activity are different in different populations;<sup>14</sup> the prevalence of *CYP2D6*\*4 (null allele) has been found to be higher in Caucasians<sup>15</sup> than in Asians<sup>16</sup> or Thais,<sup>17</sup> while that of *CYP2D6*\*10 (reduced allele) has been found to be higher in Asians,<sup>16</sup> including Thais,<sup>18</sup> than in Caucasians<sup>15</sup> and *CYP3A5*\*3 (null allele) is the major allele in Caucasians<sup>15</sup> and Asians,<sup>16</sup> including Thais.<sup>18</sup> Previous research has suggested that the low activities of CYP2D6 and CYP3A5 enzymes account for 25%–55% and 40%–50% of the polymorphisms, respectively, in Thai breast cancer patients.<sup>17–19</sup> Early researches suggested that *CYP2D6*\*10/\*10 patients had shorter disease-free survival than heterozygous *CYP2D6*\*10 ( $p=0.046$ )<sup>17</sup> and lower END concentrations than those patients with *CYP2D6*\*1/\*10 and *CYP2D6*\*1/\*1 ( $p=0.045$ )<sup>19</sup> among Thai breast cancer patients; however, the result was not completely generalized to target population because of the limited sample size.<sup>19</sup> Furthermore, the associations between *CYP3A5* polymorphisms and levels of TAM and its metabolites have never been explored in Thai breast cancer patients, even though a high prevalence of *CYP3A5* incomplete allele (*CYP3A5*\*3) has been suggested in a previous study.<sup>18</sup> The purpose of the present study was to determine the associations of *CYP2D6* and *CYP3A5* polymorphisms and the concentrations of TAM and its metabolites in large numbers of Thai breast cancer patients undergoing TAM treatment.

## Patients and methods

### Patients and samples preparation

A total of 134 Thai breast cancer patients undergoing TAM treatment were recruited from the Thai Tamoxifen Project.<sup>18,20</sup> In brief, the patients took 20 mg of TAM once daily for at least 2 months to ensure a steady-state concentration and

visited the outpatient clinic at King Chulalongkorn Memorial Hospital between February and March 2015. All patients were 18 years or older, with normal hepatic and renal functions (aspartate aminotransferase and alanine aminotransferase  $\leq 2$  upper normal limit, serum creatinine  $\leq 1.2$  mg/dL) in the previous 4 weeks.<sup>18,20</sup> Medication nonadherence was evaluated through self-reporting. Medication records were screened for drug–drug interactions by a clinical pharmacist. Patients who reported  $<80\%$  adherence, showed an evident drug–drug interaction or were diagnosed for psychiatric illness/cognitive impairment were excluded from the study. A sample size calculation was performed using G\*Power version 3.1 program<sup>21</sup> using the priori method<sup>22</sup> with type-I errors 0.05 (two-tailed) and type-II errors 0.2. Ten milliliters of whole blood was drawn from each patient by a professional nurse and stored in a Vacutainer® (K<sub>2</sub>EDTA [di-potassium salts of ethylenediaminetetraacetic acid]; 10 mL) tube (BD, Franklin Lakes, NJ, USA).<sup>18</sup> The DNA extraction for assessing *CYP2D6* and *CYP3A5* polymorphisms and the determination of the polymorphisms have been described in previous research.<sup>18</sup>

The plasma section was separated from the collected whole blood and stored at  $-80^{\circ}\text{C}$  until use. Then 10  $\mu\text{L}$  of internal standard (IS; Mexiletine 5 mg/mL; Sigma-Aldrich, Singapore) was added to 1 mL of plasma sample, followed by 1 mL of acetonitrile (RCI Labscan, Bangkok, Thailand) and 500  $\mu\text{L}$  of methanol (Fisher Scientific, Loughborough, UK) in a 15 mL centrifuge tube (Corning Incorporated, Corning, NY, USA). The tube was capped and vortexed for 10 minutes and subsequently centrifuged twice at 3,000 rpm ( $4^{\circ}\text{C}$ ) for 30 and 10 minutes, respectively. The supernatant was filtered through a 0.22  $\mu\text{m}$  nylon filter and thereafter derivatized using an ultraviolet lamp at a wavelength of 366 nm for 20 minutes before being injected into a high-performance liquid chromatography (HPLC) column.

### Quantification of TAM and its metabolites

HPLC system with a fluorescent (FLU) detector: The concentrations of TAM and its metabolites concentrations in plasma were quantified using reverse-phase HPLC with a FLU detector. The HPLC-FLU method validation and the plasma extraction protocol were modified from the methods developed by Zhu et al<sup>23</sup> and Areepium et al.<sup>19</sup> The HPLC-FLU system was set as follows: Prostar (model 363) with autosampler (model 410) and column oven (model 510) with fluorescence detector and Varian Star software (Agilent Technologies, Santa Clara, CA, USA), column: Luna 5U C18 (2) 100 A, 250 $\times$ 4.6 mm ( $35^{\circ}\text{C}$ ; Phenomenex, Torrance, CA, USA), mobile phase: 1% trimethylamine and methanol (19:81 %V/V) with flow

rate 1.1 mL/min. TAM and metabolites standards: TAM (Fluka, Singapore), (*E/Z*)-4OHT (Fluka), NDMT (Sigma-Aldrich) and (*E/Z*)-END (Sigma-Aldrich) and IS: mexiletine (Sigma-Aldrich).

### Standards and chemicals

The methanolic standard stock solutions of TAM, NDMT, END, 4OHT and mexiletine (IS) were prepared using powder dissolution to obtain 5 mg/mL of mexiletine, 0.01 mg/mL of END and 4OHT, 0.1 mg/mL of TAM and 0.3 mg/mL of NDMT. The working solutions were prepared from each stock solution with a sufficient volume of methanol to obtain six non-zero standard solutions containing TAM (25, 50, 100, 500, 750 and 1,000 ng/mL), NDMT (25, 50, 100, 500, 750 and 1,000 ng/mL), END (5, 10, 50, 75, 100 and 300 ng/mL) and 4OHT (2.5, 5.0, 7.5, 12.5, 25 and 50 ng/mL). All stock solutions were stored at  $-20^{\circ}\text{C}$  and protected from light. Trimethylamine HPLC grade (Sigma-Aldrich), methanol HPLC grade (Fisher Scientific), acetonitrile HPLC grade (RCI Labscan) and ultrapure analytical grade type I water were used for the mobile phase and plasma extraction.

### Method validation and calibration curve

The chromatogram of a blank sample (six sources of plasma) with the IS was compared with the chromatograms of TAM 750 ng/mL, NDMT 750 ng/mL, END 75 ng/mL and 4OHT 25 ng/mL in blank plasma to assess the selectivity of each metabolite. The retention times of IS, TAM, NDMT, END and 4OHT were 5.0, 26.1, 19.6, 6.9 and 7.3 minutes, respectively. The recovery of those metabolites was  $118.0\% \pm 12.0\%$ ,  $123.7\% \pm 7.5\%$ ,  $130.7\% \pm 17.5\%$  and  $98.0\% \pm 13.1\%$  for TAM 750 ng/mL, NDMT 750 ng/mL, END 100 ng/mL and 4OHT 100 ng/mL, respectively, and  $122.3\% \pm 23.8\%$ ,  $109.0\% \pm 23.8\%$ ,  $72.3\% \pm 13.4\%$  and  $108.4\% \pm 20.1\%$  for TAM 50 ng/mL, NDMT 100 ng/mL, END 50 ng/mL and 4OHT 50 ng/mL, respectively. The coefficient of determination ( $R^2$ ) of those calibration curves was 0.991, 0.995, 0.990 and 0.991 for TAM, NDMT, END and 4OHT, respectively. The accuracy of detection of TAM 100 ng/mL, NDMT 750 ng/mL, END 75 ng/mL and 4OHT 25 ng/mL was 85.0–115.0, 637.5–862.5, 63.8–86.3 and 21.3–28.8 ng/mL, respectively. The coefficients of variation of TAM, NDMT, END and 4OHT were 11.6%, 12.4%, 14.0% and 19.0%, respectively. The concentration of TAM and its metabolites in the plasma samples was determined from the constructed calibration curves and is presented in ng/mL. The metabolic ratios (MR) of TAM (ng/mL)/NDMT (ng/mL): MR (TAM-NDMT),

NDMT (ng/mL)/END (ng/mL): MR (NDMT-END), TAM (ng/mL)/4OHT (ng/mL): MR (TAM-4OHT) and 4OHT (ng/mL)/END (ng/mL): MR (4OHT-END) were calculated for use in hypothesis testing.

### Data analysis

Kolmogorov–Smirnov and Shapiro–Wilk tests were performed to test for normality. Mean and standard deviation (SD) were used to describe normally distributed data, while median and interquartile range were used to describe non-normally distributed data. The genetic data were described as genotypes (*CYP2D6*\*1, *CYP2D6*\*2, *CYP2D6*\*4, *CYP2D6*\*10, *CYP3A5*\*1 and *CYP3A5*\*3) and phenotypes (extensive metabolizer [EM], intermediate metabolizer [IM] and poor metabolizer [PM]). The phenotypes were classified using a conventional method which is described in a previous report.<sup>18</sup> Briefly, the genotypes were firstly divided into two groups (complete and incomplete alleles). The complete alleles included all wild-type alleles (*CYP2D6*\*1, *CYP2D6*\*2 and *CYP3A5*\*1), while the incomplete alleles consisted of *CYP2D6*\*4 (null allele), *CYP2D6*\*10 (reduced allele) and *CYP3A5*\*3 (null allele). Secondly, the phenotype was classified as EM if at least one wild-type allele was present. It was classified as IM if at least one reduced allele was present. The rest of them were classified as PM.

Kruskal–Wallis test and Mann–Whitney *U* test were used to perform hypothesis testing. If conflicting results were produced, the interaction effect was taken into account and a simple main effects analysis<sup>24</sup> was performed to confirm the results.

### Ethical approval

This study was approved by the Institutional Review Board of the Faculty of Medicine, Chulalongkorn University (IRB No. 406/57). Written informed consent was obtained from all individual participants included in the study.

### Results

The subjects consisted of 134 breast cancer patients undergoing TAM treatment. The demographic data of the patients have been described in previous research.<sup>18</sup> In brief, these patients demonstrated stage 0–IV breast cancer. Their mean age and body mass index (BMI) were  $51.6 \pm 11.6$  years and  $24.0 \pm 4.3$ , respectively. Of these patients, 53.0% were premenopausal, 10.4% were perimenopausal and 36.6% were postmenopausal. The majority of the patients were estrogen positive/progesterone positive (71.6%). The median

duration of TAM administration was 17.2 (interquartile range 16.1) months. *CYP2D6*-EM/*CYP3A5*-EM was found in 23.1% of the patients, while 20.9% carried *CYP2D6* and *CYP3A5* incomplete alleles (*CYP2D6*\*4, *CYP2D6*\*10 or *CYP3A5*\*3).<sup>18</sup> One patient was excluded from determination of levels of TAM and its metabolites because of extremely low levels of TAM (44.9 ng/mL). The descriptive data concerning the concentration of TAM and its metabolites are shown in Table 1.

## CYP2D6 polymorphisms and TAM and its metabolites

The concentrations of TAM and its metabolites were not significantly affected by *CYP2D6* genotype or phenotype, with the exception that the concentration of TAM was found to be significantly affected by *CYP2D6* phenotype ( $p=0.031$ ; Table 2). Patients demonstrating *CYP2D6*-PM had low TAM, NDMT and END concentrations compared with those with *CYP2D6*-EM and *CYP2D6*-IM, while *CYP2D6*-IM resulted in lower END and 4OHT concentrations compared with *CYP2D6*-EM, but had higher TAM and NDMT concentrations than *CYP2D6*-EM (Table 2).

**Table 1** Descriptive data concerning the concentrations of TAM and its metabolites (ng/mL)

TAM and its metabolites (N=133)	Mean±SD	Median (IQR)	Min–max
TAM (ng/mL)	406.7±14.9	374.7 (230.2)	82.5–984.2
NDMT (ng/mL)	1,149.7±486.7	1,064.9 (599.6)	80.8–2,543.8
END (ng/mL)	68.5±4.6	54.5 (52.5)	2.3–443.8
4OHT (ng/mL)	5.8±0.3	5.0 (3.1)	2.1–21.7

**Abbreviations:** 4OHT, 4-hydroxytamoxifen; END, endoxifen; IQR, interquartile range; NDMT, N-desmethyl tamoxifen; SD, standard deviation; TAM, tamoxifen.

These inconclusive findings suggest that considering the original values of the concentrations of TAM and its metabolites might not accurately reveal the effects of *CYP2D6* polymorphisms because of the effects of several TAM-metabolizing enzymes involved in END transformation including the two subpathways for END formation (through NDMT and 4OHT).<sup>18</sup> Therefore, the concentration values of TAM and its metabolites were converted to MR to compare the baseline concentration of the input metabolite with that of the output metabolite in each subpathway of the TAM-metabolizing process. The MR (TAM-NDMT) was significantly different between different *CYP2D6* genotypes ( $p=0.000$ ) and phenotypes ( $p=0.013$ ). MR (NDMT-END) and MR (TAM-4OHT) were significantly different between different *CYP2D6* phenotypes ( $p=0.014$  and  $p=0.017$ , respectively), but were not significantly different between different *CYP2D6* genotypes ( $p=0.078$  and  $p=0.094$ , respectively), while MR (4OHT-END) was not statistically different among *CYP2D6* phenotypes or genotypes ( $p=0.594$  and  $p=0.470$ , respectively; Table 3).

## CYP3A5 polymorphisms and TAM and its metabolites

The concentrations of TAM and its metabolites were not significantly different between different *CYP3A5* genotypes or phenotypes. The MR of the metabolites was also not significantly different between different *CYP3A5* genotypes ( $p=0.307$ ,  $p=0.786$ ,  $p=0.742$  and  $p=0.642$ ) or phenotypes ( $p=0.831$ ,  $p=0.657$ ,  $p=0.508$  and  $p=0.400$ ), including those for MR (TAM/NDMT), MR (NDMT/END), MR (TAM/4OHT) and MR (4OHT/END), respectively (Table 4).

**Table 2** The concentrations of TAM and its metabolites (ng/mL) among different *CYP2D6* polymorphisms

<i>CYP2D6</i>	n	TAM	NDMT	END	4OHT
<b>Genotype</b>	<b>133</b>	<b><math>p=0.122</math></b>	<b><math>p=0.078</math></b>	<b><math>p=0.226</math></b>	<b><math>p=0.491</math></b>
<i>CYP2D6</i> *1/*1	13	429.6 (266.5)	948.5 (466.0)	69.9 (71.8)	5.9 (9.0)
<i>CYP2D6</i> *1/*10	21	324.7 (239.5)	997.8 (741.6)	61.2 (79.9)	6.5 (3.4)
<i>CYP2D6</i> *10/*10	72	375.5 (231.9)	1,095.8 (597.4)	51.1 (46.0)	4.5 (2.5)
<i>CYP2D6</i> *2/*2	5	355.7 (171.2)	1,085.9 (838.9)	56.2 (51.2)	4.6 (2.5)
<i>CYP2D6</i> *2/*10	13	283.3 (174.9)	994.6 (462.0)	46.3 (57.3)	5.4 (3.1)
<i>CYP2D6</i> *1/*2	3	481.7 (0.0)	1,170.9 (0.0)	113.5 (0.0)	5.1 (0.0)
<i>CYP2D6</i> *4/*4	2	238.4 (0.0)	803.2 (0.0)	41.9 (0.0)	6.2 (0.0)
<i>CYP2D6</i> *4/*10	4	486.5 (138.7)	1,924.6 (730.5)	48.2 (29.4)	7.0 (9.6)
<b>Phenotype</b>	<b>133</b>	<b><math>p=0.031^*</math></b>	<b><math>p=0.052</math></b>	<b><math>p=0.128</math></b>	<b><math>p=0.156</math></b>
EM	55	373.1 (224.7)	1,031.0 (503.6)	64.8 (69.3)	5.8 (3.3)
IM	76	382.9 (232.5)	1,113.1 (612.0)	50.4 (43.5)	4.6 (3.2)
PM	2	238.4 (0.0)	803.2 (0.0)	41.9 (0.0)	6.2 (0.0)

**Note:** The concentrations of TAM and its metabolites are presented as median (interquartile range).  $p=p$  value from Kruskal–Wallis test, \* $p<0.05$ .

**Abbreviations:** 4OHT, 4-hydroxytamoxifen; EM, extensive metabolizer; END, endoxifen; IM, intermediate metabolizer; NDMT, N-desmethyl tamoxifen; PM, poor metabolizer; TAM, tamoxifen.

**Table 3** The MR of TAM and its metabolites among different *CYP2D6* polymorphisms

<i>CYP2D6</i>	n	MR (TAM-NDMT)	MR (NDMT-END)	MR (TAM-4OHT)	MR (4OHT-END)
<b>Genotype</b>	<b>133</b>	<b>p=0.000*</b>	<b>p=0.078</b>	<b>p=0.094</b>	<b>p=0.470</b>
<i>CYP2D6</i> *1/*1	13	0.51 (0.23)	11.72 (22.57)	61.05 (52.86)	0.08 (0.33)
<i>CYP2D6</i> *1/*10	21	0.38 (0.11)	15.19 (20.04)	56.18 (60.29)	0.08 (0.16)
<i>CYP2D6</i> *10/*10	72	0.34 (0.09)	23.57 (25.79)	77.07 (49.80)	0.10 (0.09)
<i>CYP2D6</i> *2/*2	5	0.32 (0.09)	16.84 (13.87)	63.50 (52.33)	0.08 (0.10)
<i>CYP2D6</i> *2/*10	13	0.34 (0.10)	19.53 (20.39)	60.66 (18.32)	0.11 (0.06)
<i>CYP2D6</i> *1/*2	3	0.40 (0.00)	10.96 (0.00)	92.99 (0.00)	0.04 (0.00)
<i>CYP2D6</i> *4/*4	2	0.30 (0.00)	21.56 (0.00)	40.48 (0.00)	0.17 (0.00)
<i>CYP2D6</i> *4/*10	4	0.27 (0.10)	39.96 (33.71)	69.34 (119.84)	0.15 (0.30)
<b>Phenotype</b>	<b>133</b>	<b>p=0.013*</b>	<b>p=0.014*</b>	<b>p=0.017*</b>	<b>p=0.594</b>
EM	55	0.10 (0.03)	15.19 (15.55)	61.05 (34.92)	0.08 (0.13)
IM	76	0.09 (0.02)	24.06 (26.15)	76.80 (49.8)	0.10 (0.09)
PM	2	0.08 (0.00)	21.56 (0.00)	40.48 (0.00)	0.17 (0.00)

**Note:** The concentrations of TAM and its metabolites are presented as median (interquartile range).  $p$ = $p$  value from Kruskal–Wallis test, \* $p$ <0.05.

**Abbreviations:** 4OHT, 4-hydroxytamoxifen; EM, extensive metabolizer; END, endoxifen; IM, intermediate metabolizer; MR, metabolic ratios; NDMT, N-desmethyl tamoxifen; PM, poor metabolizer; TAM, tamoxifen.

**Table 4** The concentrations and MR of TAM and its metabolites among different *CYP3A5* polymorphisms

<i>CYP3A5</i>	n	TAM (ng/mL)	NDMT (ng/mL)	END (ng/mL)	4OHT (ng/mL)
<b>Genotype</b>	<b>133</b>	<b>p=0.771</b>	<b>p=0.680</b>	<b>p=0.844</b>	<b>p=0.223</b>
<i>CYP3A5</i> *1/*1	18	363.76 (199.20)	1,057.48 (682.44)	51.12 (67.14)	4.37 (2.35)
<i>CYP3A5</i> *1/*3	64	364.20 (249.26)	1,049.24 (501.11)	53.42 (54.21)	5.03 (2.92)
<i>CYP3A5</i> *3/*3	51	384.02 (235.09)	1,085.89 (724.63)	56.22 (49.41)	5.39 (3.18)
<b>Phenotype</b>	<b>133</b>	<b>p=0.493</b>	<b>p=0.451</b>	<b>p=0.813</b>	<b>p=0.100</b>
EM	82	363.76 (234.85)	1,049.24 (553.93)	53.42 (57.28)	4.80 (2.86)
PM	51	384.02 (235.09)	1,085.89 (724.63)	56.22 (49.41)	5.39 (3.18)
<i>CYP3A5</i>	n	MR (TAM-NDMT)	MR (NDMT-END)	MR (TAM-4OHT)	MR (4OHT-END)
<b>Genotype</b>	<b>133</b>	<b>p=0.307</b>	<b>p=0.786</b>	<b>p=0.742</b>	<b>p=0.642</b>
<i>CYP3A5</i> *1/*1	18	0.32 (0.10)	21.72 (30.33)	79.49 (51.14)	0.10 (0.15)
<i>CYP3A5</i> *1/*3	64	0.36 (0.10)	15.45 (23.37)	70.34 (41.05)	0.08 (0.09)
<i>CYP3A5</i> *3/*3	51	0.34 (0.09)	23.26 (19.97)	63.23 (47.45)	0.10 (0.08)
<b>Phenotype</b>	<b>133</b>	<b>p=0.831</b>	<b>p=0.657</b>	<b>p=0.508</b>	<b>p=0.400</b>
EM	82	0.36 (0.10)	15.89 (25.49)	71.04 (42.44)	0.08 (0.12)
PM	51	0.34 (0.09)	23.26 (19.97)	63.23 (47.45)	0.10 (0.08)

**Notes:** The concentrations of TAM and its metabolites are presented as median (interquartile range).  $p$ = $p$  value from Kruskal–Wallis test.

**Abbreviations:** 4OHT, 4-hydroxytamoxifen; EM, extensive metabolizer; END, endoxifen; IM, intermediate metabolizer; MR, metabolic ratios; NDMT, N-desmethyl tamoxifen; PM, poor metabolizer; TAM, tamoxifen.

## CYP2D6 with CYP3A5 polymorphisms and TAM and its metabolites

The concentrations of TAM and its metabolites were significantly different between different *CYP2D6* phenotypes (Table 2), but these differences were not significant when both *CYP2D6* and *CYP3A5* were combined in the analysis ( $p=0.265$ ; Table 5). The NDMT, END and 4OHT concentrations were not significantly different between different combined *CYP2D6* and *CYP3A5* phenotypes ( $p=0.114$ ,  $p=0.244$  and  $p=0.224$ , respectively; Table 5). The MR (TAM-NDMT) and MR (NDMT-END) were significantly different between different combined *CYP2D6* and *CYP3A5* phenotypes ( $p=0.032$  and  $p=0.026$ , respectively), while the MR (TAM-4OHT) and MR (4OHT-END) were not significantly different between different combined *CYP2D6* and *CYP3A5*

phenotypes ( $p=0.079$  and  $p=0.622$ , respectively). The MR of TAM and its metabolites between different combined *CYP2D6* and *CYP3A5* polymorphisms are presented in Table 5.

## Discussion

The patients consisted of 134 Thai women with breast cancer who demonstrated all stages of breast cancers and were both premenopausal and postmenopausal. Saladores et al suggested that a combination of genetic factors (*CYP2C9*, *CYP2C19* and *CYP3A5*) and nongenetic factors (age and BMI) produced a 2.8% contribution to MR (NDMT-END),<sup>25</sup> and Lien et al reported that age was positively correlated to TAM, NDMT and END concentrations.<sup>26</sup> Therefore, in the present study, the associations between these factors (age and BMI) and the concentration of TAM and its metabolites were



**Table 5** The MR of TAM and its metabolites between different combinations of *CYP2D6* and *CYP3A5* polymorphisms

Combined phenotype	n (133)	TAM (ng/mL), p=0.265	NDMT (ng/mL), p=0.114	END (ng/mL), p=0.244	4OHT (ng/mL), p=0.224
<i>CYP2D6</i> (EM)– <i>CYP3A5</i> (EM)	31	373.13 (266.69)	1,030.96 (487.88)	72.86 (67.52)	5.31 (2.76)
<i>CYP2D6</i> (EM)– <i>CYP3A5</i> (PM)	24	365.23 (173.35)	1,030.41 (738.27)	58.71 (66.85)	6.25 (2.80)
<i>CYP2D6</i> (IM)– <i>CYP3A5</i> (EM)	51	358.31 (231.36)	1,094.70 (630.68)	46.56 (46.50)	4.70 (2.62)
<i>CYP2D6</i> (IM)– <i>CYP3A5</i> (PM)	25	425.63 (227.09)	1,193.56 (610.65)	57.47 (42.03)	4.44 (3.22)
<i>CYP2D6</i> (PM)– <i>CYP3A5</i> (PM)	2	238.36 (0.00)	803.19 (0.00)	41.86 (0.00)	6.23 (0.00)
Combined phenotype	n (133)	MR TAM-NDMT (p=0.032*)	MR NDMT-END (p=0.026*)	MR TAM-4OHT (p=0.079)	MR 4OHT-END (p=0.622)
<i>CYP2D6</i> (EM)– <i>CYP3A5</i> (EM)	31	0.38 (0.19)	11.83 (7.06)	65.09 (42.45)	0.07 (0.08)
<i>CYP2D6</i> (EM)– <i>CYP3A5</i> (PM)	24	0.34 (0.11)	18.06 (27.87)	60.29 (28.26)	0.11 (0.17)
<i>CYP2D6</i> (IM)– <i>CYP3A5</i> (EM)	51	0.34 (0.09)	23.84 (27.62)	75.47 (46.76)	0.10 (0.14)
<i>CYP2D6</i> (IM)– <i>CYP3A5</i> (PM)	25	0.35 (0.08)	24.28 (17.3)	78.98 (68.10)	0.10 (0.07)
<i>CYP2D6</i> (PM)– <i>CYP3A5</i> (PM)	2	0.30 (0.00)	21.56 (0.00)	40.48 (0.00)	0.17 (0.00)

**Notes:** The concentration of TAM and its metabolites are presented in median (interquartile range). *p*=*p* value from Kruskal–Wallis test, \**p*<0.05.

**Abbreviations:** 4OHT, 4-hydroxytamoxifen; EM, extensive metabolizer; END, endoxifen; IM, intermediate metabolizer; MR, metabolic ratios; NDMT, N-desmethyl tamoxifen; PM, poor metabolizer; TAM, tamoxifen.

explored to determine other possible effects. Age showed a significant nonparametric correlation to MR (TAM-4OHT; *p*=0.026) and BMI showed a significant nonparametric correlation to MR (TAM-NDMT; *p*=0.020) and MR (TAM-4OHT; *p*=0.023; data not shown). However, the distribution of these factors (BMI and age) was not statistically different between *CYP2D6*, *CYP3A5* and combined *CYP2D6* and *CYP3A5* phenotypes (data not shown) which indicated the same distribution of age and BMI among those analyzed phenotype groups.

One patient was excluded from the gene polymorphisms–concentration association analysis because of an extremely low concentration of TAM (44.9 ng/mL) which was lower than 20% of the median concentration. This low concentration might result from nonadherence or other unknown factors. The patient interview nonadherence screening method may overestimate medication adherence in some instances. Saladores et al reported the use of plasma TAM concentration as a criterion for medication adherence screening in breast cancer patients and they found that 39/587 of patients were excluded from the analysis based on this adherence screening method.<sup>25</sup>

The concentrations of TAM and its metabolites were not significantly different between the combined *CYP2D6* and *CYP3A5* phenotypes (Table 5), even though TAM concentration was significantly different between the *CYP2D6* phenotypes (Table 2) which might have resulted from the effect of several enzymes involved in the TAM metabolic pathway.<sup>1,2,8–10</sup> However, the same baseline concentration of TAM was present before applying the gene polymorphisms–concentrations association analysis. Since there are several TAM metabolic subpathways and several enzymes are involved in each subpathway,<sup>1,2,8–10,18</sup> the MR of these metabolites were

used to adjust the baseline concentration of each metabolite formed in each TAM metabolic subpathway. The MR (TAM-NDMT) and MR (NDMT-END) were significantly different between different combined *CYP2D6* and *CYP3A5* phenotypes (*p*=0.032 and *p*=0.026, respectively; Table 5) and the *CYP2D6* phenotype (*p*=0.013 and *p*=0.014, respectively; Table 3), but were not significantly different between different *CYP3A5* phenotypes (*p*=0.831 and *p*=0.657, respectively; Table 4). These results revealed the main effect of the *CYP2D6* phenotype on MR (TAM-NDMT) and MR (NDMT-END) without the effect of *CYP3A5* phenotype on those metabolites. On the contrary, the MR (TAM-4OHT) was not significantly different between the combined *CYP2D6* and *CYP3A5* phenotypes (*p*=0.079; Table 5) or the *CYP3A5* phenotype (*p*=0.508; Table 4), but was significantly different between the *CYP2D6* phenotypes (*p*=0.017; Table 3). These conflicting results indicate the potential interaction between the *CYP2D6* and *CYP3A5* phenotypes in affecting MR (TAM-4OHT); therefore, the interaction between *CYP2D6* and *CYP3A5* was considered. According to the nonparametric tests that were used to perform the hypotheses testing in this research, the usual interaction test was of limited use. Instead, the trend graphs of the *CYP2D6* and *CYP3A5* phenotypes were plotted to see the possible interaction effect and the simple main effect test<sup>24</sup> was subsequently performed to determine the effect of *CYP2D6* and *CYP3A5* on the MR (TAM-4OHT). No significant differences were detected when the *CYP2D6* or *CYP3A5* phenotypes were fixed in these simple main effect analyses, which confirmed that neither the *CYP2D6* nor the *CYP3A5* phenotypes affected the MR (TAM-4OHT; Table 6). However, the significance level between the different *CYP2D6* phenotypes in the *CYP3A5*-PM group

**Table 6** Simple main effect analysis of *CYP2D6* and *CYP3A5* phenotypes on MR (TAM-4OHT)

Gene polymorphisms		n	p	Gene polymorphisms		n	p
CYP3A5-EM	CYP2D6-EM	31	0.176	CYP2D6-EM	CYP3A5-EM	31	0.671
	CYP2D6-IM	51			CYP3A5-PM	24	
	CYP2D6-PM	0		CYP2D6-IM	CYP3A5-EM	51	
CYP3A5-PM	CYP2D6-EM	24	0.056		CYP3A5-PM	25	0.778
	CYP2D6-IM	25		CYP2D6-PM	CYP3A5-EM	0	
	CYP2D6-PM	2			CYP3A5-PM	2	

**Notes:** The concentrations of TAM and its metabolites are presented as median (interquartile range).  $p=p$  value from Kruskal–Wallis test.


**Abbreviations:** 4OHT, 4-hydroxytamoxifen; EM, extensive metabolizer; IM, intermediate metabolizer; MR, metabolic ratios; PM, poor metabolizer; TAM, tamoxifen.

was such that the null hypothesis was nearly rejected, which implied that it is possible that the *CYP2D6* polymorphisms might affect MR (TAM-4OHT) in the *CYP3A5*-PM group and might be responsible for the statistically significant results concerning MR (TAM-4OHT) between the different *CYP2D6* phenotypes (Table 3).

The MR (4OHT-END) was not significantly different between the combined *CYP2D6* and *CYP3A5* phenotype ( $p=0.622$ ; Table 5), the *CYP2D6* phenotype ( $p=0.594$ ; Table 3) or the *CYP3A5* phenotype ( $p=0.400$ ; Table 4), which suggested that *CYP2D6* or *CYP3A5* phenotype did not affect the MR (4OHT-END).

These results suggested that the effect of the *CYP2D6* phenotype on MR (TAM-NDMT) and MR (NDMT-END) does not involve effects from the *CYP3A5* phenotype on these metabolites. These results correspond to those of a study by Mürdter et al which found that the *CYP2D6* phenotype was associated with MR (NDMT-END;  $p<10^{-16}$ ),<sup>9</sup> another study by Saladores et al which suggested that *CYP2D6* made 53% of the contribution toward an MR (NDMT-END)<sup>25</sup> and previous studies by Lim et al<sup>16</sup> and Fernández-Santander et al<sup>15</sup> which suggested no association between *CYP3A5* polymorphisms and plasma concentrations of TAM and its metabolites.

In Figure 1, the *CYP2D6* and *CYP3A5* phenotypes are represented by the *CYP2D6* and *CYP3A5* polymorphisms, respectively, while the MR (TAM-NDMT) and MR (TAM-4OHT) are represented by the NDMT and 4OHT concentrations, respectively, and the MR (NDMT-END) is represented by the END concentrations through the NDMT subpathway (Figure 1) and the MR (4OHT-END) is represented by the END concentrations through the 4OHT subpathway (Figure 1). These MR analyses were useful for discriminating END concentrations through the NDMT subpathway, which is the main pathway for END formation from another minor subpathway through 4OHT. In conclusion, this study suggests that the effects of the *CYP2D6* polymorphisms on NDMT and END concentrations occur through the NDMT subpathway,



**Figure 1** TAM metabolic pathways and their interaction with *CYP2D6* and *CYP3A5* polymorphisms.

**Note:** \*Shows significant differences.

**Abbreviations:** 4OHT, 4-hydroxytamoxifen; END, endoxifen; NDMT, N-desmethyl tamoxifen; TAM, tamoxifen.

but the *CYP3A5* polymorphisms did not produce these effects on the concentrations of TAM and its metabolites.

## Limitations

Firstly, a comedication list was collected from a hospital database and patient medical records were used for drug–drug interaction screening without plasma drug level determination, which might lead to over- or underestimation of the drug–drug interaction problem. Secondly, medication adherence was assessed using a face-to-face interview with the patients without other medication adherence assessment tools, which might lead to overestimation of patient medication adherence. Thirdly, plasma concentrations of TAM and its metabolites were determined using HPLC with a fluorescence detector. Some extrapolations for NDMT concentration (78 patients) and END concentration (1 patient) need to be taken into account when reporting the accuracy of the quantification of these metabolites. However, the relevant hypotheses testing results should not be affected by these limitations, according to the nonparametric analyses which were based on the rank sum test rather than their actual values. Finally, genes encoding other metabolizing enzymes that are involved in the TAM metabolic pathway, such as *CYP2C9*, *CYP2C19* or *CYP3A4*, might need to be explored to yield more concise findings.

These results can be further applied to identify a high-risk patient group for potential ineffective TAM treatment,

in terms of their genetic background resulting in low concentrations of its active metabolites. This information could be used to improve medication plans by adjusting TAM dosage based on individual genetic factors. However, the associations between the polymorphisms, plasma concentrations of TAM and its metabolites and true clinical outcomes should be confirmed before these results can be applied in clinical practice.

## Acknowledgments

This research was funded by the thesis grant for doctoral degree student of the National Research Council of Thailand (NRCT; 2015) and the 90th Anniversary of Chulalongkorn University Scholarship (2015).

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Klein DJ, Thorn CF, Desta Z, Flockhart DA, Altman RB, Klein TE. PharmGKB summary: tamoxifen pathway, pharmacokinetics. *Pharmacogenet Genomics*. 2013;23(11):643–647.
- Saladores PH, Precht JC, Schroth W, Brauch H, Schwab M. Impact of metabolizing enzymes on drug response of endocrine therapy in breast cancer. *Expert Rev Mol Diagn*. 2013;13(4):349–365.
- Rae JM, Drury S, Hayes DF, et al. CYP2D6 and UGT2B7 genotype and risk of recurrence in tamoxifen-treated breast cancer patients. *J Natl Cancer Inst*. 2012;104(6):452–460.
- Karle J, Bolbrinker J, Vogl S, et al. Influence of CYP2D6-genotype on tamoxifen efficacy in advanced breast cancer. *Breast Cancer Res Treat*. 2013;139(2):553–560.
- Regan MM, Leyland-Jones B, Bouzyk M, et al. CYP2D6 Genotype and tamoxifen response in postmenopausal women with endocrine-responsive breast cancer: the breast international group 1–98 trial. *J Natl Cancer Inst*. 2012;104(6):441–451.
- Westbrook K, Stearns V. Pharmacogenomics of breast cancer therapy: an update. *Pharmacol Ther*. 2013;139(1):1–11.
- Binkhorst L, Mathijssen RHJ, Ghobadi Moghaddam-Helmantel IM, et al. Quantification of tamoxifen and three of its phase-I metabolites in human plasma by liquid chromatography/triple-quadrupole mass spectrometry. *J Pharm Biomed Anal*. 2011;56(5):1016–1023.
- Kiyotani K, Mushiroda T, Nakamura Y, Zembutsu H. Pharmacogenomics of tamoxifen: roles of drug metabolizing enzymes and transporters. *Drug Metab Pharmacokinet*. 2012;27(1):122–131.
- Mürdter TE, Schroth W, Bacchus-Gerybadze L, et al. Activity levels of tamoxifen metabolites at the estrogen receptor and the impact of genetic polymorphisms of phase I and II enzymes on their concentration levels in plasma. *Clin Pharmacol Ther*. 2011;89(5):1–10.
- Barginear MF, Jaremkó M, Peter I, et al. Increasing tamoxifen dose in breast cancer patients based on CYP2D6 genotypes and endoxifen levels: effect on active metabolite isomers and the antiestrogenic activity score. *Clin Pharmacol Ther*. 2011;90(4):605–611.
- Zafra-Ceres M, de Haro T, Farez-Vidal E, et al. Influence of CYP2D6 polymorphisms on serum levels of tamoxifen metabolites in Spanish women with breast cancer. *Int J Med Sci*. 2013;10(7):932–937.
- Ratain MJ, Nakamura Y, Cox NJ. CYP2D6 genotype and tamoxifen activity: understanding interstudy variability in methodological quality. *Clin Pharmacol Ther*. 2013;94(2):185–187.
- Hertz DL, McLeod HL, Irvin Jr WJ. Tamoxifen and CYP2D6: a contradiction of data. *Oncologist*. 2012;17(5):620–630.
- Teh LK, Bertilsson L. Pharmacogenomics of CYP2D6: molecular genetics, interethnic differences and clinical importance. *Drug Metab Pharmacokinet*. 2012;27(1):55–67.
- Fernández-Santander A, Gaibar M, Novillo A, et al. Relationship between genotypes Sult1a2 and Cyp2d6 and tamoxifen metabolism in breast cancer patients. *PLoS One*. 2013;8(7):e70183.
- Lim JS, Chen XA, Singh O, et al. Impact of CYP2D6, CYP3A5, CYP2C9 and CYP2C19 polymorphisms on tamoxifen pharmacokinetics in Asian breast cancer patients. *Br J Clin Pharmacol*. 2011;71(5):737–750.
- Chamnanphon M, Pechatanan K, Sirachainan E, et al. Association of CYP2D6 and CYP2C19 polymorphisms and disease-free survival of Thai post-menopausal breast cancer patients who received adjuvant tamoxifen. *Pharmacogenomics Pers Med*. 2013;6(1):37–48.
- Charoenchokthavee W, Panomvana D, Sriuranpong V, Areepium N. Prevalence of CYP2D6\*2, CYP2D6\*4, CYP2D6\*10, and CYP3A5\*3 in Thai breast cancer patients undergoing tamoxifen treatment. *Breast Cancer (Dove Med Press)*. 2016;8:149–155.
- Areepium N, Panomvana D, Rungwanonchai P, Sathaporn S, Voravud N. Effects of CYP2D6 and UGT2B7 polymorphisms on pharmacokinetics of tamoxifen in Thai breast cancer patients. *Breast Cancer (Dove Med Press)*. 2013;5:73–78.
- Charoenchokthavee W, Ayudhya DP, Sriuranpong V, Areepium N. Effects of SULT1A1 copy number variation on estrogen concentration and tamoxifen-associated adverse drug reactions in premenopausal Thai breast cancer patients: a preliminary study. *Asian Pac J Cancer Prev*. 2016;17(4):1851–1855.
- Faul F, Erdfelder E, Lang A-G, Buchner A. G\*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods*. 2007;39(2):175–191.
- Cohen Jacob. *Statistical Power Analysis for the Behavioral Sciences*. 2nd ed. Hillsdale, NJ: Lawrence Erlbaum Associates; 1988.
- Zhu YB, Zhang Q, Zou JJ, Yu CX, Xiao DW. Optimizing high-performance liquid chromatography method with fluorescence detection for quantification of tamoxifen and two metabolites in human plasma: application to a clinical study. *J Pharm Biomed Anal*. 2008;46(2):349–355.
- Kirk RE. *Experimental Design: Procedures for the Behavioral Sciences*. 4th ed. Thousand Oaks, CA: SAGE; 2013.
- Saladores P, Mürdter T, Eccles D, et al. Tamoxifen metabolism predicts drug concentrations and outcome in premenopausal patients with early breast cancer. *Pharmacogenomics J*. 2014;15(1):84–94.
- Lien EA, Soiland H, Lundgren S, et al. Serum concentrations of tamoxifen and its metabolites increase with age during steady-state treatment. *Breast Cancer Res Treat*. 2013;141(2):243–248.

### Breast Cancer - Targets and Therapy

### Publish your work in this journal

Breast Cancer - Targets and Therapy is an international, peer-reviewed open access journal focusing on breast cancer research, identification of therapeutic targets and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient.

Submit your manuscript here: <https://www.dovepress.com/breast-cancer---targets-and-therapy-journal>

Dovepress

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.