

Response to Association of Triglyceride-Glucose Index with Risk of Large for Gestational Age: A Prospective Cohort Study [Letter]

Rongyan Wei, Yuechou Nong

Department of Endocrinology and Metabolism, The Guangxi Academy of Medical Sciences, The People's Hospital of Guangxi Zhuang Autonomous Region, Nanning, Guangxi, 530021, People's Republic of China

Correspondence: Yuechou Nong, Email yuechou_gx@163.com

Dear editor

Endocrine and metabolic disorders play a crucial role in adverse pregnancy outcomes, which is currently a research hotspot. We noticed Lin et al¹ recently published a paper entitled “Association of Triglyceride-Glucose Index with Risk of Large for Gestational Age: A Prospective Cohort Study” in *Diabetes Metabolic Syndrome and Obesity-Targets and Therapy*. They delved into the relationship between the TyG index and the risk of large for gestational age (LGA) based on a birth cohort. They found that a higher TyG index in the first trimester was independently associated with a higher risk of LGA. Logically, a high TyG index in the first trimester may serve as an early predictor for LGA.

While we appreciate the novel work, certain aspects of the current study still need further improvement to align with clinical practice. First, there is a lack of the differential diagnosis of Hashimoto's thyroiditis (HT), especially subclinical hypothyroidism. We note that the author excluded hypothyroidism and hyperthyroidism in Table 5 but did not exclude Hashimoto's thyroiditis, which is particularly closely related to LGA in patients with TSH > 2.5 mIU/L. Maternal subclinical hypothyroidism in early pregnancy (TSH > 2.5 mIU/L) was closely associated with increased odds for LGA in male newborns (OR = 1.95, 95% CI = 1.22–3.11).² Additionally, a prospective cohort study with 175,112 women in China showed a J-shaped relationship between TSH and LGA ($P < 0.001$), in which TSH within 0.91 to 1.82 mIU/L was the potential safe range for preconception women.³ Actually, over 30% of women with hypothyroidism (Hypo-HT) and euthyroidism-HT (Eu-HT) have TSH greater than 2.5 mIU/L in early pregnancy, 19.7% of women with Hypo-HT and 10.1% of women with Eu-HT have TSH higher than 4.0 mIU/L ($P < 0.05$).⁴ To prevent/reduce the risk of thyroid insufficiency inducing adverse pregnancy outcomes such as LGA during early pregnancy, preconception thyroid function optimization for women with Hashimoto thyroiditis (HT) through levothyroxine sodium is strongly suggested by guidelines.⁵ Second, there is a lack of evaluation data on serum ferritin levels. Serum ferritin concentration prior to pregnancy or in the first trimester can affect thyroid function, which is associated with adverse pregnancy outcomes, including LGA.⁶

To sum up, only by improving the above deficiencies can the research results be more dependable.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Lin L, Lin J, Yang F, et al. Association of triglyceride-glucose index with risk of large for gestational age: a prospective cohort study. *Diabetes Metabol Syndr Obes.* 2023;2023(16):3837–3846. doi:10.2147/DMSO.S436611
2. Vrijkotte TGM, Hrudevy EJ, Twickler MB. Early maternal thyroid function during gestation is associated with fetal growth, particularly in male newborns. *J Clin Endocrinol Metab.* 2017;102(3):1059–1066. doi:10.1210/je.2016-3452
3. Du H, Wu D, Zhou X, et al. Preconception TSH and adverse pregnancy outcomes in china: a nationwide prospective cohort study. *J Clin Endocrinol Metab.* 2022;107(7):e2770–e2776. doi:10.1210/clinem/dgac208

4. Moleti M, Alibrandi A, Di Mauro M, et al. Preconception thyrotropin levels and thyroid function at early gestation in women with Hashimoto thyroiditis. *J Clin Endocrinol Metab.* 2023;108(7):e464–e473. doi:10.1210/clinem/dgac748
5. Bekkering GE, Agoritsas T, Lytvyn L, et al. Thyroid hormones treatment for subclinical hypothyroidism: a clinical practice guideline. *BMJ.* 2019;365:l2006. doi:10.1136/bmj.l2006
6. Li X, Zhang Z, Cai W, et al. Association between serum ferritin levels in early pregnancy and thyroid function and pregnancy outcomes in Chinese population. *Int J Women's Health.* 2023;15:1951–1957. doi:10.2147/IJWH.S436651

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Diabetes, Metabolic Syndrome and Obesity 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Diabetes, Metabolic Syndrome and Obesity editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Diabetes, Metabolic Syndrome and Obesity

Dovepress

Publish your work in this journal

Diabetes, Metabolic Syndrome and Obesity is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert opinion and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-journal>

<https://doi.org/10.2147/DMSO.S454536>