

Sociodemographic Profile: A Forgotten Factor in Temporomandibular Disorders? A Scoping Review

Ivo Henrique Alexandrino Carapinha¹, Giancarlo De la Torre Canales¹⁻³, Rodrigo Lorenzi Poluha⁴, Mariana Barbosa Câmara-Souza³, Nikolaos Christidis², Malin Ernberg², Andre Mariz de Almeida¹, Ana Cristina Garcia de Matos Manso¹

¹Egas Moniz Center for Interdisciplinary Research (CiiEM); Egas Moniz School of Health & Science, Almada, Portugal; ²Division of Oral Diagnostics and Rehabilitation, Department of Dental Medicine, Karolinska Institutet, Huddinge, Sweden; ³Department of Dentistry, Ingá University Center, Uningá, Paraná, Brazil; ⁴Department of Dentistry, State University of Maringá, Paraná, Brazil

Correspondence: Giancarlo De la Torre Canales, Tel +351 913025948, Email gcanales@egasmoniz.edu.pt

Abstract: The literature on Temporomandibular Disorders (TMD) incidence commonly reports sociodemographic factors such as gender and age. However, the role and prevalence of other sociodemographic factors in TMD are not well defined. Therefore, this scoping review aimed to report the prevalence of sociodemographic factors in TMD patients. A systematic search was conducted in the PubMed and Web of Science databases to identify clinical trials in adult populations, using the Research Diagnostic Criteria for TMD (RDC/TMD) or the Diagnostic Criteria for TMD (DC/TMD) and reporting sociodemographic data in TMD patients. Twenty-seven studies meeting the criteria were included in this review. The most commonly reported sociodemographic factors assessed in the included studies were age, race, education, job, income, and marital status. TMD prevalence was observed to be higher among younger and divorced individuals among the included studies. However, conflicting results were found for education level, and employment was not considered a risk factor for TMD. Although this review has methodological limitations, it suggests an association between TMD incidence and certain sociodemographic factors; nevertheless, further studies are needed to establish this relationship more conclusively.

Keywords: temporomandibular joint disorders, sociodemographic factors, orofacial pain

Introduction

Temporomandibular disorders (TMDs) are the most prevalent group and the most common source of chronic pain in the orofacial area.^{1,2} They are characterized by pain in the temporomandibular joints (TMJ) and masticatory muscles, noises of the TMJ during mandibular function, and by restriction in jaw movements.^{1,3,4} Painful TMDs have been shown to be biopsychosocial and multifactorial disorders; hence, there is not a singular cause that can explain the onset of painful TMDs.⁵ Notwithstanding, the psychological profile, state of pain amplification and general health and global symptoms have been proposed to play an important role in the etiology of painful TMDs,⁵⁻⁷ which are probably adjusted by gene expression and affected by environmental contributions such as social and demographic factors.⁶

In addition, the most common sociodemographic factor reported by the literature related with TMDs incidence is the gender frequency, which is higher in females than in males, and age prevalence, being greatest among people between 18 and 44 years old.^{5,8} However, while other sociodemographic factors disparities like race, social and ethnicity have been associated with clinical pain experiences in many health conditions,⁹⁻¹¹ their role in TMDs is not clear, especially regarding TMDs onset.⁶ In the same way, discrepancies have been described in subgroups with different socioeconomic status. Further, it seems that educational attainment and factors associated with moving to another country are less important predictors, even though could influence TMDs pain.^{12,13}

Most clinical trials in the field of TMDs use the Research Diagnostic Criteria for TMD (RDC/TMD),¹⁴ or its updated version, the Diagnostic Criteria for TMD (DC/TMD),¹⁵ as a diagnostic tool that also assesses the sociodemographic profile of

patients. Intriguingly, even though they assess the sociodemographic profile of the included patients, these data are rarely reported. Both the RDC/TMD and the DC/TMD include questions regarding age, race, sex, marital status, ethnicity, level of schooling and family income, which as explained before, could influence the onset of TMDs.^{14,15} Therefore, since the actual knowledge about the role of sociodemographic factors is scant, but merits attention for both scientific and public health reasons, it is important to summarize the current findings to provide a basis to construct TMDs patients sociodemographic profile as well as suggestions for future researches. Taking this into account, the aim of the present scoping review is to report the prevalence of sociodemographic factors in TMDs patients using the RDC/TMD and DC/TMD.

Materials and Methods

Search Strategy and Selection Criteria

During February 2023, a search of the literature was performed using the following Medical Subjects Headings (MeSH) and related terms: [Temporomandibular Joints disorders] OR [Temporomandibular disorders] OR [TMD] AND [sociodemographic factors] OR [sociodemographic] AND [Research Diagnostic Criteria for Temporomandibular Disorders] OR [RDC/TMD] AND [Diagnostic Criteria for Temporomandibular Disorders] OR [DC/TMD] in the National Library of Medicine Medline (PubMed) and Web of Science databases to identify a list of potential papers for inclusion in this scoping review. In addition, search expansion strategies were also used to identify other potentially relevant citations (ie, hands-on search in private libraries and reference lists of the included articles).

The inclusion criteria were limited to: (1) studies written in English published from January 1992 to February 2023, (2) articles on adult populations (if adults and teenagers/children were included in the same study, just adult data was considered for the analysis), (3) clinical trials focused on reporting sociodemographic data in TMD patients, (4) studies using the RDC/TMD and DC/TMD as diagnostic tools. Manuscripts using different diagnostics tools, not reporting the criteria for TMD diagnosis, reporting data on TMD prevalence only in teenagers/children, or unrelated to the review aim and case reports were excluded.

Assessment of Papers

Two of the authors (I.C. and G.D.C.) independently reviewed the titles and abstracts of all articles, and potential articles were obtained in full text for careful read to check the eligibility. In all cases of disagreement between reviewers regarding the potential inclusion of an article or data interpretation, a third author (R.P.) was involved. Then, after final selection and inclusion of articles for the review, another author (M.B.C.S.) performed data extraction based on the Population Exposure Comparison Outcome (PECO) strategy.¹⁶ The population (“P”) was described including the sample size. The exposure (“E”) concerned information on the study design, diagnostic tool used, and TMD diagnoses. The comparison (“C”) included if data for the control group if present depending on the study design. The outcome (“O”) was reported in terms of sociodemographic frequency data.

Results

Overview

The literature search identified 316 articles, of which eight were overlapping articles retrieved in both databases. From the 308 articles screened by title and abstract, 181 were read in full for eligibility. Of these, 151 were excluded for not fulfilling the inclusion criteria, and three were excluded since it was not possible to retrieve the full text. Therefore, a total of 27 manuscripts were included in the review (Figure 1).

The included articles covered a wide spectrum of populations of different sex, age and ethnic background. The age of the subjects varied from ≥ 15 years to ≤ 80 years, and the sample size ranged from 15 to 4289 participants with and without TMD. Regarding the sex distribution, a predominance of females was found (Table 1).

Summary of the Studies

Most of the included studies aimed to investigate the role of sociodemographic characteristics and their association with TMDs. Besides sociodemographic aspects, such as age, race, education, job, income, and marital status, two manuscripts




Figure 1 Flow diagram of the literature search strategy.

investigated features related to height, weight, and BMI. Further, psychological characteristics and parafunctional habits were investigated in most studies but are not presented in this scoping review since they do not relate to the present aim (Table 2).

The environment in which each study was conducted differed substantially from each other. While there were epidemiological studies aiming to access the influence of socioeconomic features, some papers were developed to investigate specific situations, such as patients with Parkinson's disease, COVID-19 pandemic, headaches, etc. Thus,

Table 1 Characteristics of the Included Studies

Study First Author, Year	Study Design	Diagnostic Criteria	Population
Velly, 2002 ¹⁷ (Canada)	Case-Control	RDC/TMD	N = 59 (43 W, 16 M, mean age = 36 y) with Disc Displacement (75% with reduction, 12% without reduction, 13% with arthralgia osteoarthritis-osteoarthritis - restrictions during jaw movement, 7%). Control group: 100 patients (64 W, 36 M, mean age = 36 y).
Gesch, 2004 ¹⁸ (Germany)	Cross-sectional observational	No mention to DC or RDC/TMD exam [†]	4289 patients (2180 W, 2109 M).
Bernhardt, 2005 ¹⁹ (Germany)	Population-based cross-sectional	No mention if DC or RDC/TMD exam [†]	4255 subjects (2165 W, 2090 M).
Mundt, 2005 ²⁰ (Germany)	Population-based, randomized, cross sectional	No mention if DC or RDC/TMD exam [†]	2963 subjects (1493 W, 1470 M).
Casanova-Rosado, 2006 ²¹ (Mexico)	Cross-sectional	RDC/TMD	506 patients (274 W, 232 M, mean age = 17.2 ± 2.7 y)
Selaimen, 2007 ²² (Brazil)	Analytical, case-control	RDC/TMD	72 patients (all women, mean age = 32.4 ± 12.1 y), diagnosed with myofascial pain, with or without limited opening and arthralgia. Control group: 30 patients (all women, mean age = 38.7 ± 14.4 y).
Ommerborn, 2008 ²³ (Germany)	Prospective observational clinical	RDC/TMD	125 patients (80 W, 45 M, mean age = 49.24 ± 15.73 y)
Quinteromarmol-Juárez, 2008 ²⁴ (Mexico)	Comparative cross-sectional	RDC/TMD	130 patients (70 W, 51 M, mean age = 34 ± 10 years), with or without TMD.
Slade, 2011 ²⁵ (USA)	Case-control	RDC/TMD	185 patients (155W, 30M) with chronic TMD Control group: 1633 pain-free volunteers (925W, 708M).
Reissmann, 2012 ²⁶ (Germany)	Case-control	RDC/TMD	70 patients (57 W, 13 M, mean age = 41.9 ± 15.6 y), with at least one pain-related diagnosis Control group: 868 subjects (493 W, 375 M, mean age = 40.4 ± 11.8 y) - general-population adults without any pain-related TMD.
Blanco-Hungría, 2012 ²⁷ (Spain)	Cross-sectional	RDC/TMD	748 patients (624 W, 124 M, mean age = 53.28 ± 27.76 y) with some of the following signs or symptoms: mandibular or TMJ pain, limitation or restriction during oral aperture or lateralization, or joint sounds with or without pain.
Slade, 2013 ⁵ (USA)	Community-based prospective cohort	RDC/TMD	2737 participants (1630 W, 1107 M) without history of TMD.
Blanco-Aguilera, 2014 ²⁸ (Spain)	Cross-sectional epidemiological	RDC/TMD	407 patients (365 W, 42 M) with at least one of signs and symptoms as: pain in the jaw or TMJs, restricted or limited range of motion when opening or closing the mouth or lateral excursions of the jaw, and joint sounds (with or without pain).
Lei, 2015 ²⁹ (China)	Cohort	RDC/TMD	510 patients (387 W, 123 M, mean age = 31.06 ± 14.40 y) with TMD.

(Continued)

Table 1 (Continued).

Study First Author, Year	Study Design	Diagnostic Criteria	Population
Dahan, 2016 ³⁰ (Canada)	Comparative cross-sectional, multicenter	RDC/TMD	180 patients (149 W, 31 M, mean age = 42.8 ± 1.2 y), with chronic TMD.
Su, 2017 ³¹ (Netherlands)	Cross-sectional	DC/TMD	320 patients with TMD (250 W, 70 M, mean age = 43.2 ± 14.6 years).
Di Paolo, 2017 ³² (Italy)	Retrospective cohort	DC/TMD	929 patients with TMD.
Santiago, 2019 ³³ (USA)	Case-control	RDC/TMD	124 patients (all women, mean age = 40.3 ± 14.9 y) with Myofascial TMD (27.4% with muscle pain only – n = 34, 72.6% with muscle and joint pain – n = 90).
Miller, 2019 ³⁴ (USA)	Cross-sectional Multicenter study Ω	RDC/TMD	846 participants with chronic TMD (652 W, 194 M, mean age = 29.0 ± 7.8 years) aged 18–44 years.
Balik, 2019 ³⁵ (Turkey)	Cross-sectional	RDC/TMD	104 patients (64 W, 40 M, mean age = 33.46 ± 10.51 y) with chronic TMD pain for at least 6 months.
Arikan, 2019 ³⁶ (Turkey)	Cross-sectional	RDC/TMD	77 patients with TMD (59 W, 18 M, mean age = 32.69 ± 13.64 years).
Resende, 2020 ³⁷ (Brazil)	Nonpaired, controlled case	RDC/TMD	120 patients (80 W, 40 M, mean age = 33.29 ± 13.68 years) with and without TMD.
de Caxias, 2021 ³⁸ (Brazil)	Epidemiological cross-sectional and analytic	DC/TMD	2301 patients (1513 W, 537 M, 2 Non-binary, mean age = 41.4 ± 13.64 y), practicing social isolation during the COVID-19 pandemic. 249 patients (150W, 99M), not practicing social isolation during COVID-19 pandemic.
Delgado-Delgado, 2021 ³⁹ (Spain)	Observational	DC/TMD	59 participants with (n = 45) and without (n = 14) TMD (35 F, 24 M, mean age = 28.0 ± 10.1 years).
Tavares, 2021 ⁴⁰ (Brazil)	Cross-sectional cohort cut	RDC/TMD	15 patients (5 W, 10 M, mean age = 69.0 years), elderly people with Parkinson's disease associated with TMD and depression.
Lei, 2021 ⁴¹ (China)	Cross-sectional	DC/TMD	1079 patients (856 W, 223 M, mean age = 29.6 ± 14.2 y) with TMD.
Mendonça, 2022 ⁴² (Brazil)	Cross-sectional	RDC/TMD	41 patients (all women, mean age = 26.83 ± 7.54 y), diagnosed with at least one painful TMD, who presented for treatment before and during COVID-19 pandemic.

Notes: [†]Followed the guidelines of the Academy of Orofacial Pain. ^ΩUniversity of Buffalo, NY; University of Florida Gainesville, FL; University of Maryland in Baltimore, MD; and University of North Carolina at Chapel Hill, NC.

the results of the 27 papers retrieved differed substantially in most aspects. Nevertheless, it was possible to summarize the findings according to the frequency of reported results (Table 2).

In general, no statistically significant difference was reported by the included studies between men and women regarding sociodemographic factors. The differences found were that women demonstrated higher prevalence of TMDs and pain-related impairments. Only one study reported that men were more likely to have higher pain-related disability than women.³¹

Diagnosed TMDs were higher among younger individuals, while those aged above 50 years showed a lower prevalence. Conversely, the prevalence of non-painful TMDs was also higher in the 20–30 years old individuals, while older individuals were more likely to experience high-impact pain.

Table 2 Summary of the Findings from the Included Studies (PECO)

Study First Author, Year	Exposure (Sociodemographic Factors)	Comparisons	Outcomes
Velly, 2002 ¹⁷ (Canada)	Age categories (years) 18–27: n = 23 28–37: n = 18 38–45: n = 5 46–60: n = 13 Race Non-white: n = 10 White: n = 49 Education Less than University: n = 30 University: n = 29 Job Household: n = 4 Employed: n = 37 Not employed: n = 18 Household income (Can\$/year) < 30,000: n = 29 30,000–60,000: n = 17 > 60,000: n = 8 Non-reported: n = 5 Marital status Married: n = 22 No partner (single, divorced, widowed): n = 36	Age categories (years) 18–27: n = 25 28–37: n = 24 38–45: n = 25 46–60: n = 26 Race on-white: n = 123 White: n = 77 Education Less than University: n = 57 University: n = 43 Job Household: n = 10 Employed: n = 55 Not employed: n = 18 Household income (Can\$/year) < 30,000: n = 55 30,000–60,000: n = 30 > 60,000: n = 7 Non-reported: n = 8 Marital status Married: n = 55 No partner (single, divorced, widowed): n = 44	- Gender was not associated with Disc Displacement (DD). - Age levels from 28 to 37 (OR = 0.84; 95% CI: 0.37–1.88), from 38 to 45 (OR = 0.25; 95% CI: 0.07–1.10), and greater than 46 (OR = 0.56; 95% CI: 0.23–1.30) were not associated with DD. - Race, education, and household income were not related to DD.
Gesch, 2004 ¹⁸ (Germany)	Age (years) 20–30: n = 592 30–40: n = 760 40–50: n = 748 50–60: n = 789 60–70: n = 789 > 70: n = 611 Race - Not reported Education Less than University: n = 3586 University: n = 703 Job - Not reported Household income (US\$) < 875 (lower): n = 939 875–2000: n = 2453 > 2000: n = 897 Marital status - Not reported	No comparison group.	- Women (OR = 2.3) compared with men, or subjects aged 30 to 40 years (OR = 1.6) or 40 to 50 years (OR = 1.4), compared with the age group of 20 to 30 years displayed more than one clinical sign of TMD. - Higher education was associated with more TMD signs (OR = 1.4). - Associations between age or sex and TMD were found. Being woman was a clinically relevant sociodemographic risk marker regarding TMD signs (OR = 2.3).

Bernhardt, 2005 ¹⁹ (Germany)	Age categories (years) 20–29: n = 591 30–39: n = 758 40–49: n = 747 50–59: n = 787 60–69: n = 789 70–79: n = 583 Race - Not reported Education - Not reported Job - Not reported Household income - Not reported Marital status - Not reported	Participants with Headache (287W, 98M) Age categories (years) 20–29: n = 57 30–39: n = 76 40–49: n = 90 50–59: n = 74 60–69: n = 56 70–79: n = 32 Race - Not reported Education - Not reported Job - Not reported Household income - Not reported Marital status - Not reported	- Women had a significantly higher risk for developing frequent headache compared to men. - Being 50 years and older showed a decreased risk of suffering from frequent headache. - Women showed a clear dose/response effect for one to three painful muscles (OR = 2.10) and for four or more (OR = 3.47) painful muscles. - Of the entire sample, 1.3% reported pain in the masticatory muscles, which was significantly correlated with frequent headache in women (OR = 2.37). - Pain upon palpation of the TMJ was only related to headache in women upon lateral palpation.
Mundt, 2005 ²⁰ (Germany)	Presence of Pain Muscle Tenderness and/or Pain (n = 416, 276W, 140M) Age categories (years) 35–44: n = 72W; 35M 45–54: n = 54W; 20M 55–64: n = 91W; 39M 65–74: n = 56W; 46M Race - Not reported Education Less than University: n = 138W; 74M University: n = 138W; 66M Job - Not reported Income - Not reported Marital status Married: n = 189W; 109M No partner (single, divorced or widowed): n = 87W; 31M TMJ Tenderness and/or Pain (n = 159, 109W, 50M) Age categories (years) 35–44: n = 30W; 15M 45–54: n = 28W; 11M 55–64: n = 34W; 14M 65–74: n = 17W; 10M Race - Not reported Education Less than University: n = 48FW 22M University: n = 61W; 28M Job - Not reported Income - Not reported Marital status Married: n = 845W; 1091M No partner (single, divorced or widowed): n = 372W; 239M	Absence of Pain Muscle Tenderness and/or Pain (n = 2547, 1217W, 1330M) Age categories (years) 35–44: n = 317W; 323M 45–54: n = 341W; 318M 55–64: n = 326W; 366M 65–74: n = 233W; 329M Race - Not reported Education Less than University: n = 526W; 617M University: n = 691W; 713M Job - Not reported Income - Not reported Marital status Married: n = 845W; 1091M No partner (single, divorced or widowed): n = 372W; 239M TMJ Tenderness and/or Pain (n = 2804, 1384W, 1420M) Age categories (years) 35–44: n = 359W; 343M 45–54: n = 370W; 327M 55–64: n = 383W 391M 65–74: n = 272W; 359M Race - Not reported Education Less than University: n = 616W; 669M University: n = 768W; 751M Job - Not reported Income - Not reported Marital status Married: n = 71W; 43M No partner (single, divorced or widowed): n = 38W; 7M	- Lower muscle tenderness was observed in men aged 45 to 54 years compared to other age groups ($P = 0.062$). - For both TMD signs, the differences in proportions of other age groups were not significant according to bivariate tests. - Individuals with school education of up to 9 years were more likely to have muscle tenderness or pain (men, $P = 0.024$). - Women who bruxed also showed a significant difference in muscle tenderness (25.2% vs 35.1%, $P = 0.001$). - Sociodemographic data – such as age, marital status, and school education modified the observed effects.

(Continued)

Table 2 (Continued).

Study First Author, Year	Exposure (Sociodemographic Factors)	Comparisons	Outcomes
Casanova-Rosado, 2006 ²¹ (Mexico)	Age categories (years) Females: mean age = 17.35 ± 2.67 Males: mean age = 17.05 ± 2.71 Race - Not reported Education - Not reported Job - Not reported Household income - Not reported Marital status - Not reported	No comparison group.	- TMD prevalence = 46.1% (n = 233). - Women have higher prevalence of TMD (52.9 vs 37.9%) than men ($P < 0.01$), and more likely to have TMD than men (OR = 1.8). - The mean age was higher within the group with TMD (17.6 ± 2.9 vs 16.9 ± 2.5) than those without TMD ($P < 0.01$). - Pain diagnosis was associated with age (OR = 1.2; 95% CI = 1.1–1.3) and female sex (OR = 2.3; 95% CI = 1.2–4.5).
Selaimen, 2007 ²² (Brazil)	Age categories - Not reported Race - Not reported Education Less than University: n = 61 University: n = 11 Job Employed: n = 25 Unemployed: n = 47 Household income Up to 5 minimum wage: n = 45 > 5 minimum wage: n = 27 Marital status Married: n = 43 No partner (single, divorced or widowed): n = 29	Age categories - Not reported Race - Not reported Education Less than University: n = 23 University: n = 7 Job Employed: n = 21 Unemployed: n = 9 Household income Up to 5 minimum wage: n = 22 > 5 minimum wage: n = 8 Marital status Married: n = 20 No partner (single, divorced or widowed): n = 10	- Unemployment and age did show statistically significant differences between TMD patients and non-pain subjects.
Ommerborn, 2008 ²³ (Germany)	TMD patients (n = 60, 49W, 11M, mean age = 43.45 ± 14.01 y) Age categories - Not reported Race - Not reported Education Less than University: n = 39 University: n = 21 Job Employed: n = 43.3% Unemployed: n = 48.4% Retired: n = 8.3% Household income - Not reported Marital status Married: n = 34 No partner (single, divorced, widowed): n = 26	Non-TMD participants (n = 65, 31W, 34M, mean age = 53.03 ± 14.09 y) Age categories - Not reported Race - Not reported Education Less than University: n = 48 University: n = 17 Job Employed: n = 20% Unemployed: n = 49.2% Retired: n = 30.8% Household income - Not reported Marital status Married: n = 42 No partner (single, divorced, widowed): n = 23	- Statistically significant differences were found with respect to age and gender. - Education and marital status showed no significant differences between both groups.

Quinteromarmol-Juárez, 2008 ²⁴ (Mexico)	With TMD (n = 65, 48W, 17M) Age categories - Not reported Race - Not reported Education (years) - 14.00 ± 3.60 Job - Not reported Household income (Mex\$) Average: 5178.00 ± 4786.00 Marital status Married: n = 45 No partner (single, divorced, widowed): n = 20	Without TMD (n = 65, 37W, 28M) Age categories - Not reported Race - Not reported Education (years) - 14.10 ± 3.30 Job - Not reported Household income (Mex\$) Average: 4546.00 ± 4257.00 Marital status Married: n = 43 No partner (single, divorced, widowed): n = 22	- Statistically significant differences were found when comparing by sex ($P < 0.04$). - The statistical data showed that there was no difference between demographic data (age, years of study and income) and TMD.
Slade, 2011 ²⁵ (USA)	Age categories (years) 18–24: n = 72 25–34: n = 60 35–44: n = 53 Race White (non-hispanic): n = 145 Non-whites: n = 40 Education - Not reported Job - Not reported Household income - Not reported Marital status - Not reported	Age categories (years) 18–24: n = 838 25–34: n = 451 35–44: n = 344 Race White (non-hispanic): n = 839 Non-whites: n = 794 Education - Not reported Job - Not reported Household income - Not reported Marital status - Not reported	- Odds of TMD increased across successively older age groups. - Women had more than three times the odds of TMD as males. - Relative to non-Hispanic Whites, other racial groups had lower odds of TMD (OR = 0.2). - Higher educational attainment was associated with greater odds of TMD (OR = 2). - There was a conspicuous lack of association between TMD and income, satisfaction with socioeconomic position, and health insurance coverage.
Reissmann, 2012 ²⁶ (Germany)	Age group - Not reported Race - Not reported Education Less than University: n = 46 University: n = 23 Job - Not reported Household income - Not reported Marital status - Not reported	Age group - Not reported Race - Not reported Education Less than University: n = 490 University: n = 372 Job - Not reported Household income - Not reported Marital status - Not reported	- TMD patients were significantly more often women ($P < 0.001$) and had a lower level of education ($P < 0.05$) than the general population subjects. - No statistically significant differences in age between both groups.

(Continued)

Table 2 (Continued).

Study First Author, Year	Exposure (Sociodemographic Factors)	Comparisons	Outcomes
Blanco-Hungria, 2012 ²⁷ (Spain)	Age categories (years) 16–29: n = 151 30–40: n = 176 41–60: n = 142 ≥ 61: n = 279 Race - Not reported Education Less than University: n = 564 University: n = 184 Job Employed: n = 373 Unemployed: n = 369 Household income 6000: n = 181 6000–15,000: n = 306 15,000–24,000: n = 211 > 24,000: n = 50 Marital status Married: n = 473 No partner (single, divorced or widowed): n = 275	No comparison group.	- The characteristic pain intensity (CPI) score was almost 15 points higher in women than in men (55.73 and 40.91 respectively). - The patients with a lesser educational level yielded higher CPI scores (pain intensity increase of over 5 points), while those with a higher education level yielded comparatively lower pain scores. - Marital status: divorced patients reported higher intensity pain ($P < 0.05$), followed by married subjects. Married or divorced status implied a pain intensity increment of 11.8 and 23.3 points, respectively. - No statistically significant relationship between pain intensity and age group, occupation at the time of the study (employed or otherwise), or income was found.
Slade, 2013 ⁵ (USA)	Age categories (years) 18–24: n = 1421 25–34: n = 736 35–44: n = 580 Race White: n = 1448 Black/African American: n = 766 Asian: n = 256 Hispanic: n = 178 Other: n = 89 Education Less than University: n = 1538 University: n = 1164 Not reported: n = 35 Job - Not reported Household income (US\$) ≤ 20,000/year: n = 421 20,000 – < 40,000: n = 493 40,000 – < 80,000: n = 583 ≥ 80,000: n = 624 Non-reported: n = 616 Marital status Married: n = 539 No partner (single, divorced, widowed): n = 2156 Not reported: n = 42	No comparison group.	TMD incidence was positively associated with age, whereas women had only slightly greater incidence than men (hazard ratio = 1.30). Compared to whites, Asians had lower TMD incidence whereas African Americans had greater incidence, although the latter was attenuated somewhat after adjusting for satisfaction with socioeconomic circumstances. - First-onset TMD increased according to age, from 2.5%/year among 18- to 24-year-olds to 4.5%/year among 35- to 44-year-olds. - Marital status was not significantly associated with TMD incidence.

Blanco-Aguilera, 2014 ²⁸ (Spain)	Age (years) Women mean age: 42.15 ± 14.63 Men mean age: 41.48 ± 17.28 Age group (years) 16–29: n = 98 30–40: n = 106 41–60: n = 154 ≥ 61: n = 49 Race - Not reported Education (years) - Not reported Job - Not reported Household income - Not reported Marital status Married: n = 216 No partner (single, divorced, widowed): n = 186 Not reported: n = 5	No comparison group.	Significant association for gender, age, marital status, and pain duration.
Lei, 2015 ²⁹ (China)	Myofascial Pain (n = 128, 103W, 25M) Age (years) - 33.59 ± 15.30 Race - Not reported Education Less than University: n = 51 University: n = 77 Job - Not reported Household income - Not reported Marital status - Not reported	Non-myofascial Pain (n = 382, 284W, 98M) Age (years) - 30.22 ± 13.99 Race - Not reported Education Less than University: n = 251 University: n = 131 Job - Not reported Household income - Not reported Marital status - Not reported	The myofascial pain group had the average age significantly higher than those of non-myofascial pain group. - No significant difference in sex distribution was observed between the two groups.
Dahan, 2016 ³⁰ (Canada)	Myofascial TMD (n = 121, 102W, 19M) Age (years) - 42.5 ± 1.4 Race - Not reported Education - Not reported Job Part-time employed: n = 21 Full-time employed: n = 64 Unemployed: n = 25 Retired (on disability): n = 11 Household income - Not reported Marital status Married: n = 55 No partner (single, divorced, widowed): n = 66	Non-myofascial TMD (n = 59, 47F, 12M) Age (years) - 43.4 ± 2.1 Race - Not reported Education - Not reported Job Part-time employed: n = 11 Full-time employed: n = 28 Unemployed: n = 19 Retired (on disability): n = 1 Household income - Not reported Marital status Married: n = 37 No partner (single, divorced, widowed): n = 22	- No sociodemographic differences were observed between the myofascial TMD (m-TMD) and non-myofascial TMD (n-TMD) groups.

(Continued)

Table 2 (Continued).

Study First Author, Year	Exposure (Sociodemographic Factors)	Comparisons	Outcomes
Su, 2017 ³¹ (Netherlands)	Age (years) Female mean age: 43.4 ± 14.5 Male mean age: 42.1 ± 15.2 Race - Not reported Education (years) - Not reported Job - Not reported Household income - Not reported Marital status - Not reported	No comparison group.	- Higher pain intensity tended to be associated with younger age. - Women were more likely to have higher pain intensity than men. - Men were more likely to have higher pain-related disability than females and age was not associated with pain-related disability.
Di Paolo, 2017 ³² (Italy)	Headache (n = 625) Age categories (years) ≤ 15: n = 10 16–25: n = 142 26–40: n = 182 41–50: n = 166 51–60: n = 72 61–70: n = 37 > 70: n = 6 Race - Not reported Education - Not reported Job - Not reported Household income - Not reported Marital status - Not reported	Without Headache (n = 304) Age categories (years) ≤ 15: n = 1 16–25: n = 73 26–40: n = 53 41–50: n = 45 51–60: n = 69 61–70: n = 51 > 70: n = 12 Race - Not reported Education - Not reported Job - Not reported Household income - Not reported Marital status - Not reported	- Sociodemographic factors did not show a statistically significant correlation in either group. - In the Headache group, the age classes most frequently involved were 26–40 (n = 182) and 41–50 (n = 166).
Santiago, 2019 ³³ (USA)	Age (years) TMD muscle pain: 40.3 ± 15.5 TMD muscle + joint pain: 40.3 ± 14.7 Race Hispanic: TMD muscle pain: n = 5 TMD muscle + joint pain: n = 22 Education (years) TMD muscle pain: mean study time: 15.4 ± 2.3, n = 34 TMD muscle + joint pain: mean study time: 15.5 ± 2.3, n = 88 Job - Not reported Household income (US\$) > 15,000: TMD muscle pain: n = 24 TMD muscle + joint pain: n = 65 Marital status - Not reported	Control group (n = 46, all women) Age (years) - 36.1 ± 13.5 Race - Hispanic: n = 9 Education (years) - 15.7 ± 2.3 Job - Not reported Household income (US\$) > 15,000: n = 32 Marital status - Not reported	- No significant differences were found between muscle pain and muscle + joint pain groups on demographic variables or comorbid fibromyalgia.

Miller, 2019 ³⁴ (USA)	Low Impact Pain, n = 563 (437 W, 126 M) Age categories (years) 18–24: n = 229 25–34: n = 205 35–44: n = 129 Race White: n = 429 Black/African American: n = 49 Asian: 29 Hispanic: 38 Other: 18 Education - Not reported Job - Not reported Household income - Not reported Marital status - Not reported Study site University of North Carolina: n = 177 University of Buffalo: n = 122 University of Florida: n = 173 University of Maryland: n = 91	High Impact Pain, n = 283 (215 W, 68 M) Age categories (years) 18–24: n = 85 25–34: n = 100 35–44: n = 98 Race White: n = 175 Black/African American: n = 71 Asian: 9 Hispanic: 18 Other: 10 Education - Not reported Job - Not reported Household income - Not reported Marital status - Not reported Study site University of North Carolina: n = 62 University of Buffalo: n = 79 University of Florida: n = 64 University of Maryland: n = 78	- Older people were more likely to experience high-impact pain. - Sex distribution by impact was very similar. There were more women than men in the sample of cases, with the 3:1 female to male ratio observed in both low- and high-impact groups. - Gender had no effect on pain impact. - Black/African American people were more likely to experience high-impact pain compared to other racial/ethnic categories and had 3.5 times the odds of having high-impact pain compared to whites (AUC = 0.34, 95% CI (2.4, 5.2)). - Black/African American people were older than participants from other racial groups. - People identified as Asian, Hispanic, other or multiple racial/ethnic groups did not have elevated estimates of high-impact pain.
Balik, 2019 ³⁵ (Turkey)	Age categories - Not reported Race - Not reported Education (years) ≤ 8: n = 50 > 9: n = 54 Job Employed (full or part-time): n = 57 Unemployed: n = 47 Household income - Not reported Marital status Married: n = 51 No partner (single, divorced or widowed): n = 53	No comparison group.	Significant differences were found in the functional limitation subscale in terms of educational level ($P = 0.036$), and employment status ($P = 0.042$). - There were no differences found in the physical and mental component summary scores and its subscales in terms of socio-demographic variables. - Weak correlations were found in age/role limitations related to emotional problems ($r = -0.203$, $P < 0.05$); age/vitality ($r = -0.243$, $P < 0.05$); age/social functioning ($r = -0.229$, $P < 0.05$). - Lower educational level (Odds Ratio = 0.08, 95% Confidence Interval = 0.01 to 0.56), was found to be one of the most important predictors for higher pain-related disability.

(Continued)

Table 2 (Continued).

Study First Author, Year	Exposure (Sociodemographic Factors)	Comparisons	Outcomes
Arikan, 2019 ³⁶ (Turkey)	Age (years) Muscle disorders, mean age: 25.52 ± 7.61 (16W, 9M) Disc Displacement, mean age: 24.59 ± 7.82 (21W, 6M) Other common diseases, mean age: 48.6 ± 8.62 (22W, 3M) Height (cm) Muscle disorders: 167.96 ± 10.29 Disc Displacement: 165.52 ± 6.9 Other common diseases: 165.13 ± 8.92 Weight (kg) Muscle disorders: 65.44 ± 12.57 Disc Displacement: 65.26 ± 13.69 Other common diseases: 70.36 ± 12.92 Body Mass Index (kg/m²) Muscle disorders: 23.18 ± 3.69 Disc Displacement: 23.66 ± 4.05 Other common diseases: 25.77 ± 4.33 Race - Not reported Education (years) - Not reported Job - Not reported Household income - Not reported Marital status - Not reported	No comparison group.	Mean age of the Other common diseases group were significantly higher than the ones in Muscle disorders and Disc displacement groups ($P < 0.001$). Higher the age, increase the expectation of osteoarthritis and osteoarthritis. - There was no difference among the three groups in weight, height and body mass index (BMI).
Resende, 2020 ³⁷ (Brazil)	TMD, n = 60 (48W, 12M) Age categories - Not reported Race - Not reported Education - Not reported Job Employed: n = 20 Unemployed: n = 39 Household income - Not reported Marital status Married: n = 30 No partner (single, divorced or widowed): n = 28	Without TMD, n = 60 (32W, 28M) Age categories - Not reported Race - Not reported Education - Not reported Job Employed: n = 8 Unemployed: n = 52 Household income - Not reported Marital status Married: n = 45 No partner (single, divorced or widowed): n = 15	Among patients with TMD, 60% were women ($P = 0.002$), 65.1% were single ($P = 0.009$), and 71.4% were employed ($P = 0.008$). - Sociodemographic data showed an association with TMD: being woman (OR = 3.5), being employed (OR = 3.3; $P = 0.008$), and do not have partner (OR = 2.8; $P = 0.009$).

de Caxias, 2021 ³⁸ (Brazil)	Practice of social isolation Age categories - Not reported Race - Not reported Education Current studying: n = 431 Not studying: n = 1621 Job Worker: n = 1460 Not employed: n = 592 Household income - Not reported Marital status - Not reported Social class (household gross monthly income – Brazilian currency) A (> R\$ 15,760): n = 259 B (> 7880): n = 667 C (> 3152): n = 724 D (> 1576): n = 305 E (< 1576): n = 97	No Practice of social isolation Age categories - Not reported Race - Not reported Education Current studying: n = 31 Not studying: n = 318 Job Worker: n = 212 Not employed: n = 37 Household income - Not reported Marital status - Not reported Social class (household gross monthly income) A (> R\$ 15,760): n = 24 B (> R\$ 7880): n = 81 C (> R\$ 3152): n = 99 D (> R\$ 1576): n = 30 E (< R\$ 1576): n = 15	- Gender was associated with “pain/stiffness in the jaw on awakening” ($P = 0.037$), “change of pain during jaw habits” ($P = 0.034$) and “perception of change in the situations mentioned in the TMD-Pain Screener” ($P = 0.020$), “depression” ($P = 0.012$), “anxiety” ($P = 0.006$) and “impact of the event” ($P = 8.3E-11$). No associations were found between “gender” and “presence of pain in the jaw and temporalis” ($P = 0.070$), “chewing hard food” (question 3.a from the TMD-Pain Screener) ($P = 0.735$), “opening or moving the jaw” (question 3.b from the TMD-Pain Screener) ($P = 0.708$), “other jaw activities” (question 3.d from the TMD-Pain Screener) ($P = 0.101$), nor “presence of pain for three months” ($P = 0.102$). - Men presented a 28% lesser chance of having pain/stiffness in the jaw on awakening, were 1.34 times more likely to have changes of pain during jaw habits and were 1.23 times more likely to perceive changes of situations mentioned in the TMD-Pain Screener.
Delgado-Delgado, 2021 ³⁹ (Spain)	Age (years) Women mean age: 27.1 ± 8.9 Men mean age: 29.2 ± 11.6 Height (m) Women: 1.63 ± 0.03 Men: 1.80 ± 0.02 Weight (kg) Women: 59.4 ± 4.5 Men: 80.7 ± 5.8 Body Mass Index (kg/m^2) Women: 22.2 ± 1.2 Men: 24.9 ± 1.6 Race - Not reported Education (years) - Not reported Job - Not reported Household income - Not reported Marital status - Not reported	No comparison group.	- TMD was not associated with sociodemographic features. - Earlier age was associated with presence of parafunctional disorders ($P < 0.05$). Anxiety, a pain predictor, was negatively associated with height and weight ($P < 0.05$). - Neither awake nor sleep bruxism was associated with none of the variables assessed.

(Continued)

Table 2 (Continued).

Study First Author, Year	Exposure (Sociodemographic Factors)	Comparisons	Outcomes
Tavares, 2021 ⁴⁰ (Brazil)	Age group (years) 60–69: n = 7 70–79: n = 7 ≥ 80: n = 1 Race Black: n = 4 White: n = 4 Yellow: n = 1 Brown: n = 5 Other: n = 1 Education Less than University: n = 8 University: n = 7 Job - Not reported Household income < 1 minimum wage: n = 2 1–2 minimum wage: n = 8 3–4 minimum wage: n = 4 5–10 minimum wage: n = 1 >60000 Can\$/year: n = 8 Non-reported: n = 5 Marital status Married: n = 9 No partner (single, divorced, widowed): n = 6	No comparison group.	- The sociodemographic profile of elderly people who had Parkinson's and associated TMD and depression were of the male sex, married or with a partner, on a low income, with nine or more years of schooling, and moderate stage of the disease.
Lei, 2021 ⁴¹ (China)	Painful TMD (n = 519, mean age = 40.0 ± 16.6 y) Age categories - Not reported Race - Not reported Education - Not reported Job - Not reported Household income - Not reported Marital status - Not reported	Non-painful TMD (n = 560, mean age = 25.9 ± 11.3 y) Age categories - Not reported Race - Not reported Education - Not reported Job - Not reported Household income - Not reported Marital status - Not reported	- Participants with painful TMDs were often older. - Higher prevalence of nonpainful TMD in younger subjects.

Mendonça, 2022 ⁴² (Brazil)	Age categories (years) < 30: n = 27 ≥ 30: n = 14 Race White: n = 23 Black: n = 1 Yellow: n = 3 Brown: n = 14 Education Less than University: n = 29 University: n = 12 Job Employed: n = 24 Unemployed (student): n = 17 Household income (minimum wage) < 1: n = 6 1–1.9: n = 5 2–2.9: n = 7 3–5: n = 11 > 6: n = 12 Marital status Married: n = 4 No partner (single, divorced or widowed): n = 37	Before and during COVID-19 pandemic. - Same sociodemographic characteristics, changes related to Job (occupation during COVID-19) Unaltered: n = 10 Home-office: n = 6 No occupation: n = 25	- Before pandemic (T1), subject's occupation was associated with OHIP-14 global score, physical pain, and physical disability domains. - During pandemic (T2), age was associated with OHIP-14 global scores, physical pain, psychological discomfort, and psychological disability domains.
---------------------------------------	--	---	--

As for race, no significant difference was found in most of the evaluated studies, except for one in which Black/African American people were more likely to experience high-impact pain compared to other racial/ethnic categories.³⁴ When it comes to education, two articles reported that higher educational level, could be a risk factor for TMDs, because it may be associated with having more stressful jobs.^{18,25} Conversely, three studies demonstrated that a lower level of education, was associated with a higher pain degree.^{20,27,35} Further, being employed or not, in general, was not a risk factor for TMDs.

Considering the marital status, no difference was usually found, but when reported it was considered that divorced individuals reported higher intensity pain ($p < 0.05$), followed by married ones.

Discussion

The influence of sociodemographic factors on the development, presentation, and treatment outcomes of chronic painful conditions is widely acknowledged.^{43–45} Understanding these factors can inform the creation of tailored prevention programs and personalized treatment methods, ultimately enhancing patient outcomes and improving the quality of life for individuals with TMDs. This scoping review highlights age, gender, marital status, education level, and profession as the most frequently studied sociodemographic factors.

Age was investigated in every study in this review. While TMDs can manifest in any age group, research indicates that the incidence rises during adolescence and young adulthood, peaking between 20 and 40 years of age. This heightened prevalence may be linked to hormonal fluctuations, psychosocial stress, and an increased vulnerability to injury or trauma during this life stage.^{8,46–48} With aging, structural changes occur in the joints and surrounding tissues. Wear on joint cartilage, bone remodeling, and tissue degeneration can impact the function and stability of the TMJ, potentially contributing to TMDs such as disc displacement, as noted in Velly et al case–control study.^{17,49–51}

Similar to age, gender was a focal point across all studies in this review. Extensive evidence highlights gender disparities in the prevalence of TMDs, with women exhibiting a higher prevalence compared to men in diverse populations and geographic regions (Table 1). Women tend to perceive oral health, particularly orofacial pain, more negatively than men.²⁸ This gender bias is influenced by hormonal factors, anatomical variations in the TMJ and muscles, psychosocial factors, and differences in pain reporting and mechanisms.^{31,52–54} A comprehensive understanding of gender-specific aspects in TMDs can contribute to the development of personalized treatment strategies and targeted interventions to enhance TMDs management.

Marital status was investigated in only 14 studies included in this review. Unlike other health conditions, such as coronary diseases, where the impact of marital status is well documented, the consideration of this factor in the realm of TMDs remains limited. Research in coronary diseases shows that a satisfying marital relationship reduces biological, lifestyle, and psychosocial risk factors associated with disease development.⁵⁵ However, in the context of TMDs, only a few studies have explored this variable. The current findings indicate that individuals who are divorced, separated, or widowed exhibit a higher prevalence of TMDs compared to those who are married or single. This association may stem from the psychosocial ramifications of relationship dissolution.^{20,37}

In general, a patient's level of education is a significant factor, as higher education tends to correlate with greater self-care and motivation to seek specialized help for addressing pathologies, particularly painful ones.⁵⁶ The current review reveals that studies examining educational levels indicate that individuals with lower academic degrees or fewer years of education may face an elevated risk of developing TMDs and experiencing more severe symptoms. Health behaviors, psychosocial factors, pain perception, and treatment adherence are all potential mediators in the relationship between educational level and TMDs.^{19,26,27,35}

The link between a patient's profession and TMDs has been a focal point in research investigating occupational factors and their potential influence on TMDs prevalence, risk factors, and symptomatology. In this review, numerous studies have shed light on the correlation between profession and TMDs. Various occupational factors have been identified as potential risk factors for the development of TMDs, with variations based on the nature of the profession. Specific professions have been highlighted as potentially having a higher prevalence of TMDs or increased risk factors. These include occupations involving prolonged or repetitive use of the masticatory system, such as playing musical instruments or professions requiring extensive speaking or voice use. Furthermore, professions characterized by high levels of stress, such as teaching, computer office work, healthcare, military service, and emergency services, may also exhibit an increased risk of TMDs.^{57–61}

In this sociodemographic study segment, parallels can be drawn with other chronic painful conditions, revealing striking similarities. For example, both fibromyalgia and chronic migraine conditions tend to disproportionately affect young women with lower levels of education, thus adversely impacting their personal relationships.^{62–64} These shared patient profile similarities also indicate that individuals with TMD could potentially benefit from successful treatment strategies like those used for other conditions, and vice versa.

Finally, all results must be interpreted cautiously due to the methodological limitations of the review. While systematic reviews are considered the highest level of scientific evidence, the wide variability in objectives and methodologies of the included studies made it infeasible in this case. Hence, the decision to conduct a scoping review was based on its capacity to map and provide an overview of the research field, allowing for a flexible analysis of the addressed studies. Moving forward, when there are scientific articles with relevant methodology, a systematic review on the subject should also be conducted. Furthermore, we strongly recommend that future clinical studies on individuals with TMDs include data collection on the not commonly reported sociodemographic factors such as education, job, income, and marital status. By doing that it will allow to explore the actual relationship of sociodemographic factors, especially for individuals with painful TMDs. As a final remark, the results of this review aimed to contribute to the understanding of the influence of various sociodemographic factors on TMD occurrence, but were also an attempt to characterize the individuals suffering from TMD. In addition, by characterizing the individuals with TMD regarding their sociodemographic variables, their psychosocial variables and correlate them to TMDs, especially painful TMDs, might provide clinicians with tools and categorizations that will help them setting up individualized treatment plans based on their condition but also based on sociodemographic and psychosocial factors and variables affecting and causing the condition. That would in turn improve treatment outcome and prognosis, while it will reduce individual pain and suffering.

Conclusion

Given the conflicting results of the included studies on TMD individuals sociodemographic profiles, it can be inferred that young women with lower educational levels and without a partner were the most susceptible to experiencing TMD signs and symptoms.

Data Sharing Statement

Datasets related to this article will be available upon request to the corresponding author.

Acknowledgments

We acknowledge the Foundation for Science and Technology, I.P., under the project UIDB/04585/2020.

Funding

This study was financed by national funds through the FCT—Foundation for Science and Technology, I.P., under the project UIDB/04585/2020. The funding source was not involved in study design, collection, analysis and interpretation of data, decision to publish, or preparation of the manuscript.

Disclosure

The authors declare no conflicts of interest in this work.

References

1. List T, Jensen RH. Temporomandibular disorders: old ideas and new concepts. *Cephalalgia*. 2017;37(7):692–704. doi:10.1177/0333102416686302
2. Fillingim RB, Wallace MR, Herbstman DM, Ribeiro-Dasilva M, Staud R. Genetic contributions to pain: a review of findings in humans. *Oral Dis*. 2008;14(8):673–682. doi:10.1111/j.1601-0825.2008.01458.x
3. Durham J. Temporomandibular disorders (TMD): an overview. *Oral Surg*. 2008;1(2):60–68. doi:10.1111/j.1752-248X.2008.00020.x
4. Progiante PS, Pattussi MP, Lawrence HP, Goya S, Grossi PK, Grossi ML. Prevalence of temporomandibular disorders in an adult Brazilian community population using the research diagnostic criteria (Axes I and II) for temporomandibular disorders (the maringá study). *Int J Prosthodont*. 2015;28(6):600–609. doi:10.11607/ijp.4026
5. Slade GD, Fillingim RB, Sanders AE, et al. Summary of findings from the OPPERA prospective cohort study of incidence of first-onset temporomandibular disorder: implications and future directions. *J Pain*. 2013;14(12 Suppl):T116–T124. doi:10.1016/j.jpain.2013.09.010

6. Maixner W, Diatchenko L, Dubner R, et al. Orofacial pain prospective evaluation and risk assessment study – the OPPERA study. *J Pain*. 2011;12 (11 Suppl):T4–T11.e2. doi:10.1016/j.jpain.2011.08.002
7. Bair E, Gaynor S, Slade GD, et al. Identification of clusters of individuals relevant to temporomandibular disorders and other chronic pain conditions: the OPPERA study. *Pain*. 2016;157(6):1266–1278. doi:10.1097/j.pain.0000000000000518
8. Lövgren A, Häggman-Henrikson B, Visscher CM, Lobbezoo F, Marklund S, Wänman A. Temporomandibular pain and jaw dysfunction at different ages covering the lifespan – a population based study. *Eur J Pain*. 2016;20(4):532–540. doi:10.1002/ejp.755
9. Edwards CL, Fillingim RB, Keefe F. Race, ethnicity and pain. *Pain*. 2001;94(2):133–137. doi:10.1016/S0304-3959(01)00408-0
10. Edwards RR, Fillingim RB, Yamauchi S, et al. Effects of gender and acute dental pain on thermal pain responses. *Clin J Pain*. 1999;15(3):233–237. doi:10.1097/00002508-199909000-00011
11. Green CR, Anderson KO, Baker TA, et al. The unequal burden of pain: confronting racial and ethnic disparities in pain. *Pain Med*. 2003;4 (3):277–294. doi:10.1046/j.1526-4637.2003.03034.x
12. Von Korff M, Dworkin SF, Le Resche L, Kruger A. An epidemiologic comparison of pain complaints. *Pain*. 1988;32(2):173–183. doi:10.1016/0304-3959(88)90066-8
13. Visscher CM, Ligthart L, Schuller AA, et al. Comorbid disorders and sociodemographic variables in temporomandibular pain in the general Dutch population. *J Oral Facial Pain Headache*. 2015;29(1):51–59. doi:10.11607/ofph.1324
14. Dworkin SF, LeResche L. Research diagnostic criteria for temporomandibular disorders: review, criteria, examinations and specifications, critique. *J Craniomandib Disord*. 1992;6(4):301–355.
15. Schiffman E, Ohrbach R, Truelove E, et al. Diagnostic Criteria for Temporomandibular Disorders (DC/TMD) for clinical and research applications: recommendations of the international RDC/TMD consortium network and orofacial pain special interest group. *J Oral Facial Pain Headache*. 2014;28(1):6–27. doi:10.11607/jop.1151
16. Morgan RL, Whaley P, Thayer KA, Schünemann HJ. Identifying the PECO: a framework for formulating good questions to explore the association of environmental and other exposures with health outcomes. *Environ Int*. 2018;121(Pt 1):1027–1031. doi:10.1016/j.envint.2018.07.015
17. Velly AM, Gornitsky M, Philippe P. A case-control study of temporomandibular disorders: symptomatic disc displacement. *J Oral Rehabil*. 2002;29(5):408–416. doi:10.1046/j.1365-2842.2002.00913.x
18. Gesch D, Bernhardt O, Kocher T, John U, Hensel E, Alte D. Association of malocclusion and functional occlusion with signs of temporomandibular disorders in adults: results of the population-based study of health in Pomerania. *Angle Orthod*. 2004;74(4):512–520. doi:10.1043/0003-3219(2004)074<0512:AOMAFO>2.0.CO;2
19. Bernhardt O, Gesch D, Schwahn C, et al. Risk factors for headache, including TMD signs and symptoms, and their impact on quality of life. Results of the Study of Health in Pomerania (SHIP). *Quintessence Int*. 2005;36(1):55–64.
20. Mundt T, Mack F, Schwahn C, et al. Gender differences in associations between occlusal support and signs of temporomandibular disorders: results of the population-based Study of Health in Pomerania (SHIP). *Int J Prosthodont*. 2005;18(3):232–239.
21. Casanova-Rosado JF, Medina-Solis CE, Vallejos-Sánchez AA, Casanova-Rosado AJ, Hernández-Prado B, Ávila-burgos L. Prevalence and associated factors for temporomandibular disorders in a group of Mexican adolescents and youth adults. *Clin Oral Investig*. 2006;10(1):42–49. doi:10.1007/s00784-005-0021-4
22. Selaimen CM, Jeronimo JC, Brilhante DP, Lima EM, Grossi PK, Grossi ML. Occlusal risk factors for temporomandibular disorders. *Angle Orthod*. 2007;77(3):471–477. doi:10.2319/0003-3219(2007)077[0471:ORFFTD]2.0.CO;2
23. Ommerborn MA, Hugger A, Kruse J, et al. The extent of the psychological impairment of prosthodontic outpatients at a German University Hospital. *Head Face Med*. 2008;4:23. doi:10.1186/1746-160X-4-23
24. Quinteromarmol-Juárez M, Espinosa-de Santillana IA, Martínez-Torres J, Vargas-García HA. Trastornos temporomandibulares y funcionamiento familiar. *Rev Med Inst Mex Seguro Soc*. 2008;46(5):473–478.
25. Slade GD, Bair E, By K, et al. Study methods, recruitment, socio-demographic findings and demographic representativeness in the OPPERA study. *J Pain*. 2011;12(11 Suppl):T12–T26. doi:10.1016/j.jpain.2011.08.001
26. Reissmann DR, John MT, Schierz O, Seedorf H, Doering S. Stress-related adaptive versus maladaptive coping and temporomandibular disorder pain. *J Orofac Pain*. 2012;26(3):181–190.
27. Blanco-Hungria A, Rodríguez-Torronteras A, Blanco-Aguilera A, et al. Influence of sociodemographic factors upon pain intensity in patients with temporomandibular joint disorders seen in the primary care setting. *Med Oral Patol Oral Cir Bucal*. 2012;17(6):e1034–e1041. doi:10.4317/medoral.17576
28. Blanco-Aguilera A, Blanco-Hungria A, Biedma-Velázquez L, et al. Application of an oral health-related quality of life questionnaire in primary care patients with orofacial pain and temporomandibular disorders. *Med Oral Patol Oral Cir Bucal*. 2014;19(2):e127–e135. doi:10.4317/medoral.19061
29. Lei J, Liu MQ, Yap AU, Fu KY. Sleep disturbance and psychologic distress: prevalence and risk indicators for temporomandibular disorders in a Chinese population. *J Oral Facial Pain Headache*. 2015;29(1):24–30. doi:10.11607/ofph.1301
30. Dahan H, Shir Y, Nicolau B, Keith D, Allison P. Self-reported migraine and chronic fatigue syndrome are more prevalent in people with myofascial vs nonmyofascial temporomandibular disorders. *J Oral Facial Pain Headache*. 2016;30(1):7–13. doi:10.11607/ofph.1550
31. Su N, Lobbezoo F, van Wijk A, van der Heijden GJ, Visscher CM, van der Heijden GJMG. Associations of pain intensity and pain-related disability with psychological and socio-demographic factors in patients with temporomandibular disorders: a cross-sectional study at a specialised dental clinic. *J Oral Rehabil*. 2017;44(3):187–196. doi:10.1111/joor.12479
32. Di Paolo C, D’Urso A, Papi P, et al. Temporomandibular disorders and headache: a retrospective analysis of 1198 patients. *Pain Res Manag*. 2017;2017:3203027. doi:10.1155/2017/3203027
33. Santiago V, Raphael K. Absence of joint pain identifies high levels of sleep masticatory muscle activity in myofascial temporomandibular disorder. *J Oral Rehabil*. 2019;46(12):1161–1169. doi:10.1111/joor.12853
34. Miller VE, Poole C, Golightly Y, et al. Characteristics associated with high-impact pain in people with temporomandibular disorder: a cross-sectional study. *J Pain*. 2019;20(3):288–300. doi:10.1016/j.jpain.2018.09.007
35. Balik A, Peker K, Ozdemir-Karatas M. Comparisons of measures that evaluate oral and general health quality of life in patients with temporomandibular disorder and chronic pain. *Cranio*. 2019;39(4):310–320. doi:10.1080/08869634.2019.1622869

36. Arikan H, Sertel M, Bas B. Evaluation of the musculoskeletal systems and kinesiophobia of the individuals with temporomandibular disorders. *Eurasian J Med Oncol*. 2019;3(2):132–138. doi:10.14744/ejmo.2019.22613
37. Resende CMBM, Rocha LGDDS, Paiva RP, et al. Relationship between anxiety, quality of life, and sociodemographic characteristics and temporomandibular disorder. *Oral Surg Oral Med Oral Pathol Oral Radiol*. 2020;129(2):125–132. doi:10.1016/j.oooo.2019.10.007
38. de Caxias FP, Athayde FRF, Januzzi MS, Pinheiro LV, Turcio KHL. Impact event and orofacial pain amid the COVID-19 pandemic in Brazil: a cross-sectional epidemiological study. *J Appl Oral Sci*. 2021;29:e20210122. doi:10.1590/1678-7757-2021-0122
39. Delgado-Delgado R, Iriarte-álvarez N, Valera-Calero JA, Centenera-Centenera MB, Garnacho-Garnacho VE, Gallego-Sendarrubias GM. Association between temporomandibular disorders with clinical and sociodemographic features: an observational study. *Int J Clin Pract*. 2021;75(5):e13961. doi:10.1111/ijcp.13961
40. Tavares RB, Oliveira JS, Faccio PF, Coriolano MGWS, Asano NMJ, Lins CCSA. Sociodemographic profile of elderly people with temporomandibular disorder and depression in combination with parkinson's disease. *Pesqui Bras Odontopediatria Clin Integr*. 2021;21:e0169. doi:10.1590/pboci.2021.026
41. Lei J, Yap AU, Zhang M, Fu KY. Temporomandibular disorder subtypes, emotional distress, impaired sleep, and oral health-related quality of life in Asian patients. *Oral Epidemiol*. 2021;49(6):543–549. doi:10.1111/cdoe.12643
42. Mendonça AKR, Fontoura LPG, Rocha TDD, et al. Influence of the COVID-19 pandemic on pain and oral health-related quality of life in women with temporomandibular disorder. *Dental Press J Orthod*. 2022;27(3):e2220422. doi:10.1590/2177-6709.27.3.e2220422.oar
43. Mills SEE, Nicolson KP, Smith BH. Chronic pain: a review of its epidemiology and associated factors in population-based studies. *Br J Anaesth*. 2019;123(2):e273–e283. doi:10.1016/j.bja.2019.03.023
44. Joseph V, Huo J, Cook R, et al. Sociodemographic and clinical characteristics associated with worst pain intensity among cancer patients. *Pain Manag Nurs*. 2022;23(4):424–429. doi:10.1016/j.pmn.2021.11.006
45. Atzeni F, Alciati A, Bazzichi L, et al. Sociodemographic factors in fibromyalgia: results from the Italian fibromyalgia registry. *Clin Exp Rheumatol*. 2022;40(6):1183–1188. doi:10.55563/clinexpheumatol/64963d
46. Kuttilla M, Niemi PM, Kuttilla S, Alanen P, Le Bell Y. TMD treatment need in relation to age, gender, stress, and diagnostic subgroup. *J Orofac Pain*. 1998;12(1):67–74.
47. Manfredini D, Piccotti F, Ferronato G, Guarda-Nardini L. Age peaks of different RDC/TMD diagnoses in a patient population. *J Dent*. 2010;38(5):392–399. doi:10.1016/j.jdent.2010.01.006
48. Zheng Y, Zhu R, Xiao C, et al. Age and gender, but not pain are associated with pressure pain thresholds in patients with temporomandibular disorders: a cross-sectional study. *J Pain Res*. 2023;16:2205–2216. doi:10.2147/JPR.S414276
49. Stocum DL, Roberts WE. Part I: development and physiology of the temporomandibular joint. *Curr Osteoporos Rep*. 2018;16(4):360–368. doi:10.1007/s11914-018-0447-7
50. Roberts WE, Stocum DL. Temporomandibular Joint (TMJ)—regeneration, degeneration, and adaptation. *Curr Osteoporos Rep*. 2018;16(4):369–379. doi:10.1007/s11914-018-0462-8
51. Ilgunas A, Häggman-Henrikson B, Visscher CM, et al. The longitudinal relationship between jaw catching/locking and pain. *J Dent Res*. 2023;102(4):383–390. doi:10.1177/00220345221138532
52. Fillingim RB, King CD, Ribeiro-Dasilva MC, Rahim-Williams B, Riley JL. Sex, gender, and pain: a review of recent clinical and experimental findings. *J Pain*. 2009;10(5):447–485. doi:10.1016/j.jpain.2008.12.001
53. Bueno CH, Pereira DD, Pattussi MP, Grossi PK, Grossi ML. Gender differences in temporomandibular disorders in adult populational studies: a systematic review and meta-analysis. *J Oral Rehabil*. 2018;45(9):720–729. doi:10.1111/joor.12661
54. Gałczyńska-Rusin M, Pobudek-Radzikowska M, Gawriolek K, Czajka-Jakubowska A. Gender-related biomechanical properties of masseter muscle among patients with self-assessment of bruxism: a comparative study. *J Clin Med*. 2022;11(3):845. doi:10.3390/jcm11030845
55. Gallo LC, Troxel WM, Matthews KA, Kuller LH. Marital status and quality in middle-aged women: associations with levels and trajectories of cardiovascular risk factors. *Health Psychol*. 2003;22(5):453–463. doi:10.1037/0278-6133.22.5.453
56. Zajacova A, Rogers RG, Grodsky E, Grol-Prokopczyk H. The relationship between education and pain among adults aged 30–49 in the United States. *J Pain*. 2020;21(11–12):1270–1280. doi:10.1016/j.jpain.2020.03.005
57. Chisnoiu AM, Picos AM, Popa S, et al. Factors involved in the etiology of temporomandibular disorders – a literature review. *Chujul Med*. 2015;88(4):473–478. doi:10.15386/cjmed-485
58. Jang J-Y, Kwon J-S, Lee DH, Bae J-H, Kim ST. Clinical signs and subjective symptoms of temporomandibular disorders in instrumentalists. *Yonsei Med J*. 2016;57(6):1500–1507. doi:10.3349/ymj.2016.57.6.1500
59. Eli I, Zigler-Garburg A, Winocur E, et al. Temporomandibular disorders and bruxism among sex workers – a cross sectional study. *J Clin Med*. 2022;11(22):6622. doi:10.3390/jcm11226622
60. Bragatto MM, Bevilacqua-Grossi D, Regalo SC, Sousa JD, Chaves TC. Associations among temporomandibular disorders, chronic neck pain and neck pain disability in computer office workers: a pilot study. *J Oral Rehabil*. 2016;43(5):321–332. doi:10.1111/joor.12377
61. Shigeishi H. Association of temporomandibular disorder with occupational visual display terminal use. *Biomed Rep*. 2016;5(1):7–10. doi:10.3892/br.2016.669
62. Mur Martí T, Llordés Llordés M, Custal Jordà M, López Juan G, Martínez Pardo S. Perfil de pacientes con fibromialgia que acuden a los centros de atención primaria en Terrassa [Profile of patients with fibromyalgia who attend primary care centers in Terrassa]. *Reumatol Clin*. 2017;13(5):252–257. doi:10.1016/j.reuma.2016.05.008
63. Buse DC, Manack AN, Fanning KM, et al. Chronic migraine prevalence, disability, and sociodemographic factors: results from the American migraine prevalence and prevention study. *Headache*. 2012;52(10):1456–1470. doi:10.1111/j.1526-4610.2012.02223.x
64. Buse DC, Manack A, Serrano D, Turkel C, Lipton RB. Sociodemographic and comorbidity profiles of chronic migraine and episodic migraine sufferers. *J Neurol Neurosurg Psychiatry*. 2010;81(4):428–432. doi:10.1136/jnnp.2009.192492

Journal of Pain Research**Dovepress****Publish your work in this journal**

The Journal of Pain Research is an international, peer reviewed, open access, online journal that welcomes laboratory and clinical findings in the fields of pain research and the prevention and management of pain. Original research, reviews, symposium reports, hypothesis formation and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-pain-research-journal>