

An Analysis of Preoperative Inflammatory Indicators That Influence the Drainage Tube Retention Time in Patients with Breast Cancer Surgery

Qi Li^{1,*}, Cong Gao¹, Xinrui Zhao¹, Jiahui Li¹, Qinghong Shen¹, Li Chen^{2,*}

¹Department of Breast Surgical Oncology, National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, 100021, People's Republic of China; ²Department of Thyroid and Breast Surgery, Tongji Hospital, Tongji Medical College of Huazhong University of Science and Technology, Wuhan, Hubei, 430030, People's Republic of China

*These authors contributed equally to this work

Correspondence: Qi Li, Department of Breast Surgical Oncology, National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, 100021, People's Republic of China, Email 1589898345@qq.com; Li Chen, Department of Thyroid and Breast Surgery, Tongji Hospital, Tongji Medical College of Huazhong University of Science and Technology, Wuhan, Hubei, 430030, People's Republic of China, Email cswtth@163.com

Objective: The study was aimed to investigate the influence factor between preoperative inflammatory indicators and drainage tube retention time in patients with breast cancer.

Methods: This retrospective study enrolled 121 patients with breast cancer who were undergoing surgery between October 2020 and June 2021. The enumeration data were used the Chi-square test, and the measurement data were used the *t*-test analysis. The univariate and multivariate logistic regression models were performed to access the risk factors for affecting drainage tube retention time in patients with breast cancer. The receiver operating characteristic curve (ROC) was performed to test the prediction effect of the model.

Results: Through the median extraction time of postoperative drainage tube retention time, all patients were divided into two groups: drainage tube retention time (DTRT) < 13 (d) and drainage tube retention time (DTRT) ≥ 13 (d). The results showed that type of surgery, total lymph nodes (TLN), pathological T stage, NLR were related to the drainage tube retention time ($P < 0.05$). Moreover, the univariate and multivariate logistic regression analysis performed that Hb, type of surgery, pathological T stage, chest wall drainage tube, NRI were the independent risk predictors of affecting drainage tube retention time. Furthermore, a significant correlation existed between NRI and drainage tube retention at different times ($P < 0.05$).

Conclusion: NRI is an independent risk factor for postoperative drainage tube extraction time and can effectively predict the probability of drainage tube retention time. Thus, it can also provide personalized nursing intervention for patients with breast cancer after drainage tube retention time and the rehabilitation process.

Keywords: breast cancer, inflammatory indicator, drainage tube, operation, nutritional indicator

Introduction

Breast cancer (BC) is the most common public health threat to females across the global.¹ In recent decades, the number of patients with breast cancer shows an upward trend with each passing year, and the age of onset inclines to become younger in average age.² At the moment, the treatment of breast cancer primarily includes surgery, chemotherapy, radiotherapy, targeted therapy, and endocrine therapy.³ The surgical operation of breast cancer comprises mastectomy and breast-conserving surgery. As a result of mastectomy, patients have extensive trauma, and the drainage tubes are routinely placed after operation.⁴ Nowadays, clinical pathways are commonly designed as a standardized tool for perioperative management of patients with breast cancer who received surgical operations.⁵ Through the long-term

clinical practice and experience of surgical operation in our department, the drainage tube retention time is about two weeks, and the incision cicatrized time is about three weeks, respectively.

It has been found that inflammatory cell and inflammatory response affects tumor proliferation, invasion, metastasis by transforming the tumor immunization microenvironment.⁶ The common peripheral blood inflammatory factors contain white blood cell (W), lymphocyte (L), monocyte (M), neutrophils (N), platelet (P), C-reactive protein (CRP), and form the derived ratio, for example, neutrophil-to-lymphocyte ratio (NLR), derived neutrophil-to-lymphocyte ratio (dNLR), monocyte-to-lymphocyte ratio (MLR), platelet-to-lymphocyte ratio (PLR), are used to forecast the prognosis of tumors.^{7–10}

There are many invasive operations during the operation. The large wound and the length of the operations may bring incision infection complications for the patients, and then cause infection in other parts of the body. Moreover, it also affects the treatment effect, and brings serious physical and mental injury to cancer patients.^{11,12} Drainage tubes are routinely placed after breast cancer surgery. The drainage volume and drainage time after surgical operations will affect postoperative rehabilitation and functional exercise. The purpose of the current study is aimed at the influence factor between preoperative inflammatory indicators and drainage tube retention time in patients with breast cancer, and to provide personalized nursing intervention and recover limb function.

Materials and Methods

Patients' Selection and Data Collection

From October 2020 to June 2021, this retrospective study involved 121 patients with breast cancer at Cancer Hospital Chinese Academy of Medical Sciences. All patients were female and underwent surgery. After mastectomy and axillary dissection, two drainage tubes were put in vacuum aspiration: one at the chest wall and the other at the axilla. After breast-conserving surgery and axillary dissection, one drainage tube was placed to provide suction under negative pressure at the axilla. Moreover, surgical dressing and elastic bandages were applied to eliminate the dead chambers in the chest wall and axilla. In addition, dressings and elastic bandages were used to remove dead chambers in the chest wall and armpit after the operation. This study was approved by the ethics committee of Cancer Hospital Chinese Academy of Medical Sciences and was conducted in accordance with the amended Declaration of Helsinki. And all enrolled patients signed informed consent.

Inclusion Criteria and Exclusion Criteria

The inclusion criteria were as follows: 1) Patients were diagnosed with breast cancer by pathology; 2) Surgical treatment of unilateral breast; 3) All patients had indwelling drainage tube after the operation, and with chest wall drainage tube and/or axillary drainage tube. The exclusion criteria were as follows: 1) with metastasis or other tumors; 2) without indwelling drainage tube after surgery; 3) complicated with chronic diseases and difficult to control, for instance, hypertension or diabetes mellitus.

Inflammatory Index

We analyzed the common inflammatory parameters, for instance, white blood cell (W), red blood cell (R), hemoglobin (Hb), neutrophils (N), lymphocyte (L), monocyte (M), platelet (P), and the complex index, such as neutrophil-to-lymphocyte ratio (NLR), derived neutrophil-to-lymphocyte ratio (dNLR), monocyte-to-lymphocyte ratio (MLR), platelet-to-lymphocyte ratio (PLR), systemic inflammation response index (SIRI), systemic immune-inflammation index (SII), prognostic nutritional index (PNI), nutritional risk index (NRI), albumin-to-fibrinogen ratio (AFR), HALP, controlling nutritional status, and breast immune prognostic index (BIPI).

The dNLR was defined as neutrophil count/(white blood cell count – neutrophil count). The systemic inflammation response index (SIRI) was an indicator that combined the neutrophil, monocyte, and lymphocyte and defined as neutrophil count \times monocyte count/lymphocyte count. The systemic immune-inflammation index (SII) was an indicator that combined the platelet, neutrophil, and lymphocyte and defined as platelet count \times neutrophil count/lymphocyte count. The PNI was defined as albumin level (g/L) + 5 \times total lymphocyte count (10^9 /L). The NRI was defined as $1.519 \times$

albumin level (g/L) + 0.417×(present weight/usual weight×100). The HALP was determined by hemoglobin level (g/L) × albumin level (g/L) × lymphocyte count (10⁹/L)/platelet count (10⁹/L). The controlling nutritional status score (CONUT) based on serum albumin, total cholesterol concentration, and total peripheral lymphocyte counts. The breast immune prognostic index (BIPI) was an indicator that combined the serum lactate dehydrogenase (LDH) level and the derived neutrophil-to-lymphocyte ratio (dNLR).

Statistical Analyses

All statistical analyses were conducted using GraphPad Prism 8.0 software and RStudio software version 3.6.0. The enumeration data were used the Chi-square test, and the measurement data were used the *t*-test analysis. The univariate and multivariate logistic regression models were performed to access the risk factors for affecting drainage tube retention time in patients with breast cancer. The nomogram was validated predicting the risk for drainage tube retention time. Two-sided *P* values less than 0.05 were considered statistically significant.

Results

Clinicopathological Characteristics of All Enrolled Patients

One hundred and twenty-one patients with breast cancer who received surgical treatment were selected as the study subjects from October 2020 to June 2021. The median drainage duration of chest wall suction drain (DDCSD) of all patients was 13 days, and the median drainage duration of the axillary suction drain (DDASD) of all patients was 13 days, respectively. Based on the postoperative drainage tube retention time, which included drainage duration of chest wall suction drain (DDCSD) and drainage duration of the axillary suction drain (DDASD), we choose the median DDCSD as the cutoff value for drainage tube retention time. And all cases were divided into two groups: drainage tube retention time (DTRT) <13 (d) and drainage tube retention time (DTRT) ≥13 (d). The results showed that type of surgery, total lymph nodes (TLN), pathological T stage were related to the drainage tube retention time (*P* < 0.05). All detailed information could be found in Table 1.

Table 1 Clinicopathological Features of Patients with Breast Cancer in the Present Study

n	Level	Overall	DTRT<13 (d)	DTRT ≥13 (d)	p
		N=121	N=67	N=54	
Age (median [IQR])		46.00 [38.00, 56.00]	46.00 [38.00, 57.00]	45.00 [38.25, 55.00]	0.624
Weight (median [IQR])		60.00 [54.00, 66.00]	59.00 [52.00, 65.00]	60.50 [55.00, 66.00]	0.383
Height (median [IQR])		1.61 [1.58, 1.65]	1.61 [1.58, 1.65]	1.62 [1.58, 1.65]	0.738
BMI (median [IQR])		22.83 [20.58, 25.78]	21.99 [20.56, 25.86]	23.21 [21.02, 25.75]	0.274
Healthy lower arm circumference (median [IQR])		19.00 [17.50, 24.00]	19.00 [18.00, 23.50]	19.50 [17.50, 23.88]	0.863
Healthy upper arm circumference (median [IQR])		26.00 [24.00, 28.00]	26.00 [24.00, 28.00]	26.50 [24.00, 28.00]	0.586
Affected side lower arm circumference (median [IQR])		19.00 [18.00, 23.50]	19.00 [18.00, 23.25]	19.00 [17.62, 23.38]	0.712
Affected side upper arm circumference (median [IQR])		26.00 [24.00, 28.00]	26.00 [24.00, 28.00]	26.25 [24.00, 28.00]	0.615
Chest wall drainage tube (median [IQR])		13.00 [10.00, 18.00]	10.00 [6.00, 12.00]	18.00 [15.25, 21.00]	<0.001
Axillary drainage tube (median [IQR])		13.00 [10.00, 17.00]	12.00 [9.00, 13.00]	16.00 [13.25, 20.00]	<0.001
Marital status (%)	Married	107 (88.4)	60 (89.6)	47 (87.0)	0.592
	Unmarried	8 (6.6)	5 (7.5)	3 (5.6)	
	Divorce	5 (4.1)	2 (3.0)	3 (5.6)	
	Widowed	1 (0.8)	0 (0.0)	1 (1.9)	
Occupation (%)	Mental worker	101 (83.5)	57 (85.1)	44 (81.5)	0.511
	Manual worker	1 (0.8)	0 (0.0)	1 (1.9)	
	Others	19 (15.7)	10 (14.9)	9 (16.7)	
Type of surgery (%)	MRM	31 (25.6)	13 (19.4)	18 (33.3)	0.006
	M+SLNB	33 (27.3)	23 (34.3)	10 (18.5)	
	BCS+SLNB	15 (12.4)	13 (19.4)	2 (3.7)	
	BCS+ALND	7 (5.8)	4 (6.0)	3 (5.6)	
	BR	35 (28.9)	14 (20.9)	21 (38.9)	

(Continued)

Table I (Continued).

n	Level	Overall	DTRT<13 (d)	DTRT ≥13 (d)	p
		N=121	N=67	N=54	
Tumor size (%)	≤2cm	54 (44.6)	34 (50.7)	20 (37.0)	0.306
	>2 and <5cm	58 (47.9)	29 (43.3)	29 (53.7)	
	≥5cm	9 (7.4)	4 (6.0)	5 (9.3)	
TLN (median [IQR])		7.00 [4.00, 19.00]	6.00 [3.00, 17.00]	8.00 [5.00, 22.75]	0.049
PLN (median [IQR])		0.00 [0.00, 1.00]	0.00 [0.00, 1.00]	0.00 [0.00, 1.75]	0.683
Histologic type (%)	Noninvasive carcinoma	18 (14.9)	12 (17.9)	6 (11.1)	0.479
	Invasive special carcinoma	5 (4.1)	2 (3.0)	3 (5.6)	
	Invasive nonspecific carcinoma	98 (81.0)	53 (79.1)	45 (83.3)	
Histologic grade (%)	0	6 (5.0)	3 (4.5)	3 (5.6)	0.628
	I	35 (28.9)	22 (32.8)	13 (24.1)	
	II	62 (51.2)	34 (50.7)	28 (51.9)	
	III	18 (14.9)	8 (11.9)	10 (18.5)	
Pathological T stage (%)	Tis/T0	28 (23.1)	16 (23.9)	12 (22.2)	0.050
	T I	56 (46.3)	37 (55.2)	19 (35.2)	
	T2	36 (29.8)	14 (20.9)	22 (40.7)	
	T3	1 (0.8)	0 (0.0)	1 (1.9)	
Pathological N stage (%)	N0	82 (67.8)	47 (70.1)	35 (64.8)	0.891
	N1	22 (18.2)	12 (17.9)	10 (18.5)	
	N2	13 (10.7)	6 (9.0)	7 (13.0)	
	N3	4 (3.3)	2 (3.0)	2 (3.7)	
Pathological TNM stage (%)	Tis/T0	27 (22.3)	16 (23.9)	11 (20.4)	0.746
	I	40 (33.1)	24 (35.8)	16 (29.6)	
	II	37 (30.6)	19 (28.4)	18 (33.3)	
	III	17 (14.0)	8 (11.9)	9 (16.7)	
Molecular subtype (%)	Luminal A	38 (31.4)	25 (37.3)	13 (24.1)	0.416
	Luminal B HER2+	56 (46.3)	30 (44.8)	26 (48.1)	
	Luminal B HER2-	18 (14.9)	9 (13.4)	9 (16.7)	
	HER2 enriched	4 (3.3)	1 (1.5)	3 (5.6)	
	Triple negative	5 (4.1)	2 (3.0)	3 (5.6)	
ER status (%)	Negative	28 (23.1)	16 (23.9)	12 (22.2)	1.000
	Positive	93 (76.9)	51 (76.1)	42 (77.8)	
PR status (%)	Negative	24 (19.8)	14 (20.9)	10 (18.5)	0.923
	Positive	97 (80.2)	53 (79.1)	44 (81.5)	
HER2 (%)	Negative	99 (81.8)	57 (85.1)	42 (77.8)	0.425
	Positive	22 (18.2)	10 (14.9)	12 (22.2)	
Ki-67 (%)	Negative	41 (33.9)	27 (40.3)	14 (25.9)	0.142
	Positive	80 (66.1)	40 (59.7)	40 (74.1)	
AR (%)	Negative	29 (24.0)	20 (29.9)	9 (16.7)	0.140
	Positive	92 (76.0)	47 (70.1)	45 (83.3)	
CK5/6 (%)	Negative	114 (94.2)	62 (92.5)	52 (96.3)	0.625
	Positive	7 (5.8)	5 (7.5)	2 (3.7)	
E-cad (%)	Negative	35 (28.9)	22 (32.8)	13 (24.1)	0.393
	Positive	86 (71.1)	45 (67.2)	41 (75.9)	
EGFR (%)	Negative	104 (86.0)	56 (83.6)	48 (88.9)	0.567
	Positive	17 (14.0)	11 (16.4)	6 (11.1)	
P53 (%)	Negative	50 (41.3)	30 (44.8)	20 (37.0)	0.500
	Positive	71 (58.7)	37 (55.2)	34 (63.0)	
TOP2A (%)	Negative	53 (43.8)	35 (52.2)	18 (33.3)	0.058
	Positive	68 (56.2)	32 (47.8)	36 (66.7)	
Lymph vessel invasion (%)	Negative	99 (81.8)	55 (82.1)	44 (81.5)	1.000
	Positive	22 (18.2)	12 (17.9)	10 (18.5)	
Neural invasion (%)	Negative	104 (86.0)	57 (85.1)	47 (87.0)	0.964
	Positive	17 (14.0)	10 (14.9)	7 (13.0)	

Abbreviations: BMI, body mass index; TLN, Total lymph nodes; PLN, Positive lymph nodes; ER, estrogen receptor; PR, progesterone receptor; HER2, Human Epidermal Growth Factor Receptor 2; AR, androgen receptor; CK5/6, Cytokeratin 5/6; E-cad, E-cadherin; EGFR, Epidermal Growth Factor Receptor; TOP2A, topoisomerase 2A.

The Effect of the Drainage Tube Retention Time for Inflammatory Parameters

The common inflammatory cells included LDH, ALB, CRP, CHOL, FIB, W, R, Hb, N, L, M, P; and the complex inflammatory index, for instance, NLR, dNLR, MLR, PLR, SIRI, SII, PNI, NRI, AFR, HALP; and the complex inflammatory scores, such as CONUT and BIPI. The cutoff values of inflammatory cells and complex inflammatory index or scores were determined by the ROC curve. Moreover, [Figures S1](#) and [S2](#) were developed to show the ROC curves for these complex inflammatory indexes or scores. [Table 2](#) illustrates the comparison of preoperative clinical inflammatory evaluation indexes between the two groups. The results indicated that NRI was associated with drainage tube retention time ($P<0.05$).

Table 2 Comparison of Preoperative Inflammatory Parameters with the Drainage Tube Retention Time

n	Level	Overall	DTRT<13 (d)	DTRT ≥13 (d)	p
		N=121	N=67	N=54	
LDH (median [IQR])		158.10 [141.10, 180.00]	158.10 [141.05, 178.00]	158.55 [141.85, 181.88]	0.741
ALB (median [IQR])		44.40 [42.80, 46.60]	43.90 [42.80, 46.35]	45.30 [42.85, 46.98]	0.370
CRP (median [IQR])		0.02 [0.00, 0.08]	0.03 [0.00, 0.08]	0.02 [0.00, 0.08]	0.688
CHOL (median [IQR])		4.83 [4.25, 5.53]	4.80 [4.24, 5.68]	4.86 [4.30, 5.46]	0.942
FIB (median [IQR])		2.73 [2.46, 3.08]	2.77 [2.48, 3.04]	2.65 [2.28, 3.10]	0.289
W (median [IQR])		5.68 [5.02, 6.94]	5.64 [5.03, 6.50]	5.84 [5.02, 7.08]	0.406
R (median [IQR])		4.45 [4.21, 4.69]	4.54 [4.28, 4.70]	4.39 [4.08, 4.69]	0.194
Hb (median [IQR])		133.00 [128.00, 140.00]	135.00 [129.50, 140.00]	132.00 [122.25, 139.00]	0.057
N (median [IQR])		3.56 [2.87, 4.46]	3.56 [2.89, 4.37]	3.57 [2.84, 4.78]	0.778
L (median [IQR])		1.69 [1.37, 2.15]	1.69 [1.38, 2.15]	1.70 [1.36, 2.14]	0.956
M (median [IQR])		0.30 [0.24, 0.36]	0.29 [0.24, 0.36]	0.30 [0.25, 0.36]	0.444
P (median [IQR])		252.00 [208.00, 283.00]	245.00 [209.00, 284.00]	255.50 [208.00, 282.75]	0.802
NLR (%)	<1.86	44 (36.4)	22 (32.8)	22 (40.7)	0.479
	≥1.86	77 (63.6)	45 (67.2)	32 (59.3)	
dNLR (%)	<1.48	45 (37.2)	22 (32.8)	23 (42.6)	0.360
	≥1.48	76 (62.8)	45 (67.2)	31 (57.4)	
MLR (%)	<0.12	24 (19.8)	14 (20.9)	10 (18.5)	0.923
	≥0.12	97 (80.2)	53 (79.1)	44 (81.5)	
PLR (%)	<159.3	77 (63.6)	40 (59.7)	37 (68.5)	0.417
	≥159.3	44 (36.4)	27 (40.3)	17 (31.5)	
SIRI (%)	<0.63	62 (51.2)	38 (56.7)	24 (44.4)	0.246
	≥0.63	59 (48.8)	29 (43.3)	30 (55.6)	
SII (%)	<262.5	14 (11.6)	5 (7.5)	9 (16.7)	0.198
	≥262.5	107 (88.4)	62 (92.5)	45 (83.3)	
PNI (%)	<55.58	88 (72.7)	53 (79.1)	35 (64.8)	0.121
	≥55.58	33 (27.3)	14 (20.9)	19 (35.2)	
NRI (%)	<109.7	44 (36.4)	31 (46.3)	13 (24.1)	0.020
	≥109.7	77 (63.6)	36 (53.7)	41 (75.9)	
AFR (%)	<17.36	78 (64.5)	48 (71.6)	30 (55.6)	0.100
	≥17.36	43 (35.5)	19 (28.4)	24 (44.4)	
HALP (%)	<34.87	44 (36.4)	27 (40.3)	17 (31.5)	0.417
	≥34.87	77 (63.6)	40 (59.7)	37 (68.5)	
CONUT (%)	<1	69 (57.0)	40 (59.7)	29 (53.7)	0.633
	≥1	52 (43.0)	27 (40.3)	25 (46.3)	
BIPI (%)	Good	108 (89.3)	61 (91.0)	47 (87.0)	0.412
	Intermediate	12 (9.9)	5 (7.5)	7 (13.0)	
	Poor	1 (0.8)	1 (1.5)	0 (0.0)	

Abbreviations: LDH, lactic dehydrogenase; ALB, albumin; CRP, C-reactive protein; CHOL, cholesterol; FIB, fibrinogen; W, white blood cell; R, red blood cell; Hb, hemoglobin; N, neutrophils; L, lymphocyte; M, monocyte; P, platelet; NLR, neutrophil-to-lymphocyte ratio; dNLR, derived neutrophil-to-lymphocyte ratio; MLR, monocyte-to-lymphocyte ratio; PLR, platelet-to-lymphocyte ratio; SIRI, systemic inflammation response index; SII, systemic immune-inflammation index; PNI, prognostic nutritional index; NRI, nutritional risk index; AFR, albumin-to-fibrinogen ratio; HALP, hemoglobin × albumin × lymphocyte / platelet; CONUT, controlling nutritional status; BIPI, breast immune prognostic index.

Univariate and Multivariate Logistic Regression Analysis of Factors Affecting Drainage Tube Retention Time in Patients with Breast Cancer

According to the univariate and multivariate logistic regression analysis, the results showed that Hb, type of surgery, pathological T stage, chest wall drainage tube, NRI were independent predictors of affecting drainage tube retention time in patients with breast cancer (Table 3). A nomogram was constructed based on the results of the multivariate Logistic regression analysis to predict the risk for drainage tube retention time in patients with breast cancer (Figure 1).

Table 3 Univariate and Multivariate Logistic Regression Analysis of Factors Affecting Drainage Tube Retention Time in Patients with Breast Cancer

n	Level	Univariate				Multivariate			
		OR	95% CI low	95% CI high	p	OR	95% CI low	95% CI high	p
Age	<46	I (Reference)			0.935				
	≥46	0.971	0.473	1.991					
Marital status	Married	I (Reference)			0.668				
	Unmarried + Divorce + Widowed	1.277	0.410	3.974					
Occupation	Mental worker	I (Reference)			0.597				
	Manual worker + Others	1.295	0.491	3.425					
Weight	<60	I (Reference)			0.372				
	≥60	1.389	0.677	2.875					
Height	<1.61	I (Reference)			0.837				
	≥1.61	0.927	0.451	1.905					
BMI	<22.83	I (Reference)			0.082				
	≥22.83	1.906	0.926	3.980					
LDH	<158.10	I (Reference)			0.935				
	≥158.10	0.971	0.473	1.991					
ALB	<44.40	I (Reference)			0.168				
	≥44.40	1.662	0.810	3.453					
CRP	<0.02	I (Reference)			0.627				
	≥0.02	0.837	0.406	1.715					
CHOL	<4.83	I (Reference)			0.776				
	≥4.83	1.110	0.541	2.280					
FIB	<2.73	I (Reference)			0.329				
	≥2.73	0.699	0.338	1.433					
W	<5.68	I (Reference)			0.516				
	≥5.68	1.269	0.619	2.614					
R	<4.45	I (Reference)			0.181				
	≥4.45	0.611	0.294	1.253					
Hb	<133	I (Reference)			0.036	I (Reference)			0.035
	≥133	0.457	0.217	0.943		0.401	0.168	0.926	
N	<3.56	I (Reference)			0.966				
	≥3.56	0.985	0.480	2.022					

(Continued)

Table 3 (Continued).

n	Level	Univariate				Multivariate			
		OR	95% CI low	95% CI high	p	OR	95% CI low	95% CI high	p
L	<1.69	1 (Reference)			0.904				
	≥1.69	1.045	0.510	2.147					
M	<0.30	1 (Reference)			0.144				
	≥0.30	1.719	0.834	3.591					
P	<252	1 (Reference)			0.776				
	≥252	1.110	0.541	2.280					
Type of surgery	Mastectomy	1 (Reference)			0.019	1 (Reference)	0.071	0.883	0.040
	Breast-conserving surgery	0.278	0.086	0.759		0.273			
Tumor size	<2	1 (Reference)			0.133				
	≥2	1.752	0.848	3.676					
Histologic type	Ductal	1 (Reference)			0.556				
	Lobular + Others	1.321	0.529	3.441					
Histologic grade	0+I	1 (Reference)			0.376				
	II+III	1.414	0.661	3.079					
Pathological T stage	Tis/T0	1 (Reference)			0.011	1 (Reference)	1.155	44.961	0.039
	T1+T2+T3	2.809	1.278	6.363		6.684			
Pathological N stage	I	1 (Reference)			0.533				
	2	1.276	0.592	2.752					
Pathological TNM stage	I	1 (Reference)			0.287				
	2	1.481	0.720	3.070					
TLN	<7	1 (Reference)			0.394				
	≥7	1.367	0.667	2.824					
PLN	<1	1 (Reference)			0.816				
	≥1	1.095	0.506	2.358					
Molecular subtype	Luminal A + Luminal B (HER2+) + Luminal B (HER2-)	1 (Reference)			0.181				
	HER2 enriched + Triple negative	2.667	0.668	13.140					
ER status	Negative	1 (Reference)			0.830				
	Positive	1.098	0.470	2.618					
PR status	Negative	1 (Reference)			0.745				
	Positive	1.162	0.473	2.939					
HER2	Negative	1 (Reference)			0.304				
	Positive	1.629	0.643	4.203					
Ki-67	Negative	1 (Reference)			0.099				
	Positive	1.929	0.894	4.290					
AR	Negative	1 (Reference)			0.095				
	Positive	2.128	0.897	5.372					
CK5/6	Negative	1 (Reference)			0.388				
	Positive	0.477	0.066	2.313					

(Continued)

Table 3 (Continued).

n	Level	Univariate				Multivariate			
		OR	95% CI low	95% CI high	p	OR	95% CI low	95% CI high	p
E-cad	Negative	I (Reference)			0.292				
	Positive	1.542	0.696	3.518					
EGFR	Negative	I (Reference)			0.406				
	Positive	0.636	0.206	1.803					
P53	Negative	I (Reference)			0.391				
	Positive	1.378	0.665	2.892					
TOP2A	Negative	I (Reference)			0.039	I (Reference)			0.226
	Positive	2.188	1.051	4.656		1.945	0.672	5.920	
Lymph vessel invasion	Negative	I (Reference)			0.931				
	Positive	1.042	0.404	2.638					
Neural invasion	Negative	I (Reference)			0.758				
	Positive	0.849	0.288	2.382					
Healthy lower arm circumference	<19	I (Reference)			0.655				
	≥19	1.178	0.575	2.423					
Healthy upper arm circumference	<26	I (Reference)			0.329				
	≥26	1.431	0.698	2.955					
Affected side lower arm circumference	<19	I (Reference)			0.966				
	≥19	1.016	0.495	2.085					
Affected side upper arm circumference	<26	I (Reference)			0.368				
	≥26	1.393	0.678	2.879					
Chest wall drainage tube	<13	I (Reference)			0.000	I (Reference)			0.000
	≥13	92.188	29.083	374.069		127.427	25.938	1139.723	
Axillary drainage tube	<13	I (Reference)			0.000	I (Reference)			0.480
	≥13	5.298	2.441	12.090		1.872	0.311	11.822	
NLR	<1.86	I (Reference)			0.370				
	≥1.86	0.711	0.336	1.498					
dNLR	<1.48	I (Reference)			0.271				
	≥1.48	0.659	0.312	1.383					
MLR	<0.12	I (Reference)			0.745				
	≥0.12	1.162	0.473	2.939					
PLR	<159.3	I (Reference)			0.317				
	≥159.3	0.681	0.317	1.438					
SIRI	<0.63	I (Reference)			0.181				
	≥0.63	1.638	0.798	3.398					
SII	<262.5	I (Reference)			0.124				
	≥262.5	0.403	0.117	1.248					
PNI	<55.58	I (Reference)			0.082				
	≥55.58	2.055	0.918	4.698					

(Continued)

Table 3 (Continued).

n	Level	Univariate				Multivariate			
		OR	95% CI low	95% CI high	p	OR	95% CI low	95% CI high	p
NRI	<109.7	1 (Reference)			0.013	1 (Reference)			0.025
	≥109.7	2.716	1.256	6.116		2.837	1.164	7.292	
AFR	<17.36	1 (Reference)			0.068				
	≥17.36	2.021	0.954	4.345					
HALP	<34.87	1 (Reference)			0.317				
	≥34.87	1.469	0.695	3.159					
CONUT	<1	1 (Reference)			0.508				
	≥1	1.277	0.619	2.645					
BIPI	Good	1 (Reference)			0.481				
	Intermediate + Poor	1.514	0.473	4.990					

Abbreviations: BMI, body mass index; LDH, lactic dehydrogenase; ALB, albumin; CRP, C-reactive protein; CHOL, cholesterol; FIB, fibrinogen; WBC, white blood cell; R, red blood cell; Hb, hemoglobin; N, neutrophils; L, lymphocyte; M, monocyte; P, platelet; TLN, Total lymph nodes; PLN, Positive lymph nodes; ER, estrogen receptor; PR, progesterone receptor; HER2, Human Epidermal Growth Factor Receptor 2; AR, androgen receptor; CK5/6, Cytokeratin 5/6; E-cad, E-cadherin; EGFR, Epidermal Growth Factor Receptor; TOP2A, topoisomerase 2A; NLR, neutrophil-to-lymphocyte ratio; dNLR, derived neutrophil-to-lymphocyte ratio; MLR, monocyte-to-lymphocyte ratio; PLR, platelet-to-lymphocyte ratio; SIRI, systemic inflammation response index; SII, systemic immune-inflammation index; PNI, prognostic nutritional index; NRI, nutritional risk index; AFR, albumin-to-fibrinogen ratio; HALP, hemoglobin × albumin × lymphocyte / platelet; CONUT, controlling nutritional status; BIPI, breast immune prognostic index.

The Useful Factors by Univariate Logistic Regression Analysis for Drainage Tube at Different Times

Through the univariate Logistic regression analysis, the results indicated that Hb, type of surgery, pathological T stage, TOP2A, chest wall drainage tube, axillary drainage tube, NRI were independent predictors of affecting drainage tube retention time in patients with breast cancer. There were significant differences in the type of surgery, pathological T stage, chest wall drainage tube, NRI by removing the drainage tube at different times (Table 4).




Figure 1 Nomogram constructed by the multivariate Logistic regression analysis. ***Statistical significance of P value.

Table 4 The Useful Factors by Univariate Logistic Regression Analysis for Drainage Tube at Different Times

n	Level	Overall	DTRT≤7 (d)	DTRT≤14 (d)	DTRT≤21 (d)	DTRT>21 (d)	p
		121	24	48	38	11	
Hb (%)	<133	61 (50.4)	10 (41.7)	21 (43.8)	23 (60.5)	7 (63.6)	0.271
	≥133	60 (49.6)	14 (58.3)	27 (56.2)	15 (39.5)	4 (36.4)	
Type of surgery (%)	Mastectomy	98 (81.0)	10 (41.7)	44 (91.7)	34 (89.5)	10 (90.9)	<0.001
	Breast-conserving surgery	23 (19.0)	14 (58.3)	4 (8.3)	4 (10.5)	1 (9.1)	
Pathological T stage (%)	Tis/T0	84 (69.4)	20 (83.3)	37 (77.1)	21 (55.3)	6 (54.5)	0.041
	T1+T2+T3	37 (30.6)	4 (16.7)	11 (22.9)	17 (44.7)	5 (45.5)	
TOP2A (%)	Negative	53 (43.8)	13 (54.2)	23 (47.9)	14 (36.8)	3 (27.3)	0.341
	Positive	68 (56.2)	11 (45.8)	25 (52.1)	24 (63.2)	8 (72.7)	
Chest wall drainage tube (%)	<13	63 (52.1)	18 (75.0)	41 (85.4)	4 (10.5)	0 (0.0)	<0.001
	≥13	58 (47.9)	6 (25.0)	7 (14.6)	34 (89.5)	11 (100.0)	
Axillary drainage tube (%)	<13	70 (57.9)	12 (50.0)	31 (64.6)	20 (52.6)	7 (63.6)	0.557
	≥13	51 (42.1)	12 (50.0)	17 (35.4)	18 (47.4)	4 (36.4)	
NRI (%)	<109.7	44 (36.4)	13 (54.2)	19 (39.6)	11 (28.9)	1 (9.1)	0.047
	≥109.7	77 (63.6)	11 (45.8)	29 (60.4)	27 (71.1)	10 (90.9)	

Abbreviations: Hb, hemoglobin; TOP2A, topoisomerase 2A; NRI, nutritional risk index.

Discussion

After the operation, the complications of breast cancer mainly include subcutaneous effusion, wound infection, delayed healing, bleeding, and skin flap necrosis.^{13,14} The subcutaneous effusion and wound infection cause a huge gap among the wounds, the flap and the surface of the wound cannot be appressed effectively. This will affect the quality of life and survival time of breast cancer patients. The drainage tubes are usually to prevent the complications of subcutaneous effusion and infection. The drainage tube removal time is determined on the basis of drainage volume and color, and the postoperative recovery condition. The preoperative and postoperative nursing care of breast cancer patients is a comprehensive procedure. The drainage tube retention time and nutritional status may influence the wound healing time, and this may be related to the poor nutritional status and immune function after the operation. It is very critical to control basic diseases (diabetes mellitus, hypertension), strengthen nutritional support, and enhance immunity.

Inflammation is a critical component of the tumor microenvironment (TME) and is an indispensable participant in the development, progression, and metastasis of cancer.¹⁵ The TME is largely orchestrated by inflammatory cells and also selected some signal molecules of the innate immune system.¹⁶ It is an attractive strategy to make a profound study of inflammation for cancer prevention and treatment. A number of reports have confirmed that in many types of cancers, for instance, digestive tract cancers, lung cancer, liver cancer, and breast cancer, the inflammatory reactions are abnormal, and related to the prognosis of tumors.^{17–24} Yet, the relationship between the duration of suction drainage and the inflammatory reactions has been rarely studied.

This study systematically analyzed the effects of common inflammatory cells and the composite inflammatory indexes on the postoperative drainage tube extraction time. Our drainage tube retention time is determined by the median DDCSD time. The results showed that the type of surgery, total lymph nodes, pathological T stage, NRI were related to the drainage tube retention time. Moreover, the univariate and multivariate Logistic regression analysis performed that Hb, type of surgery, pathological T stage, chest wall drainage tube, NRI were the independent risk predictors of affecting drainage tube retention time. Furthermore, we also found that a significant correlation existed between NRI and drainage tube retention at different times. These results go a step further to suggest that NRI is an important risk factor affecting drainage tube retention time and emphasizes the major impact of nutrition on breast cancer patients.

The NRI is a clinical biological index that combines the strength of two nutritional indicators-albumin and weight loss.²⁵ Serum albumin is an important indicator of nutritional status and immunological functioning. The albumin level decreases upon the occurrence of inflammatory reactions. Albumin can be used as a non-specific host defense substance and used to fight against various toxic metabolites during infection, so as to reduce the harmful substances in the

body.^{26,27} The doctors can identify patients who are more likely to retain suction drainage by the serum albumin levels and take early interventions. Obesity has been proved to be a risk factor for postoperative recovery for cancer patients.²⁸ Owing to the special anatomical position of the breast, the use of high-frequency electric knives in operation leads to the liquefaction of adipose tissue for obese women with rich subcutaneous adipose tissue. After the operation, it is easy to have an insufficient level of blood supply, which could result in subcutaneous effusion, and subsequent delays in the extraction time of the drainage tube. The blockage of the drainage tube by adipose tissue leads to the flap stays in a free state. The flap cannot establish a normal blood supply with the thorax, which further leads to flap necrosis.

Observing the nature of drainage fluid is an important part of nursing after breast surgery. The drainage volume and color are ideal clinical observation indicators that can reflect the incision exudation. The drainage tube retention time can be determined by the flow and the color of the drainage tube decreasing. At the same time, it can also assist clinicians to understand the condition of the healing of the incision, so as to formulate the next diagnosis and treatment plan. Negative pressure drainage can form continuous negative pressure, make the wound cavity narrow, and then make the flap close to the chest wall and armpit. On the one hand, continuous negative pressure suction can promote the formation of the capillary, provide sufficient blood supply and establish blood circulation.²⁹ On the other hand, continuous negative pressure suction can reduce skin tension, promote wound healing, and prevent flap necrosis.²⁹ At the same time, continuous negative pressure suction can also effectively prevent the spread of bacteria into the incision, and prevent the spread of bacteria into the incision.³⁰

The difficulty in predicting the drainage tube retention time in these patients poses an uncertainty that may complicate the development of a suitable clinical pathway. This study had some limitations. Firstly, this study had insufficient data, and bring about a bias. Secondly, the patients are from a single research site, which means that the findings of our study may not be applicable to other research contexts. Furthermore, follow-up data are not available.

Conclusions

The preoperative inflammatory indicators are related to the drainage tube retention time in patients with breast cancer. NRI is an independent risk factor for postoperative drainage tube extraction time and can provide personalized nursing intervention of patients with breast cancer after drainage tube retention time and rehabilitation process.

Data Sharing Statement

The data that support the findings of this study are available from the corresponding author (Li Chen) upon reasonable request.

Ethics Statement

This study was approved by the ethics committee of Cancer Hospital Chinese Academy of Medical Sciences and was conducted in accordance with the amended Declaration of Helsinki.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

This research was supported by grants from the Hubei Province Postdoctoral Innovation Research Post Fund Project (No.0106540096), Open Fund for the Key Laboratory of Organ Transplantation of Ministry of Education and National Health Commission (No.2021QYKF03), Tongji Hospital Cultivation Project (No.2022B03), Chen Xiao-ping Foundation for the Development of Science and Technology of Hubei province, Youth Science Special Fund (No.CXPJJH123001-2308).

Disclosure

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

- Li N, Deng Y, Zhou L, et al. Global burden of breast cancer and attributable risk factors in 195 countries and territories, from 1990 to 2017: results from the Global Burden of Disease Study 2017. *J Hematol Oncol*. 2019;12(1):140. PMID: 31864424; PMCID: PMC6925497. doi:10.1186/s13045-019-0828-0
- Radecka B, Litwiniuk M. Breast cancer in young women. *Ginekol Pol*. 2016;87(9):659–663. PMID: 27723074. doi:10.5603/GP.2016.0062
- Shien T, Iwata H. Adjuvant and neoadjuvant therapy for breast cancer. *Jpn J Clin Oncol*. 2020;50(3):225–229. PMID: 32147701. doi:10.1093/jjco/hyz213
- Chen CF, Lin SF, Hung CF, Chou P. Risk of infection is associated more with drain duration than daily drainage volume in prosthesis-based breast reconstruction: a cohort study. *Medicine*. 2016;95(49):e5605. PMID: 27930584; PMCID: PMC5266056. doi:10.1097/MD.0000000000005605
- Héquet D, Huchon C, Baffert S, et al. Preoperative clinical pathway of breast cancer patients: determinants of compliance with EUSOMA quality indicators. *Br J Cancer*. 2017;116(11):1394–1401. PMID: 28441385; PMCID: PMC5520093. doi:10.1038/bjc.2017.114
- Mantovani A, Allavena P, Sica A, Balkwill F. Cancer-related inflammation. *Nature*. 2008;454(7203):436–444. PMID: 18650914. doi:10.1038/nature07205
- Borsig L, Wolf MJ, Roblek M, Lorentzen A, Heikenwalder M. Inflammatory chemokines and metastasis--tracing the accessory. *Oncogene*. 2014;33(25):3217–3224. Epub 2013 Jul 15. PMID: 23851506. doi:10.1038/ncr.2013.272
- Dong X, Liu C, Yuan J, et al. Prognostic roles of neutrophil-to-lymphocyte ratio and stromal tumor-infiltrating lymphocytes and their relationship in locally advanced triple-negative breast cancer treated with neoadjuvant chemotherapy. *Breast Care*. 2021;16(4):328–334. PMID: 34602938; PMCID: PMC8436630. doi:10.1159/000509498
- Morrison L, Laukkanen JA, Ronkainen K, Kurl S, Kauhanen J, Toriola AT. Inflammatory biomarker score and cancer: a population-based prospective cohort study. *BMC Cancer*. 2016;16:80. PMID: 26860264; PMCID: PMC4748611. doi:10.1186/s12885-016-2115-6
- Xu M, Chen R, Liu L, et al. Systemic immune-inflammation index and incident cardiovascular diseases among middle-aged and elderly Chinese adults: the Dongfeng-Tongji cohort study. *Atherosclerosis*. 2021;323:20–29. PMID: 33773161. doi:10.1016/j.atherosclerosis.2021.02.012
- Browne JP, Jeevan R, Gulliver-Clarke C, Pereira J, Caddy CM, van der Meulen JHP. The association between complications and quality of life after mastectomy and breast reconstruction for breast cancer. *Cancer*. 2017;123(18):3460–3467. Epub 2017 May 17. PMID: 28513834. doi:10.1002/cncr.30788
- Yfantis A, Sarafis P, Moisoglou I, et al. How breast cancer treatments affect the quality of life of women with non-metastatic breast cancer one year after surgical treatment: a cross-sectional study in Greece. *BMC Surg*. 2020;20(1):210. PMID: 32957940; PMCID: PMC7507267. doi:10.1186/s12893-020-00871-z
- Wang Z, Han X. Clinical significance of breast-conserving surgery for early breast cancer and its impact on patient life quality of life. *J BUON*. 2019;24(5):1898–1904. PMID: 31786853.
- Palve JS, Luukkaala TH, Kääriäinen MT. Predictive risk factors of complications in different breast reconstruction methods. *Breast Cancer Res Treat*. 2020;182(2):345–354. Epub 2020 May 28. PMID: 32468337; PMCID: PMC7297836. doi:10.1007/s10549-020-05705-3
- Coussens LM, Werb Z. Inflammation and cancer. *Nature*. 2002;420(6917):860–867. PMID: 12490959; PMCID: PMC2803035. doi:10.1038/nature01322
- Chen D, Zhang X, Li Z, Zhu B. Metabolic regulatory crosstalk between tumor microenvironment and tumor-associated macrophages. *Theranostics*. 2021;11(3):1016–1030. PMID: 33391518; PMCID: PMC7738889. doi:10.7150/thno.51777
- Li Y, Shao Y, Bai L, Zhou X. Increased derived neutrophil-to-lymphocyte ratio and breast imaging-reporting and data system classification predict poor survival in patients with non-distant metastatic HER2+ breast cancer treated with neoadjuvant chemotherapy. *Cancer Manag Res*. 2018;10:3841–3847. PMID: 30288115; PMCID: PMC6161733. doi:10.2147/CMAR.S174537
- Chen L, Kong X, Wang Z, Wang X, Fang Y, Wang J. Pretreatment systemic inflammation response index in patients with breast cancer treated with neoadjuvant chemotherapy as a useful prognostic indicator. *Cancer Manag Res*. 2020;12:1543–1567. PMID: 32184659; PMCID: PMC7060771. doi:10.2147/CMAR.S235519
- Zhang Y, Sun Y, Zhang Q. Prognostic value of the systemic immune-inflammation index in patients with breast cancer: a meta-analysis. *Cancer Cell Int*. 2020;20:224. PMID: 32528232; PMCID: PMC7282128. doi:10.1186/s12935-020-01308-6
- Hua X, Long ZQ, Huang X, et al. The Value of Prognostic Nutritional Index (PNI) in predicting survival and guiding radiotherapy of patients with T1-2N1 breast cancer. *Front Oncol*. 2020;9:1562. PMID: 32083015; PMCID: PMC7002465. doi:10.3389/fonc.2019.01562
- Sasaki M, Miyoshi N, Fujino S, et al. The geriatric nutritional risk index predicts postoperative complications and prognosis in elderly patients with colorectal cancer after curative surgery. *Sci Rep*. 2020;10(1):10744. PMID: 32612136; PMCID: PMC7329855. doi:10.1038/s41598-020-67285-y
- Huang L, Mo Z, Zhang L, Qin S, Qin S, Li S. Diagnostic value of albumin to fibrinogen ratio in cervical cancer. *Int J Biol Markers*. 2020;35(2):66–73. Epub 2020 May 11. PMID: 32389031. doi:10.1177/1724600820915916
- Guo Y, Shi D, Zhang J, et al. The Hemoglobin, Albumin, Lymphocyte, and Platelet (HALP) score is a novel significant prognostic factor for patients with metastatic prostate cancer undergoing cytoreductive radical prostatectomy. *J Cancer*. 2019;10(1):81–91. PMID: 30662528; PMCID: PMC6329846. doi:10.7150/jca.27210
- Ahiko Y, Shida D, Horie T, et al. Controlling nutritional status (CONUT) score as a preoperative risk assessment index for older patients with colorectal cancer. *BMC Cancer*. 2019;19(1):946. PMID: 31690275; PMCID: PMC6833132. doi:10.1186/s12885-019-6218-8
- Sagou K, Ozeki K, Ukai S, Adachi Y, Fukushima N, Kohno A. Impact of a nutritional risk index on clinical outcomes after allogeneic hematopoietic cell transplantation. *Biol Blood Marrow Transplant*. 2019;25(11):2287–2296. Epub 2019 Jul 5. PMID: 31284071. doi:10.1016/j.bbmt.2019.07.002
- Van de Sande L, Cosyns S, Willaert W, Ceelen W. Albumin-based cancer therapeutics for intraperitoneal drug delivery: a review. *Drug Deliv*. 2020;27(1):40–53. PMID: 31858848; PMCID: PMC6968566. doi:10.1080/10717544.2019.1704945

27. Hoogenboezem EN, Duvall CL. Harnessing albumin as a carrier for cancer therapies. *Adv Drug Deliv Rev.* 2018;130:73–89. Epub 2018 Jul 27. PMID: 30012492; PMCID: PMC6200408. doi:10.1016/j.addr.2018.07.011
28. Picon-Ruiz M, Morata-Tarifa C, Valle-Goffin JJ, Friedman ER, Slingerland JM. Obesity and adverse breast cancer risk and outcome: mechanistic insights and strategies for intervention. *CA Cancer J Clin.* 2017;67(5):378–397. Epub 2017 Aug 1. PMID: 28763097; PMCID: PMC5591063. doi:10.3322/caac.21405
29. Lin W, Yang Y, Zhong W, et al. The effect of low and high vacuum drainage on the postoperative drainage of breast cancer: insights from a prospective, non-inferiority, randomized clinical trial. *Cancer Manag Res.* 2020;12:12487–12496. PMID: 33299355; PMCID: PMC7721110. doi:10.2147/CMAR.S283031
30. Matusiak D, Wichtowski M, Pieszko K, Kobylarek D, Murawa D. Is negative-pressure wound therapy beneficial in modern-day breast surgery? *Contemp Oncol.* 2019;23(2):69–73. Epub 2019 May 27. PMID: 31316287; PMCID: PMC6630394. doi:10.5114/wo.2019.85199

Breast Cancer: Targets and Therapy

Dovepress

Publish your work in this journal

Breast Cancer - Targets and Therapy is an international, peer-reviewed open access journal focusing on breast cancer research, identification of therapeutic targets and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/breast-cancer—targets-and-therapy-journal>