

IAP proteins as targets for drug development in oncology

Laurence Dubrez^{1,2}

Jean Berthelet^{1,2}

Valérie Glorian^{1,2}

¹Institut National de la Santé et de la Recherche Médicale (Inserm), Dijon, France; ²Université de Bourgogne, Dijon, France

Abstract: The inhibitors of apoptosis (IAPs) constitute a family of proteins involved in the regulation of various cellular processes, including cell death, immune and inflammatory responses, cell proliferation, cell differentiation, and cell motility. There is accumulating evidence supporting IAP-targeting in tumors: IAPs regulate various cellular processes that contribute to tumor development, such as cell death, cell proliferation, and cell migration; their expression is increased in a number of human tumor samples, and IAP overexpression has been correlated with tumor growth, and poor prognosis or low response to treatment; and IAP expression can be rapidly induced in response to chemotherapy or radiotherapy because of the presence of an internal ribosome entry site (IRES)-dependent mechanism of translation initiation, which could contribute to resistance to antitumor therapy. The development of IAP antagonists is an important challenge and was subject to intense research over the past decade. Six molecules are currently in clinical trials. This review focuses on the role of IAPs in tumors and the development of IAP-targeting molecules for anticancer therapy.

Keywords: Smac mimetics, apoptosis, antitumor therapy

Introduction: IAP family of proteins

The inhibitors of apoptosis (IAPs) constitute a family of proteins highly conserved throughout evolution. IAPs were initially discovered in baculoviruses two decades ago,¹ as potent inhibitors of apoptosis in infected insect cells. The first human homologous neuronal apoptosis inhibitory protein (NAIP) and cellular IAP 1 and 2 (cIAP1 and cIAP2) were characterized 2 years later,^{2,3} followed by X-chromosome linked IAP (XIAP),^{4,5} survivin,⁶ Apollon (also called BRUCE),⁷ melanoma IAP (ML-IAP) (also called Livin),⁸ and IAP-like protein 2 (ILP2).⁹ The IAP family is defined by the presence of one to three conserved protein motifs named a baculoviral IAP repeat (BIR). Most of them form a surface hydrophobic groove that specifically binds a conserved tetrapeptide motif, called IAP binding motif (IBM), found in the active subunits of apoptotic protease caspase-3, -7, and -9 and in cellular IAP antagonists, such as the second mitochondria-derived activator of caspases (Smac) (also named direct IAP-binding protein with low isoelectric point (pI) [DIABLO])¹⁰⁻¹³ and the high temperature requirement protein A2 (HtrA2)^{12,14} (Figure 1). The first BIR of XIAP and cIAPs does not bind IBM but rather, the signaling molecule transforming growth factor beta (TGF β)-activated kinase 1-binding protein 1 (TAB1)¹⁵ or the tumor necrosis factor (TNF) receptor (TNFR) associated factors (TRAFs),¹⁶⁻¹⁸ connecting XIAP and cIAPs with the TGF and TNF signaling pathways, respectively. In addition to the BIRs, cIAPs, XIAP, ML-IAP and ILP2 also possess a C-terminal RING (really interesting new gene)

Correspondence: Laurence Dubrez
Inserm UMR866, Faculty of Medicine,
7 Boulevard Jeanne d'Arc, 21079 Dijon
cedex, France
Tel +33 380 393 356
Fax +33 380 393 434
Email ldubrez@u-bourgogne.fr




Figure 1 The inhibition of caspases by XIAP and the regulation by Smac and Smac mimetics.

Notes: (A) Among IAPs, XIAP is a potent caspase inhibitor. XIAP is composed of three BIR domains, one UBA domain (which binds ubiquitin chains), and one C-terminal RING domain, which confers to XIAP an E3-ubiquitin ligase activity. The first BIR (BIR1) can bind to TAB1, connecting XIAP to the TGF β signaling pathway. The BIR2 and BIR3 contain a surface hydrophobic groove allowing the interaction with IBM found in caspase-3, -7, and -9 active subunits and in IAP antagonists, such as Smac or HtrA2. Moreover, the linker region upstream of BIR2 binds across the substrate binding pocket of caspase-3 and -7, and BIR3 binds the dimer interface of caspase-9, which hinder substrate accessibility and hide the catalytic residue of caspase. Smac is released from the mitochondria into the cytosol during apoptosis, after a maturation process that removes the N-terminal mitochondrial import signal and exposes the IBM to the N-extremity of the protein. Once cytosolic, Smac forms a symmetric dimer and binds the BIR2 and BIR3 IBM grooves of XIAP, preventing them from binding caspases. In a similar manner, monovalent and bivalent Smac mimetics efficiently bind the BIR2 and BIR3 surface hydrophobic grooves and abrogate XIAP-mediated caspase inhibition. (B–D) Comparison of the XIAP-BIR3 (blue) bound to the IBM of caspase-9 (ATPFQ) (orange) (pdb 1nw9); (B) The IBM (AVPI tetrapeptide) of Smac (red) (pdb 2opz); (C) The monovalent Smac mimetic SM-130 (green) (pdb 2jk7); and (D) The BIR domains of IAPs are organized in four α -helices and three β -strand sheets maintained by a zinc ion (yellow). IBMs interact with the surface hydrophobic groove of BIRs (constructed using The PyMOL Molecular Graphics System, Version 1.5.0.4 Schrödinger, LLC, New York, NY, USA).

Abbreviations: AVPI, Smac N-terminal tetrapeptide; BIR, baculoviral IAP repeat; HtrA2, high temperature requirement protein A2; IAPs, inhibitors of apoptosis; IBM, IAP binding motif; RING, really interesting new gene; SM, Smac mimetic; Smac, second mitochondria-derived activator of caspases; TAB1, TGF β -activated kinase 1-binding protein 1; TGF β , transforming growth factor beta; UBA, ubiquitin associated; XIAP, X-chromosome linked IAP; APAF-1, apoptotic peptidase activating factor.

domain conferring an E3 ligase activity in the ubiquitination or neddylation¹⁹ reactions (for review,^{20,21}).

Numerous partners of IAPs have been identified, including some caspases,^{22–24} some signaling molecules,^{25,26} some regulators of the NF- κ B: nuclear factor of kappa-light polypeptide gene enhancer in B-cell activating pathways,²⁵ some regulators of the actin cytoskeleton,²⁷ and some transcriptional regulators.^{28,29} Thus, although they were initially characterized as inhibitors of apoptosis, IAPs display additional nonapoptotic functions in the regulation of cell proliferation, cell division, cell differentiation, cell motility, and in proinflammatory and immune response (for review,^{25,26}), which could contribute to oncogenesis.

Expression of IAPs in tumors

The expression of IAPs or cellular IAP antagonists such as Smac,¹¹ HtrA2, or the septin-like mitochondrial protein, ARTS,^{30,31} were shown to be altered in a number

of human tumor samples (Table 1). Overexpression of IAPs or downregulation of the cellular IAP antagonists have been correlated with advanced progressive disease, aggressiveness, and poor prognosis or low response to treatment (Supplementary Table S1). The alterations of IAP expression can be associated or not, with gene mutations. The *baculoviral IAP repeat containing protein (BIRC)2* and *BIRC3* genes, encoding cIAP1 and cIAP2, respectively, are located on chromosome 11q21–22, a region found amplified in human hepatocarcinoma,³² mammary carcinoma,³³ medulloblastoma,³⁴ and in pancreatic,³⁵ cervical,³⁶ lung,³⁷ oral squamous cell,³⁸ and esophageal³⁹ carcinomas. Some (30%) mucosa-associated lymphoid tissue (MALT) lymphoma are associated with the chromosomal translocation t(11;18) (q21;q21) generating a chimeric protein composed of the N-terminal sequences of cIAP2 fused to the C-terminal sequence of MALT1.^{17,40} Conversely, IAPs can also display antitumoral properties in lymphocytes. The *BIRC2* and/or

Table 1 Expression of IAPs and IAP antagonists in human tumors

IAPs and cellular IAP antagonists	Tumors
IAPs:	
XIAP overexpression	Acute myeloid leukemia, ¹³⁸ B-cell chronic lymphocytic leukemia, ^{139,140} bladder carcinoma, ¹⁴¹ breast carcinoma, ¹⁴² cervical carcinoma, ¹⁴³ colorectal cancer, ^{144,145} hepatocarcinoma, ^{146,147} melanoma, ¹⁴⁸ non-small cell lung cancer, ^{149,150} ovarian cancer, ¹¹⁹ prostate carcinoma, ^{151,152} renal carcinoma, ^{153–155} thyroid carcinoma ¹⁵⁶
cIAP1 and cIAP2 overexpression (amplicon 11q21-22)	Cervical cancer, ³⁶ esophageal squamous cell carcinoma, ³⁹ hepatocarcinoma, ³² medulloblastoma, ³⁴ non-small and small cell lung cancer, ^{37,150} oral squamous cell carcinoma, ³⁸ pancreatic cancer ¹⁵⁷
cIAP1 overexpression independently from amplicon 11q21-22	B-cell chronic lymphocytic leukemia, ^{139,158,159} bladder carcinoma, ⁸³ cervical carcinoma, ^{143,36} chronic lymphocytic leukemia, ¹³⁹ colorectal cancer, ¹⁶⁰ head and neck squamous cell carcinoma, ⁸² non-small and small cell lung cancer, ¹⁵⁰ prostate carcinoma, ¹⁵² squamous carcinoma of tongue ¹⁶¹
cIAP2 overexpression independently from amplicon 11q21-22	Breast cancer, ¹⁶² cervical carcinoma, ¹⁴³ chronic lymphocytic leukemia, ^{139,159} colorectal carcinoma, ¹⁶⁰ prostate carcinoma ¹⁵²
cIAP2/MALT chimeric protein t(11,18)(q21, q21)	MALT myeloma ^{163,164}
cIAP1/cIAP2 inactivation	Multiple myeloma ^{41,165}
ML-IAP overexpression	Acute myeloid leukemia, ¹⁶⁶ childhood acute lymphoblastic leukemia, ¹⁶⁷ bladder carcinoma, ¹⁶⁸ colorectal carcinoma, ¹⁶⁹ gastric cancer, ¹⁷⁰ melanoma, ¹⁷¹ neuroblastoma, ¹⁷² osteosarcoma, ¹⁷³ renal cell carcinoma, ^{174,175} testicular cancer ¹⁷⁶
cIAP antagonists:	
Smac downregulation	Acute myeloid leukemia, ¹⁷⁷ bladder carcinoma, ¹⁷⁸ breast carcinoma, ¹⁷⁹ cervical carcinoma ¹⁸⁰ chronic lymphocytic leukemia, ¹³⁹ colorectal carcinoma, ¹⁸¹ endometrioid endometrial cancer, ¹⁸² esophageal carcinoma, ¹¹² lung cancer, ¹⁸³ rectal adenocarcinoma, ¹⁴⁵
Smac overexpression	Bladder cancer, ¹⁸⁴ gastric adenocarcinoma, ¹⁸⁵ renal adenocarcinoma, ^{154,176,186}
HtrA2 overexpression	Endometrial cancer, ¹⁸⁷ ovarian cancer, ¹⁸⁸ prostate carcinoma, ^{189,190} renal carcinoma, ¹⁸⁶ stomach cancer, ¹⁹¹ thyroid cancer ¹⁹²
HtrA2 downregulation	Endometrial cancer, ^{187,193} ovarian cancer, ¹⁹⁴ testicular cancer
ARTS overexpression	Astrocytoma ¹⁹⁵
ARTS downregulation	Acute myeloid leukemia ¹⁹⁶
Ratio IAP/IAP antagonists:	
Increased XIAP/Smac	Renal adenocarcinoma ¹⁵⁴
Reduced XIAP/Smac	Gastric carcinoma ¹⁸⁵
Increased cIAP1/HtrA2 and cIAP1/Smac	Chronic lymphocytic leukemia ¹³⁹

Abbreviations: ARTS, septin-like mitochondrial protein; cIAP, cellular IAP; HtrA2, high temperature requirement protein A2; IAPs, inhibitors of apoptosis; MALT, mucosa-associated lymphoid tissue; ML-IAP, melanoma IAP; Smac, second mitochondria-derived activator of caspases; XIAP, X-chromosome linked IAP.

BIRC3 genes were found to be mutated in some multiple myeloma samples,^{41,42} and the *BIRC4* encoding XIAP in X-linked lymphoproliferative disease.⁴³ The expression and functions of the atypical IAP survivin in tumors, and the development of specific survivin-targeted therapy were recently reviewed by Coumar et al⁴⁴ and won't be discussed here.

Role of IAPs in cancer

IAPs as apoptotic regulators

IAPs were first characterized as inhibitors of apoptosis because of their ability to bind caspases. Indeed, cIAPs, XIAP and ML-IAP can bind caspase-3, -7, and -9 via the BIRs^{10,11,45,46} and can induce their ubiquitination or neddylation via the RING domain.^{19,22–24} The influence of the ubiquitination is still not very well established, triggering degradative or nondegradative consequences,^{22–24} while the neddylation of

caspase-7, by XIAP, inhibits its activity.¹⁹ In addition, XIAP is able to directly inhibit the enzymatic activity of caspases (Figure 1). The XIAP BIR3 binds the dimer interface of caspase-9, and the linker region upstream of BIR2 binds across the substrate binding pocket of caspase-3 and -7, which hinder substrate accessibility and hide the catalytic residue of caspases.^{47–49} The capacity of XIAP to inhibit caspase activity could account for the resistance of cancer cells to antitumor therapy. Indeed, DNA-damaging treatments, such as ionizing irradiations, induce a translational upregulation of XIAP as a consequence of the presence of an internal ribosome entry site (IRES)-dependent translation mechanism, which results in the resistance of carcinoma cells to radiation-induced apoptosis.^{50,51}

IAPs can also inhibit cell death at an earlier step, preventing the assembly of caspase-8- or -10-activating platforms. Caspase-8 and -10 are initiator caspases recruited

by the adaptor Fas-associated death domain protein (FADD) in multiprotein complexes, which provide the proximity required for caspase homodimerization and self-activation (for review,⁵²). These molecular platforms are assembled either in response to the engagement of death receptor from the TNFR superfamily (in which case, these are referred to as death-inducing signaling complex [DISC] and complex II)^{52,53} or in response to genotoxic stress, tumor necrosis factor-like weak inducer of apoptosis (TWEAK) engagement, or toll-like

receptor (TLR) 3 stimulation (in which case, they are referred to as Ripoptosome).^{54,55} Complex II^{50,51} and Ripoptosome^{52,53} share, in addition to the caspase and the adaptor FADD, the serine/threonine kinase receptor interacting protein (RIP) (Figure 2). cIAPs and XIAP are potent regulators of proteins from the RIP family, catalyzing the conjugation of ubiquitin chains that control either protein degradation or signal transduction pathways⁵⁶⁻⁶² (Figure 2). In the absence of cIAPs, non-ubiquitinated RIP1 promotes (through its kinase




Figure 2 Mechanisms of action of Smac mimetics.

Notes: IAPs are important regulators of NF-κB-activating signaling pathways. Upon TNFα stimulation, cIAPs and RIP1 are recruited to TNFR1 via the adaptors TRADD and TRAF2. cIAPs trigger self-ubiquitination and ubiquitination of RIP1. These ubiquitin chains serve as a scaffold for the recruitment of IKK, TAK, and LUBAC. Once activated, IKK complex triggers phosphorylation of IκBα, which is then degraded by the UPS and releases NF-κB dimer, which promotes the transcription of target genes. In the noncanonical pathway of NF-κB activation, cIAPs promote the degradative ubiquitination of NIK and associated TRAF proteins and prevent the activation of the IKK complex required for transcription factor activation. On the other hand, cIAPs and XIAP prevent the assembly of RIP1-containing initiator caspase-activating complexes, named complex-II and Ripoptosome; and XIAP inhibits, through a direct interaction via BIR2 and BIR3, the activity of caspase-3, -7, and -9. SMs bind to the BIR domains of cIAPs and stimulate their E3-ubiquitin ligase activity. This results first, in the ubiquitination of RIP1, leading to the canonical NF-κB activation, and second, in the rapid autoubiquitination and subsequent proteasome-mediated degradation of cIAPs. Depletion of cIAPs releases NIK, resulting in the noncanonical activation of NF-κB, and NF-κB target gene expression, including TNFα, MCP-1, and IL-6. TNFα engages TNFR1 via an autocrine pathway. In the absence of cIAP1, stimulation of TNFR1 triggers the assembly of the secondary RIP1-containing cytoplasmic complex (complex II), leading to cell death. SM-mediated IAP depletion can also favor the formation of the Ripoptosome, leading to cell death.

Abbreviations: BIR, baculoviral IAP repeat; cIAP, cellular IAP; IκBα, nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha; IAPs, inhibitors of apoptosis; IL, interleukin; IKK, IκB kinase complex; LUBAC, linear ubiquitin chain assembly complex; MCP-1, monocyte chemoattractant protein; NF-κB, nuclear factor kappa-light-chain-enhancer of activated B cells; NIK, NF-κB-inducing kinase; RIP1, receptor interacting protein 1; SM, Smac mimetic; Smac, second mitochondria-derived activator of caspases; TAK, TGFβ-activated kinase; TGFβ, transforming growth factor beta; TNF, tumor necrosis factor; TNFR1, tumor necrosis factor receptor 1; TRADD, TNFR1-associated death domain; TRAF, TNFR associated factor; UBA, ubiquitin proteasome system; XIAP, X-chromosome linked IAP; TAB, TAK1-binding partners; HOIL, heme-oxidized IRP2 ligase-1; HOIP, HOIL-IL-interaction protein; NEMO, nuclear factor-κB (NF-κB) essential modulator; UPS, ubiquitin-proteasome system.

activity) the assembly of the caspase-activating platforms that leads to cell death^{56,62} (Figure 2). Thus, cIAPs inhibit RIP1-containing caspase-activating platform assembly, either by promoting the ubiquitin-proteasome-mediated degradation of the components of the Ripoptosome⁵⁴ or by inducing a nondegradative ubiquitination of RIP1, which inhibits the cell death complex assembly and promotes survival-signaling pathway transduction.^{56,59,63}

IAPs as cell-signaling regulators

The role of IAPs in the regulation of the NF- κ B-activating signaling pathways is well documented (for review,^{25,26}). NF- κ B is a transcription factor induced by the stimulation of antigen or cytokine receptors, by the recognition of microbiological patterns by the TLRs, the nucleotide-binding oligomerization domain-containing proteins (NODs), or the NOD-like receptors (NLRs), or in response to intracellular injuries, such as DNA damage or reactive oxygen species. NF- κ B contributes to the adaptive response of cells, by mediating the expression of the proinflammatory molecules that counter microbial invasion and by promoting the expression of genes involved in cell survival, cell differentiation, and cell proliferation.⁶⁴ The transcription factor consists of heterodimers formed by one Rel subunit (RelA [also called p65], RelB, or c-Rel) and one NF- κ B subunit (the p50 subunit of NF- κ B1 or the p52 subunit of NF- κ B2). In resting cells, the p50/RelA dimer is sequestered into the cytoplasm by the inhibitor of κ B (I κ B) proteins. Upon stimulation of the cell surface or intracellular receptors, or DNA damage, p50/RelA is released as a consequence of the degradation of NF- κ B inhibitor alpha (I κ B- α) and translocated into the nucleus to stimulate proinflammatory gene transcription (Figure 2). Degradation of I κ B- α requires its phosphorylation by the I κ B kinase (IKK) complex, which is activated by ubiquitination by the linear ubiquitin chain assembly complex (LUBAC) and by phosphorylation by TGF β -activated protein kinase 1 (TAK1)⁶⁴ (Figure 2). cIAPs and XIAP promote the steric proximity of TAK1, LUBAC, and IKK complex. In the TNF-R1-signaling pathway, cIAPs are recruited along with RIP1 to the receptor⁶¹ and trigger self-ubiquitination and the nondegradative polyubiquitination of RIP1^{56,57,66} (Figure 2), and in NOD2-mediated inflammatory signaling, XIAP and cIAPs mediate the conjugation of ubiquitin chains to RIP2.^{67–69} These ubiquitin chains serve as a scaffold for the recruitment and activation of the signaling complexes leading to IKK activation^{56,61,68,70} (Figure 2). cIAPs can also modulate NF- κ B activation by catalyzing the monoubiquitination of the IKK component NF- κ B essential modulator (NEMO),

which is required for IKK activation,^{71,72} and XIAP promotes the activation of TAK1 and the steric proximity of TAK1 and IKK complex⁷¹ during TGF β and myelin basic protein (MBP) receptor signaling, or in response to DNA damage.^{15,71,73–75} A second NF- κ B-activating signaling pathway, named the noncanonical pathway, involves NF- κ B-inducing kinase (NIK), which catalyzes the phosphorylation of IKK α . In turn, IKK α induces the phosphorylation of the p100 NF- κ B2 precursor, leading to its proteolytic activation into active p52 NF- κ B2 (Figure 2). cIAPs prevent the noncanonical activation of NF- κ B by mediating the ubiquitination and the proteasomal-mediated degradation of NIK^{70,76–79} (Figure 2). Mutations in cIAP-encoding genes leading to NIK stabilization and chronic NF- κ B activation could facilitate B cell malignancy and lymphomagenesis, as observed in some multiple myelomas that harbor mutations in the cIAP1- or cIAP2-encoding genes^{41,42} and as observed in MALT lymphoma that is associated with a chromosomal translocation t(11;18)(q21;q21), generating a chimeric protein composed of the N-terminal sequence of cIAP2 fused to the C-terminal sequence of MALT1.^{17,40,80}

Cell proliferation and migration

cIAPs are positive regulators of cell proliferation, a function correlated with the nuclear localization of the proteins.^{29,81} Interestingly, the nuclear expression of cIAP1 has been associated with advanced disease stages and poor patient prognosis in human cervical and esophageal squamous cell carcinomas and bladder cancers^{36,82,83} (Supplementary Table S1). The influence of IAPs on cell proliferation can be explained by their capacity to stimulate the activity of the c-Myc and E2F1 transcription factors, which are important regulators of cell cycle progression and cell proliferation with oncogenic properties.^{28,29} IAPs have also been involved in the regulation of the invasive properties of mammalian cancer cells, as recently reviewed.⁸⁴

Targeting IAPs in cancer therapy

Targeting IAPs in tumors is an important challenge and several strategies have been explored, including the use of antisense oligonucleotides and antagonist molecules. A synthetic antisense oligonucleotide to XIAP, named AEG35156, was developed by Aegera Therapeutics Inc (Montreal, QC, Canada).⁸⁵ It demonstrated promising efficiency in the pre-clinical studies. It induced a decrease of XIAP expression in tumor cell lines and tumor xenograft models, and sensitized cells to various standard chemotherapeutic agents and Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand

(TRAIL) receptor agonists.⁸⁶ AEG35156 entered into clinical trials (<http://www.clinicaltrials.gov/>) in 2005, and to date, ten Phase 1,2, or 1/2 clinical trials have been completed in solid tumors and in acute myeloid leukemia (AML) (Table 2) (for review,^{86,87}). In the trials, AEG35156 appeared to accumulate in the liver and to have efficiently downregulated XIAP messenger ribonucleic acid (mRNA) in peripheral blood mononuclear cells and hepatocytes. AEG35156 is generally well tolerated except when administered in repeated high doses. Promising results were obtained with AEG35156 used as a single agent in solid tumors⁸⁸ and in combination with cytarabine and idarubicin in AML⁸⁹ in the Phase 1 studies, but it failed to show any significant antitumoral activity in the randomized Phase 2 studies in pancreatic adenocarcinoma, when combined with Gemcitabine,⁹⁰ or in AML, when it was given in combination with cytarabine and idarubicin.⁹¹

The structural characterization of the interaction of XIAP with caspases, or with Smac, or the drosophila Smac homologs has provided very potent tools for the design of synthetic IAP antagonists aiming to inhibit the capacity of XIAP to neutralize caspases.^{11,92-94} The surface hydrophobic groove of IAP BIRs binds the IBM found in the N-terminal

of the active subunits of caspase-3, -7, and -9 and exposed by activating proteolytic processing.^{10,11} Cellular IAP antagonists also own an IBM.¹⁰⁻¹³ During the apoptotic process, the Smac IBM is exposed as a consequence of the cleavage of the mitochondria-targeting signal, and matured Smac is released from the mitochondria into the cytosol.¹⁰⁻¹² The tetrapeptide Ala-Val-Pro-Ile (AVPI) IBM motif of Smac inserts into the XIAP BIR2 and BIR3-caspase interaction pocket and abrogates XIAP-mediated caspase inhibition^{93,95,96} (Figure 1). The Smac N-terminal peptide was also derived to produce cell permeable peptides and was shown to mimic the activity of Smac and to sensitize human cancer cell lines to diverse chemotherapeutic agents, including etoposide, teniposide, cisplatin, paclitaxel, 7-ethyl-10-hydroxycamptothecin (SN-38), and TRAIL agonists.⁹⁷⁻¹⁰⁰ In xenograft models, a Smac-derived peptide, made permeable by linking to the shuttle peptide trans-activation of transcription (TAT) from HIV, enhanced the antitumoral effect of TRAIL in glioma,⁹⁹ and a polyarginine-conjugated Smac peptide was shown to sensitize non-small cell lung carcinoma cells to cisplatin,⁹⁸ with little toxicity to normal tissues. The pharmacological

Table 2 AEG35156 XIAP antisense oligonucleotide in clinical trials (<http://www.clinicaltrials.gov/>)

Phase	Drug combination	Start date	Condition	Observations	Ref
Phase I	Docetaxel	07/2006	Adult solid tumor		
Phase I/2	Cytarabine and idarubicin	08/2006	Refractory or relapsed acute myeloid leukemia (24 patients)	<ul style="list-style-type: none"> • Generally well tolerated • Toxicity included two cases of neuropathy in patients having received multiple AEG35156 doses • Achieved target knockdown • Clinical evidence of antitumoral activity 	89 197
Phase I	Docetaxel	09/2006	Adult solid tumor		
Phase I	Single agent	10/2006	Advanced cancer (22 patients)	<ul style="list-style-type: none"> • Well tolerated • Evidence of efficiency (decreased XIAP mRNA in peripheral blood mononuclear cells) • Clinical evidence of antitumoral activity in patients with refractory lymphoma, melanoma, and breast cancer 	88
Phase I/2	Gemcitabine	11/2007	Metastatic pancreatic adenocarcinoma (14 patients)	<ul style="list-style-type: none"> • Toxicities include neutropenia, thrombocytopenia, peripheral neuropathy, fatigue, ascites, and nausea/vomiting • Failed to show significant antitumoral activity 	90
Phase I/2	Paclitaxel	11/2007	Mammary carcinoma		
Phase I/2	Carboplatin and paclitaxel	11/2007	Non-small cell lung carcinoma		
Phase I/2	Single agent	10/2008	Refractory chronic lymphocytic leukemia and indolent B-cell lymphomas		
Phase I/2	Sorafenib	04/2009	Advanced hepatocellular carcinoma		
Phase 2	Cytarabine and idarubicin	11/2009	Refractory or relapsed acute myeloid leukemia (27 patients)	<ul style="list-style-type: none"> • Well tolerated • Did not improve rates of remission 	91

Abbreviations: RNA, ribonucleic acid; XIAP, X-chromosome linked IAP; IAPs, inhibitors of apoptosis; mRNA, messenger RNA.

properties of such Smac-derived peptides were not good enough to merit consideration of these molecules as therapeutic agents; however, they provided the bases for the structure-based design of IAP antagonists named Smac mimetics (SMs). Several approaches were used, including the screening of peptide or peptidomimetic libraries,^{101,102} and the structure-based design of conformationally constrained SMs^{103,104} (Figure 3). Considerable efforts were invested to improve the affinity of the compounds to the IAP BIR domains, to improve their ability to antagonize IAPs,^{104–107} to improve cellular delivery and activity (ie, their capacity to induce apoptosis or to sensitize to apoptotic agents), and to improve their *in vivo* stability and bioavailability. The preclinical assays demonstrated their capacity to inhibit tumor growth in multiple solid

tumors,^{102,107,108} acute lymphoblastic leukemia (ALL),¹⁰⁸ and multiple myeloma¹⁰⁹ xenograft models and to sensitize cells to TRAIL, proteasome inhibitors, B-cell lymphoma protein 2 (Bcl-2) family-targeting compounds, and more conventional therapeutic agents, such as radiation, melphalan, or cisplatin.^{103,109–114} Importantly, these compounds were well tolerated by animals and did not display toxicity against normal lymphocytes and bone marrow stromal cells¹⁰⁹ or normal mammary epithelial cells.¹¹⁵ The analysis of binding affinity revealed that similarly to Smac,^{93,95,96} SMs can bind to XIAP-BIR2, preventing XIAP-caspase-7 and -3 binding, and to XIAP-BIR3, abrogating the XIAP-mediated inhibition of caspase-9. Structural and biochemical studies of the apoptotic activity of Smac cellular protein revealed, first, that it forms a symmetric dimer;^{93,94} second, that




Figure 3 Structure of the Smac N-terminal tetrapeptide (AVPI) and SMs used in clinical trials.

Notes: AT-406: CAS RN 1071992-99-8; GDC-0152: CAS RN 873652-48-3; LCL161: CAS RN 1005342-46-0; and Birinapant: CAS RN 1260251-31-7.

Abbreviations: CAS RN, CAS Registry Number®; SM, Smac mimetic; Smac, second mitochondria-derived activator of caspases.

Table 3 SMs in clinical trials

Compound	Structure	Preclinical assays in animals	Clinical trial	Conditions
AT-406 (SM-406 – Debio 1143)	Monovalent	<ul style="list-style-type: none"> Inhibited tumor growth and sensitized cells to carboplatin in ovarian cancer xenograft model¹¹⁹ Inhibited tumor growth in breast tumor xenograft model with no sign of toxicity¹²⁰ Increased chemo- and radiotherapy sensibility in head and neck squamous cell carcinoma tumor xenograft model¹²¹ 	Phase I	Advanced solid tumors and lymphoma
Ascenta therapeutics/ debiopharm SA	Bivalent	<ul style="list-style-type: none"> Tumor growth arrest or inhibition in patient-derived primary pancreatic cancer explant model¹⁹⁹ Remission in acute lymphoblastic leukemia xenograft models²⁰⁰ 	Phase I	Combination with daunorubicin and cytarabine in patients with poor-risk acute myelogenous leukemia Refractory solid tumors or lymphoma → Well tolerated with no dose limiting toxicities, potent and sustained target inhibition, apoptotic pathway activation in tumor and antitumoral activity in colon cancer and melanoma ²⁰³
Birinapant (TL-32711)	Bivalent	<ul style="list-style-type: none"> Delayed the tumor growth and increases survival in combination with ionizing radiation in a glioblastoma multiform model in mice²⁰¹ 	Phase I/2	Combination chemotherapy (doxorubicin, paclitaxel, carboplatin, gemcitabine, irinotecan, docetaxel) in advanced and metastatic solid tumors
TetraLogic pharmaceuticals	Bivalent	<ul style="list-style-type: none"> Inhibited tumor growth in combination with the immunomodulatory agents IFNα or GM-CSF in a kidney carcinoma xenograft model²⁰² 	Phase I/2	Acute myelogenous leukemia, myelodysplastic syndrome and acute lymphoblastic leukemia
GDC-0917 Genentech	Monovalent	<ul style="list-style-type: none"> Inhibits tumor growth in breast cancer xenograft without affecting normal mammary epithelial cells²⁰⁴ 	Phase I	Combination with gemcitabine in patients with advanced solid tumor
GDC-0152 Genentech	Monovalent	<ul style="list-style-type: none"> Inhibits tumor growth in breast cancer xenograft without affecting normal mammary epithelial cells²⁰⁴ 	Phase 1/2 Phase I	Advanced ovarian, fallopian tube and peritoneal cancers Combination with 5-azacytidine in myelodysplastic syndrome Refractory solid tumors or lymphoma
HGS1029 (AEG-40826)	Bivalent	<ul style="list-style-type: none"> Induces an increased systemic level of cytokines and chemokines (TNFα and MCP-1), a systemic inflammatory response and hepatic injury when IV administered in dogs;²⁰⁵ such effects were not observed in human¹³³ 	Phase I	Locally advanced or metastatic solid malignancies, or non-Hodgkin's lymphoma without leukemic phase → Well tolerated, no signs of a systemic inflammatory response
Human Genome Sciences LCL161	Bivalent	<ul style="list-style-type: none"> Delays tumor growth in multiple solid tumor xenograft models as a single agent but is ineffective in acute lymphoblastic leukemia xenograft models¹⁰⁸ 	Phase I Phase I	Advanced solid tumors and refractory lymphoid malignancies Relapsed or refractory lymphoid malignancies
Novartis pharmaceuticals	Monovalent	<ul style="list-style-type: none"> Antitumor activity in combination with chemotherapy against a range of solid tumors including primary models of breast cancer (Novartis website)* Inhibits tumor growth in combination with a Bcl-2 inhibitor in hepatocellular carcinoma xenograft models¹¹⁰ Inhibits tumor growth and prolongs survival in combination with adeno-associated virus bacteriophage-TNFα in melanoma xenograft models¹¹¹ 	Phase I Phase 2	Solid tumors → Well tolerated ¹³⁵ Combination with weekly paclitaxel in patients with advanced solid tumor Combination with weekly paclitaxel in patients with breast cancer

Note: *Novartis website: <http://www.novartis.com/research/pipeline/c1161.jsp>
Abbreviations: GM-CSF, granulocyte-macrophage colony-stimulating factor; IFN, interferon; IV, intravenous; TNF, tumor necrosis factor; MCP-1, monocyte chemoattractant protein; Smac, second mitochondria-derived activator of caspases; SMs, Smac mimetics; Bcl-2, B-cell lymphoma 2.

dimerization is essential for Smac function,^{93,116} and third, that the capacity of Smac to abrogate XIAP-mediated caspase inhibition required the binding to both BIR2 and BIR3.⁹⁵ Overall, these observations support the conclusion that compounds targeting both BIR domains could be more efficient as XIAP antagonists and lead to the development of bivalent small molecules containing two Smac AVPI IBM motif mimetics.¹¹⁷ As expected, these compounds appeared to be more potent than their monovalent counterparts, in antagonizing XIAP and in activating caspases.^{104,117,118} Like the monovalent versions, the bivalent molecules either inhibited tumor growth or sensitized cells to both conventional and nonconventional anticancer therapies in the preclinical assays and did not display toxicity to normal human primary cells;^{104,117–121} however, unlike the monovalent molecules, the bivalent SMs are not orally bioavailable. To date, more than 50 applications for patents related to IAP antagonists have been filed (for review,¹²²), and six SMs have entered human clinical trials (<http://www.clinicaltrials.gov/>) for the treatment of cancer (described in Table 3).

Mechanisms of action of SMs

As expected, SMs abrogate XIAP-mediated caspase inhibition and therefore increase caspase-3 and -7 activities (Figure 1). However, in addition to binding XIAP BIRs, SMs also bind the BIR domains of ML-IAP, cIAP1 and cIAP2.^{56,104,105,115,118} SMs stimulate the E3-ubiquitin ligase activity of cIAPs, which results in the ubiquitination of RIP1, leading in turn, to canonical NF- κ B activation and the rapid autoubiquitination and subsequent proteasome-mediated degradation of cIAPs^{123–126} (Figure 2). Depletion of cIAPs abolishes the cIAP-mediated ubiquitination and degradation of NIK and induces canonical activation of NF- κ B. In turn, NF- κ B induces the expression of proinflammatory cytokines and chemokines, including TNF α , which can trigger cell death by an autocrine pathway.^{66,125–127} Furthermore, depletion of cIAPs favors the assembly of the RIP1-containing cytoplasmic cell death complexes, such as complex II and Ripoptosome, resulting in cell death in some sensitive cancer cells, or in the sensitization to TNF α or DNA-damaging chemotherapeutic agents^{54,55} (Figure 2). SMs exert their activity through XIAP and cIAPs and both effects are required for their maximal antitumoral activity.^{128–130} Indeed, IAP antagonists displaying a high and selective affinity for cIAPs over XIAP appeared less potent than pan-IAP antagonists in promoting cancer cell death¹²⁹ and in sensitizing cancer cells to TRAIL.¹³¹

As a consequence of cIAP degradation and NF- κ B activation, the administration of SMs such as LCL161, GDC-0152, and HGS1029, resulted in the upregulation of cytokines and chemokines,^{132–134} including TNF α , monocyte chemoattractant protein (MCP)-1, interleukin (IL)-7, IL-6, and interferon (IFN) γ .¹³⁴ MCP-1 was used as a clinical biomarker for SMs efficiency in clinical programs.^{133,135} The analysis of the proinflammatory characteristics of cellular Smac-induced cell death suggests that the proinflammatory response elicited by SMs could activate the adaptive antitumor immune response in cancers.¹³⁶ In dogs, intravenous (IV) administration of GDC-0152 induced an acute systemic inflammatory response with lung and hepatic injury, which are consistent with TNF- α mediated toxicity;¹³⁴ however, a similar TNF- α -driven inflammatory response was not observed in humans.¹³³ Although the first clinical trials did not reveal extensive toxicity of SMs when orally or intravenously administered, additional analysis of the consequences of cytokine and chemokine secretion are required. Because osteoclast differentiation and function are stimulated by activation of the noncanonical NF- κ B pathway and because osteoclasts are susceptible to TNF-mediated death, Yang et al analyzed the influence of SMs on bone metastasis and demonstrated that SMs stimulated osteoporosis and specifically enhanced metastasis in bone.¹³⁷

Conclusion

SMs are a very promising new class of anticancer therapeutics. Results from preclinical studies have demonstrated an acceptable safety profile and some signs of antitumoral activity, in their use as a single agent or in combination with conventional or nonconventional therapies, such as dead receptor agonists, Bcl-2, or kinase-targeting therapies. The first clinical trials demonstrated a good tolerance and target inhibition. Ongoing and future clinical trials will determine the safety, appropriate indications, and drugs combinations. It will be important to determine the level and the site of production of TNF α and other cytokines and the consequences of cytokine production for tumoral and non-tumoral cells. Since IAPs are involved in the regulation of various cellular functions, it will be interesting to target specific IAP functions in order to limit possible adverse impacts. The consequences of SMs on the immune system *in vivo* and the use of cIAPs as potential therapeutic targets for inflammatory or immune disorders are still important questions that need to be addressed.

Acknowledgments

Our work is supported by grants from the “Comité de Côte d’Or de la Ligue contre le Cancer,” from the “Association pour la recherche sur le Cancer (ARC),” and from the “Conseil Régional de Bourgogne.” JB received a fellowship from the “Ministère de l’Enseignement Supérieur et de la Recherche” of France, and VG received a fellowship from the “Ligue Nationale contre le Cancer.”

Disclosure

The authors report no conflicts of interest in this work.

References

- Crook NE, Clem RJ, Miller LK. An apoptosis-inhibiting baculovirus gene with a zinc finger-like motif. *J Virol*. 1993;67(4):2168–2174.
- Rothe M, Pan MG, Henzel WJ, Ayres TM, Goeddel DV. The TNFR2-TRAF signaling complex contains two novel proteins related to baculoviral inhibitor of apoptosis proteins. *Cell*. 1995;83(7):1243–1252.
- Roy N, Mahadevan MS, McLean M, et al. The gene for neuronal apoptosis inhibitory protein is partially deleted in individuals with spinal muscular atrophy. *Cell*. 1995;80(1):167–178.
- Duckett CS, Nava VE, Gedrich RW, et al. A conserved family of cellular genes related to the baculovirus iap gene and encoding apoptosis inhibitors. *EMBO J*. 1996;15(11):2685–2694.
- Uren AG, Pakusch M, Hawkins CJ, Puls KL, Vaux DL. Cloning and expression of apoptosis inhibitory protein homologs that function to inhibit apoptosis and/or bind tumor necrosis factor receptor-associated factors. *Proc Natl Acad Sci U S A*. 1996;93(10):4974–4978.
- Ambrosini G, Adida C, Altieri DC. A novel anti-apoptosis gene, survivin, expressed in cancer and lymphoma. *Nat Med*. 1997;3(8):917–921.
- Chen Z, Naito M, Hori S, Mashima T, Yamori T, Tsuruo T. A human IAP-family gene, apollon, expressed in human brain cancer cells. *Biochem Biophys Res Commun*. 1999;264(3):847–854.
- Vucic D, Stennicke HR, Pisabarro MT, Salvesen GS, Dixit VM. ML-IAP, a novel inhibitor of apoptosis that is preferentially expressed in human melanomas. *Curr Biol*. 2000;10(21):1359–1366.
- Richter BW, Mir SS, Eiben LJ, et al. Molecular cloning of ILP-2, a novel member of the inhibitor of apoptosis protein family. *Mol Cell Biol*. 2001;21(13):4292–4301.
- Srinivasula SM, Hegde R, Saleh A, et al. A conserved XIAP-interaction motif in caspase-9 and Smac/DIABLO regulates caspase activity and apoptosis. *Nature*. 2001;410(6824):112–116.
- Liu Z, Sun C, Olejniczak ET, et al. Structural basis for binding of Smac/DIABLO to the XIAP BIR3 domain. *Nature*. 2000;408(6815):1004–1008.
- Hegde R, Srinivasula SM, Zhang Z, et al. Identification of Omi/HtrA2 as a mitochondrial apoptotic serine protease that disrupts inhibitor of apoptosis protein-caspase interaction. *J Biol Chem*. 2002;277(1):432–438.
- Shi Y. A conserved tetrapeptide motif: potentiating apoptosis through IAP-binding. *Cell Death Differ*. 2002;9(2):93–95.
- Verhagen AM, Silke J, Ekert PG, et al. HtrA2 promotes cell death through its serine protease activity and its ability to antagonize inhibitor of apoptosis proteins. *J Biol Chem*. 2002;277(1):445–454.
- Lu M, Lin SC, Huang Y, et al. XIAP induces NF-kappaB activation via the BIR1/TAB1 interaction and BIR1 dimerization. *Mol Cell*. 2007;26(5):689–702.
- Samuel T, Welsh K, Lober T, Togo SH, Zapata JM, Reed JC. Distinct BIR domains of cIAP1 mediate binding to and ubiquitination of tumor necrosis factor receptor-associated factor 2 and second mitochondrial activator of caspases. *J Biol Chem*. 2006;281(2):1080–1090.
- Varfolomeev E, Wayson SM, Dixit VM, Fairbrother WJ, Vucic D. The inhibitor of apoptosis protein fusion c-IAP2.MALT1stimulates NF-kappaB activation independently of TRAF1 AND TRAF2. *J Biol Chem*. 2006;281(39):29022–29029.
- Mace PD, Smits C, Vaux DL, Silke J, Day CL. Asymmetric recruitment of cIAPs by TRAF2. *J Mol Biol*. 2010;400(1):8–15.
- Broemer M, Tenev T, Rigbolt KT, et al. Systematic in vivo RNAi analysis identifies IAPs as NEDD8-E3 ligases. *Mol Cell*. 2010;40(5):810–822.
- Vucic D, Dixit VM, Wertz IE. Ubiquitylation in apoptosis: a post-translational modification at the edge of life and death. *Nat Rev Mol Cell Biol*. 2011;12(7):439–452.
- Broemer M, Meier P. Ubiquitin-mediated regulation of apoptosis. *Trends Cell Biol*. 2009;19(3):130–140.
- Huang HK, Joazeiro CA, Bonfoco E, Kamada S, Leverson JD, Hunter T. The inhibitor of apoptosis, cIAP2, functions as a ubiquitin-protein ligase and promotes in vitro monoubiquitination of caspases 3 and 7. *J Biol Chem*. 2000;275(35):26661–26664.
- Suzuki Y, Nakabayashi Y, Takahashi R. Ubiquitin-protein ligase activity of X-linked inhibitor of apoptosis protein promotes proteasomal degradation of caspase-3 and enhances its anti-apoptotic effect in Fas-induced cell death. *Proc Natl Acad Sci U S A*. 2001;98(15):8662–8667.
- Choi YE, Butterworth M, Malladi S, Duckett CS, Cohen GM, Bratton SB. The E3 ubiquitin ligase cIAP1 binds and ubiquitinates caspase-3 and -7 via unique mechanisms at distinct steps in their processing. *J Biol Chem*. 2009;284(19):12772–12782.
- Gyrd-Hansen M, Meier P. IAPs: from caspase inhibitors to modulators of NF-kappaB, inflammation and cancer. *Nat Rev Cancer*. 2010;10(8):561–574.
- Beug ST, Cheung HH, LaCasse EC, Korneluk RG. Modulation of immune signalling by inhibitors of apoptosis. *Trends Immunol*. 2012;33(11):535–545.
- Oberoi TK, Dogan T, Hocking JC, et al. IAPs regulate the plasticity of cell migration by directly targeting Rac1 for degradation. *EMBO J*. 2011;31(1):14–28.
- Xu L, Zhu J, Hu X, et al. c-IAP1 cooperates with Myc by acting as a ubiquitin ligase for Mad1. *Mol Cell*. 2007;28(5):914–922.
- Cartier J, Berthelet J, Marivin A, et al. Cellular inhibitor of apoptosis protein-1 (cIAP1) can regulate E2F1 transcription factor-mediated control of cyclin transcription. *J Biol Chem*. 2011;286(30):26406–26417.
- Gottfried Y, Rotem A, Lotan R, Steller H, Larisch S. The mitochondrial ARTS protein promotes apoptosis through targeting XIAP. *EMBO J*. 2004;23(7):1627–1635.
- García-Fernández M, Kissel H, Brown S, et al. Sept4/ARTS is required for stem cell apoptosis and tumor suppression. *Genes Dev*. 2010;24(20):2282–2293.
- Zender L, Spector MS, Xue W, et al. Identification and validation of oncogenes in liver cancer using an integrative oncogenomic approach. *Cell*. 2006;125(7):1253–1267.
- Cheng L, Zhou Z, Flesken-Nikitin A, et al. Rb inactivation accelerates neoplastic growth and substitutes for recurrent amplification of cIAP1, cIAP2 and Yap1 in sporadic mammary carcinoma associated with p53 deficiency. *Oncogene*. 2010;29(42):5700–5711.
- Reardon DA, Michalkiewicz E, Boyett JM, et al. Extensive genomic abnormalities in childhood medulloblastoma by comparative genomic hybridization. *Cancer Res*. 1997;57(18):4042–4047.
- Bashyam MD, Bair R, Kim YH, et al. Array-based comparative genomic hybridization identifies localized DNA amplifications and homozygous deletions in pancreatic cancer. *Neoplasia*. 2005;7(6):556–562.
- Imoto I, Tsuda H, Hirasawa A, et al. Expression of cIAP1, a target for 11q22 amplification, correlates with resistance of cervical cancers to radiotherapy. *Cancer Res*. 2002;62(17):4860–4866.
- Dai Z, Zhu WG, Morrison CD, et al. A comprehensive search for DNA amplification in lung cancer identifies inhibitors of apoptosis cIAP1 and cIAP2 as candidate oncogenes. *Hum Mol Genet*. 2003;12(7):791–801.

38. Snijders AM, Schmidt BL, Fridlyand J, et al. Rare amplicons implicate frequent deregulation of cell fate specification pathways in oral squamous cell carcinoma. *Oncogene*. 2005;24(26):4232–4242.
39. Imoto I, Yang ZQ, Pimkhaokham A, et al. Identification of cIAP1 as a candidate target gene within an amplicon at 11q22 in esophageal squamous cell carcinomas. *Cancer Res*. 2001;61(18):6629–6634.
40. Garrison JB, Samuel T, Reed JC. TRAF2-binding BIR1 domain of c-IAP2/MALT1 fusion protein is essential for activation of NF-kappaB. *Oncogene*. 2009;28(13):1584–1593.
41. Keats JJ, Fonseca R, Chesi M, et al. Promiscuous mutations activate the noncanonical NF-kappaB pathway in multiple myeloma. *Cancer Cell*. 2007;12(2):131–144.
42. Annunziata CM, Davis RE, Demchenko Y, et al. Frequent engagement of the classical and alternative NF-kappaB pathways by diverse genetic abnormalities in multiple myeloma. *Cancer Cell*. 2007;12(2):115–130.
43. Filipovich AH, Zhang K, Snow AL, Marsh RA. X-linked lymphoproliferative syndromes: brothers or distant cousins? *Blood*. 2010;116(18):3398–3408.
44. Coumar MS, Tsai FY, Kanwar JR, Sarvagalla S, Cheung CH. Treat cancers by targeting survivin: just a dream or future reality? *Cancer Treat Rev*. [Epub February 28, 2013.]
45. Eckelman BP, Drag M, Snipas SJ, Salvesen GS. The mechanism of peptide-binding specificity of IAP BIR domains. *Cell Death Differ*. 2008;15(5):920–928.
46. Tenev T, Zachariou A, Wilson R, Ditzel M, Meier P. IAPs are functionally non-equivalent and regulate effector caspases through distinct mechanisms. *Nat Cell Biol*. 2005;7(1):70–77.
47. Scott FL, Denault JB, Riedl SJ, Shin H, Rensatus M, Salvesen GS. XIAP inhibits caspase-3 and -7 using two binding sites: evolutionarily conserved mechanism of IAPs. *EMBO J*. 2005;24(3):645–655.
48. Shiozaki EN, Chai J, Rigotti DJ, et al. Mechanism of XIAP-mediated inhibition of caspase-9. *Mol Cell*. 2003;11(2):519–527.
49. Riedl SJ, Rensatus M, Schwarzenbacher R, et al. Structural basis for the inhibition of caspase-3 by XIAP. *Cell*. 2001;104(5):791–800.
50. Holcik M, Yeh C, Korneluk RG, Chow T. Translational upregulation of X-linked inhibitor of apoptosis (XIAP) increases resistance to radiation induced cell death. *Oncogene*. 2000;19(36):4174–4177.
51. Gu L, Zhu N, Zhang H, Durden DL, Feng Y, Zhou M. Regulation of XIAP translation and induction by MDM2 following irradiation. *Cancer Cell*. 2009;15(5):363–375.
52. Mace PD, Riedl SJ. Molecular cell death platforms and assemblies. *Curr Opin Cell Biol*. 2010;22(6):828–836.
53. Micheau O, Tschopp J. Induction of TNF receptor I-mediated apoptosis via two sequential signaling complexes. *Cell*. 2003;114(2):181–190.
54. Tenev T, Bianchi K, Darding M, et al. The Ripoptosome, a signaling platform that assembles in response to genotoxic stress and loss of IAPs. *Mol Cell*. 2011;43(3):432–448.
55. Feoktistova M, Geserick P, Kellert B, et al. cIAPs block ripoptosome formation, a RIP1/caspase-8 containing intracellular cell death complex differentially regulated by cFLIP isoforms. *Mol Cell*. 2011;43(3):449–463.
56. Bertrand MJ, Milutinovic S, Dickson KM, et al. cIAP1 and cIAP2 facilitate cancer cell survival by functioning as E3 ligases that promote RIP1 ubiquitination. *Mol Cell*. 2008;30(6):689–700.
57. Dynek JN, Goncharov T, Dueber EC, et al. c-IAP1 and UbcH5 promote K11-linked polyubiquitination of RIP1 in TNF signalling. *EMBO J*. 2010;29(24):4198–4209.
58. Park SM, Yoon JB, Lee TH. Receptor interacting protein is ubiquitinated by cellular inhibitor of apoptosis proteins (c-IAP1 and c-IAP2) in vitro. *FEBS Lett*. 2004;566(1–3):151–156.
59. Geserick P, Hupe M, Moulin M, et al. Cellular IAPs inhibit a cryptic CD95-induced cell death by limiting RIP1 kinase recruitment. *J Cell Biol*. 2009;187(7):1037–1054.
60. Vince JE, Wong WW, Gentle I, et al. Inhibitor of apoptosis proteins limit RIP3 kinase-dependent interleukin-1 activation. *Immunity*. 2012;36(2):215–227.
61. Haas TL, Emmerich CH, Gerlach B, et al. Recruitment of the linear ubiquitin chain assembly complex stabilizes the TNF-R1 signaling complex and is required for TNF-mediated gene induction. *Mol Cell*. 2009;36(5):831–844.
62. O'Donnell MA, Legarda-Addison D, Skountzos P, Yeh WC, Ting AT. Ubiquitination of RIP1 regulates an NF-kappaB-independent cell-death switch in TNF signaling. *Curr Biol*. 2007;17(5):418–424.
63. Vanlangenakker N, Vanden Berghe T, Bogaert P, et al. cIAP1 and TAK1 protect cells from TNF-induced necrosis by preventing RIP1/RIP3-dependent reactive oxygen species production. *Cell Death Differ*. 2011;18(4):656–665.
64. Hayden MS, Ghosh S. NF-kappaB, the first quarter-century: remarkable progress and outstanding questions. *Genes Dev*. 2012;26(3):203–234.
65. Shih VF, Tsui R, Caldwell A, Hoffmann A. A single NFkB system for both canonical and non-canonical signaling. *Cell Res*. 2011;21(1):86–102.
66. Varfolomeev E, Goncharov T, Fedorova AV, et al. c-IAP1 and c-IAP2 are critical mediators of tumor necrosis factor alpha (TNFalpha)-induced NF-kappaB activation. *J Biol Chem*. 2008;283(36):24295–24299.
67. Bertrand MJ, Doiron K, Labbé K, Korneluk RG, Barker PA, Saleh M. Cellular inhibitors of apoptosis cIAP1 and cIAP2 are required for innate immunity signaling by the pattern recognition receptors NOD1 and NOD2. *Immunity*. 2009;30(6):789–801.
68. Damgaard RB, Nachbur U, Yabal M, et al. The ubiquitin ligase XIAP recruits LUBAC for NOD2 signaling in inflammation and innate immunity. *Mol Cell*. 2012;46(6):746–758.
69. Krieg A, Correa RG, Garrison JB, et al. XIAP mediates NOD signaling via interaction with RIP2. *Proc Natl Acad Sci U S A*. 2009;106(34):14524–14529.
70. Varfolomeev E, Goncharov T, Maecker H, et al. Cellular inhibitors of apoptosis are global regulators of NF-kB and MAPK activation by members of the TNF family of receptors. *Sci Signal*. 2012;5(216):ra22.
71. Jin HS, Lee DH, Kim DH, Chung JH, Lee SJ, Lee TH. cIAP1, cIAP2, and XIAP act cooperatively via nonredundant pathways to regulate genotoxic stress-induced nuclear factor-kappaB activation. *Cancer Res*. 2009;69(5):1782–1791.
72. Hinz M, Stilmann M, Arslan SC, Khanna KK, Dittmar G, Scheidereit C. A cytoplasmic ATM-TRAF6-cIAP1 module links nuclear DNA damage signaling to ubiquitin-mediated NF-kB activation. *Mol Cell*. 2010;40(1):63–74.
73. Hofer-Warbinek R, Schmid JA, Stehlik C, Binder BR, Lipp J, de Martin R. Activation of NF-kappa B by XIAP, the X chromosome-linked inhibitor of apoptosis, in endothelial cells involves TAK1. *J Biol Chem*. 2000;275(29):22064–22068.
74. Birkey Reffey S, Wurthner JU, Parks WT, Roberts AB, Duckett CS. X-linked inhibitor of apoptosis protein functions as a cofactor in transforming growth factor-beta signaling. *J Biol Chem*. 2001;276(28):26542–26549.
75. Sanna MG, da Silva Correia J, Ducrey O, et al. IAP suppression of apoptosis involves distinct mechanisms: the TAK1/JNK1 signaling cascade and caspase inhibition. *Mol Cell Biol*. 2002;22(6):1754–1766.
76. Zarnegar BJ, Wang Y, Mahoney DJ, et al. Noncanonical NF-kappaB activation requires coordinated assembly of a regulatory complex of the adaptors cIAP1, cIAP2, TRAF2 and TRAF3 and the kinase NIK. *Nat Immunol*. 2008;9(12):1371–1378.
77. Vallabhapurapu S, Matsuzawa A, Zhang W, et al. Nonredundant and complementary functions of TRAF2 and TRAF3 in a ubiquitination cascade that activates NIK-dependent alternative NF-kappaB signaling. *Nat Immunol*. 2008;9(12):1364–1370.
78. Feltham R, Moulin M, Vince JE, et al. Tumor necrosis factor (TNF) signaling, but not TWEAK (TNF-like weak inducer of apoptosis)-triggered cIAP1 (cellular inhibitor of apoptosis protein 1) degradation, requires cIAP1 RING dimerization and E2 binding. *J Biol Chem*. 2010;285(23):17525–17536.
79. Mao AP, Li S, Zhong B, et al. Virus-triggered ubiquitination of TRAF3/6 by cIAP1/2 is essential for induction of interferon-beta (IFN-beta) and cellular antiviral response. *J Biol Chem*. 2010;285(13):9470–9476.

80. Rosebeck S, Madden L, Jin X, et al. Cleavage of NIK by the API2-MALT1 fusion oncoprotein leads to noncanonical NF-kappaB activation. *Science*. 2011;331(6016):468–472.
81. Samuel T, Okada K, Hyer M, Welsh K, Zapata JM, Reed JC. cIAP1 Localizes to the nuclear compartment and modulates the cell cycle. *Cancer Res*. 2005;65(1):210–218.
82. Tanimoto T, Tsuda H, Imazeki N, et al. Nuclear expression of cIAP-1, an apoptosis inhibiting protein, predicts lymph node metastasis and poor patient prognosis in head and neck squamous cell carcinomas. *Cancer Lett*. 2005;224(1):141–151.
83. Che X, Yang D, Zong H, et al. Nuclear cIAP1 overexpression is a tumor stage- and grade-independent predictor of poor prognosis in human bladder cancer patients. *Urol Oncol*. 2012;30(4):450–456.
84. Fulda S. Regulation of cell migration, invasion and metastasis by IAP proteins and their antagonists. *Oncogene*. [Epub March 11, 2013.]
85. McManus DC, Lefebvre CA, Cherton-Horvat G, et al. Loss of XIAP protein expression by RNAi and antisense approaches sensitizes cancer cells to functionally diverse chemotherapeutics. *Oncogene*. 2004;23(49):8105–8117.
86. Lacasse EC. Pulling the plug on a cancer cell by eliminating XIAP with AEG35156. *Cancer Lett*. 2013;332(2):215–224.
87. Katragadda L, Carter BZ, Borthakur G. XIAP antisense therapy with AEG 35156 in acute myeloid leukemia. *Expert Opin Investig Drugs*. 2013;22(5):663–670.
88. Dean E, Jodrell D, Connolly K, et al. Phase I trial of AEG35156 administered as a 7-day and 3-day continuous intravenous infusion in patients with advanced refractory cancer. *J Clin Oncol*. 2009;27(10):1660–1666.
89. Schimmer AD, Estey EH, Borthakur G, et al. Phase I/II trial of AEG35156 X-linked inhibitor of apoptosis protein antisense oligonucleotide combined with idarubicin and cytarabine in patients with relapsed or primary refractory acute myeloid leukemia. *J Clin Oncol*. 2009;27(28):4741–4746.
90. Mahadevan D, Chalasani P, Rensvold D, et al. Phase I trial of AEG35156 an antisense oligonucleotide to XIAP plus gemcitabine in patients with metastatic pancreatic ductal adenocarcinoma. *Am J Clin Oncol*. 2013;36(3):239–243.
91. Schimmer AD, Herr W, Hanel M, et al. Addition of AEG35156 XIAP antisense oligonucleotide in reinduction chemotherapy does not improve remission rates in patients with primary refractory acute myeloid leukemia in a randomized phase II study. *Clin Lymphoma Myeloma Leuk*. 2011;11(5):433–438.
92. Chai J, Yan N, Huh JR, et al. Molecular mechanism of Reaper-Grim-Hid-mediated suppression of DIAP1-dependent Dronc ubiquitination. *Nat Struct Biol*. 2003;10(11):892–898.
93. Chai J, Du C, Wu JW, Kyin S, Wang X, Shi Y. Structural and biochemical basis of apoptotic activation by Smac/DIABLO. *Nature*. 2000;406(6798):855–862.
94. Wu G, Chai J, Suber TL, et al. Structural basis of IAP recognition by Smac/DIABLO. *Nature*. 2000;408(6815):1008–1012.
95. Huang Y, Rich RL, Myszkowski DG, Wu H. Requirement of both the second and third BIR domains for the relief of X-linked inhibitor of apoptosis protein (XIAP)-mediated caspase inhibition by Smac. *J Biol Chem*. 2003;278(49):49517–49522.
96. Du C, Fang M, Li Y, Li L, Wang X. Smac, a mitochondrial protein that promotes cytochrome c-dependent caspase activation by eliminating IAP inhibition. *Cell*. 2000;102(1):33–42.
97. Arnt CR, Chiorean MV, Heldebrant MP, Gores GJ, Kaufmann SH. Synthetic Smac/DIABLO peptides enhance the effects of chemotherapeutic agents by binding XIAP and cIAP1 in situ. *J Biol Chem*. 2002;277(46):44236–44243.
98. Yang L, Mashima T, Sato S, et al. Predominant suppression of apoptosome by inhibitor of apoptosis protein in non-small cell lung cancer H460 cells: therapeutic effect of a novel polyarginine-conjugated Smac peptide. *Cancer Res*. 2003;63(4):831–837.
99. Fulda S, Wick W, Weller M, Debatin KM. Smac agonists sensitize for Apo2L/TRAIL- or anticancer drug-induced apoptosis and induce regression of malignant glioma in vivo. *Nat Med*. 2002;8(8):808–815.
100. Guo F, Nimmanapalli R, Parawithana S, et al. Ectopic overexpression of second mitochondria-derived activator of caspases (Smac/DIABLO) or cotreatment with N-terminus of Smac/DIABLO peptide potentiates epothilone B derivative-(BMS 247550) and Apo-2L/TRAIL-induced apoptosis. *Blood*. 2002;99(9):3419–3426.
101. Franklin MC, Kadkhodayan S, Ackerly H, et al. Structure and function analysis of peptide antagonists of melanoma inhibitor of apoptosis (ML-IAP). *Biochemistry*. 2003;42(27):8223–8231.
102. Oost TK, Sun C, Armstrong RC, et al. Discovery of potent antagonists of the antiapoptotic protein XIAP for the treatment of cancer. *J Med Chem*. 2004;47(18):4417–4426.
103. Sun H, Nikolovska-Coleska Z, Yang CY, et al. Structure-based design of potent, conformationally constrained Smac mimetics. *J Am Chem Soc*. 2004;126(51):16686–16687.
104. Li L, Thomas RM, Suzuki H, De Brabander JK, Wang X, Harran PG. A small molecule Smac mimic potentiates TRAIL- and TNFalpha-mediated cell death. *Science*. 2004;305(5689):1471–1474.
105. Zobel K, Wang L, Varfolomeev E, et al. Design, synthesis, and biological activity of a potent Smac mimetic that sensitizes cancer cells to apoptosis by antagonizing IAPs. *ACS Chem Biol*. 2006;1(8):525–533.
106. Sharma SK, Straub C, Zawel L. Development of peptidomimetics targeting IAPs. *Int J Pept Res Ther*. 2006;12(1):21–32.
107. Sun H, Nikolovska-Coleska Z, Lu J, et al. Design, synthesis, and evaluation of a potent, cell-permeable, conformationally constrained second mitochondria derived activator of caspase (Smac) mimetic. *J Med Chem*. 2006;49(26):7916–7920.
108. Houghton PJ, Kang MH, Reynolds CP, et al. Initial testing (stage 1) of LCL161, a SMAC mimetic, by the Pediatric Preclinical Testing Program. *Pediatr Blood Cancer*. 2012;58(4):636–639.
109. Chauhan D, Neri P, Velankar M, et al. Targeting mitochondrial factor Smac/DIABLO as therapy for multiple myeloma (MM). *Blood*. 2007;109(3):1220–1227.
110. Chen KF, Lin JP, Shiao CW, et al. Inhibition of Bcl-2 improves effect of LCL161, a SMAC mimetic, in hepatocellular carcinoma cells. *Biochem Pharmacol*. 2012;84(3):268–277.
111. Yuan Z, Syrkin G, Adema A, et al. Blockade of inhibitors of apoptosis (IAPs) in combination with tumor-targeted delivery of tumor necrosis factor- α leads to synergistic antitumor activity. *Cancer Gene Ther*. 2013;20(1):46–56.
112. Xu Y, Zhou L, Huang J, et al. Role of Smac in determining the chemotherapeutic response of esophageal squamous cell carcinoma. *Clin Cancer Res*. 2011;17(16):5412–5422.
113. Ziegler DS, Keating J, Kesari S, et al. A small-molecule IAP inhibitor overcomes resistance to cytotoxic therapies in malignant gliomas in vitro and in vivo. *Neuro Oncol*. 2011;13(8):820–829.
114. Keating J, Tsoli M, Hallahan AR, Ingram WJ, Haber M, Ziegler DS. Targeting the inhibitor of apoptosis proteins as a novel therapeutic strategy in medulloblastoma. *Mol Cancer Ther*. 2012;11(12):2654–2663.
115. Flygare JA, Beresini M, Budha N, et al. Discovery of a potent small-molecule antagonist of inhibitor of apoptosis (IAP) proteins and clinical candidate for the treatment of cancer (GDC-0152). *J Med Chem*. 2012;55(9):4101–4113.
116. Flanagan L, Sebastia J, Delgado ME, Lennon JC, Rehm M. Dimerization of Smac is crucial for its mitochondrial retention by XIAP subsequent to mitochondrial outer membrane permeabilization. *Biochim Biophys Acta*. 2011;1813(5):819–826.
117. Sun H, Nikolovska-Coleska Z, Lu J, et al. Design, synthesis, and characterization of a potent, nonpeptide, cell-permeable, bivalent Smac mimetic that concurrently targets both the BIR2 and BIR3 domains in XIAP. *J Am Chem Soc*. 2007;129(49):15279–15294.
118. Lu J, Bai L, Sun H, et al. SM-164: a novel, bivalent Smac mimetic that induces apoptosis and tumor regression by concurrent removal of the blockade of cIAP-1/2 and XIAP. *Cancer Res*. 2008;68(22):9384–9393.
119. Brunckhorst MK, Lerner D, Wang S, Yu Q. AT-406, an orally active antagonist of multiple inhibitor of apoptosis proteins, inhibits progression of human ovarian cancer. *Cancer Biol Ther*. 2012;13(9):804–811.

120. Cai Q, Sun H, Peng Y, et al. A Potent and orally active antagonist (SM-406/AT-406) of multiple inhibitor of apoptosis proteins (IAPs) in clinical development for cancer treatment. *J Med Chem.* 2011;54(8): 2714–2726.
121. Fulda S. Inhibitor of apoptosis (IAP) proteins as therapeutic targets for radiosensitization of human cancers. *Cancer Treat Rev.* 2012;38(6): 760–766.
122. Flygare JA, Fairbrother WJ. Small-molecule pan-IAP antagonists: a patent review. *Expert Opin Ther Pat.* 2010;20(2):251–267.
123. Feltham R, Bettjeman B, Budhidarmo R, et al. Smac mimetics activate the E3 ligase activity of cIAP1 protein by promoting RING domain dimerization. *J Biol Chem.* 2011;286(19):17015–17028.
124. Dueber EC, Schoeffler AJ, Lingel A, et al. Antagonists induce a conformational change in cIAP1 that promotes autoubiquitination. *Science.* 2011;334(6054):376–380.
125. Varfolomeev E, Blankenship JW, Wayson SM, et al. IAP antagonists induce autoubiquitination of c-IAPs, NF-kappaB activation, and TNFalpha-dependent apoptosis. *Cell.* 2007;131(4):669–681.
126. Vince JE, Wong WW, Khan N, et al. IAP antagonists target cIAP1 to induce TNFalpha-dependent apoptosis. *Cell.* 2007;131(4):682–693.
127. Petersen SL, Wang L, Yalcin-Chin A, et al. Autocrine TNFalpha signaling renders human cancer cells susceptible to Smac-mimetic-induced apoptosis. *Cancer Cell.* 2007;12(5):445–456.
128. Cohen F, Alick B, Elliott LO, et al. Orally bioavailable antagonists of inhibitor of apoptosis proteins based on an azabicyclooctane scaffold. *J Med Chem.* 2009;52(6):1723–1730.
129. Ndubaku C, Varfolomeev E, Wang L, et al. Antagonism of c-IAP and XIAP proteins is required for efficient induction of cell death by small-molecule IAP antagonists. *ACS Chem Biol.* 2009;4(7):557–566.
130. Galbán S, Hwang C, Rumble JM, et al. Cytoprotective effects of IAPs revealed by a small molecule antagonist. *Biochem J.* 2009;417(3): 765–771.
131. Allensworth JL, Sauer SJ, Lyerly HK, Morse MA, Devi GR. Smac mimetic Birinapant induces apoptosis and enhances TRAIL potency in inflammatory breast cancer cells in an IAP-dependent and TNF-alpha-independent mechanism. *Breast Cancer Res Treat.* 2013;137(2): 359–371.
132. Knights AJ, Fucikova J, Pasam A, Koernig S, Cebon J. Inhibitor of apoptosis protein (IAP) antagonists demonstrate divergent immunomodulatory properties in human immune subsets with implications for combination therapy. *Cancer Immunol Immunother.* 2013;62(2): 321–335.
133. Wong H, Budha NR, West K, et al. Dogs are more sensitive to antagonists of inhibitor of apoptosis proteins than rats and humans: a translational toxicokinetic/toxicodynamic analysis. *Toxicol Sci.* 2012;130(1): 205–213.
134. Erickson RI, Tarrant J, Cain G, et al. Toxicity profile of small-molecule IAP antagonist GDC-0152 is linked to TNF- α pharmacology. *Toxicol Sci.* 2013;131(1):247–258.
135. Infante JR, Porter D, Sen SK, et al. A phase I study of LCL161, an oral IAP inhibitor, in patients with advanced cancer. Paper presented at: 101st Annual Meeting of the American Association for Cancer Research; April 17–21, 2010; Washington, DC. Philadelphia, PA: AACR; 2010. Abstract nr 2775.
136. Emeagi PU, Van Lint S, Goyvaerts C, et al. Proinflammatory characteristics of SMAC/DIABLO-induced cell death in antitumor therapy. *Cancer Res.* 2012;72(6):1342–1352.
137. Yang C, Davis JL, Zeng R, et al. Antagonism of inhibitor of apoptosis proteins increases bone metastasis via unexpected osteoclast activation. *Cancer Discov.* 2013;3(2):212–223.
138. Tamm I, Richter S, Scholz F, et al. XIAP expression correlates with monocytic differentiation in adult de novo AML: impact on prognosis. *Hematol J.* 2004;5(6):489–495.
139. Grzybowska-Izydorzycz O, Cebula B, Robak T, Smolewski P. Expression and prognostic significance of the inhibitor of apoptosis protein (IAP) family and its antagonists in chronic lymphocytic leukaemia. *Eur J Cancer.* 2010;46(4):800–810.
140. Hussain AR, Uddin S, Ahmed M, et al. Prognostic significance of XIAP expression in DLBCL and effect of its inhibition on AKT signalling. *J Pathol.* 2010;222(2):180–190.
141. Li M, Song T, Yin ZF, Na YQ. XIAP as a prognostic marker of early recurrence of nonmuscular invasive bladder cancer. *Chin Med J (Engl).* 2007;120(6):469–473.
142. Zhang Y, Zhu J, Tang Y, et al. X-linked inhibitor of apoptosis positive nuclear labeling: a new independent prognostic biomarker of breast invasive ductal carcinoma. *Diagn Pathol.* 2011;6:49.
143. Liu SS, Tsang BK, Cheung AN, et al. Anti-apoptotic proteins, apoptotic and proliferative parameters and their prognostic significance in cervical carcinoma. *Eur J Cancer.* 2001;37(9):1104–1110.
144. Xiang G, Wen X, Wang H, Chen K, Liu H. Expression of X-linked inhibitor of apoptosis protein in human colorectal cancer and its correlation with prognosis. *J Surg Oncol.* 2009;100(8):708–712.
145. Moussata D, Amara S, Siddeek B, et al. XIAP as a radioresistance factor and prognostic marker for radiotherapy in human rectal adenocarcinoma. *Am J Pathol.* 2012;181(4):1271–1278.
146. Shi YH, Ding WX, Zhou J, et al. Expression of X-linked inhibitor-of-apoptosis protein in hepatocellular carcinoma promotes metastasis and tumor recurrence. *Hepatology.* 2008;48(2):497–507.
147. Augello C, Caruso L, Maggioni M, et al. Inhibitors of apoptosis proteins (IAPs) expression and their prognostic significance in hepatocellular carcinoma. *BMC Cancer.* 2009;9:125.
148. Hiscutt EL, Hill DS, Martin S, et al. Targeting X-linked inhibitor of apoptosis protein to increase the efficacy of endoplasmic reticulum stress-induced apoptosis for melanoma therapy. *J Invest Dermatol.* 2010;130(9):2250–2258.
149. Ferreira CG, van der Valk P, Span SW, et al. Expression of X-linked inhibitor of apoptosis as a novel prognostic marker in radically resected non-small cell lung cancer patients. *Clin Cancer Res.* 2001;7(8): 2468–2474.
150. Ferreira CG, van der Valk P, Span SW, et al. Assessment of IAP (inhibitor of apoptosis) proteins as predictors of response to chemotherapy in advanced non-small-cell lung cancer patients. *Ann Oncol.* 2001;12(6): 799–805.
151. Seligson DB, Hongo F, Huerta-Yopez S, et al. Expression of X-linked inhibitor of apoptosis protein is a strong predictor of human prostate cancer recurrence. *Clin Cancer Res.* 2007;13(20):6056–6063.
152. Krajewska M, Krajewski S, Banares S, et al. Elevated expression of inhibitor of apoptosis proteins in prostate cancer. *Clin Cancer Res.* 2003;9(13):4914–4925.
153. Ramp U, Krieg T, Caliskan E, et al. XIAP expression is an independent prognostic marker in clear-cell renal carcinomas. *Hum Pathol.* 2004;35(8): 1022–1028.
154. Yan Y, Mahotka C, Heikaus S, et al. Disturbed balance of expression between XIAP and Smac/DIABLO during tumour progression in renal cell carcinomas. *Br J Cancer.* 2004;91(7):1349–1357.
155. Mizutani Y, Nakanishi H, Li YN, et al. Overexpression of XIAP expression in renal cell carcinoma predicts a worse prognosis. *Int J Oncol.* 2007;30(4):919–925.
156. Gu LQ, Li FY, Zhao L, et al. BRAFV600E mutation and X-linked inhibitor of apoptosis expression in papillary thyroid carcinoma. *Thyroid.* 2009;19(4):347–354.
157. Esposito I, Kleeff J, Abiatari I, et al. Overexpression of cellular inhibitor of apoptosis protein 2 is an early event in the progression of pancreatic cancer. *J Clin Pathol.* 2007;60(8):885–895.
158. Vallat L, Magdelénat H, Merle-Béral H, et al. The resistance of B-CLL cells to DNA damage-induced apoptosis defined by DNA microarrays. *Blood.* 2003;101(11):4598–4606.
159. Silva KL, Vasconcellos DV, Castro ED, Coelho AM, Linden R, Maia RC. Apoptotic effect of fludarabine is independent of expression of IAPs in B-cell chronic lymphocytic leukemia. *Apoptosis.* 2006;11(2): 277–285.
160. Ponnelle T, Chapusot C, Martin L, et al. Subcellular expression of c-IAP1 and c-IAP2 in colorectal cancers: relationships with clinicopathological features and prognosis. *Pathol Res Pract.* 2003;199(11):723–731.

161. Qi S, Mogi S, Tsuda H, et al. Expression of cIAP-1 correlates with nodal metastasis in squamous cell carcinoma of the tongue. *Int J Oral Maxillofac Surg*. 2008;37(11):1047–1053.
162. Stanculescu A, Bembinster LA, Borgen K, Bergamaschi A, Wiley E, Frasar J. Estrogen promotes breast cancer cell survival in an inhibitor of apoptosis (IAP)-dependent manner. *Horm Cancer*. 2010;1(3):127–135.
163. Akagi T, Motegi M, Tamura A, et al. A novel gene, MALT1 at 18q21, is involved in t(11;18)(q21;q21) found in low-grade B-cell lymphoma of mucosa-associated lymphoid tissue. *Oncogene*. 1999;18(42):5785–5794.
164. Dierlamm J, Baens M, Wlodarska I, et al. The apoptosis inhibitor gene API2 and a novel 18q gene, MLT, are recurrently rearranged in the t(11;18)(q21;q21) associated with mucosa-associated lymphoid tissue lymphomas. *Blood*. 1999;93(11):3601–3609.
165. Annunziata CM, Davis RE, Demchenko Y, et al. Frequent engagement of the classical and alternative NF-kappaB pathways by diverse genetic abnormalities in multiple myeloma. *Cancer Cell*. 2007;12(2):115–130.
166. El-Mesallamy HO, Hegab HM, Kamal AM. Expression of inhibitor of apoptosis protein (IAP) livin/BIRC7 in acute leukemia in adults: correlation with prognostic factors and outcome. *Leuk Res*. 2011;35(12):1616–1622.
167. Choi J, Hwang YK, Sung KW, et al. Expression of Livin, an antiapoptotic protein, is an independent favorable prognostic factor in childhood acute lymphoblastic leukemia. *Blood*. 2007;109(2):471–477.
168. Gazzaniga P, Gradilone A, Giuliani L, et al. Expression and prognostic significance of LIVIN, SURVIVIN and other apoptosis-related genes in the progression of superficial bladder cancer. *Ann Oncol*. 2003;14(1):85–90.
169. Wang X, Xu J, Ju S, Ni H, Zhu J, Wang H. Livin gene plays a role in drug resistance of colon cancer cells. *Clin Biochem*. 2010;43(7–8):655–660.
170. Wang TS, Ding QQ, Guo RH, et al. Expression of livin in gastric cancer and induction of apoptosis in SGC-7901 cells by shRNA-mediated silencing of livin gene. *Biomed Pharmacother*. 2010;64(5):333–338.
171. Nachmias B, Ashhab Y, Bucholtz V, et al. Caspase-mediated cleavage converts Livin from an antiapoptotic to a proapoptotic factor: implications for drug-resistant melanoma. *Cancer Res*. 2003;63(19):6340–6349.
172. Kim DK, Alvarado CS, Abramowsky CR, et al. Expression of inhibitor-of-apoptosis protein (IAP) livin by neuroblastoma cells: correlation with prognostic factors and outcome. *Pediatr Dev Pathol*. 2005;8(6):621–629.
173. Nedelcu T, Kubista B, Koller A, et al. Livin and Bcl-2 expression in high-grade osteosarcoma. *J Cancer Res Clin Oncol*. 2008;134(2):237–244.
174. Kempkensteffen C, Hinz S, Christoph F, et al. Expression of the apoptosis inhibitor livin in renal cell carcinomas: correlations with pathology and outcome. *Tumour Biol*. 2007;28(3):132–138.
175. Haferkamp A, Bedke J, Vetter C, et al. High nuclear Livin expression is a favourable prognostic indicator in renal cell carcinoma. *BJU Int*. 2008;102(11):1700–1706.
176. Kempkensteffen C, Hinz S, Krause H, et al. Expression of splicing variants of the inhibitor of apoptosis livin in testicular germ cell tumors. *Tumour Biol*. 2008;29(2):76–82.
177. Pluta A, Wrzesien-Kus A, Cebula-Obrzut B, et al. Influence of high expression of Smac/DIABLO protein on the clinical outcome in acute myeloid leukemia patients. *Leuk Res*. 2010;34(10):1308–1313.
178. Mizutani Y, Katsuoka Y, Bonavida B. Low circulating serum levels of second mitochondria-derived activator of caspase (Smac/DIABLO) in patients with bladder cancer. *Int J Oncol*. 2012;40(4):1246–1250.
179. Pluta P, Cebula-Obrzut B, Ehemann V, et al. Correlation of Smac/DIABLO protein expression with the clinico-pathological features of breast cancer patients. *Neoplasma*. 2011;58(5):430–435.
180. Arellano-Llamas A, Garcia FJ, Perez D, et al. High Smac/DIABLO expression is associated with early local recurrence of cervical cancer. *BMC Cancer*. 2006;6:256.
181. Endo K, Kohnoe S, Watanabe A, et al. Clinical significance of Smac/DIABLO expression in colorectal cancer. *Oncol Rep*. 2009;21(2):351–355.
182. Dobrzycka B, Terlikowski SJ, Bernaczyk PS, et al. Prognostic significance of smac/DIABLO in endometrioid endometrial cancer. *Folia Histochem Cytobiol*. 2011;48(4):678–681.
183. Sekimura A, Konishi A, Mizuno K, et al. Expression of Smac/DIABLO is a novel prognostic marker in lung cancer. *Oncol Rep*. 2004;11(4):797–802.
184. Mizutani Y, Katsuoka Y, Bonavida B. Prognostic significance of second mitochondria-derived activator of caspase (Smac/DIABLO) expression in bladder cancer and target for therapy. *Int J Oncol*. 2010;37(2):503–508.
185. Shibata T, Mahotka C, Wethkamp N, Heikus S, Gabbert HE, Ramp U. Disturbed expression of the apoptosis regulators XIAP, XAF1, and Smac/DIABLO in gastric adenocarcinomas. *Diagn Mol Pathol*. 2007;16(1):1–8.
186. Kempkensteffen C, Hinz S, Christoph F, et al. Expression levels of the mitochondrial IAP antagonists Smac/DIABLO and Omi/HtrA2 in clear-cell renal cell carcinomas and their prognostic value. *J Cancer Res Clin Oncol*. 2008;134(5):543–550.
187. Ouyang YQ, zur Hausen A, Orłowska-Volk M, et al. Expression levels of hnRNP G and hTra2-beta1 correlate with opposite outcomes in endometrial cancer biology. *Int J Cancer*. 2011;128(9):2010–2019.
188. Yang X, Xing H, Gao Q, et al. Regulation of HtrA2/Omi by X-linked inhibitor of apoptosis protein in chemoresistance in human ovarian cancer cells. *Gynecol Oncol*. 2005;97(2):413–421.
189. Hu XY, Xu YM, Chen XC, Ping H, Chen ZH, Zeng FQ. Immunohistochemical analysis of Omi/HtrA2 expression in prostate cancer and benign prostatic hyperplasia. *APMIS*. 2006;114(12):893–898.
190. Hu X, Chen X, Ping H, Chen Z, Zeng F, Lu G. Immunohistochemical analysis of Omi/HtrA2 expression in prostate cancer and benign prostatic hyperplasia. *J Huazhong Univ Sci Technolog Med Sci*. 2005;25(6):671–673.
191. Lee SH, Lee JW, Kim HS, et al. Immunohistochemical analysis of Omi/HtrA2 expression in stomach cancer. *APMIS*. 2003;111(5):586–590.
192. Zurawa-Janicka D, Kobiela J, Galczynska N, et al. Changes in expression of human serine protease HtrA1, HtrA2 and HtrA3 genes in benign and malignant thyroid tumors. *Oncol Rep*. 2012;28(5):1838–1844.
193. Narkiewicz J, Lapinska-Szumczyk S, Zurawa-Janicka D, Skorko-Glonek J, Emerich J, Lipinska B. Expression of human HtrA1, HtrA2, HtrA3 and TGF-beta1 genes in primary endometrial cancer. *Oncol Rep*. 2009;21(6):1529–1537.
194. Narkiewicz J, Klasa-Mazurkiewicz D, Zurawa-Janicka D, Skorko-Glonek J, Emerich J, Lipinska B. Changes in mRNA and protein levels of human HtrA1, HtrA2 and HtrA3 in ovarian cancer. *Clin Biochem*. 2008;41(7–8):561–569.
195. Gottfried Y, Voldavsky E, Yodko L, Sabo E, Ben-Itzhak O, Larisch S. Expression of the pro-apoptotic protein ARTS in astrocytic tumors: correlation with malignancy grade and survival rate. *Cancer*. 2004;101(11):2614–2621.
196. Elhasid R, Sahar D, Merling A, et al. Mitochondrial pro-apoptotic ARTS protein is lost in the majority of acute lymphoblastic leukemia patients. *Oncogene*. 2004;23(32):5468–5475.
197. Carter BZ, Mak DH, Morris SJ, et al. XIAP antisense oligonucleotide (AEG35156) achieves target knockdown and induces apoptosis preferentially in CD34+38- cells in a phase 1/2 study of patients with relapsed/refractory AML. *Apoptosis*. 2011;16(1):67–74.
198. Viertl D, Perillo-Adamer F, Rigotti S, et al. The SMAC-mimetic Debio 1143 efficiently enhanced chemo and radiotherapy in head and neck squamous cell carcinoma models. Paper presented at: 140th Annual Meeting of the American Association for Cancer Research; April 6–10, 2013; Washington, DC. Philadelphia, PA: AACR; 2013. Abstract nr 2055.

199. Ma WW, Zhang H, Hylander B, et al. TL32711, a novel Smac mimetic, exerts significant antitumor efficacy in primary pancreatic adenocarcinoma model. Paper presented at: 103rd Annual Meeting of the American Association for Cancer Research; March 31–April 4, 2012; Chicago, IL. Philadelphia, PA: AACR; 2012. Abstract nr 1939.
200. Smith MA, Carol H, Evans K, et al. Birinapant (TL32711), a small molecule Smac mimetic, induces regressions in childhood acute lymphoblastic leukemia (ALL) xenografts that express TNF α and synergizes with TNF α in vitro – a report from the Pediatric Preclinical Testing Program (PPTP). *Blood*. 2012;120:3565.
201. Cerna JD, Donna C, Naoko T, Norman C, M. AM, Stephen Y. Radiosensitization of GBM by a novel peptidomimetic of the second mitochondria-derived activator of caspases (SMAC). Paper presented at: 104th Annual Meeting of the American Association for Cancer Research; April 6–10, 2013; Washington, DC. Philadelphia, PA: AACR; 2013. Abstract nr 1599.
202. Benetatos CA, Burns JM, Borden EC, et al. The Smac mimetic Birinapant synergistically induces apoptosis in combination with type I interferons and GM-CSF. Paper presented at: 104th Annual Meeting of the American Association for Cancer Research; April 6–10, 2013; Washington, DC. Philadelphia, PA: AACR; 2013. Abstract 3336.
203. Amaravadi RK, Schilder RJ, Dy GK, et al. Phase I study of the Smac mimetic TL32711 in adult subjects with advanced solid tumors and lymphoma to evaluate safety, pharmacokinetics, pharmacodynamics, and antitumor activity. Paper presented at: 102nd Annual Meeting of the American Association for Cancer Research; April 2–6, 2011; Orlando, FL. Philadelphia, PA: AACR; 2010. Abstract nr LB-406.
204. Flygare JA, Beresini M, Budha N, et al. Discovery of a potent small-molecule antagonist of inhibitor of apoptosis (IAP) proteins and clinical candidate for the treatment of cancer (GDC-0152). *J Med Chem*. 2012;55(9):4101–4113.
205. Erickson RI, Tarrant J, Cain G, et al. Toxicity profile of small-molecule IAP antagonist GDC-0152 is linked to TNF- α pharmacology. *Toxicol Sci*. 2013;131(1):247–258.

Supplementary material

Table S1 Role of IAPs in cancer

	Cohort	Observations
XIAP overexpression		
AML	92 ¹	Associated with poor cytogenetics ¹
	78 ²	Inversely correlated with overall survival ^{1,2} Correlated with sensitivity to anticancer drugs (cytarabine) ²
BCLL	100 ³	Correlated with Ki-67 proliferation index and progressive disease; inverse correlation with overall survival ³
	301 ⁴	Associated with poor clinical outcome ⁴
Bladder carcinoma	176 ⁵	Independent prognostic factor for early recurrence of invasive cancers Correlated with poor differentiation Inversely correlated with recurrence-free survival
Breast carcinoma	102 ⁶	Nuclear expression Independent negative prognostic factor for overall survival
Cervical carcinoma	77 ⁷	
Colorectal cancer	96 ⁸	Cytoplasmic expression Independent negative prognostic factor Correlated with tumor dedifferentiation, invasion, stage, and lower disease-free and overall survival ⁸
	38 ⁹	Correlated with resistance to irradiation ⁹
Hepatocellular carcinoma	69 ¹⁰	Associated with shorter survival ^{10,11} and increased risk of relapse and metastasis ¹¹
	192 ¹¹	The cytoplasmic expression is an independent negative prognostic factor ¹¹
Melanoma	55 ¹²	Correlated with advanced tumor stage and inversely correlated with patient survival
NSCLC	144 ¹³	Mainly expressed in the cytoplasm Independent positive prognostic factor for survival in resected patients Correlated with longer overall survival
	55 ¹⁴	Inversely correlated with proliferation Ki-67 proliferation index Cytoplasmic expression No correlation with chemotherapy or radiotherapy
Ovarian cancer		AT-406-induced apoptosis is correlated with its ability to downregulate XIAP expression ¹⁵
Prostate carcinoma	226, ¹⁶ 691 ¹⁷	Deregulation of XIAP occurs early in the pathogenesis of prostate cancer ¹⁷ Independent predictor of tumor recurrence ¹⁶
	145 ¹⁸	Independent negative prognostic factor ¹⁸
Renal carcinoma	66 ¹⁹	Correlated with tumor grade and advanced tumor stage ^{18,19,20}
	109 ²⁰	Inversely correlated with patient survival ^{18,19,20}
Thyroid carcinoma	72 ²¹	
cIAP1 and cIAP2 overexpression associated with amplicon 11q21-22		
Cervical cancer	70 ²²	Nuclear expression correlated with low overall survival
ESC	42 ²³	Correlated with resistance to cisplatin/camptotecin
Hepatocarcinoma	25 ²⁴	
Mammary carcinoma	25 ²⁵	
Medulloblastoma	17 ²⁶	
NSCLC and SCLC	25 ²⁷	
	55 ¹⁴	Cytoplasmic expression No correlation with chemotherapy or radiotherapy ¹⁴
Pancreatic cancer	22, ²⁸ 33 ²⁹	Inversely correlated with patient survival
cIAP1 overexpression independent from 11q21-22 amplicon		
AML		Associated with resistance to several anticancer drugs ²
B-cell CLL	22, ³⁰ 30 ³¹	Correlated with resistance to irradiation ³⁰ No correlation with fludarabine sensitivity ³¹
	102 ³²	Nuclear expression correlated with proliferation index (Ki-67), tumor stage, and grade Inversely correlated with overall survival and recurrence free-survival
Cervical carcinoma	70 ²²	Nuclear expression Correlated with the resistance to irradiation Inversely correlated with overall survival and recurrence-free survival
	100 ³	Correlated with advanced tumor stage
Colorectal cancer	46 ³³	Nuclear expression

(Continued)

Table S1 (Continued)

	Cohort	Observations
HNSCC	55 ³⁴	Nuclear expression correlated with metastasis, advanced stage, and poor patient prognosis
NSCLC and SCLC	55 ¹⁴	Nuclear expression No correlation with chemotherapy or radiotherapy
Prostate carcinoma	691 ¹⁷	Inversely correlated with refractory disease
Squamous carcinoma of tongue	75 ³⁵	Nuclear and cytoplasmic expression Correlated with metastasis
cIAP1/cIAP2 inactivation		
Multiple myeloma	155 ^{36,37}	
c-IAP1/ HtrA2		
c-IAP1/Smac DIABLO		
CLL	100 ³	Correlated with a better response to treatment (cladribine, cyclophosphamide, fludarabine)
c-IAP2 overexpression independent of t(11q21)		
Breast cancer	144 ³⁸	
Cervical carcinoma	77 ⁷	
CLL	100 ³ 30 ³¹	Associated with progressive disease Cytoplasmic expression No correlation with fludarabine sensitivity
Colorectal cancer	46 ³³	Cytoplasmic expression
Prostate carcinoma	691 ¹⁷	Deregulation of c-IAP2 occurs early in the pathogenesis of prostate cancer Correlated with tumor stage and with refractory disease
c-IAP2/MALT chimeric protein t(11,18)(q21, q21)		
MALT myeloma	5 ^{39,40}	
ML-IAP overexpression		
AML	34 ⁴¹	Inversely correlated with overall survival
Adults ALL	34 ⁴¹	Inverse correlation with relapse-free survival and overall survival
Childhood ALL	222 ⁴²	Correlated with relapse-free survival
Bladder cancer	30 ⁴³	Correlated with relapse-free survival
Colorectal cancer		Correlated with resistance to etoposide, vincristine, 5-fluorouracil ⁴⁴
Gastric cancer	40 ⁴⁵	Correlated with metastasis and dedifferentiation
Melanoma	27 ⁴⁶	Resistance to etoposide
Neuroblastoma	68 ⁴⁷	Associated with MYCN amplification → inversely correlated with patient survival
Osteosarcoma	29 ⁴⁸	Nuclear expression: inverse correlation with overall survival
Renal cell carcinoma	152, ⁴⁹ 204 ⁵⁰	Nuclear expression ⁵⁰
Testicular cancer	131 ⁵¹	Correlated with dedifferentiation
Smac downregulation		
AML	71 ⁵²	Correlated with response to chemotherapy
Bladder cancer	173 ⁵³	Inversely correlated with advanced tumor stage and tumor grade (serum)
Breast cancer	62 ⁵⁴	Inversely correlated with tumor stage
CLL	100 ³	Inversely correlated with advanced tumor stage
Cervical carcinoma	86 ⁵⁵	Inversely correlated with local recurrence
Colorectal carcinoma	121 ⁵⁶	Inversely correlated with metastasis and advanced tumor stage Correlated with patient survival
Endometrioid endometrial cancer	76 ⁵⁷	Inversely correlated with tumor grade and correlated with longer disease-specific survival
Esophageal carcinoma	86 ⁵⁸	Inversely correlated with chemoresistance
Lung cancer	88 ⁵⁹	Inversely correlated with advanced tumor stage
Rectal adenocarcinoma	38 ⁹	Correlated with resistance to irradiation
Smac overexpression		
Bladder cancer	75 ⁶⁰	Correlated with postoperative recurrence-free period
Gastric adenocarcinoma	46 ⁶¹	Correlated with advanced tumor stage
Renal carcinoma	66, ¹⁹ 85 ⁶²	Correlated with advanced tumor stage
XIAP/ Smac		
Gastric adenocarcinoma	46 ⁶¹	Low XIAP/ Smac ratio
Renal carcinoma	66 ¹⁹	High XIAP/ Smac ratio is correlated with advanced tumor stage

(Continued)

Table S1 (Continued)

	Cohort	Observations
HtrA2 overexpression		
Endometrial cancer	139 ⁶³	Nuclear HtrA2 expression is elevated in poorly differentiated and lymph node metastatic cancer Nuclear HtrA2 expression is an independent prognostic factor for endometrial cancer progression-free survival
Ovarian cancer	64	Cytoplasmic HtrA2 expression increased in cisplatin-resistant cells
Prostate carcinoma	105, ⁶⁵ 61 ⁶⁶	Correlated with tumor grade and dedifferentiation ⁶⁵
Renal carcinoma	85 ⁶²	Correlated with recurrence-free and tumor-specific survival
Stomach cancer	60 ⁶⁷	
Thyroid cancer	68	
HtrA2 downregulation		
Endometrial cancer	124 ⁶⁹	
Ovarian cancer	79 ⁷⁰	
ARTS overexpression		
Astrocytoma	72 ⁷¹	Correlation with tumor grade and higher rate of apoptosis
ARTS downregulation		
AML	72	Resistance to 5-azacytidine

Abbreviations: ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; ARTS, septin-like mitochondrial protein; BCLL, B-cell chronic lymphocytic leukemia; cIAP1, cellular inhibitors of apoptosis; CLL, chronic lymphocytic leukemia; DIABLO, IAP-binding protein with low pI; ESC, esophageal squamous cell carcinoma; HNSCC, head and neck squamous cell carcinomas; HtrA2, high temperature requirement protein A2; IAP, inhibitors of apoptosis; NSCLC, non-small cell lung cancer; SCLC, small cell lung cancer; MALT, mucosa-associated lymphoid tissue; ML-IAP, melanoma IAP; SLL, small lymphocytic lymphoma; Smac, second mitochondria-derived activator of caspases; XIAP, X-chromosome linked IAP; pI, isoelectric point.

References

- Tamm I, Richter S, Scholz F, et al. XIAP expression correlates with monocytic differentiation in adult de novo AML: impact on prognosis. *Hematol J*. 2004;5(6):489–495.
- Tamm I, Kornblau SM, Segall H, et al. Expression and prognostic significance of IAP-family genes in human cancers and myeloid leukemias. *Clin Cancer Res*. 2000;6(5):1796–1803.
- Grzybowska-Izydorczyk O, Cebula B, Robak T, Smolewski P. Expression and prognostic significance of the inhibitor of apoptosis protein (IAP) family and its antagonists in chronic lymphocytic leukaemia. *Eur J Cancer*. 2010;46(4):800–810.
- Hussain AR, Uddin S, Ahmed M, et al. Prognostic significance of XIAP expression in DLBCL and effect of its inhibition on AKT signalling. *J Pathol*. 2011;222(2):180–190.
- Li M, Song T, Yin ZF, Na YQ. XIAP as a prognostic marker of early recurrence of nonmuscular invasive bladder cancer. *Chin Med J (Engl)*. 2007;120(6):469–473.
- Zhang Y, Zhu J, Tang Y, et al. X-linked inhibitor of apoptosis positive nuclear labeling: a new independent prognostic biomarker of breast invasive ductal carcinoma. *Diagn Pathol*. 2011;6:49.
- Liu SS, Tsang BK, Cheung AN, et al. Anti-apoptotic proteins, apoptotic and proliferative parameters and their prognostic significance in cervical carcinoma. *Eur J Cancer*. 2001;37(9):1104–1110.
- Xiang G, Wen X, Wang H, Chen K, Liu H. Expression of X-linked inhibitor of apoptosis protein in human colorectal cancer and its correlation with prognosis. *J Surg Oncol*. 2009;100(8):708–712.
- Moussata D, Amara S, Siddeek B, et al. XIAP as a radioresistance factor and prognostic marker for radiotherapy in human rectal adenocarcinoma. *Am J Pathol*. 2012;181(4):1271–1278.
- Augello C, Caruso L, Maggioni M, et al. Inhibitors of apoptosis proteins (IAPs) expression and their prognostic significance in hepatocellular carcinoma. *BMC Cancer*. 2009;9:125.
- Shi YH, Ding WX, Zhou J, et al. Expression of X-linked inhibitor-of-apoptosis protein in hepatocellular carcinoma promotes metastasis and tumor recurrence. *Hepatology*. 2008;48(2):497–507.
- Hiscutt EL, Hill DS, Martin S, et al. Targeting X-linked inhibitor of apoptosis protein to increase the efficacy of endoplasmic reticulum stress-induced apoptosis for melanoma therapy. *J Invest Dermatol*. 2010;130(9):2250–2258.
- Ferreira CG, van der Valk P, Span SW, et al. Expression of X-linked inhibitor of apoptosis as a novel prognostic marker in radically resected non-small cell lung cancer patients. *Clin Cancer Res*. 2001;7(8):2468–2474.
- Ferreira CG, van der Valk P, Span SW, et al. Assessment of IAP (inhibitor of apoptosis) proteins as predictors of response to chemotherapy in advanced non-small-cell lung cancer patients. *Ann Oncol*. 2001;12(6):799–805.
- Brunckhorst MK, Lerner D, Wang S, Yu Q. AT-406, an orally active antagonist of multiple inhibitor of apoptosis proteins, inhibits progression of human ovarian cancer. *Cancer Biol Ther*. 2012;13(9):804–811.
- Seligson DB, Hongo F, Huerta-Yepez S, et al. Expression of X-linked inhibitor of apoptosis protein is a strong predictor of human prostate cancer recurrence. *Clin Cancer Res*. 2007;13(20):6056–6063.
- Krajewska M, Krajewski S, Banares S, et al. Elevated expression of inhibitor of apoptosis proteins in prostate cancer. *Clin Cancer Res*. 2003;9(13):4914–4925.
- Ramp U, Krieg T, Caliskan E, et al. XIAP expression is an independent prognostic marker in clear-cell renal carcinomas. *Hum Pathol*. 2004;35(8):1022–1028.
- Yan Y, Mahotka C, Heikaus S, et al. Disturbed balance of expression between XIAP and Smac/DIABLO during tumour progression in renal cell carcinomas. *Br J Cancer*. 2004;91(7):1349–1357.
- Mizutani Y, Nakanishi H, Li YN, et al. Overexpression of XIAP expression in renal cell carcinoma predicts a worse prognosis. *Int J Oncol*. 2007;30(4):919–925.
- Gu LQ, Li FY, Zhao L, et al. BRAFV600E mutation and X-linked inhibitor of apoptosis expression in papillary thyroid carcinoma. *Thyroid*. 2009;19(4):347–354.
- Imoto I, Tsuda H, Hirasawa A, et al. Expression of cIAP1, a target for 11q22 amplification, correlates with resistance of cervical cancers to radiotherapy. *Cancer Res*. 2002;62(17):4860–4866.
- Imoto I, Yang ZQ, Pimkhaokham A, et al. Identification of cIAP1 as a candidate target gene within an amplicon at 11q22 in esophageal squamous cell carcinomas. *Cancer Res*. 2001;61(18):6629–6634.
- Zender L, Spector MS, Xue W, et al. Identification and validation of oncogenes in liver cancer using an integrative oncogenomic approach. *Cell*. 2006;125(7):1253–1267.

25. Cheng L, Zhou Z, Flesken-Nikitin A, et al. Rb inactivation accelerates neoplastic growth and substitutes for recurrent amplification of cIAP1, cIAP2 and Yap1 in sporadic mammary carcinoma associated with p53 deficiency. *Oncogene*. 2010;29(42):5700–5711.
26. Reardon DA, Michalkiewicz E, Boyett JM, et al. Extensive genomic abnormalities in childhood medulloblastoma by comparative genomic hybridization. *Cancer Res*. 1997;57(18):4042–4047.
27. Dai Z, Zhu WG, Morrison CD, et al. A comprehensive search for DNA amplification in lung cancer identifies inhibitors of apoptosis cIAP1 and cIAP2 as candidate oncogenes. *Hum Mol Genet*. 2003;12(7):791–801.
28. Bashyam MD, Bair R, Kim YH, et al. Array-based comparative genomic hybridization identifies localized DNA amplifications and homozygous deletions in pancreatic cancer. *Neoplasia*. 2005;7(6):556–562.
29. Esposito I, Kleeff J, Abiatari I, et al. Overexpression of cellular inhibitor of apoptosis protein 2 is an early event in the progression of pancreatic cancer. *J Clin Pathol*. 2007;60(8):885–895.
30. Vallat L, Magdelénat H, Merle-Béral H, et al. The resistance of B-CLL cells to DNA damage-induced apoptosis defined by DNA microarrays. *Blood*. 2003;101(11):4598–4606.
31. Silva KL, Vasconcellos DV, Castro ED, Coelho AM, Linden R, Maia RC. Apoptotic effect of fludarabine is independent of expression of IAPs in B-cell chronic lymphocytic leukemia. *Apoptosis*. 2006;11(2):277–285.
32. Che X, Yang D, Zong H, et al. Nuclear cIAP1 overexpression is a tumor stage- and grade-independent predictor of poor prognosis in human bladder cancer patients. *Urol Oncol*. 2012;30(4):450–456.
33. Ponnelle T, Chapusot C, Martin L, et al. Subcellular expression of c-IAP1 and c-IAP2 in colorectal cancers: relationships with clinicopathological features and prognosis. *Pathol Res Pract*. 2003;199(11):723–731.
34. Tanimoto T, Tsuda H, Imazeki N, et al. Nuclear expression of cIAP-1, an apoptosis inhibiting protein, predicts lymph node metastasis and poor patient prognosis in head and neck squamous cell carcinomas. *Cancer Lett*. 2005;224(1):141–151.
35. Qi S, Mogi S, Tsuda H, et al. Expression of cIAP-1 correlates with nodal metastasis in squamous cell carcinoma of the tongue. *Int J Oral Maxillofac Surg*. 2008;37(11):1047–1053.
36. Keats JJ, Fonseca R, Chesi M, et al. Promiscuous mutations activate the noncanonical NF-kappaB pathway in multiple myeloma. *Cancer Cell*. 2007;12(2):131–144.
37. Annunziata CM, Davis RE, Demchenko Y, et al. Frequent engagement of the classical and alternative NF-kappaB pathways by diverse genetic abnormalities in multiple myeloma. *Cancer Cell*. 2007;12(2):115–130.
38. Stanculescu A, Bembinsler LA, Borgen K, Bergamaschi A, Wiley E, Frasor J. Estrogen promotes breast cancer cell survival in an inhibitor of apoptosis (IAP)-dependent manner. *Horm Cancer*. 2010;1(3):127–135.
39. Akagi T, Motegi M, Tamura A, et al. A novel gene, MALT1 at 18q21, is involved in t(11;18)(q21;q21) found in low-grade B-cell lymphoma of mucosa-associated lymphoid tissue. *Oncogene*. 1999;18(42):5785–5794.
40. Dierlamm J, Baens M, Wlodarska I, et al. The apoptosis inhibitor gene API2 and a novel 18q gene, MLT, are recurrently rearranged in the t(11;18)(q21;q21) associated with mucosa-associated lymphoid tissue lymphomas. *Blood*. 1999;93(11):3601–3609.
41. El-Mesallamy HO, Hegab HM, Kamal AM. Expression of inhibitor of apoptosis protein (IAP) livin/BIRC7 in acute leukemia in adults: correlation with prognostic factors and outcome. *Leuk Res*. 2011;35(12):1616–1622.
42. Choi J, Hwang YK, Sung KW, et al. Expression of Livin, an antiapoptotic protein, is an independent favorable prognostic factor in childhood acute lymphoblastic leukemia. *Blood*. 2007;109(2):471–477.
43. Gazzaniga P, Gradilone A, Giuliani L, et al. Expression and prognostic significance of LIVIN, SURVIVIN and other apoptosis-related genes in the progression of superficial bladder cancer. *Ann Oncol*. 2003;14(1):85–90.
44. Wang X, Xu J, Ju S, Ni H, Zhu J, Wang H. Livin gene plays a role in drug resistance of colon cancer cells. *Clin Biochem*. 2010;43(7–8):655–660.
45. Wang TS, Ding QQ, Guo RH, et al. Expression of livin in gastric cancer and induction of apoptosis in SGC-7901 cells by shRNA-mediated silencing of livin gene. *Biomed Pharmacother*. 2010;64(5):333–338.
46. Nachmias B, Ashhab Y, Bucholtz V, et al. Caspase-mediated cleavage converts Livin from an antiapoptotic to a proapoptotic factor: implications for drug-resistant melanoma. *Cancer Res*. 2003;63(19):6340–6349.
47. Kim DK, Alvarado CS, Abramowsky CR, et al. Expression of inhibitor-of-apoptosis protein (IAP) livin by neuroblastoma cells: correlation with prognostic factors and outcome. *Pediatr Dev Pathol*. 2005;8(6):621–629.
48. Nedelcu T, Kubista B, Koller A, et al. Livin and Bcl-2 expression in high-grade osteosarcoma. *J Cancer Res Clin Oncol*. 2008;134(2):237–244.
49. Kempkensteffen C, Hinz S, Christoph F, et al. Expression of the apoptosis inhibitor livin in renal cell carcinomas: correlations with pathology and outcome. *Tumour Biol*. 2007;28(3):132–138.
50. Haferkamp A, Bedke J, Vetter C, et al. High nuclear Livin expression is a favourable prognostic indicator in renal cell carcinoma. *BJU Int*. 2008;102(11):1700–1706.
51. Kempkensteffen C, Hinz S, Krause H, et al. Expression of splicing variants of the inhibitor of apoptosis livin in testicular germ cell tumors. *Tumour Biol*. 2008;29(2):76–82.
52. Pluta P, Wrzesien-Kus A, Cebula-Obrzut B, et al. Influence of high expression of Smac/DIABLO protein on the clinical outcome in acute myeloid leukemia patients. *Leuk Res*. 2010;34(10):1308–1313.
53. Mizutani Y, Katsuoka Y, Bonavida B. Low circulating serum levels of second mitochondria-derived activator of caspase (Smac/DIABLO) in patients with bladder cancer. *Int J Oncol*. 2012;40(4):1246–1250.
54. Pluta P, Cebula-Obrzut B, Ehemann V, et al. Correlation of Smac/DIABLO protein expression with the clinico-pathological features of breast cancer patients. *Neoplasia*. 2011;58(5):430–435.
55. Arellano-Llamas A, Garcia FJ, Perez D, et al. High Smac/DIABLO expression is associated with early local recurrence of cervical cancer. *BMC Cancer*. 2006;6:256.
56. Endo K, Kohnoe S, Watanabe A, et al. Clinical significance of Smac/DIABLO expression in colorectal cancer. *Oncol Rep*. 2009;21(2):351–355.
57. Dobrzycka B, Terlikowski SJ, Bernaczyk PS, et al. Prognostic significance of smac/DIABLO in endometrioid endometrial cancer. *Folia Histochem Cytobiol*. 2011;48(4):678–681.
58. Xu Y, Zhou L, Huang J, et al. Role of Smac in determining the chemotherapeutic response of esophageal squamous cell carcinoma. *Clin Cancer Res*. 2011;17(16):5412–5422.
59. Sekimura A, Konishi A, Mizuno K, et al. Expression of Smac/DIABLO is a novel prognostic marker in lung cancer. *Oncol Rep*. 2004;11(4):797–802.
60. Mizutani Y, Katsuoka Y, Bonavida B. Prognostic significance of second mitochondria-derived activator of caspase (Smac/DIABLO) expression in bladder cancer and target for therapy. *Int J Oncol*. 2010;37(2):503–508.
61. Shibata T, Mahotka C, Wethkamp N, Heikaus S, Gabbert HE, Ramp U. Disturbed expression of the apoptosis regulators XIAP, XAF1, and Smac/DIABLO in gastric adenocarcinomas. *Diagn Mol Pathol*. 2007;16(1):1–8.
62. Kempkensteffen C, Hinz S, Christoph F, et al. Expression levels of the mitochondrial IAP antagonists Smac/DIABLO and Omi/HtrA2 in clear-cell renal cell carcinomas and their prognostic value. *J Cancer Res Clin Oncol*. 2008;134(5):543–550.
63. Ouyang YQ, zur Hausen A, Orlowska-Volk M, et al. Expression levels of hnRNP G and hTra2-beta1 correlate with opposite outcomes in endometrial cancer biology. *Int J Cancer*. 2011;128(9):2010–2019.
64. Yang X, Xing H, Gao Q, et al. Regulation of HtrA2/Omi by X-linked inhibitor of apoptosis protein in chemoresistance in human ovarian cancer cells. *Gynecol Oncol*. 2005;97(2):413–421.
65. Hu XY, Xu YM, Chen XC, Ping H, Chen ZH, Zeng FQ. Immunohistochemical analysis of Omi/HtrA2 expression in prostate cancer and benign prostatic hyperplasia. *APMIS*. 2006;114(12):893–898.

66. Hu X, Chen X, Ping H, Chen Z, Zeng F, Lu G. Immunohistochemical analysis of Omi/HtrA2 expression in prostate cancer and benign prostatic hyperplasia. *J Huazhong Univ Sci Technolog Med Sci.* 2005;25(6):671–673.
67. Lee SH, Lee JW, Kim HS, et al. Immunohistochemical analysis of Omi/HtrA2 expression in stomach cancer. *APMIS.* 2003;111(5):586–590.
68. Zurawa-Janicka D, Kobiela J, Galczynska N, et al. Changes in expression of human serine protease HtrA1, HtrA2 and HtrA3 genes in benign and malignant thyroid tumors. *Oncol Rep.* 2012;28(5):1838–1844.
69. Narkiewicz J, Lapinska-Szumczyk S, Zurawa-Janicka D, Skorko-Glonek J, Emerich J, Lipinska B. Expression of human HtrA1, HtrA2, HtrA3 and TGF-beta1 genes in primary endometrial cancer. *Oncol Rep.* 2009;21(6):1529–1537.
70. Narkiewicz J, Klasa-Mazurkiewicz D, Zurawa-Janicka D, Skorko-Glonek J, Emerich J, Lipinska B. Changes in mRNA and protein levels of human HtrA1, HtrA2 and HtrA3 in ovarian cancer. *Clin Biochem.* 2008;41(7–8):561–569.
71. Gottfried Y, Voldavsky E, Yodko L, Sabo E, Ben-Itzhak O, Larisch S. Expression of the pro-apoptotic protein ARTS in astrocytic tumors: correlation with malignancy grade and survival rate. *Cancer.* 2004; 101(11):2614–2621.
72. Elhasid R, Sahar D, Merling A, et al. Mitochondrial pro-apoptotic ARTS protein is lost in the majority of acute lymphoblastic leukemia patients. *Oncogene.* 2004;23(32):5468–5475.

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress