

# Bilateral septic knee arthritis after treatment of an ectopic pregnancy with methotrexate

Fatih Karaaslan<sup>1</sup>  
Musa Uğur Mermerkaya<sup>1</sup>  
Emre Yurdakul<sup>2</sup>  
Özlem Tanın<sup>3</sup>

<sup>1</sup>Department of Orthopaedics and Traumatology, Faculty of Medicine, Bozok University, Yozgat, Turkey;

<sup>2</sup>Department of Orthopaedics and Traumatology, Osmaniye State Hospital, Osmaniye, Turkey;

<sup>3</sup>Department of Gynecology and Obstetrics, Faculty of Medicine, Bozok University, Yozgat, Turkey

**Abstract:** We describe a case of bilateral septic knee arthritis that followed the administration of methotrexate for nonsurgical termination of an ectopic pregnancy. We believe that methotrexate is a relatively effective alternative to surgery in patients with early unruptured tubal pregnancies, but suggest that it be used with caution in view of possible septic events.

**Keywords:** bilateral septic arthritis, methotrexate, ectopic pregnancy, immunosuppression

## Introduction

Methotrexate (MTX) is a folic acid antagonist and cell cycle-specific antimetabolite. While still an important component of hematological and solid tumor treatment regimens and ectopic pregnancies, low-dose MTX is now used to treat many autoimmune disorders, including Crohn's disease, rheumatoid arthritis, and psoriasis. MTX-related hematological toxicity includes leukopenia, thrombocytopenia, megaloblastic anemia, and pancytopenia.<sup>1,2</sup> In the circulation, 50% of MTX is protein-bound. It exhibits particular affinity for hepatocytes, myeloid precursors, red blood cells, and fibroblasts. It is converted to its active polyglutamate metabolite, which persists for months and allows weekly dosing. Excretion is mostly renal and, to a lesser extent, biliary.<sup>3</sup> MTX is regarded as an important, beneficial drug, but side effects should be monitored carefully.

Septic arthropathy is an unusual complication of pregnancy. The causative organism is typically *Staphylococcus aureus*. It can quickly lead to irreversible arthropathy and disability. Factors predisposing to septic arthritis pertinent to the gravida include diabetes, rheumatoid arthritis, drug abuse, a prior intra-articular steroid injection, or trauma.<sup>4</sup> Bilateral simultaneous knee septic arthritis was identified in this case, which has, to our knowledge, not been reported before.

## Case report

A 25-year-old female, gravida 3, was admitted to the outpatient gynecology clinic complaining of abdominal pain and vaginal bleeding after 6 weeks of amenorrhea. Her previous menstrual cycles had been regular at 30 days, with average flow and no dysmenorrhea.

She had tenderness in the right lower quadrant of the abdomen, with normal vital signs. Transvaginal ultrasonography showed an empty uterine cavity and a gestational sac in the right adnexa, with no fetal cardiac activity, and there was no intra-abdominal free fluid. Laboratory tests showed an hCG level of 585 mIU/mL.

A single dose of MTX (50 mg/m<sup>2</sup>) was administered intramuscularly on day 1, and the hCG was 19.1 (minute 5) and 9.8 mIU/mL, respectively, on days 4 and 7.

Correspondence: Fatih Karaaslan  
Department of Orthopaedics and Traumatology, Faculty of Medicine, Bozok University, 18 Madrasa Adnan Menderes, Boulevard, Yozgat 66200, Turkey  
Email [fkaraaslan@gmail.com](mailto:fkaraaslan@gmail.com)

The patient's liver function tests were normal and an additional dose of MTX was given. Subsequently, the hCG levels decreased. After the treatment, however, she developed pain and swelling of both knees.

On examination, she was afebrile and appeared well. Both knees were swollen and tender. A blood count revealed a hemoglobin of 12 g/dL (normal range 12–16 g/dL), total white blood cells at 13,600/L (normal range 3,600–10,000/L), and platelets at 576,000/L (normal range 130,000–400,000/L). The erythrocyte sedimentation rate was 100 mm/h (normal range 0–20 mm/h), and the C-reactive protein (CRP) 43 mg/L (normal range 0–7 mg/L). A direct radiographic evaluation was normal (Figure 1). She had no comorbidities. Joint ultrasonography showed anterior soft tissue swelling with a thickened synovium and fluid collections measuring 1.0×0.9×2 cm in the right knee and 1.1×0.9×1.8 cm in the left. She was diagnosed with bilateral septic arthritis and underwent arthroscopic debridement and washout.

In the operating room, the puncture wound was extended about 1 cm in both directions. Pus was expressed with manual manipulation. A moderate amount of pus remained in both joints. Arthroscopic debridement and irrigation were done to break up any adhesions in the joint and to ensure that there were no pockets of pus or anything walled off, such as an abscess or phlegmon; culture antibiogram evaluation done and both joints were irrigated with approximately 3 L of saline. A Hemovac drain was placed in each joint. The wound cultures grew Gram-positive cocci (*S. aureus*) resistant to ampicillin, so vancomycin was started. The patient improved rapidly after the surgery. Active and passive movement increased to a total of 90°.

## Discussion

Ectopic pregnancy is a gynecological emergency and a serious complication of early pregnancy because of the association with maternal morbidity and mortality in the

first trimester.<sup>5</sup> Septic arthritis is an orthopedic emergency and a serious complication because of cartilage tissue damage. Factors predisposing to septic arthritis pertinent to the gravida include diabetes, rheumatoid arthritis, intravenous drug abuse, a prior intra-articular steroid injection, or trauma. Joint pain in pregnancy has a large differential diagnosis, including trauma, bacterial or viral arthropathy, collagen vascular disease, and enteropathic (inflammatory bowel disease) arthritis. Smaller, more distal joints are easier to assess than larger, more central joints. Nonetheless, joint destruction can occur with either. The patient's history and concomitant illness provide useful clues to the origins of the arthritis.<sup>6,7</sup>

Immunosuppressants have been the mainstay of treatment for various diseases for many years. Treatments have led to a change in the classical presentation of acute bone, joint, and soft tissue infections. The normal findings of severe pain and tenderness on examination can be absent or simply mimic a typical exacerbation of a chronic joint condition.<sup>8,9</sup>

We present a unique case of bilateral septic arthritis in a patient treated with MTX for ectopic pregnancy. We hope that this article will enable physicians to appreciate the changing face of septic arthritis in the modern era of immunosuppressant treatments.

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Huffman DH, Wan SH, Azarnoff DL, Hogstraten B. Pharmacokinetics of methotrexate. *Clin Pharmacol Ther.* 1973;14:572–579.
- Cheung KK, Chow KM, Szeto CC, Tai MH, Kwan BC, Li PK. Fatal pancytopenia in a hemodialysis patient after treatment with low-dose methotrexate. *J Clin Rheumatol.* 2009;15(4):177–180.
- Neves C, Jorge R, Barcelos A. A teia de toxicidade do metotrexato [The network of methotrexate toxicity]. *Acta Reumatol Port.* 2009;34:11–34. Portuguese.
- Almoujahed MO, Khatib R, Baran J. Pregnancy-associated pyogenic sacroiliitis: case report and review. *Infect Dis Obstet Gynecol.* 2003;11:53–57.
- Chen CH, Lee WL, Chiu LH, Sun HD, Liu WM, Wang PH. A cohort study to evaluate the effectiveness of laparoscopic-guided local injection of etoposide in the management of women with unruptured tubal pregnancy. *Fertil Steril.* 2011;96:654–658.
- Sharff KA, Richards EP, Townes JM. Clinical management of septic arthritis. *Curr Rheumatol Rep.* 2013;15:332.
- DeNoble PH, Gonzalez D. Septic arthritis of the shoulder following elective termination of pregnancy. *J Shoulder Elbow Surg.* 2009;18:e5–e6.
- Bourré-Tessier J, Haraoui B. Methotrexate drug interactions in the treatment of rheumatoid arthritis: a systematic review. *J Rheumatol.* 2010;37(7):1416–1421.
- Murav'ev IuV, Gridneva GI, Karateev DE, et al. [Evaluation of methotrexate effect on the acute-phase response in rheumatoid arthritis after 12-week treatment]. *Klin Med (Mosk).* 2014;92(3):59–63. Russian.



Figure 1 Anteroposterior plain radiography of both knees.

### International Medical Case Reports Journal

Dovepress

#### Publish your work in this journal

The International Medical Case Reports Journal is an international, peer-reviewed open-access journal publishing original case reports from all medical specialties. Previously unpublished medical posters are also accepted relating to any area of clinical or preclinical science. Submissions should not normally exceed 2,000 words or

4 published pages including figures, diagrams and references. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/international-medical-case-reports-journal-journal>