

# Prospective study on the overuse of blood test-guided antibiotics on patients with acute diarrhea in primary hospitals of China

Xinghui Liu<sup>1,\*</sup>

Xueke Tong<sup>2,\*</sup>

Liyin Jin<sup>3</sup>

Minghao Ha<sup>4</sup>

Feng Cao<sup>5</sup>

Fengxia Xu<sup>1</sup>

Yongbin Chi<sup>1</sup>

Denghai Zhang<sup>1</sup>

Limin Xu<sup>1</sup>

<sup>1</sup>Department of Laboratory,

<sup>2</sup>Department of Infectious Diseases, Shanghai Gongli Hospital, The Second Military Medical University,

<sup>3</sup>Department of Laboratory, Jinyang Community Health Service Center, Pudong New District, <sup>4</sup>Department of Infectious Diseases and Hepatology, The Affiliated Seventh People's Hospital of Shanghai University of Traditional Chinese Medicine,

<sup>5</sup>Department of Preventive Care, Shanghai Gongli Hospital, The Second Military Medical University, Shanghai, People's Republic of China

\*These authors contributed equally to this work

Correspondence: Limin Xu;

Denghai Zhang

Department of Clinical Laboratory, Shanghai Gongli Hospital, The Second Military Medical University, 219 Miaopu Rd, Pudong New Area, Shanghai 200135, People's Republic of China

Tel +86 21 5885 8730 ext 5236; 5177

Fax +86 21 5821 9029

Email winexlm@163.com;

shanghai\_zhang@hotmail.com

**Background:** Overuse with antibiotics in the treatment of infectious diseases has become a central focus of public health over the years. The aim of this study was to provide an up-to-date evaluation of the blood test-guided antibiotic use on patients with acute diarrhea in primary hospitals of China.

**Materials and methods:** A cross-sectional survey was conducted on 330 patients with acute diarrhea in Shanghai, People's Republic of China, from March 2013 to February 2016. These patients were treated with or without antibiotics based on the results of their blood tests, including examinations of C-reactive protein (CRP), white blood cells (WBC), and the percentage of neutrophils (Neu%). The infection types, which included bacterial, viral, and combination diarrhea, were determined by microbiological culture methods. Antibiotics used in non-bacterial diarrhea patients were considered misused and overused.

**Results:** There were significant overall differences in the clinical characteristics and blood tests between patients with diarrhea with a bacterial infection and patients with other types of infections. The patients were divided into four grading groups (0–3) according to the number of the positive results from three blood testes (CRP, WBC, and Neu%). The misuse rates of antibiotics in each group (0–3) were 81.3%, 71.1%, 72.4%, and 64.9%, respectively.

**Conclusion:** In this prospective study, the current diagnostic criteria (CRP, WBC, and Neu%) based on blood tests are not reliable in diagnosing bacterial diarrhea or guiding antibiotics use. To limit antibiotic overuse, a rapid and accurate differentiation of bacterial diarrhea from other types of diarrhea is pivotal.

**Keywords:** antibiotics overuse, acute diarrhea, primary hospitals

## Introduction

Acute diarrhea is defined as the presence of three or more abnormally loose or watery stools over a period of 24 h. The most common cause of acute diarrhea is infective microorganism, such as viruses, bacteria, fungi, or parasites.<sup>1</sup> Routine fecal cultures primarily for invasive bacterial pathogens are the major methods used to distinguish bacterial and viral infections in acute diarrhea diagnosis.<sup>2</sup> The well-known weaknesses of the stool culture method are that they are quite time-consuming, costly, and have low sensitivity. Moreover, trained technicians and appropriate equipment are required.<sup>3</sup> Therefore, in the primary hospitals in People's Republic of China, the most common tests used for the quick diagnosis and treatment guide of acute diarrhea are blood examinations.<sup>4,5</sup> The widely used diagnostic criteria include the number of white blood cells (WBC), the percentage of neutrophils (Neu%), and C-reactive protein (CRP).<sup>6</sup> Normally, the treatments will be selected directly depending on the results from these blood examinations.<sup>7</sup>

The principal components of the acute diarrhea treatments include fluid and electrolyte replacement, dietary modifications, and antibiotics therapy. Like the other infectious diseases, the irrational use of antibiotics for acute diarrhea treatment has become an outstanding problem all over the world, especially in low- and middle-income countries.<sup>8</sup> In People's Republic of China, for economical and historical reasons, antibiotics have been frequently overprescribed in clinical practice for patients with acute diarrhea.<sup>9,10</sup> Actually, the extreme overuse of antibiotics is a very serious and long-standing health issue in People's Republic of China.<sup>11</sup> More than 20 years ago, 88.9% of patients with acute diarrhea were prescribed antibiotics, with only 30.1% having clinically diagnosed bacillary dysentery.<sup>10</sup> The government has adopted approaches of both education and policy regulation to reduce the overuse of antibiotics.<sup>12,13</sup> Five years ago, antibiotics use in People's Republic of China was under control but still much higher than recommended.<sup>14,15</sup>

We aimed to study and update our knowledge about the blood examination-guided antibiotic use in emergency care of patients with acute diarrhea from the primary hospitals of People's Republic of China. Over 300 patients were divided into four groups based on their blood test results. We measured the antibiotic consumption and the misuse rate based on the stool culture examinations. Profound antibiotic overuse was discovered in patients with acute diarrhea, especially for patients who have tested negative in at least one blood examination.

## Materials and methods

### Study population

This study was carried out in 330 patients (317 adults who were defined as  $\geq 14$  years of age and 13 children) with acute diarrhea at Gongli Hospital and Jinyang Community Health Service Center, Pudong New Area, Shanghai, People's Republic of China, from March 2013 to February 2016. In this study, there was no immunosuppressed patients, no patients with sepsis or mortality associated with diarrhea and no population needed hospitalization. In this survey, acute diarrhea refers to acute infectious diarrhea. The symptoms lasted 2–24 h before first medical contact. Patients were considered to have acute diarrhea if they were passing at least three loose or watery stools in 24 h or passing at least one bloody stool (macroscopic observation revealed blood mixed with feces or inseparable from the stool), and with an illness duration of no longer than 14 days.

### Data collection

An anonymous, paper-based, descriptive questionnaire was used for collecting data. The questionnaire required

physicians to report the basic and clinical characteristics of patients (age, gender, etc.), as well as any tests and medications that they had ordered. All the patients with acute diarrhea during the study period, and who agreed to participate in the survey, were included. Four infectious disease specialists were participating in the study. They have been working with other hospital physicians to help direct care for infectious disease for 4–15 years.

### Blood tests

Blood samples for three tests (WBC count, Neu%, and CRP) were obtained on admission. WBC and differential counts were measured via the Hematology Analyzer (XS-1000, Sysmex, Kobe, Japan). WBC counts above  $10 \times 10^9/L$  represent above normal (positive). The Neu% was considered positive when  $>75\%$ . The CRP level was obtained by a Latex agglutination slide test (Humatec, Wiesbaden, Germany). CRP level above 10 mg/L was considered as elevated (positive) in our laboratory.

### Microbiological culture

Fecal samples were selected in the study by the routine enteric laboratory at the hospital. All samples included were diarrheal (adopting the shape of the container), had been stored at  $4^\circ C < 48$  h, and had sufficient fecal material to allow testing with all the assays. For the stool microbiological culture, a drop of a stool sample was resuspended in selective media for *Yersinia* (Sigma, St. Louis, MO, USA) and then inoculated on a Karmali agar plate (Oxoid, Basingstoke, Hampshire, UK) and incubated at  $37^\circ C$  in a microaerobic atmosphere. Colonies were observed daily until the third day after culture. The suspected colonies were confirmed based on motility, Gram staining, and oxidase activity. The bacterial species were identified by polymerase chain reaction (PCR) using species-specific PCR primers (Tables S1 and S2).<sup>16,17</sup> Cultures were considered negative if no bacterial or other pathogen growth was visible after 3 days of incubation. For detection of diarrhea virus, 200  $\mu L$  of supernatant was taken and used MagNA Pure LC Total Nucleic Acid isolation Kit to extract RNA by the Roche MagNA Pure LC2.0 system, and the total RNA was kept in  $-8^\circ C$  for storage. PCR methods were used to detect the target through One-Step real-time polymerase chain reaction kit (Qiagen, Hilden, Germany). The reaction condition is listed in Table S3.

### Ethics

This study was approved by the ethics committee at Gongli Hospital and Jinyang Community Health Service Center, Pudong New Area, Shanghai, People's Republic of China. Verbal informed consent was obtained from patients whose

medical data were included in the study. Participation was voluntary and no financial incentives were given. Personal identifying information was not collected from practitioners or participating patients.

## Data analysis

Data were analyzed using Statistical Package for the Social Sciences (SPSS) software (SPSS Inc., version 13.0; SPSS Inc., Chicago, IL, USA). Between-region differences were assessed for categorical variables with the Pearson's chi-square test. Continuous variables are expressed as mean  $\pm$  standard deviation (SD) with 95% confidence intervals; differences were considered statistically significant at  $P < 0.05$  or  $< 0.01$ .

## Results

### Basic characteristics of the patients with diarrhea

All patients were divided into four groups, that is, negative group (no infection), bacteria group (bacterial infection), virus group (viral infection), and combination (bacterial and viral infection). Characteristics of patients in different infection groups were examined and compared as shown in Table 1. No difference in gender, age, symptom (vomit and stomachache), or number of times diarrhea was observed in these four groups. There was a significant increase in WBC, Neu%, and number of fever cases in the bacterial group, compared to the negative group and virus group (Figure 1). Interestingly, the CRP level and the rate of fever symptom in the combination group are much lower than other groups. Together, these data indicate that patients in the bacterial group show significant signs of inflammation, compared with other patients.

### Blood tests of CRP, WBC, and Neu% in patients with diarrhea

To evaluate blood test-guided antibiotics use in the patients with diarrhea, the patients were stratified into four grading groups (from 0 to 3) based on the number of positive results (from 0 to 3) in the blood tests (CRP, WBC, and Neu%). As shown in Table 2, from grading group 0 to group 3, the percentage of no infection cases decreased from 66.3% to 35.1%, whereas the percentage of bacterial infection cases increased from 17.6% to 32.4%. Notably, the majority of these patients have either no infection or non-bacterial infection, which indicates that most of them are not qualified for antibiotics therapy. For example, even for patients with positive results from all blood tests (group 3), almost two of three of them (35.1% no infection and 29.7% viral infection) should not be treated with antibiotics. These data showed that the blood tests are not an accurate differentiation test for identifying bacterial infection in patients with diarrhea.

### Antibiotic use in patients with bacterial and non-bacterial diarrhea


To calculate the misuse rate of antibiotics, the negative group and virus group were combined into non-bacterial infection group, and bacterial group and combination group were combined into bacterial infection group in the following study. As shown in Table 3, the overall antibiotics use rates, as well as the use rates in each group, were calculated. The antibiotics use rate in non-bacterial infection groups was considered as a misuse of antibiotics. The overall misuse rate of antibiotics is 72.0%, and the misuse rate in individual grading groups (from 0 to 3) decreased from 81.3% to 64.9%. Even in group 3, the ratio of over-treated patients to properly treated patients (ratio of non-bacterial use to bacterial use) is nearly double (1.8).

**Table 1** Characteristics of patients in different infection groups

Variables	Infection groups (total n=330)				Overall P-value
	Negative (n=168)	Bacteria (n=68)	Virus (n=85)	Combination (n=9)	
Male, %	82, 48.8%	35, 51.5%	45, 52.9%	6, 66.7%	0.720
Age, years	43.88 $\pm$ 17.86	42.46 $\pm$ 17.26	45.13 $\pm$ 16.44	44.56 $\pm$ 24.28	0.828
Vomiting, %	38, 22.6%	17, 25.0%	32, 37.6%	3, 33.3%	0.079
Stomachache, %	82, 48.8%	37, 54.4%	40, 47.1%	4, 44.4%	0.806
Fever, %	22, 13.1%	23, 33.8%	15, 17.6%	0, 0.0%	0.001**
CRP (mg/L)	11.31 $\pm$ 24.41	12.70 $\pm$ 20.40	11.94 $\pm$ 14.24	5.97 $\pm$ 10.20	0.869
WBC ( $10^9/L$ )	8.84 $\pm$ 3.32	11.12 $\pm$ 4.44	8.98 $\pm$ 3.02	10.00 $\pm$ 3.86	0.000**
Neu%	73.57 $\pm$ 12.84	80.28 $\pm$ 8.78	76.76 $\pm$ 14.27	78.91 $\pm$ 12.65	0.002**
Symptom duration (h)	17 $\pm$ 8	13 $\pm$ 8	15 $\pm$ 7	15 $\pm$ 8	0.124
Diarrhea (number of times)	6.21 $\pm$ 4.07	5.99 $\pm$ 2.50	6.81 $\pm$ 3.84	6.00 $\pm$ 2.55	0.519

**Notes:** Infection groups were classified by microbiological culture examination. Combination group includes patients with both bacteria and virus detected. P, Pearson's chi-square test. Data presented as n % or mean  $\pm$  standard deviation. \*\* $P < 0.01$ .

**Abbreviations:** CRP, C-reactive protein; Neu%, percentage of neutrophils; WBC, white blood cells.



**Figure 1** Baseline laboratory results for the study cohort. **Notes:** (A) The percentage of neutrophils in blood work tests of each group. (B) The number of WBC in each group. (C) The value of CRP in each group. Infection groups were classified by microbiological culture examination. Combination group means both bacteria and virus were detected. \*\*P<0.01. **Abbreviations:** CRP, C-reactive protein; Neu%, percentage of neutrophils; WBC, white blood cells.

Hence, we concluded that this retrospective study showed a profound misuse of antibiotics in the patients with acute diarrhea, which was directly guided by the results from the blood tests of WBC, Neu%, and CRP.

### Discussion

In this retrospective cohort study, we observed a profound misuse of antibiotics in patients with acute diarrhea who were treated based on blood testing (CRP, WBC, and Neu%). Although the results of the blood tests displayed a significant difference between patients with bacterial diarrhea and other groups of patients, these tests failed to prove their accuracy in differentiating between bacterial diarrhea and non-bacterial diarrhea.

As a common symptom faced by emergency doctors in People’s Republic of China, acute diarrhea and the intestinal secretory processes can be activated by infection from bacteria,

viruses, or other pathogens. It was reported that the most common enteral infectious agents in People’s Republic of China from 2010 to 2015 include 7 bacteria, 5 viruses, and 3 protozoa. Diarrheagenic *Escherichia coli* was identified as the most predominant pathogen in diarrhea cases.<sup>18,19</sup> In our study, the most frequent causes of infectious diarrhea were virus (*Vibrio*) and bacteria (*Salmonella*). In consistent with our findings, it seems that the epidemiology change during the several past years is virus becoming a more frequent cause than bacteria in Chinese as reported by citywide hospitals in big cities. For different infections, there were different treatments and the doctors in the primary hospitals and institutes determine the chance of bacterial infection using three criteria: elevated Neu%, WBC count, and CRP level. Doctors are more likely to prescribe antibiotics if more of these criteria

**Table 2** Blood test results in different infection groups

Grading group	N (%) in each infection group				Total
	Negative	Bacteria	Virus	Combination	
0	69 (66.3)	12 (11.5)	20 (19.2)	3 (2.9)	104
1	40 (42.6)	20 (21.3)	31 (33.0)	3 (3.2)	94
2	46 (48.4)	24 (25.3)	23 (24.2)	2 (2.1)	95
3	13 (35.1)	12 (32.4)	11 (29.7)	1 (2.7)	37

**Note:** Grading groups were classified by the total number of positive results in tests of CPR, WBC, and Neu%.

**Abbreviations:** CRP, C-reactive protein; Neu%, percentage of neutrophils; WBC, white blood cells.

**Table 3** Antibiotic use rate in each infection group

Grading group	N (%) of antibiotic use			Ratio (non-bacterial use/bacterial use)
	Non-bacterial infection (overuse)	Bacterial infection (proper use)	Total	
0	26 (81.3)	6 (18.8)	32	4.3
1	54 (71.1)	22 (28.9)	76	2.5
2	63 (72.4)	24 (27.6)	87	2.6
3	24 (64.9)	13 (35.1)	37	1.8
Overall	167 (72.0)	65 (28.0)	232	2.6

**Notes:** Non-bacterial infection includes negative and virus groups; bacterial infection includes bacteria and combination groups. Antibiotic use in non-bacterial infection group is considered as misuse of antibiotics.

are met as they are directly related to bacterial infection. However, in our current findings, these indicators showed low accuracy in identifying bacterial infectious diarrhea.

The Neu% and WBC count are two criteria used to diagnose bacterial infection. Neutrophils are responsible for the most rapid immune response and are the immune system's first line of defense.<sup>20</sup> Upon bacterial infection, pathogen-associated molecular patterns of the bacteria, such as lipopolysaccharides and poly (I:C), may directly stimulate the infected tissue or local immune cells through toll-like receptors.<sup>21</sup> This strong inflammatory stimulation would directly induce the production of cytokines and chemokines, among them are several CXC chemokines.<sup>22–24</sup> These two criteria seem to be specific for bacterial infection. However, recent discoveries showed that viral infection can also induce elevated peripheral neutrophil count and percentage through similar mobilization mechanisms. The neutrophils are also recruited to the virus-infected tissue.<sup>25–27</sup> Thus, analyzing the neutrophil percentage and WBC count cannot clearly distinguish bacterial infection from viral infection. Moreover, the elevated WBC count is dependent on neutrophil mobilization. This makes these two criteria interdependent and should not be evaluated as two separated criteria.

Another criterion the Chinese clinical practitioners use to diagnose bacterial infection is the elevation of CRP. CRP is a typical acute-phase protein. CRP can directly bind to and opsonize bacteria. It also orchestrates the immune response.<sup>28</sup> Importantly, many studies showed that extremely high CRP is related to acute bacterial infection but not viral infection. This makes it a good candidate for a criterion that can distinguish between bacterial and viral infection.<sup>29–31</sup> However, in our study, we did not find an extremely high average CRP concentration in the bacterial diarrhea infections (Figure 1). This disappointing discovery may be due to the dynamic pattern of CRP expression. As we know, CRP was the acute-phase protein. Plasma CRP is produced by hepatocytes and study showed that in healthy adults the median concentration of CRP was 0.8 mg/L and the 99th centile is 10 mg/L and after an acute-phase stimulus, CRP values may increase to 10,000-fold.<sup>32</sup> We noticed that in acute bacterial diarrhea, drastically elevated CRP concentration can only be found within 20 h after the infection (data not shown), while in our clinical practice the blood testing is often conducted outside of this timeframe. This may explain that why CRP elevation may not be found in most of patients with bacterial diarrhea. Due to this fact, a moderate CRP concentration is an unreliable criterion for differentiating between viral diarrhea and bacterial diarrhea. Another cause that leads to misdiagnose is

when the patients have both bacterial and viral infections, and the CRP level and other physical symptom are much lower, instead of higher, as we observed in this study. Actually, in most of the combined infection patients, the viral and bacterial infections usually occur simultaneously,<sup>33</sup> which make the typical features of the bacterial infection disappear.

In brief, the imprecise nature of these diagnostic criteria may have led to many misdiagnoses of bacterial diarrhea. This is reflected in our data, which shows that an extremely high percentage of antibiotics overuse occurred because the doctors relied solely on these tests. This is not the only contributing factor to antibiotics overuse, however.

Previous studies showed that Chinese health literacy has been low and antibiotics have been considered a stronger, faster, and higher quality medicine for treatment.<sup>34</sup> Pressure from patients and the financial concerns are two of major causes that conspire to encourage the dangerous overuse of antibiotics.<sup>8</sup> The antimicrobial resistance of antibiotics is commonly increasing in People's Republic of China, whereas the rate varies on different infections. For instance, studies indicated that the antimicrobial resistance of levofloxacin is 53.4% in *Shigella* infections.<sup>35</sup> In the *Campylobacter jejuni* infection, the resistance to cefuroxime was increased to 93% in 2010 while it was 0% in the 1990s.<sup>36</sup> Actually, the preventive usage of antibiotics is beneficial in immunocompromised patients,<sup>37,38</sup> which only counts for a small amount of patients with infectious diseases. For instance, only ~5%–8% of acute salmonellosis is in general population.<sup>39</sup> It is suggested that most patients should be prescribed with antibiotics only after confirmation of bacterial infection. However, confirmation of bacterial infection with culture analysis usually takes days, which is an unrealistic amount of time to wait to provide a treatment. Thus, Chinese medical practitioners utilized an inexact method to prescribe antibiotics, taking into account only the patients' symptoms and quick lab tests. When information about antibiotics became available to the general public in People's Republic of China, the elderly and parents of small children with limited health insurance preferred the use of antibiotics treatment because it was an inexpensive option. In addition, monetary incentives also contribute to clinicians prescribing antibiotics excessively. This is especially true of primary care clinicians as monetary incentives from drug prescriptions are a major source of income for them.<sup>34</sup>

In summary, the higher rate of antibiotic prescription for patients with acute diarrhea by primary care clinicians may be explained by a combination of inaccurate criteria for diagnosis, lack of sufficient training, and stronger economic incentives. Practitioners should be cautious about



overprescribing antibiotics and rely on other testing methods for treatment. Medical professionals, as well as government personnel working in the health sector, need to update their diagnostic and treatment guidelines based on the latest scientific evidence. The remedy of this situation requires that educational awareness programs be conducted on a regular basis to educate doctors and patients about the dangers of overuse and abuse of antibiotics.

Overall, we found that using these three blood tests to diagnose bacterial diarrhea and guide antibiotics treatment is not reliable. More accurate and efficient diagnostic criteria for identifying bacterial diarrhea are needed in People's Republic of China to reduce antibiotic overuse and improve the efficiency of the clinical management of diarrhea.

## Acknowledgments

This study was supported by Key Discipline Construction Project of Pudong Health Bureau of Shanghai (No: PWZx2014-03) and Shanghai Municipal Commission of Health and Family Planning Fund (No: 20134396). We thank SinoScript for proofreading this manuscript.

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Farthing M, Salam MA, Lindberg G, et al. Acute diarrhea in adults and children: a global perspective. *J Clin Gastroenterol*. 2013;47(1):12–20.
- Pawlowski SW, Warren CA, Guerrant R. Diagnosis and treatment of acute or persistent diarrhea. *Gastroenterology*. 2009;136(6):1874–1886.
- Guerrant RL, Shields DS, Thorson SM, Schorling JB, Groschel DH. Evaluation and diagnosis of acute infectious diarrhea. *Am J Med*. 1985;78(6B):91–98.
- Wang D, Fang L, Xiao S. Porcine epidemic diarrhea in China. *Virus Res*. 2016;226:7–13.
- Zhu M, Zhang J, Gao Y, Wang Z. [Present research on disease burden of diarrhea illness]. *Wei Sheng Yan Jiu*. 2008;37(1):126–128. Chinese.
- Gonzalez MD, Wilen CB, Burnham CA. Markers of intestinal inflammation for the diagnosis of infectious gastroenteritis. *Clin Lab Med*. 2015;35(2):333–344.
- Radlovic N, Lekovic Z, Vuletic B, Radlovic V, Simic D. [Acute diarrhea in children]. *Srp Arh Celok Lek*. 2015;143(11–12):755–762. Slovak.
- Li Y. China's misuse of antibiotics should be curbed. *BMJ*. 2014;348:g1083.
- Hou FQ, Wang Y, Li J, Wang GQ, Liu Y. Management of acute diarrhea in adults in China: a cross-sectional survey. *BMC Public Health*. 2013;13:41.
- Ke B, Ran L, Wu S, et al. Survey of physician diagnostic and treatment practices for patients with acute diarrhea in Guangdong province, China. *Foodborne Pathog Dis*. 2012;9(1):47–53.
- Mao W, Vu H, Xie Z, Chen W, Tang S. Systematic review on irrational use of medicines in China and Vietnam. *PLoS One*. 2015;10(3):e0117710.
- Xiao Y, Zhang J, Zheng B, Zhao L, Li S, Li L. Changes in Chinese policies to promote the rational use of antibiotics. *PLoS Med*. 2013;10(11):e1001556.
- Zhang Y, Harvey K. Rational antibiotic use in China: lessons learnt through introducing surgeons to Australian guidelines. *Australia New Zealand Health Policy*. 2006;3:5.
- Dong L, Yan H, Wang D. Antibiotic prescribing patterns in village health clinics across 10 provinces of Western China. *J Antimicrob Chemother*. 2008;62(2):410–415.
- Dong L, Yan H, Wang D. Drug prescribing indicators in village health clinics across 10 provinces of Western China. *Fam Pract*. 2011;28(1):63–67.
- Yu XF, Pan JC, Meng DM, Wang HQ, Zhang W, Zheng W. [Multiplex real-time PCR detecting *Salmonella*, *Shigella* and diarrheagenic *Escherichia coli*]. *Zhonghua yu fang yi xue za zhi*. 2007;41(6):461–465. Chinese.
- Wang J, Xu Z, Niu P, et al. A two-tube multiplex reverse transcription PCR assay for simultaneous detection of viral and bacterial pathogens of infectious diarrhea. *Biomed Res Int*. 2014;2014:648520.
- Zhang SX, Yang CL, Gu WP, et al. Case-control study of diarrheal disease etiology in individuals over 5 years in southwest China. *Gut Pathog*. 2016;8:58.
- Wei ZQ, Chang HL, Li YF, Xu XB, Zeng M. [Clinical epidemiology and antimicrobial resistance of nontyphoidal *Salmonella* enteric infections in children: 2012–2014]. *Zhonghua Er Ke Za Zhi*. 2016;54(7):489–495. Chinese.
- Kolaczowska E, Kubes P. Neutrophil recruitment and function in health and inflammation. *Nat Rev Immunol*. 2013;13(3):159–175.
- Zheng J, Meng J, Zhao S, Singh R, Song W. *Campylobacter*-induced interleukin-8 secretion in polarized human intestinal epithelial cells requires *Campylobacter*-secreted cytolethal distending toxin- and toll-like receptor-mediated activation of NF- $\kappa$ B. *Infect Immun*. 2008;76(10):4498–4508.
- Amulic B, Cazalet C, Hayes GL, Metzler KD, Zychlinsky A. Neutrophil function: from mechanisms to disease. *Annu Rev Immunol*. 2012;30:459–489.
- Eash KJ, Greenbaum AM, Gopalan PK, Link DC. CXCR2 and CXCR4 antagonistically regulate neutrophil trafficking from murine bone marrow. *J Clin Invest*. 2010;120(7):2423–2431.
- Chang B, Xu MJ, Zhou Z, et al. Short- or long-term high-fat diet feeding plus acute ethanol binge synergistically induce acute liver injury in mice: an important role for CXCL1. *Hepatology*. 2015;62(4):1070–1085.
- Galani IE, Andreaskos E. Neutrophils in viral infections: current concepts and caveats. *J Leukoc Biol*. 2015;98(4):557–564.
- Drescher B, Bai F. Neutrophil in viral infections, friend or foe? *Virus Res*. 2013;171(1):1–7.
- Jenne CN, Wong CH, Zemp FJ, et al. Neutrophils recruited to sites of infection protect from virus challenge by releasing neutrophil extracellular traps. *Cell Host Microbe*. 2013;13(2):169–180.
- Zhou Z, Xu MJ, Gao B. Hepatocytes: a key cell type for innate immunity. *Cell Mol Immunol*. 2016;13(3):301–315.
- Sasaki K, Fujita I, Hamasaki Y, Miyazaki S. Differentiating between bacterial and viral infection by measuring both C-reactive protein and 2'-5'-oligoadenylate synthetase as inflammatory markers. *J Infect Chemother*. 2002;8(1):76–80.
- Haran JP, Beaudoin FL, Suner S, Lu S. C-reactive protein as predictor of bacterial infection among patients with an influenza-like illness. *Am J Emerg Med*. 2013;31(1):137–144.
- Korppi M, Kroger L. C-reactive protein in viral and bacterial respiratory infection in children. *Scand J Infect Dis*. 1993;25(2):207–213.
- Pepys MB, Hirschfield GM. C-reactive protein: a critical update. *J Clin Invest*. 2003;111(12):1805–1812.
- DuPont HL. Persistent diarrhea: a clinical review. *JAMA*. 2016;315(24):2712–2723.
- Reynolds L, McKee M. Factors influencing antibiotic prescribing in China: an exploratory analysis. *Health Policy*. 2009;90(1):32–36.
- Zhang CL, Liu QZ, Wang J, Chu X, Shen LM, Guo YY. Epidemic and virulence characteristic of *Shigella* spp. with extended-spectrum cephalosporin resistance in Xiaoshan district, Hangzhou, China. *BMC Infect Dis*. 2014;14:260.

36. Hou FQ, Sun XT, Wang GQ. Clinical manifestations of *Campylobacter jejuni* infection in adolescents and adults, and change in antibiotic resistance of the pathogen over the past 16 years. *Scand J Infect Dis*. 2012; 44(6):439–443.
37. Kuruvilla M, de la Morena MT. Antibiotic prophylaxis in primary immune deficiency disorders. *J Allergy Clin Immunol Pract*. 2013;1(6): 573–582.
38. Miao R, Huang S, Zhou Z, et al. Targeted disruption of MCP1/ Zc3h12a results in fatal inflammatory disease. *Immunol Cell Biol*. 2013;91(5):368–376.
39. Onwuezobe IA, Oshun PO, Odigwe CC. Antimicrobials for treating symptomatic non-typhoidal *Salmonella* infection. *Cochrane Database Syst Rev*. 2012;11:CD001167.

## Supplementary materials

**Table S1** Primer sequences for bacteria identification

Gene name (bacteria type)	Primers (5'-3')	Production length (bp)
<i>invA</i> ( <i>Salmonella</i> )	F: AGG TGA CAC TAT AGA ATA R: GTA CGA CTC ACT ATA GGG A	323
<i>ipaH</i> ( <i>Shigella</i> and <i>EIEC</i> )	F: GTT CCT TGA CCG CCT TTC CGA TAC CGT C R: GCC GGT CAG CCA CCC TCT GAG AGT AC	619
<i>stx1</i> ( <i>EHEC</i> )	F: AAA TCG CCA TTC GTT GAC TAC TTC T R: TGC CAT TCT GGC AAC TCG CGA TGC A	370
<i>stx2</i> ( <i>EHEC</i> )	F: CAG TCG TCA CTC ACT GGT TTC ATC AC R: GGA TAT TCT CCC CAC TCT GAC ACC	283
<i>eaeA</i> ( <i>EHEC</i> and <i>EPEC</i> )	F: TCA ATG CAG TTC CGT TAT CAG TT R: GTA AAG TCC GTT ACC CCA ACC TG	482
<i>aggR</i> ( <i>EAEC</i> )	F: CTA TAC ACA AAA GAA CCA AGC R: ACA GAA TCG TCA GCA TCA GC	254
<i>lt</i> ( <i>ETEC</i> )	F: TCT CTA TGT GCA TAC GGA GC R: CCA TAC TGA TTG CCG CAA	322
<i>mapA</i> ( <i>Campylobacter jejuni</i> )	F: AGG TGA CAC TAT AGA ATA R: GTA CGA CTC ACT ATA GGG A	627

**Table S2** Primer sequences for virus identification

Virus type	Primers and probes	Sequence
<i>Rotavirus</i>	NVP3-F	ACCATCTTCACRTRACCCTC
	NVP3-R	GGTCACATAACGCCCC
	NVP3-Probe	ATGAGCACAATAGTTAAAAGCTAACACTGTCAA
<i>Adenovirus</i>	Adeno-F	GCCACGGTGGGGTTTCTAAACTT
	Adeno-R	GCCCCAGTGGTCTTACATGCACATC
	Adeno-Probe	TGCACCAGACCCGGGCTCAGGTACTCCGA
<i>Astrovirus</i>	AV1	CCGAGTAGGATCGAGGGT
	AV2	GCTTCTGATTAATCAATTTTAA
	Avs	CTTTTCTGTCTCTGTTTAGATTATTTTAATCACC
<i>Sapovirus</i>	SV1	GCTGTTTCTACTGGTGCA
	SV2	GGCATCCTGTCTTTCCAAGCA
	SVP	CCAATCTAATGTCCCTGAGGCAATACGTAA
<i>Norovirus</i>	COG1F	CGTTGGATGCGCTTTTCATGA
	COG1R	CTTAGACGCCATCATCATTTAC
	Probe	
<i>Vibrio</i>	RING1-TP	AGATTGCGATCTCCTGTCCA
	TLHF	AGGTGACACTATAGAATA
	TLHR	GTACGACTCACTATAGGGA
	Probe	



**Table S3** The polymerase chain reaction condition of diarrhea pathogens

Pathogen type	Temperature (°C)	Time	Cycles
Bacteria and <i>Vibrio</i>	50	30 min	
	95	15 min	
	90	30 s	10
	56	30 s	
	72	60 s	
	90	30 s	10
	68	30 s	
	72	60 s	
	90	30 s	20
	50	30 s	
	72	60 s	
	72	3 min	
<i>Rotavirus</i>	50	30 min	45
	95	15 min	
	95	15 s	
	60	1 min	
	50	45 min	
<i>Astrovirus</i>	95	15 min	45
	94	15 s	
	55	1 min	
<i>Sapovirus</i>	50	30 min	40
	95	15 min	
	94	15 s	
	55	34 s	
<i>Norovirus</i>	50	30 min	45
	95	15 min	
	95	10 s	
	60	1 min	
<i>Adenovirus</i>	95	15 min	45
	95	15 s	
	60	1 min	

### Patient Preference and Adherence

Dovepress

### Publish your work in this journal

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focuses on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize

clinical outcomes for existing disease states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/patient-preference-and-adherence-journal>