

# Impact of disability on different domains of health-related quality of life in the noninstitutionalized general population

Javier Mar<sup>1,2</sup>  
Isabel Larrañaga<sup>3</sup>  
Arantzazu Arrospide<sup>1</sup>  
José María Begiristain<sup>3</sup>

<sup>1</sup>Research Unit, Hospital Alto Deba, Navarra 16, Mondragón, Spain;

<sup>2</sup>Clinical Management Unit, Hospital Alto Deba, Navarra 16, Mondragón, Spain; <sup>3</sup>Gipuzkoa Health Authority, Basque Government, Sancho El Sabio 35, Donostia-San Sebastián, Spain

**Abstract:** The relationship between disability and health related quality of life (HRQL) is complex because of the role that function plays in the measurement, and certain points need to be dealt with in greater detail when the analysis is applied to the different dimensions of HRQL. The purpose of this study was to assess the impact of disability on different domains of quality of life. Variables were drawn from the 2002 Basque Country Health Survey. Logistic regression models were constructed to measure the adjusted effect of disability on quality of life as determined by the SF-36 questionnaire. The models were adjusted for sociodemographic variables and the presence of comorbidity. The greatest difference between disabled and nondisabled subjects was found in the physical components of the questionnaire. Odds Ratios (ORs) for a suboptimal level of quality of life as determined by functional status were much higher for the physical than for the mental dimensions. Adjustment for different variables showed a decrease in ORs in the physical, mental, and social dimensions. This study establishes a significant relationship between disability and the loss of quality of life in both physical and mental dimensions of SF-36, irrespective of the associated disease.

**Keywords:** quality of life, disability, health status, SF-36

## Introduction

The impact of disease in a population has traditionally been measured by its effect on the mortality rate and thus on the decrease in life expectancy. This approach has been adequate in an epidemiologic context characterized by acute diseases with high mortality rates.<sup>1</sup> However, the determinants of health in developed countries, such as population aging, have changed, resulting in an increased prevalence of diseases causing disability. The list of diseases according to their burden is evidence of this new scenario,<sup>2,3</sup> and stroke and dementia are among the leading conditions because of their great impact in terms of disability.<sup>4</sup> However, society and the medical community still have not made its care a social priority, and research for those diseases that generate disability does not receive funds according to the burden imposed.<sup>5</sup> The lack of knowledge often fosters the tacit assumption that there is no problem, especially when compared to the information available about life lost due to premature mortality.<sup>4</sup>

Therefore, the evaluation of diseases, besides mortality, should include measurement of long-term consequences such as disability and, at a broader level, health related quality of life (HRQL). The term “HRQL” refers to the physical, psychological, and social domains of health, which are distinct areas that are influenced by a person’s experiences, beliefs, expectations, and perceptions.<sup>6</sup> The World Health Organization defines ‘disability’ as a restriction or lack of ability to perform an activity in the manner

Correspondence: Javier Mar  
Unidad de Gestión Sanitaria, Hospital Alto Deba, Avda Navarra, 20500, Mondragón, Gipuzkoa, Spain  
Tel +34 943 035 300  
Fax +34 943 035 314  
Email javier.marmedina@osakidetza.net

or within the range considered normal for a human being.<sup>7</sup> This term does not include the subjective perceptions that determine the HRQL of a person. In the field of outcome research both concepts are used separately, and impairments appear to be better reflected in disability measures than in HRQL instruments.<sup>8,9</sup> Nevertheless, the differentiation of both concepts is complex because of the role played by function in the measurement of health and disability.<sup>9</sup> Some authors have noted that new measures of perceived health status are needed to avoid the confusion between function and health.<sup>9</sup> This approach needs to deal with certain points in greater detail when the analysis is applied to the different dimensions of HRQL. Physical scales are scored according to the performance of activities of daily living, but items that produce mental and social scales are less closely related to the functional status.<sup>10,11</sup> The new framework for the analysis of functioning, disability, and health proposed by the World Health Organization differentiates the concepts of disability and health and assumes that a disabled can be a healthy person.<sup>7</sup>

Given the parallel use of the two concepts, it would be important to ascertain the relationship of disability separately within the physical, mental, and social domains of HRQL. From this perspective we aimed to check the theoretical model by assessing the influence of disability in the different domains of HRQL in the noninstitutionalized general population of the Basque Country.

## Methods

Data from a noninstitutionalized population were used to conduct a descriptive study of the HRQL as perceived by a disabled population subset, versus the HRQL perceived by a nondisabled subset. The disabled population was defined as subjects with any disability that resulted from a process of gradual change from diseases and lesions first affecting organ function and ultimately the individual. Disability may thus be defined as the inability to independently perform activities of daily living within a given environment.<sup>12</sup> Disabled people depend on their families or the community to carry out basic or instrumental activities of daily living. Questionnaires that evaluate the level of autonomy in different daily tasks estimate the degree of disability.<sup>13</sup>

## Sample (Basque Country Health Survey)

The 2002 Basque Country Health Survey (BCHS) served as the information source. This survey is conducted every 5 years to collect information about the health and social characteristics of the noninstitutionalized population of

the Basque Country. The surveyed sample consisted of 8415 subjects representative of the Basque Country population. The authors of the survey have provided a more detailed description of their methods elsewhere.<sup>14</sup>

## Variables and measurements

Items related to the limitation of activities of daily living, quality of life (SF-36 questionnaire), and presence of disease were drawn from the BCHS. Social and demographic variables were also processed: sex, age, marital status, educational level, socioeconomic group, confidential support, and affective support. The presence of disease was recorded based on the subjects' self-report, and it was coded according to the International Classification of Diseases (ICD-9-MC). For the analysis, the 9 most relevant disease groups were selected: stroke; ischemic heart disease; cancer; diabetes; respiratory, other circulatory, musculoskeletal, neurological diseases and lesions; and trauma. These groups were included in our analysis because the presence of chronic conditions plays an important role in the relationship between functional status and HRQL.<sup>15,16</sup>

The functional capacity for activities of daily living was assessed by recording 10 basic activities: eating, bathing, using the toilet, getting out of bed, rising from a chair and walking, adequate nutrition, washing, dressing, putting shoes on, and spending the night alone. Individuals were asked if they were able to carry out each activity. Ten points were assigned for complete ability, 5 points for partial autonomy for each activity, and 0 when they were totally unable to carry out the activity. Partial autonomy was that when provided with the choice, the individual would prefer to perform the task with help. The resulting total value ranged from 0 to 100, and subjects with a value of 95 or less were defined as disabled. The same criterion has been used in other studies and means that a person is not capable, totally or partially, to accomplish some activity. Individuals with a score of 100 were defined as nondisabled.<sup>17,18</sup>

The SF-36 Health Survey (SF-36), a standardized questionnaire consisting of 36 items that assess subjects' health, is one of the tools most commonly used internationally to measure HRQL.<sup>10</sup> By running a clustering algorithm, eight scales (dimensions) with values ranging from 0 (minimum health perceived) to 100 (maximum health perceived) are obtained. Two summary variables, a physical component summary (PCS) and a mental component summary (MCS), are calculated from these dimensions. The summary variables are standardized to have a mean of 50 and a standard deviation of 10 in the general population.

## Analysis

The first step in our study was calculating the SF-36 variables (eight dimensions and two summary variables) in disabled subjects and in the nondisabled population. The extent of the difference between disabled and nondisabled was measured by the effect size (the difference between the means divided by the standard deviation from the nondisabled sample).<sup>19,20</sup> Cohen's benchmarks were used to classify the magnitude of effect sizes [25]: below 0.20 was not significant; between 0.20 and 0.50 was small; between 0.50 and 0.80 was moderate; and above 0.80 large. The association between functional status and the presence of different diseases was measured by calculating the crude odds ratios (OR) of the disease declaration in disabled versus nondisabled subjects.<sup>21</sup>

Given the disparity in sex, age, social support variables, and presence of chronic conditions in the nondisabled and disabled populations, we decided to apply a statistical approach to adjust the OR values.<sup>15,22,23</sup> We analyzed the differences in HRQL between disabled and nondisabled subjects by logistic regression models, because linear regression modeling presented a problem of lack of linearity and normality in the scores of SF-36 dimensions that we planned to use as dependent variables.<sup>24</sup> Guallar-Castillón et al used the suboptimal health index to analyze gender differences by recoding the value of each SF-36 dimension, depending on whether it was below the maximum value of 100.<sup>23</sup> In this study, the same method was applied and the mean of the general population was used as a threshold for good quality of life (values below the mean were recorded as suboptimal HRQL and values over the mean as optimal HRQL). Logistic regression models were used to calculate the ORs of being in suboptimal health for the different SF-36 dimensions, and summaries in subjects with disability versus subjects without disability. Crude ORs were calculated first, followed by adjusted ORs. The first model was adjusted for sociodemographic variables (age, sex, and education). In addition to age, sex, and education, the second OR took into account the presence of comorbidity variables (stroke; ischemic heart disease; cancer; diabetes; lesions; trauma; and respiratory, circulatory, musculoskeletal, and neurological diseases). The process of sequential calculation of ORs allowed for assessing the relative contribution of the different groups of variables to the differences between nondisabled and disabled subjects in the different scales of SF-36. We calculated the Nagelkerke  $R^2$  to quantify the contribution of the groups of variables to the explanation of the variance of the dependent variable. Nagelkerke  $R^2$  is a corrected version of the Cox  $R^2$  that covers the range from 0 to 1.

## Results

Table 1 shows the social and demographic characteristics of the sample by level of autonomy. There were more disabled subjects among women, elderly people, widowed people, those in low socioeconomic groups, and those lacking education. When the subjects were classified by the number of declared diseases, the proportion of disabled subjects was seen to increase with the number of diseases. Out of a sample of 8415 people, 303 subjects met the disability criterion, that is, 3.6% of the noninstitutionalized general population said they needed help for at least one basic activity of daily living (Table 1). Table 2 shows the presence of diseases as reported by the subjects themselves, stratified by autonomy level. Stroke and mental disorders exhibited the highest ORs associated with disability (18.54 and 15.70).

Scores and effect sizes in different SF-36 dimensions, with physical and mental summaries, and SF-6D in the nondisabled

**Table 1** Sociodemographic characteristics of the population by functional status (nondisabled/disabled)

	Nondisabled		Disabled	
	N	%	N	%
Total	8112	96.4	303	3.6
<b>Sex***</b>				
Males	3824	47.1	112	37.0
Females	4288	52.9	191	63.0
<b>Age***</b>				
0–24	1155	14.2	3	1.0
25–44	2555	31.5	22	7.3
45–64	2673	33.0	63	20.8
65–69	597	7.4	31	10.2
70 y+	1132	14.0	184	60.7
<b>Marital status***</b>				
Single	2663	32.8	47	15.5
Married or with couple	4674	57.6	137	45.2
Separated, divorced	178	2.2	5	1.7
Widow/er	597	7.4	114	37.6
<b>Education***</b>				
No schooling	299	3.7	55	18.2
Primary education	5877	72.4	228	75.2
Graduate education	1936	23.9	20	6.6
<b>Socioeconomic group*</b>				
I–II	1878	23.2	52	17.2
III	1243	15.3	48	15.8
IV–V	4991	61.5	203	67.0
<b>Number of diseases***</b>				
0	4315	53.2	29	9.6
1	1966	24.2	64	21.1
2	974	12.0	66	21.8
3	441	5.4	57	18.8
4	220	2.7	33	10.9
5	120	1.5	22	7.3
6	76	0.9	32	10.6

Notes: \* $P < 0.05$ ; \*\*\* $P < 0.0001$ .

**Table 2** Presence of chronic diseases by functional status and odds ratios of disease in disabled versus nondisabled

Disease	Nondisabled N = 8112	Disabled N = 303	Total 8415	OR (95% CI)
Stroke	66	40	106	18.54 (12.29–27.98)
Ischemic heart disease	333	65	398	6.38 (4.75–8.57)
Cancer	55	18	73	9.25 (5.36–15.96)
Diabetes	261	53	314	6.38 (4.63–8.79)
Respiratory diseases	607	27	634	1.21 (0.81–1.81)
Other circulatory diseases	1300	128	1428	3.83 (3.03–4.85)
Musculoskeletal diseases	1286	155	1441	5.56 (4.40–7.02)
Neurological diseases	315	69	384	7.30 (5.46–9.77)
Mental problems	495	153	648	15.70 (12.31–20.01)
Other diseases	2238	197	2435	4.88 (3.83–6.21)

**Abbreviations:** OR, odds ratio; CI, confidence intervals.

and disabled populations, are given in Table 3. The greatest difference between disabled and nondisabled subjects was found in the physical components of the questionnaire.

The second and third columns of Table 4 show the percentage of subjects in each sample (nondisabled and disabled) that scored lower than the mean of the general population in each domain. ORs by functional status meant how many times higher the probability was of a disabled individual being in a suboptimal level of quality of life compared with nondisabled, and this was much higher for the physical than for the mental dimensions. Adjustment for different variables showed a decrease in ORs in all cases. Nevertheless, ORs continued to be very high for physical functioning (15.6) and PCS (13.2). The MCS showed the lowest OR (1.5), and its difference from the crude OR (1.7) was small. Adjusted ORs for the other physical dimensions

were approximately 5, whereas ORs for mental dimensions were between 2 and 3. The percentages of change in the ORs produced by both models were relevant. The combined effect of the sociodemographic variables (sex, age, and education) had a negative effect in the dimension role-emotional and MCS. Table 4 also shows the percentage of explanation of the variance of the probability of being in suboptimal health by the two models. These different results by dimension receive the same comments as the figures of ORs.

## Discussion

Our understanding of the relationship between health and disability changed after the publication in 2001 of the International Classification of Function, Disability, and Health (ICF) that introduced the idea that a disabled person can live in a healthy condition.<sup>7</sup> However, in this sample of the noninstitutionalized general population, disability had a significant impact on subjects' HRQL, which is seen as a decrease in the eight dimensions of SF-36, irrespective of the associated disease. The main finding is that the effect of disability persisted in both physical and mental scales, despite adjustment for sociodemographic and social support variables, and the presence of diseases. As ICF notes, contextual factors play a key role in the manifestation of chronic conditions.<sup>7</sup> Environmental and personal factors determine how disabled people cope with their limitation in the performance of daily-living activities. In the Basque Country, the family has played a predominant role as the main support network to cover the needs of disabled people. However, this situation is changing and future health surveys in the Basque Country will show if the perceived health of disabled Basque people worsens. Therefore, a traditional strength that could be defined as the psychological resilience to disability is probably going to disappear. Resilience is defined as a dynamic process where individuals exhibit positive behavioral adaptation when they

**Table 3** Scores in the SF-36 dimensions of the nondisabled population (N = 8112) and the disabled population (N = 303), score differences, and effect sizes

	Nondisabled mean (SD)	Disabled mean (SD)	Difference	Effect size
Physical functioning	88.8 (17.7)	36.0 (30.5)	52.8*	3.0
Role-physical	88.1 (29.0)	39.5 (45.5)	48.6*	1.7
Pain	78.5 (24.3)	45.7 (31.4)	32.8*	1.3
Perceived health	66.7 (18.3)	37.7 (21.6)	29.0*	1.6
Vitality	66.3 (19.1)	42.9 (23.6)	23.3*	1.2
Social functioning	89.9 (18.0)	57.5 (33.5)	32.4*	1.8
Role-emotional	92.0 (24.2)	72.9 (43.3)	19.1*	0.8
Mental health	73.3 (17.2)	57.6 (23.0)	15.8*	0.9
Physical component summary	50.2 (9.2)	26.8 (12.7)	23.5*	2.5
Mental component summary	50.1 (9.8)	45.9 (15.7)	4.2*	0.4

**Notes:** \*Significance level of the difference <0.001; SD, standard difference.

**Table 4** Percent of population with suboptimal health (worse than mean health of the population) by functional status in each dimension of the SF-36, summary indices, crude and adjusted odds ratios, and R<sup>2</sup> for poor health related quality of life in disabled vs nondisabled subjects

Dimension	Nondisabled %	Disabled %	Crude ORs (95% CI) (disabled vs nondisabled)	Model 1*		Model 2**	
				OR (95% CI)	R <sup>2</sup>	OR (95% CI)	R <sup>2</sup>
Physical functioning	27.2	94.7	47.9 (28.9–79.4)	26.5 (15.5–45.2)	0.509	15.6 (9.0–26.8)	0.413
Role-physical	17.6	68.4	10.1 (7.9–13.0)	8.3 (6.3–10.7)	0.094	5.4 (4.1–7.2)	0.186
Bodily pain	44.1	81.2	5.4 (4.1–7.3)	4.2 (3.1–5.6)	0.062	2.7 (2.0–3.8)	0.151
General health	41.2	89.0	11.6 (8.1–16.7)	7.2 (5.0–10.5)	0.118	4.5 (3.1–6.6)	0.175
Vitality	49.7	83.1	5.0 (3.7–6.7)	3.8 (2.8–5.2)	0.050	2.7 (1.9–3.7)	0.096
Social functioning	33.5	79.2	7.6 (5.7–10.0)	6.4 (4.8–8.5)	0.056	4.5 (3.3–6.1)	0.129
Role-emotional	11.5	29.7	3.3 (2.5–4.2)	3.5 (2.7–4.6)	0.033	2.3 (1.7–3.1)	0.098
Mental health	45.9	73.1	3.1 (2.5–4.1)	2.7 (2.0–3.5)	0.040	2.0 (1.5–2.6)	0.092
Physical component summary	32.7	94.9	38.6 (22.9–65.0)	22.2 (13.0–37.7)	0.206	13.2 (7.7–22.7)	0.294
Mental component summary	37.0	50.3	1.7 (1.3–2.1)	1.9 (1.5–2.4)	0.022	1.5 (1.2–2.0)	0.070

**Abbreviations/notes:** OR, odds ratio; CI, confidence interval; R<sup>2</sup>, Nagelkerke R<sup>2</sup>. \*Model 1: ORs adjusted for sociodemographic variables (age, sex, education). \*\*Model 2: OR adjusted for sociodemographic variables and comorbidity variables (stroke, ischemic heart diseases, cancer, diabetes, respiratory diseases, circulatory diseases, musculoskeletal diseases, neurological diseases, lesions, and trauma).

encounter significant adversity, trauma, tragedy, threats, or even significant sources of stress.<sup>25</sup>

These results need a different judgment for the physical, mental, and social dimensions of HRQL. As it was expected, the effect sizes and ORs associated with the physical dimensions were higher. It is especially noteworthy that the association with physical functioning can be described as huge. Nevertheless, the notable coincidence of the items used to assess disability and the questions of SF-36 that generate this scale compels us to qualify the association as partially misleading. In this sense, a detailed analysis of SF-36 items of physical functioning shows that they record the presence of a range of nine physical limitations (such as walking or climbing stairs) and only one self-care item (bathing or dressing). In this study the variable disability is based on the lack of autonomy to perform 10 activities of daily living such as eating, walking, bathing, or using the toilet. Whereas the disability score relies on the performance of some activities, HRQL is based mainly on the individual's perception of their disability.<sup>8</sup> The multidimensional approach to the measure of HRQL raises the difficulty of defining the boundary between both concepts, because in practice there is partial overlap. Therefore, this huge OR cannot be alleged to be evidence of association, because the assumption of content independence is partially broken. However, this limitation does not apply to the three other physical dimensions. Because these dimensions measure different issues, we can accept a highly significant level of association with disability. Moreover, the concept of self-care that is at the basis of the concept of disability doesn't overlap with pain, health self-perception, or work activities

that constitute the origin of the scales of role-physical, bodily pain, and general health. Even when adjusted by covariates, disabled people living in the community have a risk of pain three to five times higher than that of nondisabled people, a self-perception of poor health, and problems performing work activities. Disabled people also have worse mental health than the nondisabled population, but the impact measured by ORs and effect sizes is lower than that on physical scales. Nevertheless, when compared with the nondisabled population, the disabled are at a significantly higher risk (between 2.0 and 4.5) of perceiving poor vitality, social functioning, role-emotional, or mental health. Of these scales, social functioning shows the highest adjusted OR (4.5). This elevated figure reflects frequent interference with social activities because of physical or emotional problems. The scales related to individuals' mood perception as a level of vitality, and mental health appears less affected (adjusted ORs of 2.0 and 2.7), but they show that the disabled feel tired and depressed more often than the nondisabled group.

The SF-36 summary measures originate from the eight scales and consequently share the features described previously. The high weight of physical functioning in the PCS warns of the same association of partial lack of independence with the huge OR obtained by PCS in the logistic regression models. It is well known that the highest value on the scale for physical functioning correlates with the PCS. Our results corroborate this feature by showing that the disabled, with an adjusted OR of 13.2, have a worse score on the PCS. The disabled population's risk of a perception of poor global mental health is higher (1.5 times) than the risk in the nondisabled

population, and this difference is not only statistically significant but also epidemiologically relevant because it underscores the psychological effect of disability. Wilson and Cleary proposed a theoretical framework for understanding the relationship between the biological and physiological determinants of health, their impact in terms of functional status, and their effect on the loss of HRQL. These authors suggested that functional status has a mediating role between disease and HRQL, in what usually is referred to as 'physical domains'.<sup>16</sup> Our results also support the application of this model to the psychological and social scales of SF-36. The analysis of the percentage of the perceived health explained by the two models corroborates the previous statement. They fluctuate between 10% for mental dimensions and 40% for physical functioning. The models including comorbidity as an independent variable (model 2) produced higher levels of explanation.

Various ways to interpret differences in the scores on quality-of-life questionnaires have been proposed in the literature.<sup>19</sup> Some authors have proposed that a 10% difference should be interpreted as clinically relevant.<sup>26</sup> Accordingly, crude differences between nondisabled and disabled subjects in our study were greater than the threshold for all dimensions and the two component summaries. The effect size is an alternative way to assess impact.<sup>19,20</sup> Effect size for crude differences in our study was large ( $>0.8$ ) for PCS and the eight dimensions of SF-36, and small for MCS (0.43). Assessment of HRQL is based on the subjects' perception of their own health status, as determined by questionnaires such as SF-36. However, HRQL consists of different factors that are, in turn, related to each other. The challenge lies in quantifying the role of disability and separating it from the role of other determinants of HRQL, ie, whether disability is a factor associated with the loss of HRQL itself or only accompanies the negative effect of the presence of disease. Logistic regression models have been used in our study for this purpose. The conclusion is that disability is independently associated with HRQL deficit, notwithstanding the methodological reservations already expressed for the physical functioning. The adjusted ORs range between 2.0 and 5.4, except for those of the PCS and physical functioning, which are much higher.

The contribution of the different groups of variables for suboptimal health to differences in ORs between nondisabled and disabled subjects had a slightly different pattern. The first group of sociodemographic variables (age, sex, and educational level) roughly reduced the crude ORs of physical dimensions in a fourth and the ORs of the mental dimensions in an eighth. Significantly, the presence of

diseases produced a similar decrease in all dimensions. The final adjusted ORs of the physical scales became halved, and those of the mental scales were reduced to a third, which indicates a slightly higher association of those groups of variables with physical health. The analysis of the relative presence of those diseases in disabled people indicates that stroke and mental problems are the groups most frequently associated with disability. The probability of being disabled is 18.54 and 15.70 times higher for people declaring stroke and mental problems, respectively.

The prevalence of disability in the noninstitutionalized general population of the Basque Country was 3.6%, and it rose to 11.1% in the population more than 65 years of age. A study to assess disability in the noninstitutionalized Spanish population found that 19.1% of the subjects reported some limitation of activities of daily living.<sup>22</sup> Freedmann et al reported similar differences after measuring limitations for activities of daily living in different surveys of the American population more than 65 years of age.<sup>12</sup> In the National Health Interview Survey, disability increased from 6.4% to 8.4% from 1982 through 1993.<sup>27,28</sup> Higher rates were, however, reported in the National Long Term Care Survey, with values changing from 13% to 11% during the same period.<sup>29</sup> The lack of a standard framework to assess disability made comparison with other studies difficult. Besides, we had no figures for disabled populations residing in nursing homes in Spain. Under these conditions, no conclusions could be drawn from the various results. Therefore, we emphasize the importance of the development of standardized formulas that help to evaluate geographic and temporal differences in measuring disability.

Some limitations of this study should be discussed. The first has already been discussed and applies to the physical functioning scales. The second is that the BCHS does not include the institutionalized population because it is aimed at assessing the health status of the general population rather than the specific characteristics of the disabled population. The level of information available was therefore lower than in surveys designed for a more specific purpose, and the population with greater dependence levels was excluded, since these people are cared for in nursing homes. In addition, the cross-sectional design of the study did not allow for establishing a causal relationship between HRQL and disability, but only for relating both factors and quantifying the effect size.

## Conclusions

This study establishes a significant relationship between disability and the loss of quality of life in both physical and mental dimensions of SF-36, irrespective of the associated disease.

The effect in the mental dimensions was lower, but still relevant. The consequence for public health is that care of disabled must include not only the physical performance of activities of daily living but also the mental aspects and social activities.

## Acknowledgments

We would like to acknowledge the editorial assistance provided by Sally Ebeling. This study was supported by research grant BIO05/DCA/001 from the Department of Health and grant PI05/1703 from the Carlos III Institute.

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Murray CJ, Lopez AD. Regional patterns of disability-free life expectancy and disability-adjusted life expectancy: Global Burden of Disease Study. *Lancet*. 1997;349:1347–1352.
- Murray CJ, Lopez AD. Global mortality, disability, and the contribution of risk factors: Global Burden of Disease Study. *Lancet*. 1997;349:1436–1442.
- Murray CJ, Lopez AD. Alternative projections of mortality and disability by cause 1990–2020: Global Burden of Disease Study. *Lancet*. 1997;349:1498–1504.
- Murray CJL, Lopez AD, editors. *The Global Burden of Disease: a comprehensive assessment of mortality and disability from diseases, injuries, and risk factors in 1990 and projected to 2020*. Cambridge: Harvard University Press; 1996.
- Rothwell PM. The high cost of not funding stroke research: a comparison with heart disease and cancer. *Lancet*. 2001;357:1612–1616.
- Testa MA, Simonson DC. Assessment of quality-of-life outcomes. *N Engl J Med*. 1996;334:835–840.
- World Health Organization. *International classification of functioning, disability, and health*. Geneva, Switzerland: World Health Organization; 2001.
- Weisscher N, de Haan RJ, Vermeulen M. The impact of disease-related impairments on disability and health-related quality of life: a systematic review. *BMC Med Res Methodol*. 2007;7:24.
- Krahn GL, Fujiura G, Drum CE, Cardinal BJ, Nosek MA. The dilemma of measuring perceived health status in the context of disability. *Disabil Health J*. 2009;2:49–56.
- Ware JR, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care*. 1992;30:473–483.
- Horsman J, Furlong W, Feeny D, Torrance G. The Health Utilities Index (HUI): concepts, measurement properties and applications. *Health Qual Life Outcomes*. 2003;1:54.
- Freedman VA, Martin LG, Schoeni RF. Recent trends in disability and functioning among older adults in the United States. *JAMA*. 2002;288:3137–3146.
- Mahoney FI, Barthel DW. Functional evaluation: the Barthel index. *Md Med J*. 1965;14:61–65.
- Encuesta de Salud del País Vasco (ESCAV 2002). Departamento de Sanidad, Gobierno Vasco. [http://www.osasun.ejgv.euskadi.net/r52-20726/es/contenidos/informacion/encuesta\\_salud/es\\_4044/encues\\_salud.html](http://www.osasun.ejgv.euskadi.net/r52-20726/es/contenidos/informacion/encuesta_salud/es_4044/encues_salud.html).
- Orfila F, Ferrer M, Lamarca R, Tebe C, Domingo-Salvany A, Alonso J. Gender differences in health-related quality of life among the elderly: The role of objective functional capacity and chronic conditions. *Soc Sci Med*. 2006;63:2367–2380.
- Wilson IB, Cleary PD. Linking clinical variables with health-related quality of life. A conceptual model of patient outcomes. *JAMA*. 1995;273:59–65.
- Kwiatkowski TG, Libman RB, Frankel M, et al. Effects of tissue plasminogen activator for acute ischemic stroke at one year. *N Engl J Med*. 1999;340:1781–1787.
- Mar J, Begiristain JM, Arrozola A. Cost-effectiveness analysis of thrombolytic treatment for stroke. *Cerebrovasc Dis*. 2005;20:193–200.
- Walters SJ, Brazier JE. What is the relationship between the minimally important difference and health state utility values? The case of the SF-6D. *Health Qual Life Outcomes*. 2003;1:4.
- Kazis LE, Anderson JJ, Meenan RF. Effect sizes for interpreting changes in health status. *Med Care*. 1989;27 Suppl 3:S178–S189.
- Cohen J. *Statistical power analysis for the behavioral sciences*. New York: Academic Press; 1977.
- Graciani A, Banegas JR, Lopez-García E, Rodriguez-Artalejo F. Prevalence of disability and associated social and health-related factors among the elderly in Spain: a population-based study. *Maturitas*. 2004;48:381–392.
- Guallar-Castillon P, Redondo-Sendino A, Banegas JR, Lopez-García E, Rodriguez-Artalejo F. Differences in quality of life between women and men in the older population of Spain. *Soc Sci Med*. 2005;60:1229–1240.
- Arostegui I, Núñez-Antón V, Quintana JM. Analysis of the short form-36 (SF-36): the beta-binomial distribution approach. *Stat Med*. 2007;26(6):1318–1342.
- Luthar SS, Cicchetti D, Becker B. The construct of resilience: a critical evaluation and guidelines for future work. *Child Dev*. 2000;71:543–562.
- Juniper EF, Gordon GH, Willan A, Griffith LE. Determining a minimal important change in a disease-specific quality of life questionnaire. *J Clin Epidemiol*. 1994;47:81–87.
- Crimmins EM, Saito Y, Reynolds SR. Further evidence on recent trends in the prevalence and incidence of disability among older Americans from two sources: the LSOA and the NHIS. *J Gerontol B Psychol Sci Soc Sci*. 1997;52: S59–S71.
- Schoeni R, Freedmann VA, Wallace R. Persistent, consistent, widespread and robust? Another look at recent trends in old-age disability. *J Gerontol B Psychol Sci Soc Sci*. 2001;55:S206–S218.
- Manton KG, Gu X. Changes in the prevalence of chronic disability in the United States black and non-black population above age 65 from 1982 to 1999. *Proc Natl Acad Sci U S A*. 2001;98:6354–6359.

### ClinicoEconomics and Outcomes Research

### Publish your work in this journal

ClinicoEconomics & Outcomes Research is an international, peer-reviewed open-access journal focusing on Health Technology Assessment, Pharmacoeconomics and Outcomes Research in the areas of diagnosis, medical devices, and clinical, surgical and pharmacological intervention. The economic impact of health policy and health systems

Submit your manuscript here: <http://www.dovepress.com/clinicoeconomics-and-outcomes-research-journal>

Dovepress

organization also constitute important areas of coverage. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.