

Impact Of Spinal Cord Injury On Sleep: Current Perspectives

This article was published in the following Dove Press journal:
Nature and Science of Sleep

Abdulghani Sankari ^{1,2}
M Safwan Badr ^{1,2}
Jennifer L Martin ^{3,4}
Najib T Ayas⁵
David J Berlowitz ⁶⁻⁸

¹Department of Internal Medicine, John D. Dingell VA Medical Center, Detroit, MI, USA; ²Department of Internal Medicine, Wayne State University, Detroit, MI, USA; ³Geriatric Research, Education and Clinical Center, VA Greater Los Angeles Healthcare System, North Hills, CA, USA; ⁴Geriatric Research, Education and Clinical Center, David Geffen School of Medicine at the University of California, Los Angeles, CA, USA; ⁵Department of Medicine, The University of British Columbia, Vancouver, BC, Canada; ⁶Department of Physiotherapy, University of Melbourne, Melbourne, Australia; ⁷Department of Physiotherapy, Austin Health, Melbourne, Australia; ⁸Institute for Breathing and Sleep, Austin Health, Melbourne, Australia

Abstract: Sleep disorders are commonly encountered in people living with spinal cord injury (SCI). Primary sleep disorders such as sleep-disordered breathing (SDB), sleep-related movement disorders, circadian rhythm sleep-wake disorders, and insomnia disorder are common conditions after SCI but remain under-recognized, underdiagnosed and therefore remain untreated for a majority of patients. Sleep disturbances in people living with SCI are associated with significant impairments of daytime function and quality of life. Previous reviews have described findings related mainly to SDB but have not examined the relationship between other sleep disorders and SCI. This narrative review examines various sleep abnormalities and related functional and physical impairments in people living with SCI. It discusses new evidence pertaining to management, highlights existing limitations in the literature and recommends future directions for research.

Keywords: SCI, sleep-disordered breathing, leg movements, circadian rhythm sleep disorders, insomnia

Introduction

Sleep disorders are more common in individuals with spinal cord injury (SCI) than in the general population and likely contribute to reduced societal participation and quality of life.¹ Poor sleep quality is a consistent patient-reported outcome in populations with chronic SCI.²⁻⁶ Multiple factors may adversely influence sleep after SCI including very high rates of sleep-disordered breathing (SDB), particularly in those with high thoracic and cervical level injuries,⁷ high frequency of abnormal legs movements during wake and sleep,⁸ and poor sleep quality. The latter may be due to pain, insomnia and/or sleep-wake circadian rhythm disturbances.^{9,10} Subsequently, individuals living with SCI report daytime symptoms such as fatigue, excessive day time sleepiness,^{11,12} difficulties with concentration, and impaired quality of life. Furthermore, fatigue and excessive daytime sleepiness, have been associated with pain,¹³ depression,¹⁴ side effects of medications, spasticity,¹⁵ discomforts related to posture¹⁶ and autonomic dysfunction¹⁷ in people living with SCI. Interestingly, recent studies suggested that SCI does not in itself result in significantly impact of fatigue on health-related quality of life.¹⁸ However, approximately 40% of individuals with SCI were reported to be at risk of experiencing psychological disorder, such as depression and anxiety disorder.

In this narrative review and commentary, we examine the impact of SCI on sleep and daytime function. This paper is written to summarize current perspectives, provoke discussion and dialogue and to support the clinical practice of spinal,

Correspondence: Abdulghani Sankari
Division of Pulmonary, Critical Care and Sleep Medicine, 3990 John R, 3-Hudson, Detroit, MI 48201, USA
Tel +1 (313) 745-6033
Fax +1 (313) 745-8725
Email asankari@wayne.edu

rehabilitation, neurology, respiratory and sleep health professionals who provide care for people living with the complex co-morbidities of SCI. This paper is not intended to be a comprehensive systematic review; rather we have reviewed the existing literature exploring health risks due to the coexistence of sleep disorders and SCI. Of the recognized sleep disorders, we discuss and summarize the current evidence for the evaluation and management of the most prevalent and well-described conditions in SCI patients: sleep-disordered breathing (SDB), circadian rhythm sleep-wake disorders (CRSWD), insomnia, and periodic legs movements during sleep. Finally, we present the challenges faced in the management of these disorders and identify future directions for research.

Sleep-Disordered Breathing In SCI

Sleep-disordered breathing (SDB) is highly prevalent after SCI. While many studies noted the increased frequency of obstructive sleep apnea (OSA),^{10,19} there are reports of increased hypoventilation at sleep onset,²⁰ central sleep apnea (CSA) and narrow CO₂ reserve, a marker of increased susceptibility to CSA in cervical SCI.²¹ The severity of SDB is characterized by the apnea-hypopnea index (AHI) which is the number of times the airway narrows or closes per hour of sleep with arousals or desaturations of 3 or 4%. SDB is common in the general population of adults, with moderate to the severe disease present in approximately 10% of middle-aged (30–49 years) men, 3% of middle-aged women, and higher rates in older adults of both sexes.²² SDB seems to develop quickly after the onset of SCI; in a study by Berlowitz and colleagues, 62% of individuals with SCI assessed by full polysomnography (PSG) had SDB by four weeks post-injury. The study conservatively estimated that 10% of the cohort may have had SDB prior to the injury, indicating most developed SDB after the injurious event.²³ Chronically, SDB has a prevalence of 40–91% in SCI patients.^{2,24–28} The prevalence in people with paraplegia does not appear significantly different from what is seen in the general population; however, the literature is less comprehensive in this group owing partly to difficulties in completing diagnostic testing in patients with limited mobility.

Multiple mechanisms predispose to the development of SDB in SCI.²⁹ These include increase upper airway collapsibility,³⁰ a reduced dilator muscle responsiveness/effectiveness,³¹ a reduced arousal threshold,³² and an unstable ventilatory control system.²¹ The increased upper airway collapsibility can be due to multiple physiological

and/or anatomical factors. These include obesity (as indicated by increased specific measures of body habitus such as weight, body mass index (BMI), waist, abdominal, and neck girth) which is associated with OSA in both SCI²⁴ and the general population.³³ Obesity leads to narrowing of the upper airway by deposition in pharyngeal tissues including the tongue in both the general population and in SCI,^{22,34} in addition to the effect of reducing lung volume on decreased upper airway caliber during sleep.³⁵ In chronic SCI, most authors have observed associations between OSA prevalence and increasing age, BMI and neck circumference,^{24,25,27,36} but these relationships appear weaker in the acute, post-injury period.²³ Many individuals with SCI receive medications to control other symptoms that can affect breathing, including sedatives, muscle relaxants, and narcotics.³⁷ In addition, there is some cross-sectional evidence that loud snoring (a cardinal symptom of OSA) is associated with the use of anti-spasticity medications and obesity in SCI.³⁸

The loss of sympathetic modulation to the airways with resultant unopposed parasympathetic activation may result in bronchoconstriction in individuals with SCI,¹⁷ and in increased nasal resistance in tetraplegia.³⁹ Increased nasal resistance may contribute to upper airway narrowing by increasing negative pressure in the upper airway. There is no evidence that links bronchoconstriction to sleep disturbance in SCI and while a pilot study of the effect of nasal congestion in tetraplegia suggested a potential reduction in OSA severity,³⁹ the experiment requires replication with larger numbers and perhaps alternative decongesting medications.⁴⁰

Abnormalities of ventilatory control may also contribute to breathing instability and SDB after SCI. A recent study showed that the genioglossus reflex response to negative upper airway pressure (a protective reflex to maintain upper airway patency) was reduced in people with both OSA and SCI compared to those with OSA but without SCI. However, the mechanical corollary for this physiologic derangement is yet to be established.³¹ Also, accumulating data suggest that some patients with cervical spinal injuries may also experience CSA, perhaps related to a sleep onset hypoventilation and increased plant gain and a corresponding decrease in the CO₂ reserve.^{20,21,41} These physiologic abnormalities may predispose to upper airway collapse and consequent upper airway obstruction by reducing upper airway tone during periods of central apnea.⁴² Although these studies had small sample sizes and the underlying etiology of these ventilatory control abnormalities are unclear, it suggests the possibility that the mechanisms driving the increased prevalence and severity of SDB in

SCI may be different to those observed in the general population. Whether this may eventually lead to specifically tailored therapies for SDB in SCI is currently under research investigation.

Sleep-disordered breathing in tetraplegia may also be an etiologically bi-phasic disorder; acutely caused by the cervical SCI with partial resolution during injury recovery, only to increase again with age, weight gain, ongoing chronic intermittent hypoxia and the use of medications that compromise respiration.^{7,23,24} An association between cognitive performance and the severity of SDB has been described acutely after tetraplegic injury, impairments that persisted after adjustment for the known confounders of age and pre-injury estimates of verbal intelligence.⁴³ It is further suggested that an individual with acute tetraplegia and undiagnosed or untreated SDB may participate less in rehabilitation⁴⁴ due to sleepiness and fatigue⁴⁵ and therefore be less engaged in activities that improve quality of life and maintain functioning over time.

Consequences Of Sleep Disordered Breathing In SCI

Consistent with findings from individuals without SCI, people with chronic SCI and SDB suffer reduced cognitive function⁴⁶ that can be observed early after injury. A study of 104 people with acute tetraplegia by Schembri and colleagues found that severe OSA (AHI > 30) was associated with poorer attention, information processing, and recall compared to milder OSA.⁴³ The deficit in attention and information processing observed in those with severe OSA was large; equivalent to the expected decline after 31 additional years of aging. The substantial cortical reorganization has been well described after SCI⁴⁷⁻⁴⁹ and further research is required to understand the relative importance and time course of changes in generalized central nervous system inflammation,^{50,51} deafferentation,⁴⁹ functional connectivity,⁴⁸ and SDB.⁴³

People with SDB often report poor sleep quality and daytime sleepiness. Treatment with CPAP significantly improves daytime sleepiness in the general population with SDB.^{52,53} Likewise, untreated SDB is strongly associated with the incidence of cardiovascular and cerebrovascular disease including hypertension, strokes, and myocardial infarction.⁵⁴ The impact of SDB on cardiovascular health in SCI has not been well studied, though one study of 50 patients with tetraplegia noted increased cardiac medication use in patients with sleep apnea.²⁷ Given the high rates of cardiovascular disease in SCI patients, the associations between SDB and cardiovascular risks in SCI need to be further studied.¹²

Periodic Leg Movements

Periodic Leg Movements (PLM) are characterized by periodic episodes of repetitive and highly stereotyped limb movements, typically big toe and ankle dorsiflexion which is often accompanied by knee and hip flexion.^{8,55} The associated Restless Leg Syndrome (RLS), is characterized by uncomfortable leg sensations usually prior to sleep onset that causes an almost irresistible urge to move.⁵⁶ Both conditions can result in considerable disruption to sleep quality and are typically associated with excessive daytime sleepiness. Up to 80% of RLS patients also experience PLMS (Periodic Leg Movement syndrome), defined as five or more PLM events per hour of sleep.⁵⁷

The prevalence of periodic leg movement disorder (PLMD) appears to be markedly elevated in people with SCI. While the majority of previous samples are small and clinic-based rather than true population estimates, the prevalence in those with a lesion above T10 has been reported to range from approximately 50–100%.^{55,58–68} As detailed previously, SDB is also highly prevalent in tetraplegia,^{23,24} which precludes the diagnosis of PLMD according to the ICSD-3. For this reason, the American Association of Sleep Medicine (AASM) PLM scoring rules exclude events that occur from 0.5 seconds prior to or following a scored respiratory event. While some studies investigating PLMs excluded participants with SDB;⁵⁷ others have included participants with SDB and excluded PLM events adjacent to respiratory events as per the AASM rules,⁵⁹ and other authors made no reference to possible co-existent SDB.^{66,68} Peters et al recently collated 262 clinical and research sleep studies in people with tetraplegia.⁸ Similarly to the 2015 Prosperpio paper,⁵⁹ Peters excluded PLM events as per the AASM rules and observed an estimated prevalence (proportion with a PLM rate of > 15 events per hour) of 58%. No significant difference was observed in PLM prevalence or severity between participants with or without SDB. The arousal index was associated with the SDB severity, but not with the PLM index.⁸ Similarly, Salminen et al published a clinical case strongly suggesting SDB and PLM are independent phenomena in tetraplegia.⁵⁸ As such, it appears that PLMD is common in SCI, especially in tetraplegia and is not substantially modified in prevalence or severity by SDB.

Effect Of SCI On Circadian Rhythm Sleep-Wake Disorders (CRSWD)

There is evidence that the circadian rhythmicity of melatonin is disrupted in tetraplegia. Following a complete

cervical SCI, the afferent input of somatic and autonomic fibers below the SCI is disrupted and the efferent sympathetic innervation of the pineal gland via the superior cervical ganglion is interrupted. During the daytime, light striking the retina stimulates afferent pathways through the suprachiasmatic nuclei that in turn pass to the superior cervical ganglion and onward to the pineal gland to inhibit melatonin production. Darkness removes this inhibition and melatonin production ensues. The fibers from the suprachiasmatic nuclei travel with the sympathetic fibers which exit the spine between the first and sixth thoracic levels before ascending outside the spinal cord to the superior cervical ganglion. Fibers then pass back into the cortex and innervate the pineal gland. In 2000, Zeitzer and colleagues showed that complete cervical SCI, which cuts the suprachiasmatic nuclei to superior cervical ganglion pathway, is associated with near abolition of circadian melatonin rhythmicity and a markedly reduced circulating level.⁶⁹ These findings have been replicated by others.^{70–74} Other endocrine rhythms similarly modified by circadian influences, such as cortisol and thyroid-stimulating hormone, do not appear markedly affected by cervical SCI, however, the temperature is affected.⁷⁵ Thijsen et al observed a substantial phase advancement of temperature in tetraplegia. Core temperature phase and amplitude of change in paraplegia were not different from the non-disabled control participants. The temperature phase advance observed in tetraplegia may be yet another contributor to the poor sleep quality reported in this group.

Whether SCI affects sleep timing per se is similarly unclear. Delayed REM onset has been reported in a community survey and home PSG study of 78 people with tetraplegia.²⁴ Participants with motor and sensory complete lesions (ASIA A, $n=35$ mean REM latency in minutes of 145 ($SD=82$)) took significantly longer to attain first REM than those with incomplete lesion (AIS B-E, $n=43$, 94(58.5) minutes) although OSA was also more severe in those with complete lesions potentially confounding this finding. Further, while sleep complaints in SCI are frequent,^{6,10,19,24,76–78} and anecdotally, delayed sleep phase is reported in SCI, the relative contribution of environmental, social and physiological effects remains unclear. As detailed above,⁷⁵ temperature phase is substantially advanced in tetraplegia but it is unclear if or how this interacts with melatonin dysregulation to affect sleep. Highly prevalent PLMD and SDB further complicate symptomology, diagnosis, and management.

Effect Of SCI On Insomnia

Insomnia is highly prevalent in the general population, with estimates ranging from 15–30%, and women having a 1.4 times higher risk than men for insomnia.⁷⁹ Insomnia disorder, defined as difficulty falling or staying asleep that persists for at least three months, occurs at least three nights per week and is accompanied by daytime consequences, is likely less common, but estimates still suggest rates of 10–20% in the population.⁷⁹ A recent study using the Sleep Heart Health Study Sleep Habits Questionnaire found that 57% of SCI individuals had insomnia symptoms,⁷⁸ although it is not clear that these individuals would meet criteria for insomnia disorder as they also had significantly higher risk for SDB and had greater differences in weekday/weekend sleep which may be associated with insufficient sleep rather than insomnia disorder. This does suggest that evaluation for insomnia disorder is useful in understanding the effect of other sleep disorders experienced by individuals with SCI and can affect the choice of treatment.

Evaluation Of Sleep Disorders In Individuals With SCI

SDB is underdiagnosed in this population, and this is likely true for other sleep disorders as well.⁸⁰ Given the high prevalence of sleep disorders in SCI, any substantial sleep-related complaints should be further evaluated so that appropriate treatment can be initiated. Common symptoms include daytime fatigue/sleepiness, poor quality sleep, nocturnal awakenings, and witnessed apneas. These symptoms are not specific and further assessment is usually required to identify the correct diagnosis. The gold standard for diagnosis of SDB and periodic legs movements is full night attended PSG in the sleep laboratory; however, access to these facilities is often limited with substantial wait times.⁸⁰ In addition, many laboratories are not equipped to handle the requirements of individuals with SCI (eg, lifts, special beds for pressure care, adequate staff) representing another barrier to diagnosis. Home sleep apnea testing may provide an alternative diagnostic tool for SDB in these patients. This could include full PSG (including EEG recording) in the patients' homes, which similar to what is done in sleep laboratories (Type II studies). However, costs and resources required for these are substantial, and often not feasible outside of a research setting. Another possibility is truncated Type III or oximetry studies, which record some of the physiologic signals included in PSG but lack

others (e.g. EEG and EMG). These systems can be more easily set up in the home and have been used for SCI screening.⁸¹ The study by Bauman et al coupled a Level III device with transcutaneous carbon dioxide monitoring and observed similar sample prevalence (81%) as has been previously described, albeit at the upper end of the prevalence estimate ranges. A more recent study suggests that the incorporation of a tetraplegia-specific questionnaire with oximetry had reasonable sensitivity (77–83%) and specificity (81–88%) in detecting moderate to severe SDB.⁸⁰ The Graco et al model was compared with full PSG and as such is currently the best-validated, population-based screening approach available.

Management Of Sleep Disorders In Individuals With SCI

Treatment Of Obstructive Sleep Apnea

Treatment of SDB in people living with SCI is challenging, although substantial symptomatic improvements may be observed in many individuals. Although not studied in SCI, lifestyle interventions including exercise and weight loss⁸² should be encouraged as these have positive effects on SDB in the general population, and there is no reason to assume a lack of benefit in SCI.⁸³ The most commonly used treatment for SDB is continuous positive airway pressure (CPAP) which consists of a mask attached to the nose/face that administers a positive pressure to the upper airway preventing its collapse during sleep. CPAP is effective in eliminating obstructive events during sleep and improving oxygenation. One of the major limitations of CPAP is poor adherence, with only about 50% of patients being able to adhere to the therapy on a long-term basis.⁸⁴ Given the challenges related to arm strength/mobility in tetraplegia, and additional factors such as increased nasal congestion, it is perhaps not surprising that only 20–50% of people with chronic SCI and SDB are reported to adhere with CPAP.^{2,25,27,85,86} The less claustrophobic nature of nasal pillows may mean that they are well suited to tetraplegic users and the lower pressures required by many CPAP users with tetraplegia allows for greater flexibility in interface choice as leak is less likely at lower pressures.⁸⁷

The nature of the barriers to diagnosis and management of sleep disorders in SCI has recently been explored qualitatively through interviews with 20 specialist spinal medical practitioners from the UK, USA, Canada, Europe, and Australia.⁸⁸ Commonly reported barriers to independent

management practices were lack of skills, low confidence about capabilities, and the belief that SDB management was outside their scope of practice. The three spinal units independently managing SDB were well resourced with multidisciplinary involvement and “clinical champions” to lead the program.

The experiences of people living with SCI and comorbid SDB and the relationships of these experiences to concordance with CPAP prescription has also been reported recently.⁸⁶ Sixteen people with SCI and newly diagnosed SDB were provided with CPAP and participated in semi-structured interviews one month later. Usage was then reviewed at one, six and 12 months. The perceived balance of benefits versus burdens strongly influenced the ongoing level of use in study participants. Burden attributed to CPAP use was common and included mask discomfort and physical and emotional problems. Adherent participants were motivated by the immediate daytime benefits to mood, alertness, and reduced sleepiness. There was a tendency for patients to fail to recognize symptoms of SDB until after they were treated with CPAP. These practical challenges to the use of CPAP for SDB in tetraplegia and the uncertainty as to the real benefit of the therapy in SCI likely limit the ability of clinicians to provide evidence-based advice to their patients and for people with SCI to value the therapy.

Berlowitz et al recently assessed the effectiveness of CPAP in among individuals with acute SCI.⁸⁹ One hundred and sixty patients were randomized to CPAP or no CPAP for three months. Sleepiness was significantly improved with CPAP ($p=0.01$) though other cognitive parameters were unchanged. One issue was that CPAP adherence was relatively low in the study (2.9 hrs/night, with 21% achieving >4 hrs per night) despite only randomizing patients who were able to tolerate a three-day CPAP run-in period. The fact that sleepiness was improved with CPAP despite only partial adherence is encouraging and suggests that if adherence could be improved, more substantial benefits could be achieved.

Whether other ventilator modes, such as bilevel therapy, might be more effective in this patient group has been explored in a recent case series.⁹⁰ A combination of bilevel PAP support (for upper airway dysfunction) and bilevel PAP with volume assured pressure support (for SDB plus hypoventilation during sleep but not daytime) were used to support people with SCI (74% quadriplegia) and SDB over 12 months. If those lost to follow-up in this

cohort paper are assumed to be non-adherent, then overall PAP use of 28% at 3 months is similar to the 21% observed in the *COSAQ* (CPAP treatment for obstructive sleep apnea after acute quadriplegia) study, suggesting that the type of PAP is not a determinant of adherence.

An additional but frequently overlooked confounder to the interpretation of PAP use in clinical studies is the influence of government or privately funded diagnosis and treatment on practice. If, for example, an accident compensation system will only support one PAP device per patient, almost regardless of complexity, then clinicians and patients may well choose a more complex device. Conversely, if reimbursement requires a “step-wise” increase in complexity of treatment with evidence of failure before progression (CPAP to bi-level to controlled ventilation), then practice may well be different. Increased sleep onset ventilatory instability²⁰ and a reduced carbon dioxide reserve²¹ may well predispose people with tetraplegia to pressure support induced central apnea and sleep fragmentation. In such a case, CPAP at low pressures may be best while similarly, controlled (timed) ventilation could be indicated. We are unaware of any papers that have directly compared different PAP therapies for acceptance, usage, and efficacy in SCI as has been reported in conditions such as obesity hypoventilation syndrome.⁹¹ This paucity of literature impedes evidenced-based care and should be considered in future research priority setting exercises.

Other alternative treatments for SDB include upper airway surgery,^{92,93} hypoglossal nerve stimulation,⁹⁴ and dental appliances.⁹⁵ We are not aware of studies of these modalities in SCI, but anecdotally, some patients may respond physiologically and symptomatically and they may be considered in selected cases where risks do not outweigh the possible benefits. A recent pilot, cross-over controlled trial assessed the potential for nasal decongestion to reduce the AHI in tetraplegia.⁹⁶ No significant difference in SDB severity was observed although a trend to less severe respiratory events (more hypopneas versus apneas) was observed, similar to that observed in the general population with SDB.⁴⁰ More research into these alternative treatment measures is clearly necessary in cases of SCI.

Treatment Of Periodic Leg Movements In SCI

The origin of PLMs in the general population is generally not well understood. The clinical effectiveness of dopaminergic

agents led to the conclusion that the central brain or hypothalamic dysfunction drove the peripheral events. PLMs predominate in non-rapid eye movement (NREM) sleep in the general population, but in contrast, a number of case reports and series in SCI suggest that PLM may be observed throughout REM and without clear periodicity.^{8,55,57,58,66,68} This observation strongly suggests that PLM after SCI arises peripherally rather than centrally. Interventions which alter cortical arousal in SCI do not necessarily result in a reduction in PLM and similarly, inhibition of PLM does not necessarily reduce sleep disruption.^{56,58} Further, a recent case series evaluated PSG data in both SCI and Multiple Sclerosis from a spasticity clinic population who were typically referred for refractory “spasticity” despite often maximal anti-spasticity therapy. The study explored the possibility that spasticity was, in fact, undiagnosed and/or untreated PLM and showed a substantial reduction in PLM index and arousals from sleep in those with confirmed PLM (≥ 15 per hour of sleep) treated with a low dosage of pramipexole.⁹⁷ Taken together, these data suggest that there is likely clinical utility in SCI, especially where movements predominate in supine and during sleep, in assuming the distressing movements are PLM, not excess spasticity and treated with dopaminergic agents as first-line therapy. Polysomnography will obviously assist with differential diagnosis and more controlled trials are needed.

Treatment Of Insomnia In SCI

There are multiple challenges in treating insomnia disorder among individuals with SCI. First, pharmacological agents have a number of adverse effects that may be even more significant among individuals with SCI. For example, fall risk among those who ambulate may be elevated further, and impaired cognition may be exacerbated. As a result of these and other concerns, pharmacological therapy is recommended only after cognitive-behavioral therapy for insomnia (CBT-I) has been attempted.⁹⁸ CBT-I is a multi-component psychological treatment that includes behavioral techniques (stimulus control, sleep restriction therapy, sleep hygiene and relaxation/arousal reduction strategies) plus cognitive therapy to address sleep-related thoughts and beliefs. While no formal studies of CBT-I have been conducted in SCI, a systematic review found evidence that cognitive-behavioral therapies, in general, are helpful for anxiety, depression, adjustment, and coping problems.⁹⁹ Furthermore, there are studies of CBT-I treatments, using an adaptation of the standard behavioral techniques (e.g.,

eliminating the recommendation to get out of bed in the middle of the night if unable to sleep; addressing disease-specific cognitions) in populations with functional impairments,¹⁰⁰ and this approach may be extrapolated to SCI. Overall, SCI should not prevent people from being offered the best-available evidence-based treatment for insomnia disorder if the person is interested in a non-pharmacological approach.

There is also evidence that insomnia disorder can increase the risk for intolerance of positive airway pressure (PAP) therapy for SDB.¹⁰¹ As a result, it may be necessary to address insomnia symptoms when PAP therapy is initiated as patients with SCI already have significant challenges to PAP use due to limitations in physical function and other factors as outlined earlier.⁸⁶

Treatment Of CRSWD In SCI

Despite the clear abnormalities of circadian control in SCI and the availability of melatonin, little work has been undertaken to understand its potential therapeutic role in SCI, particularly in tetraplegia. Chronic melatonin supplementation is often used by people with blindness to counter circadian desynchronization brought on by their visual impairment and the associated jet-lag type symptoms.^{102,103} People living with tetraplegia are essentially 180 degrees out of circadian phase all of the time and as such, appropriately timed exogenous melatonin may offer unique opportunities for sleep-onset insomnia and sleep phase entrainment.

A small size pilot study⁷¹ and two subsequent randomized controlled, cross-over trials^{72,104} have demonstrated the safety of exogenous melatonin in tetraplegia. While the studies suggested some subjective improvements in sleep, both trials were underpowered statistically to demonstrate any effect on objective sleep variables. More recently, Kostovski and colleagues have shown that exogenous melatonin administration in tetraplegia can normalize clock gene expression in peripheral blood.⁷⁴ The symptomatic and longer-term physiological effects of these changes are as yet unknown.

Future Directions

Despite the increased prevalence of sleep disorders in individuals with SCI, the majority are under-recognized and untreated with several challenges in their management. Future studies could focus on the following knowledge gaps:

- Studies to determine the epidemiology and clinical manifestations of sleep disorders in individuals with

SCI are sorely needed. One particular challenge is that the parameters used for disease definition were based on observations in the general population.

- Individuals living with SCI often suffer from more than one sleep disorder, and studies on combined treatments for those with multiple sleep disorders are warranted.
- The underlying mechanisms of sleep disorders remain unclear and likely are related to physiological and behavioral changes that follow SCI. There is a critical need to identify mechanistic pathways using human as well as animal models. Such knowledge will inform the development and the testing of novel pharmacological treatments that are both effective and tolerable.
- Management of sleep disorders in this population remains a challenge given the complex clinical presentations and care needs. Likewise, there is a need to examine the positive impact of treating sleep disorders on societal participation, quality of life and function.
- The comparison of different PAP therapies for acceptance, usage, and efficacy in SCI remains unstudied.
- The effect of non-pharmacological treatments such as respiratory muscle training/exercises, dental appliances or nerve stimulation on treating SDB in SCI can inform alternative treatments when PAP therapy is not well-tolerated.
- The effect of non-pharmacological treatments such as behavioral interventions on treating insomnia and the adaptation and testing of CBT-I in SCI can lead to impactful improvements in daytime functioning and facilitate adjustment to PAP therapy in individuals with comorbid SDB and insomnia disorder.

Conclusion

The extant literature on the impact of SCI on sleep is limited and heterogeneous. While advances have been made recently on understanding the mechanisms and optimizing management of SDB after chronic SCI, other sleep disorders remain unstudied. Future studies could identify the particular expression of the disease, assess new therapies beyond traditional treatments, and determine predictors of treatment efficacy. A focus on developing standardized approaches along with measures of treatment efficacy could also provide more information on the efficacy of individual treatment components. A multidisciplinary approach to management is recommended given the overlap of sleep disorders and other co-morbid conditions in individuals

with SCI. A comprehensive management approach should include behavioral and pharmaceutical therapies, and be aligned with ongoing management and rehabilitation goals.

Acknowledgments

Dr. Sankari is supported by the (US) Department of Veterans Affairs Office of Research and Development #RX002885 and the National Heart, Lung, and Blood Institute #R21HL140447. Dr. Badr is supported by the VA Office of Research and Development, Department of Veterans Affairs #1I01RX002116, the National Heart, Lung, and Blood Institute #R01HL130552 and the Department of defense #SC150201. Dr. Ayas is supported by the Canadian Sleep and Circadian Network.

Disclosure

Dr Abdulghani Sankari reports grants from the United States Department of Veterans Affairs, grants from US Department of Defense, grants from National Institute of Health, grants from American Thoracic Society Foundation, during the conduct of the study; in addition, Dr Sankari has a patent US15706097 pending. Dr M Safwan Badr reports grants from NIH, grants from DOD, grants from the VA, during the conduct of the study; grants from NIH, grants from DOD, grants from VA, outside the submitted work. Dr Jennifer Martin reports grants from UCLA, during the conduct of the study. Dr Najib Ayas reports personal fees from bresotec, outside the submitted work. Dr David J Berlowitz reports non-financial support from Phillips Respironics, non-financial support from ResMed, outside the submitted work. This was not an industry-supported study. The opinions expressed in this article reflect those of the authors and do not necessarily represent official views of the Department of Veterans Affairs, National Institute of Health or Department of Defense. The authors report no other conflicts of interest in this work.

References

- Simpson LA, Eng JJ, Hsieh JTC, Wolfe DL, the Spinal Cord Injury Rehabilitation Evidence Research Team. The health and life priorities of individuals with spinal cord injury: a systematic review. *J Neurotrauma*. 2012;29(8):1548–1555. doi:10.1089/neu.2011.2226
- Sankari A, Bascom A, Oomman S, Badr MS. Sleep disordered breathing in chronic spinal cord injury. *J Clin Sleep Med*. 2014;10(1):65–72. doi:10.5664/jcsm.3362
- Sankari A, Martin J, Bascom A, Mitchell M, Badr M. Identification and treatment of sleep-disordered breathing in chronic spinal cord injury. *Spinal Cord*. 2015;53(2):145. doi:10.1038/sc.2014.216
- Biering-Sorensen F, Biering-Sorensen M. Sleep disturbances in the spinal cord injured: an epidemiological questionnaire investigation, including a normal population. *Spinal Cord*. 2001;39(10):505. doi:10.1038/sj.sc.3101197
- Levi R, Hultling C, Nash MS, Seiger A. The Stockholm spinal cord injury study: 1. Medical problems in a regional SCI population. *Paraplegia*. 1995;33(6):308–315.
- Jensen MP, Hirsh AT, Molton IR, Bamer AM. Sleep problems in individuals with spinal cord injury: frequency and age effects. *Rehabil Psychol*. 2009;54(3):323–331. doi:10.1037/a0016345
- Sankari A, Vaughan S, Bascom S, Martin JL, Badr MS. Sleep-disordered breathing and spinal cord injury: a state-of-the-art review. *Chest*. 2019;155(2):438–445. doi:10.1016/j.chest.2018.10.002
- Peters AEJ, van Silfhout L, Graco M, Schembri R, Thijssen D, Berlowitz DJ. Periodic limb movements in tetraplegia. *J Spinal Cord Med*. 2018;41(3):318–325. doi:10.1080/10790268.2017.1320874
- Budh CN, Hultling C, Lundeberg T. Quality of sleep in individuals with spinal cord injury: a comparison between patients with and without pain. *Spinal Cord*. 2005;43(2):85. doi:10.1038/sj.sc.3101680
- Giannoccaro MP, Moghadam KK, Pizza F, et al. Sleep disorders in patients with spinal cord injury. *Sleep Med Rev*. 2013;17s(6):399–409. doi:10.1016/j.smrv.2012.12.005
- Stepanski E, Lamphere J, Badia P, Zorick F, Roth T. Sleep fragmentation and daytime sleepiness. *Sleep*. 1984;7(1):18–26. doi:10.1093/sleep/7.1.18
- Sankari A, Badr MS. Diagnosis of sleep disordered breathing in patients with chronic spinal cord injury. *Arch Phys Med Rehabil*. 2016;97(1):176–177. doi:10.1016/j.apmr.2015.10.085
- Biering-Sorensen F, Biering-Sorensen M. Sleep disturbances in the spinal cord injured: an epidemiological questionnaire investigation, including a normal population. *Spinal Cord*. 2001;39(10):505–513. doi:10.1038/sj.sc.3101197
- Elliott TR, Frank RG. Depression following spinal cord injury. *Arch Phys Med Rehabil*. 1996;77(8):816–823. doi:10.1016/s0003-9993(96)90263-4
- de Paleville DGT, McKay WB, Folz RJ, Ovechkin AV. Respiratory motor control disrupted by spinal cord injury: mechanisms, evaluation, and restoration. *Transl Stroke Res*. 2011;2(4):463–473.
- Hammell K, Miller W, Forwell S, Forman B, Jacobsen B. Fatigue and spinal cord injury: a qualitative analysis. *Spinal Cord*. 2009;47(1):44. doi:10.1038/sc.2008.104
- Krassioukov A. Autonomic function following cervical spinal cord injury. *Respir Physiol Neurobiol*. 2009;169(2):157–164. doi:10.1016/j.resp.2009.08.003
- Wijesuriya N, Tran Y, Middleton J, Craig A. Impact of fatigue on the health-related quality of life in persons with spinal cord injury. *Arch Phys Med Rehabil*. 2012;93(2):319–324. doi:10.1016/j.apmr.2011.09.008
- Castriotta RJ, Wilde MC, Sahay S. Sleep Disorders in Spinal Cord Injury. *Sleep Med Clin*. 2012;7(4):643–653. doi:10.1016/j.jsmc.2012.10.005
- Bascom AT, Sankari A, Goshgarian HG, Badr MS. Sleep onset hypoventilation in chronic spinal cord injury. *Physiol Rep*. 2015;3(8):e12490. doi:10.14814/phy2.12490
- Sankari A, Bascom AT, Chowdhuri S, Badr MS. Tetraplegia is a risk factor for central sleep apnea. *J Appl Physiol (1985)*. 2014;116(3):345–353. doi:10.1152/jappphysiol.00731.2013
- Kim AM, Keenan BT, Jackson N, et al. Tongue fat and its relationship to obstructive sleep apnea. *Sleep*. 2014;37(10):1639–1648. doi:10.5665/sleep.4072
- Berlowitz DJ, Brown DJ, Campbell DA, Pierce RJ. A longitudinal evaluation of sleep and breathing in the first year after cervical spinal cord injury. *Arch Phys Med Rehabil*. 2005;86(6):1193–1199. doi:10.1016/j.apmr.2004.11.033
- Berlowitz DJ, Spong J, Gordon I, Howard ME, Brown DJ. Relationships between objective sleep indices and symptoms in a community sample of people with tetraplegia. *Arch Phys Med Rehabil*. 2012;93(7):1246–1252. doi:10.1016/j.apmr.2012.02.016

25. Burns SP, Kapur V, Yin KS, Buhner R. Factors associated with sleep apnea in men with spinal cord injury: a population-based case-control study. *Spinal Cord*. 2001;39(1):15–22. doi:10.1038/sj.sc.3101103
26. Leduc BE, Dagher JH, Mayer P, Bellemare F, Lepage Y. Estimated prevalence of obstructive sleep apnea-hypopnea syndrome after cervical cord injury. *Arch Phys Med Rehabil*. 2007;88(3):333–337. doi:10.1016/j.apmr.2006.12.025
27. Stockhammer E, Tobon A, Michel F, et al. Characteristics of sleep apnea syndrome in tetraplegic patients. *Spinal Cord*. 2002;40(6):286–294. doi:10.1038/sj.sc.3101301
28. Tran K, Hukins C, Geraghty T, Eckert B, Fraser L. Sleep-disordered breathing in spinal cord-injured patients: a short-term longitudinal study. *Respirology*. 2010;15(2):272–276. doi:10.1111/j.1440-1843.2009.01669.x
29. Fuller DD. How does spinal cord injury lead to obstructive sleep apnoea? *J Physiol*. 2018;596:2633. doi:10.1113/JP276162
30. Sankari A, Bascom AT, Badr MS. Upper airway mechanics in chronic spinal cord injury during sleep. *J Appl Physiol (1985)*. 2014;116(11):1390–1395. doi:10.1152/jappphysiol.00139.2014
31. Wijesuriya NS, Gainche L, Jordan AS, et al. Genioglossus reflex responses to negative upper airway pressure are altered in people with tetraplegia and obstructive sleep apnoea. *J Physiol*. 2018;596(14):2853–2864. doi:10.1113/JP275222
32. Rizwan A, Sankari A, Bascom AT, Vaughan S, Badr MS. Nocturnal swallowing and arousal threshold in individuals with chronic spinal cord injury. *J Appl Physiol*. 2018. doi:10.1152/jappphysiol.00641.2017
33. Young T, Palta M, Dempsey J, Skatrud J, Weber S, Badr S. The occurrence of sleep-disordered breathing among middle-aged adults. *N Engl J Med*. 1993;328(17):1230–1235. doi:10.1056/NEJM199304293281704
34. O'Donoghue FJ, Meaklim H, Bilston L, et al. Magnetic resonance imaging of the upper airway in patients with quadriplegia and obstructive sleep apnea. *J Sleep Res*. 2018;27(4):e12616. doi:10.1111/jsr.12616
35. Owens RL, Malhotra A, Eckert DJ, White DP, Jordan AS. The influence of end-expiratory lung volume on measurements of pharyngeal collapsibility. *J Appl Physiol*. 2009;108:445–451.
36. McEvoy RD, Mykityn I, Sajkov D, et al. Sleep apnoea in patients with quadriplegia. *Thorax*. 1995;50(6):613–619. doi:10.1136/thx.50.6.613
37. Bauman KA, Kurili A, Schotland HM, Rodriguez GM, Chiodo AE, Sitrin RG. Simplified approach to diagnosing sleep-disordered breathing and nocturnal hypercapnia in individuals with spinal cord injury. *Arch Phys Med Rehabil*. 2015;97:363–371.
38. Ayas NT, Epstein LJ, Lieberman SL, et al. Predictors of loud snoring in persons with spinal cord injury. *J Spinal Cord Med*. 2001;24(1):30–34. doi:10.1080/10790268.2001.11753552
39. Wijesuriya NS, Lewis C, Butler JE, et al. High nasal resistance is stable over time but poorly perceived in people with tetraplegia and obstructive sleep apnoea. *Respir Physiol Neurobiol*. 2017;235:27–33. doi:10.1016/j.resp.2016.09.014
40. Koutsourelakis I, Minaritzoglou A, Zakyntinos G, Vagiakis E, Zakyntinos S. The effect of nasal tramazoline with dexamethasone in obstructive sleep apnoea patients. *Eur Respir J*. 2013;42(4):1055–1063. doi:10.1183/09031936.00142312
41. Sankari A, Bascom AT, Riehani A, Badr MS. Tetraplegia is associated with enhanced peripheral chemoreflex sensitivity and ventilatory long-term facilitation. *J Appl Physiol (1985)*. 2015;119(10):1183–1193. doi:10.1152/jappphysiol.00088.2015
42. Badr MS, Toiber F, Skatrud JB, Dempsey J. Pharyngeal narrowing/occlusion during central sleep apnea. *J Appl Physiol*. 1995;78(5):1806–1815. doi:10.1152/jappphysiol.1995.78.5.1806
43. Schembri R, Spong J, Graco M, Berlowitz DJ, team Cs. Neuropsychological function in patients with acute tetraplegia and sleep disordered breathing. *Sleep*. 2017;40(2):zsw037. doi:10.1093/sleep/zsw037
44. Berlowitz DJ. Neurotrauma and sleep. *Sleep Med Rev*. 2013;17(6):391–393. doi:10.1016/j.smrv.2013.05.004
45. Berlowitz DJ, Ayas N, Barnes M, et al. Auto-titrating continuous positive airway pressure treatment for obstructive sleep apnoea after acute quadriplegia (COSAQ): study protocol for a randomized controlled trial. *Trials*. 2013;14:181. doi:10.1186/1745-6215-14-59
46. Sajkov D, Marshall R, Walker P, et al. Sleep apnoea related hypoxia is associated with cognitive disturbances in patients with tetraplegia. *Spinal Cord*. 1998;36(4):231–239.
47. Lucci G, Pisotta I, Berchicci M, et al. Proactive cortical control in spinal cord injury subjects with paraplegia. *J Neurotrauma*. 2019. doi:10.1089/neu.2018.6307
48. Athanasiou A, Terzopoulos N, Pandria N, et al. Functional brain connectivity during multiple motor imagery tasks in spinal cord injury. *Neural Plast*. 2018;2018:9354207. doi:10.1155/2018/9354207
49. Lopez-Larraz E, Montesano L, Gil-Agudo A, Minguez J, Oliviero A. Evolution of EEG motor rhythms after spinal cord injury: a longitudinal study. *PLoS One*. 2015;10(7):e0131759. doi:10.1371/journal.pone.0131759
50. Faden AI, Wu J, Stoica BA, Loane DJ. Progressive inflammation-mediated neurodegeneration after traumatic brain or spinal cord injury. *Br J Pharmacol*. 2016;173:681–691. doi:10.1111/bph.13179
51. Wu J, Stoica BA, Luo T, et al. Isolated spinal cord contusion in rats induces chronic brain neuroinflammation, neurodegeneration, and cognitive impairment. Involvement of cell cycle activation. *Cell Cycle*. 2014;13(15):2446–2458. doi:10.4161/cc.29420
52. Smith BM, Evans CT, Kurichi JE, Weaver FM, Patel N, Burns SP. Acute respiratory tract infection visits of veterans with spinal cord injuries and disorders: rates, trends, and risk factors. *J Spinal Cord Med*. 2007;30(4):355–361. doi:10.1080/10790268.2007.11753951
53. Patel SR, White DP, Malhotra A, Stanchina ML, Ayas NT. Continuous positive airway pressure therapy for treating gess in a diverse population with obstructive sleep apnea: results of a meta-analysis. *Arch Intern Med*. 2003;163(5):565–571. doi:10.1001/archinte.163.5.565
54. Ayas NT, Taylor CM, Laher I. Cardiovascular consequences of obstructive sleep apnea. *Curr Opin Cardiol*. 2016;31(6):599–605. doi:10.1097/HCO.0000000000000329
55. Dickel MJ, Renfrow SD, Moore PT, Berry RB. Rapid eye movement sleep periodic leg movements in patients with spinal cord injury. *Sleep*. 1994;17(8):733–738. doi:10.1093/sleep/17.8.733
56. Manconi M, Ferri R, Zucconi M, et al. Dissociation of periodic leg movements from arousals in restless legs syndrome. *Ann Neurol*. 2012;71(6):834–844. doi:10.1002/ana.23565
57. Ferri R, Proserpio P, Rundo F, et al. Neurophysiological correlates of sleep leg movements in acute spinal cord injury. *Clin Neurophysiol*. 2015;126(2):333–338. doi:10.1016/j.clinph.2014.05.016
58. Salminen AV, Manconi M, Rimpilä V, et al. Disconnection between periodic leg movements and cortical arousals in spinal cord injury. *J Clin Sleep Med*. 2013;9(11):1207–1209. doi:10.5664/jcsm.3174
59. Proserpio P, Lanza A, Sambusida K, et al. Sleep apnea and periodic leg movements in the first year after spinal cord injury. *Sleep Med*. 2015;16(1):59–66. doi:10.1016/j.sleep.2014.07.019
60. Esteves AM, de Mello MT, Lancellotti CL, Natal CL, Tufik S. Occurrence of limb movement during sleep in rats with spinal cord injury. *Brain Res*. 2004;1017(1–2):32–38. doi:10.1016/j.brainres.2004.05.021
61. Telles SC, Alves RC, Chadi G. Periodic limb movements during sleep and restless legs syndrome in patients with ASIA A spinal cord injury. *J Neurol Sci*. 2011;303(1–2):119–123. doi:10.1016/j.jns.2010.12.019

62. Ondo WG. Movements mimicking myoclonus associated with spinal cord pathology: is this a “pure motor restless legs syndrome”. *Tremor Other Hyperkinet Mov (N Y)*. 2012;2.
63. Telles SCL, Alves RSC, Chadi G. Spinal cord injury as a trigger to develop periodic leg movements during sleep: an evolutionary perspective. *Arq Neuropsiquiatr*. 2012;70:880–884. doi:10.1590/s0004-282x2012001100011
64. de Mello MT, Esteves AM, Tufik S. Comparison between dopaminergic agents and physical exercise as treatment for periodic limb movements in patients with spinal cord injury. *Spinal Cord*. 2004;42(4):218–221. doi:10.1038/sj.sc.3101575
65. Yokota T, Hirose K, Tanabe H, Tsukagoshi H. Sleep-related periodic leg movements (nocturnal myoclonus) due to spinal cord lesion. *J Neurol Sci*. 1991;104(1):13–18. doi:10.1016/0022-510x(91)90210-x
66. de Mello MT, Lauro FA, Silva AC, Tufik S. Incidence of periodic leg movements and of the restless legs syndrome during sleep following acute physical activity in spinal cord injury subjects. *Spinal Cord*. 1996;34(5):294–296.
67. Lee MS, Choi YC, Lee SH, Lee SB. Sleep-related periodic leg movements associated with spinal cord lesions. *Mov Disord*. 1996;11(6):719–722. doi:10.1002/mds.870110619
68. Mello MT, Silva AC, Rueda AD, Poyares D, Tufik S. Correlation between K complex, periodic leg movements (PLM), and myoclonus during sleep in paraplegic adults before and after an acute physical activity. *Spinal Cord*. 1997;35(4):248–252.
69. Zeitzer JM, Ayas NT, Shea SA, Brown R, Czeisler CA. Absence of detectable melatonin and preservation of cortisol and thyrotropin rhythms in tetraplegia. *J Clin Endocrinol Metab*. 2000;85(6):2189–2196. doi:10.1210/jcem.85.6.6647
70. Verheggen RJHM, Jones H, Nyakayiru J, et al. Complete absence of evening melatonin increase in tetraplegics. *Faseb J*. 2012;26(7):3059–3064. doi:10.1096/fj.12-205401
71. Spong J, Kennedy GA, Brown DJ, Armstrong SM, Berlowitz DJ. Melatonin supplementation in patients with complete tetraplegia and poor sleep. *Sleep Disord*. 2013;2013(Article ID 128197):8. doi:10.1155/2013/128197
72. Spong J, Kennedy GA, Tseng J, Brown DJ, Armstrong S, Berlowitz DJ. Sleep disruption in tetraplegia: a randomised, double-blind, placebo-controlled crossover trial of 3 mg melatonin. *Spinal Cord*. 2014;52(8):629–634. doi:10.1038/sc.2014.84
73. Kostovski E, Dahm AE, Mowinckel MC, et al. Circadian rhythms of homeostatic factors in tetraplegia: a double-blind, randomized, placebo-controlled cross-over study of melatonin. *Spinal Cord*. 2015;53(4):285–290. doi:10.1038/sc.2014.243
74. Kostovski E, Frigato E, Savikj M, et al. Normalization of disrupted clock gene expression in males with tetraplegia: a crossover randomized placebo-controlled trial of melatonin supplementation. *Spinal Cord*. 2018. doi:10.1038/s41393-018-0176-x
75. Thijssen DHJ, Eijssvogels TMH, Hesse M, Ballak DB, Atkinson G, Hopman MTE. The effects of thoracic and cervical spinal cord lesions on the circadian rhythm of core body temperature. *Chronobiol Int*. 2011;28(2):146–154. doi:10.3109/07420528.2010.540364
76. Berlowitz DJ, Wadsworth B, Ross J. Respiratory problems and management in people with spinal cord injury. *Breathe*. 2016;12(4):328–340. doi:10.1183/20734735.012616
77. Sankari A, Martin JL, Badr MS. Sleep disordered breathing and spinal cord injury: challenges and opportunities. *Curr Sleep Med Rep*. 2017;3(4):272–278. doi:10.1007/s40675-017-0093-0
78. Shafazand S, Anderson KD, Nash MS. Sleep complaints and sleep quality in spinal cord injury: a web-based survey. *J Clin Sleep Med*. 2019. doi:10.5664/jcsm.7760
79. Davenport PW, Chan PY, Zhang W, Chou YL. Detection threshold for inspiratory resistive loads and respiratory-related evoked potentials. *J Appl Physiol*. 2007;102(1):s276–285. doi:10.1152/jappphysiol.01436.2005
80. Graco M, Schembri R, Cross S, et al. Diagnostic accuracy of a two-stage model for detecting obstructive sleep apnoea in chronic tetraplegia. *Thorax*. 2018;73(9):864–871. doi:10.1136/thoraxjnl-2017-211131
81. Bauman KA, Kurili A, Schotland HM, Rodriguez GM, Chiodo AE, Sitrin RG. Simplified approach to diagnosing sleep-disordered breathing and nocturnal hypercapnia in individuals with spinal cord injury. *Arch Phys Med Rehabil*. 2016;97(3):363–371. doi:10.1016/j.apmr.2015.07.026
82. Shojaei MH, Alavinia SM, Craven BC. Management of obesity after spinal cord injury: a systematic review. *J Spinal Cord Med*. 2017;40(6):783–794. doi:10.1080/10790268.2017.1370207
83. Iftikhar IH, Bittencourt L, Youngstedt SD, et al. Comparative efficacy of CPAP, MADs, exercise-training, and dietary weight loss for sleep apnea: a network meta-analysis. *Sleep Med*. 2017;30:7–14. doi:10.1016/j.sleep.2016.06.001
84. Mehrtash M, Bakker JP, Ayas N. Predictors of continuous positive airway pressure adherence in patients with obstructive sleep apnea. *Lung*. 2019;197(2):115–121. doi:10.1007/s00408-018-00193-1
85. Berlowitz DJ, Spong J, Pierce RJ, Ross J, Barnes M, Brown DJ. The feasibility of using auto-titrating continuous positive airway pressure to treat obstructive sleep apnoea after acute tetraplegia. *Spinal Cord*. 2009;47(12):868–873. doi:10.1038/sc.2009.56
86. Graco M, Green SE, Tolson J, et al. Worth the effort? Weighing up the benefit and burden of continuous positive airway pressure therapy for the treatment of obstructive sleep apnoea in chronic tetraplegia. *Spinal Cord*. 2019;57(3):247–254.
87. Le Guen MC, Cistulli PA, Berlowitz DJ. Continuous positive airway pressure requirements in patients with tetraplegia and obstructive sleep apnoea. *Spinal Cord*. 2012;50(11):832–835. doi:10.1038/sc.2012.57
88. Graco M, Berlowitz DJ, Green SE. Understanding the clinical management of obstructive sleep apnoea in tetraplegia: a qualitative study using the theoretical domains framework. *BMC Health Serv Res*. 2019. doi:10.1186/s12913-019-4197-8
89. Berlowitz DJ, Schembri R, Graco M, et al. Positive airway pressure for sleep-disordered breathing in acute quadriplegia: a randomised controlled trial. *Thorax*. 2019;74(3):282–290. doi:10.1136/thoraxjnl-2018-212319
90. Brown JP, Bauman KA, Kurili A, et al. Positive airway pressure therapy for sleep-disordered breathing confers short-term benefits to patients with spinal cord injury despite widely ranging patterns of use. *Spinal Cord*. 2018. doi:10.1038/s41393-018-0077-z
91. Howard ME, Piper AJ, Stevens B, et al. A randomised controlled trial of CPAP versus non-invasive ventilation for initial treatment of obesity hypoventilation syndrome. *Thorax*. 2017;72(5):437–444. doi:10.1136/thoraxjnl-2016-208559
92. Wong KK, Marshall NS, Grunstein RR, Dodd MJ, Rogers NL. Comparing the neurocognitive effects of 40 h sustained wakefulness in patients with untreated OSA and healthy controls. *J Sleep Res*. 2008;17(3):322–330. doi:10.1111/j.1365-2869.2008.00665.x
93. Caples SM, Rowley JA, Prinsell JR, et al. Surgical modifications of the upper airway for obstructive sleep apnea in adults: a systematic review and meta-analysis. *Sleep*. 2010;33(10):1396–1407. doi:10.1093/sleep/33.10.1396
94. Strollo PJ Jr, Soose RJ, Maurer JT, et al. Upper-airway stimulation for obstructive sleep apnea. *N Engl J Med*. 2014;370(2):139–149. doi:10.1056/NEJMoa1308659
95. Ramar K, Dort LC, Katz SG, et al. Clinical practice guideline for the treatment of obstructive sleep apnea and snoring with oral appliance therapy: an update for 2015. *J Clin Sleep Med*. 2015;11(7):773–827. doi:10.5664/jcsm.4858
96. Wijesuriya NS, Eckert DJ, Jordan AS, et al. A randomised controlled trial of nasal decongestant to treat obstructive sleep apnoea in people with cervical spinal cord injury. *Spinal Cord*. 2019. doi:10.1038/s41393-019-0256-6

97. Levy J, Hartley S, Mauruc-Soubirac E, et al. Spasticity or periodic limb movements? *Eur J Phys Rehabil Med.* 2018;54(5):698–704. doi:10.23736/S1973-9087.17.04886-9
98. Qaseem A, Kansagara D, Forcica MA, Cooke M, Denberg TD. Management of chronic insomnia disorder in adults: a clinical practice guideline from the american college of physicians. *Ann Intern Med.* 2016;165(2):125–133. doi:10.7326/M15-2175
99. Mehta S, Orenczuk S, Hansen KT, et al. An evidence-based review of the effectiveness of cognitive behavioral therapy for psychosocial issues post-spinal cord injury. *Rehabil Psychol.* 2011;56(1):15–25. doi:10.1037/a0022743
100. Martin JL, Song Y, Hughes J, et al. A four-session sleep intervention program improves sleep for older adult day health care participants: results of a randomized controlled trial. *Sleep.* 2017;40(8). doi:10.1093/sleep/zsx079
101. Wallace DM, Sawyer AM, Shafazand S. Comorbid insomnia symptoms predict lower 6-month adherence to CPAP in US veterans with obstructive sleep apnea. *Sleep Breath.* 2018;22(1):5–15. doi:10.1007/s11325-017-1605-3
102. Fischer S, Smolnik R, Herms M, Born J, Fehm HL. Melatonin acutely improves the neuroendocrine architecture of sleep in blind individuals. *J Clin Endocrinol Metab.* 2003;88(11):5315–5320. doi:10.1210/jc.2003-030540
103. Skene DJ, Arendt J. Circadian rhythm sleep disorders in the blind and their treatment with melatonin. *Sleep Med.* 2007;8(6):651–655. doi:10.1016/j.sleep.2006.11.013
104. Zeitzer JM, Ku B, Ota D, Kiratli BJ. Randomized controlled trial of pharmacological replacement of melatonin for sleep disruption in individuals with tetraplegia. *J Spinal Cord Med.* 2014;37(1):46–53. doi:10.1179/2045772313Y.0000000099

Nature and Science of Sleep

Dovepress

Publish your work in this journal

Nature and Science of Sleep is an international, peer-reviewed, open access journal covering all aspects of sleep science and sleep medicine, including the neurophysiology and functions of sleep, the genetics of sleep, sleep and society, biological rhythms, dreaming, sleep disorders and therapy, and strategies to optimize healthy sleep.

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/nature-and-science-of-sleep-journal>