

Occurrence, Genetic Diversities And Antibiotic Resistance Profiles Of *Salmonella* Serovars Isolated From Chickens

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Stephen Abiola Akinola ¹
Mulunda Mwanza²
Collins Njie Ateba ¹

¹Antimicrobial Resistance and Phage Bio-Control Group, Department of Microbiology, Faculty of Natural and Agricultural Sciences, North West University, Mmabatho 2745, South Africa;

²Center for Animal Health Studies, Faculty of Natural and Agricultural Sciences, North West University, Mmabatho 2745, South Africa

Purpose: Contamination with *Salmonella* on food products and poultry in particular has been linked to foodborne infections and/or death in humans. This study investigated the occurrence, genetic diversities and antibiotic resistance profiles of *Salmonella* strains isolated from chickens.

Patients and methods: Twenty each duplicate faecal swab samples were collected from five different poultry pens of broilers, layers and indigenous chickens in the North-West Province, South Africa. Isolates identities were confirmed through amplification and sequence analysis of 16S rRNA and the *invA* gene fragments after which phylogenetic tree was constructed. *Salmonella enteritidis* (ATCC:13076TM), *Salmonella* Typhimurium (ATCC:14028TM) and *E. coli* (ATCC:25962TM) were used as positive and negative controls, respectively. The serotypes of *Salmonella* isolates were determined. Antibiotic-resistant profiles of the isolates against eleven antimicrobial agents were determined.

Results: Eighty-four (84%) of representative isolates possessed the *invA* genes. The percent occurrence and diversity of *Salmonella* subspecies in chickens were 1.81–30.9% and was highest in *Salmonella enterica subsp. enterica*. Notably, the following serotypes *Salmonella bongori* (10.09%), *Salmonella* Pullorum (1.81%), *Salmonella* Typhimurium (12.72%), *Salmonella* Weltevreden, *Salmonella* Chingola, *Salmonella* Houten and *Salmonella* Bareilly (1.81%). Isolates (96.6%) displayed multidrug resistance profiles and the identification of isolates with more than nine antibiotic resistance was a cause for concern.

Conclusion: This study indicates that isolates had pre-exposure histories to the antibiotics tested and may pose severe threats to food security and public health.

Keywords: *Salmonella*, diversity, antimicrobial resistance, phylogenetic, chickens

Introduction

Salmonella spp. are enteric pathogens that have received a lot of attention due to their ability to cause food-borne diseases and high rates of mortality amongst humans and thus were declared as agents of public health significance.^{1,2} Salmonellosis is the most common food-borne disease caused by *Salmonella* species in humans with symptoms ranging from headache, vomiting, fatigue, nausea, bloody diarrhea, gastroenteritis, and abdominal cramps and self-limiting for which often no antimicrobials are prescribed for its control.^{3–5} Hence, the pathogen is capable of causing socio-economic and public health implications to humans. *Salmonella enterica* serotype *Typhimurium* (*S.* Typhimurium) and *Salmonella enterica* serotype *Enteritidis* (*S.* Enteritidis) are considered of high

Correspondence: Collins Njie Ateba
Antimicrobial Resistance and Phage Bio-control Group, Department of Microbiology, Faculty of Natural and Agricultural Sciences, North West University, Private Bag X2046, Mmabatho 2745, South Africa
Tel +27 18 489 2720
Email collins.ateba@nwu.ac.za

health importance due to their ability to cause salmonellosis in humans and veterinary animals in both developed and developing countries of the world.

Salmonella has been highlighted as economically important zoonotic pathogens by the World Health Organisation (WHO) and the Food Agriculture Organisation (FAO) dated back to 1950s.¹ *Salmonella* spp. have been enteric pathogens co-existing with pathogens such as *Escherichia coli*, *Klebsiella* spp., and *Proteus* spp.⁶ According to Kagambèga et al,⁵ ruminants such as cattle and sheep, non-ruminants - pigs, dogs, rodents, poultry, birds, and cold-blooded animals such as fish and lizards, and humans have been implicated as reservoirs of the typhoidal and the non-typhoidal *Salmonella* species. However, Poultry and its products are the major sources of *Salmonella*-borne infection in the food chain. The ability of *Salmonella* to be transmitted from reservoirs to other animals and humans calls for concern. Thus, making its survey and control among suspected reservoirs such as chickens is necessary. The influx of many antibiotic resistance strains within the environment calls for a concern. Antibiotic resistance is currently a global problem that poses a threat to public health. Therefore, a study to investigate the occurrence and antibiotic resistance profiles of *Salmonella* among chicken whose carcasses forms a major of the South African cuisines germane.

Materials And Methods

Sample Collection

This study was conducted within farms located at Ngaka Molema Modiri District of Mafikeng, North West Province, South Africa. The study site's (Mafikeng) geographical coordinates are 25° 52' 0" South, 25° 39' 0" East. Twenty samples each were collected in duplicates from five different poultry pens, housing broiler, layer and indigenous chickens in the study area. The broilers and layers belonged to the White Leghorn breed while the indigenous belonged to Potchefstroom koekoek breed (*Gallus gallus domesticus*). Swabs from the gut were aseptically collected in duplicates from test animals and transported on ice to the laboratory for analysis within 24 hrs of collection. Ethical clearance for the study was obtained from the Mafikeng Animal Research Ethics Committee of the North West University prior to the commencement of sampling. Samples were also collected under the supervision of trained Veterinarians and Animal Health Technicians from the Centre for Animal Health Studies, North West University, South Africa.

Isolation Of Microbial Isolates

Isolation of *Salmonella* spp. from chickens was done using ISO-6579:2002 procedure.⁸ Sample pre-enrichment and enrichment were achieved in buffered peptone water and tetrathionate broth, respectively, prior to enrichment in Rappaport vassiliadis Soy (RVS) broth and incubated at 42 °C for 24 hrs. About 1 mL of the inoculated RVS broth was plated on sterile *Salmonella* - *Shigella* Agar (SSA) and was incubated aerobically at 37 °C for 18 hrs. Colonies having creamy with or without black centre on SSA were regarded as presumptive *Salmonella* isolates and were further studied. Sub-culturing was done until pure colonies were obtained.

Morphological And Biochemical Characterization Of Isolates

The morphological and biochemical tests (Gram staining, catalase, Simmons citrate test, urease and Triple sugar iron (TSI) agar) were determined as described by Ateba and Mochaiwa.⁹ Gram-negative rods and catalase-positive colonies were kept on double-strength slants and kept under -20°C for further use.

Molecular Characterisation Of Isolates

The amplification of 16S rRNA region of the bacteria was employed for the discrimination of presumptive *Salmonella* isolates. The DNA was extracted using a Fungal/Bacterial DNA extraction kit (Zymo Research Corporation, Southern California, USA) following the manufacturer's specification. The pure eluted DNA was stored at -80 °C for further analysis. The pure DNA was quantified using a Nanodrop Lite spectrophotometer (Model 1558) obtained from Thermo Scientific, USA, and the genomic DNA was quantified on a 1% agarose gel. The presence of fluorescence band when viewed under the UV Transilluminator (Biorad Gel DocTM XR+) confirmed the presence of DNA of presumptive *Salmonella* isolates.

PCR Amplification Of 16S rRNA

The 16S ribosomal RNA (16S rRNA) PCR was employed in the identification of *Salmonella* isolates.¹⁰ The amplification was done using a Biorad C1000 TouchTM Thermal Cycler. The 27F (5¹-AGAGTTTGATCCTGGCTCAG-3¹) and 1492R (5¹- GGTTACCTTGTTACGACTT - 3¹) primers used were synthesized at Inqaba Biotechnical Industries (Pty) Ltd, South Africa, having an expected

amplicon size of 1450 bp. For the polymerase chain reaction (PCR), a 25 µL reaction mix composed of 12 µL of master mix (Thermo Scientific PCR Master Mix 2X), oligonucleotides (1 µL), DNA template (4 µL) and nuclease-free water (7 µL) was used.¹⁰ The negative controls used include *Aspergillus flavus* and water as a template in the PCR assays while the positive was *Salmonella* Typhimurium ATCC 14028TM.

Amplification Of The *invA* Gene In Presumptive *Salmonella* Isolates

The *invA* gene fragment was amplified using the set of primers *invA* F (5¹-GTGAAATTATCGCCACGTTCCGGG CAA-3¹) and *invA* R 5¹-TCATCGCACCGTCAAAGG AACC -3¹) with expected amplicon size of 284 bp. Slight modifications in the annealing temperature previously reported by Ateba and Mochaiwa⁹ were used: initial denaturation (95 °C for 2 mins), denaturation (95 °C for 15 s), annealing (47.8 °C for 1 min), elongation (72 °C for 45 s) and final elongation (72 °C for 7 mins). A 25 µL reaction mix was used in the amplification, and this is composed of 12.5 µL of master mix (Thermo Scientific PCR Master Mix 2X), oligonucleotides (1 µL), DNA template (4 µL) and nuclease-free water (6.5 µL). A positive control (*Salmonella* Typhimurium ATCC 14028) and a negative control (*Escherichia coli* ATCC 25922 and non-template water) were used.

Gel Electrophoresis Of Amplicons

The molecular weight of PCR amplicons was determined by gel electrophoresis.¹¹ A DNA marker (Fermentas Life Science, Lithuania) of 1 kb was used and the gel was allowed to run at 60 volts, 400 amperes for 60 mins in 1% tris acetate ethylenediamineacetate (TAE) buffer before photographing under the UV transilluminator light (Biorad Gel DocTM XR+).

Gene Sequencing And Identification Of Isolates

The amplified product was sequenced using an automated DNA sequencer (SpectruMedix model SCE 2410) at Inqaba Biotechnical Industries (Pty) Ltd Pretoria. Resulting sequences were cleaned using the FinchTV software version 1.4.0 (Geospiza Inc.) and blasted against on the National Centre for Biotechnology Information¹² database using the Nucleotide Basic Local Alignment Search Tool (BLAST) program (<http://www.ncbi.nlm.nih.gov/BLAST>). Isolates were identified based on the highest percentage of

similarity and sequences were deposited in NCBI gene bank and accession numbers were obtained. The serotypes of presumptive *Salmonella* isolates were determined using the *Salmonella* antisera agglutination kits. Isolates were then classified into serotypes as described in the Kauffman–White *Salmonella* classification.

Phylogenetic Tree Construction

Cleaned sequences were aligned by CLUSTALW sequence alignment tool and de-gapped using Bio-Edit software package.^{13,14} To identify putative close phylogenetic relatives, multiple sequence alignments were obtained using Clustal-W against corresponding nucleotide sequences retrieved from the Gene bank. The evolutionary distance matrices were generated.¹⁵ Phylogenetic analysis was done using the neighbour joining method¹⁶ in MEGA program version 5.10.¹⁷ The bootstrap analysis was done using 1000 replications for neighbour joining. The sequences were checked for putative chimeric artefacts using the Chimera-Buster program and then manipulation and tree editing was done using the Tree View option.¹⁸ *Salmonella enterica* was used as the root to the tree.

Determination Of Antibiotics Resistance Profile Of *Salmonella* Isolates

Antibiotic sensitivity of *Salmonella* isolates was investigated against eleven antibiotics belonging to eight different classes using the disc diffusion method.¹⁹ Antibiotics used include; ampicillin (10 µg), oxy-tetracycline (30 µg), ciprofloxacin (5 µg), streptomycin (10 µg), gentamicin (10 µg), sulphamethoxazole/trimethoprim (300 µg), chloramphenicol (30 µg), erythromycin (15 µg), norfloxacin (10 µg), cephalothin (30 µg), and nalidixic acid (30 µg). Antibiotics discs were placed at an equilateral distance to each other on Muller–Hinton agar (MHA) plates and were incubated at 37 °C for 18 hrs. After incubation, zones of inhibition around the antibiotics disc were measured using a meter rule graduated in millimetres. The test was made in triplicate and the mean diameter of the inhibitory zones (IZD) were calculated. The mean IZD was determined as either susceptible, intermediate, or resistant using the Clinical and Laboratory Standards Institute²⁰ criteria. The multiple antibiotics resistance (MAR) phenotypes were recorded for isolates showing resistance to more than two antibiotics²¹ and the MAR index was calculated as shown in Equation 1.²²

$$\text{MAR} = \frac{\text{Number of resistance to antibiotics/}}{\text{Total number of antibiotics tested}} \quad (1)$$

Clustering Of Antibiotic-Resistant Patterns Of *Salmonella* Isolates

To determine the similarities and differences between *Salmonella* isolates from different sources based on their antibiotic resistance patterns, cluster analysis was done. The IZDs of *Salmonella* strains were clustered using a cluster analysis on the Statistica software package (Statsoft, USA) and a dendrogram was generated. Ward's method and the Euclidean distance method were used to generate the clusters.

Statistical Analysis

The statistical analysis of data generated was evaluated using Statistical package for Social Sciences (SPSS, version 21.0 IBM Corp., USA). The frequency and percentage of occurrence of isolates and correlations between isolates antibiotics resistance and sources were determined using Pearson's product. The cluster analysis of antibiotics sensitivity patterns of *Salmonella* isolates was evaluated through the Ward's algorithm and Euclidean distances on the Statistica software version 7.0 (Statsoft, USA). Significance and goodness of fit were evaluated at 95% confidence interval while sequence algorithms were cleaned and processed using FinchTv, Bioedit and the phylogenetic tree was constructed using the MEGA6 Software's.

Results

The morphological and biochemical characteristics of presumptive *Salmonella* isolates from chickens in Mafikeng community, South Africa, is as presented in [Supplementary material S1](#). Colonies pigment morphology ranged from pink to colourless with/without black centre, on *Salmonella Shigella* Agar. As shown in [Supplementary material A1](#), 96 percent of colonies

had a circular shape while the opacity ranged from 83.63% (opaque) and translucent (16.36%). All selected isolates were gram-negative rods having the ability to hydrolyze hydrogen peroxide in the production of catalase enzyme. Isolates showed alkalinity by a red colour pigment on slants, yellow butt with or without gas, thus signifying acid production while black pigment in butt showed hydrogen sulphide production which is typical of *Salmonella*. About 81.81% of the isolates were positive to alkalinity, 16.36% were negative while 3.63% had weak alkalinity reaction while 96.36% were able to produce acid. A number of 96.36% of the presumptive isolates had the ability to utilize citrate as a sole source of carbon and energy, while 98.18% tested negative to urease and indole production.

[Figure 1](#) presents the gel picture of 16 S rRNA amplification of representative presumptive *Salmonella* isolates from chickens. The 16 S rRNA amplification was performed twice to ensure reliability of obtained results. There was a 100% positive amplification at an expected band size of 1450 bp. There was a positive amplification of *Salmonella* Typhimurium ATCC 14028 in lane 1 while no amplification was observed in lane 14 (negative control). The positive amplification confirms the use of 27F and 1497R sets of oligonucleotides for 16 S rRNA region amplification in enteric bacteria.

Salmonella-specific PCR was conducted using the *invA* genes. As shown in [Figure 2](#), about 87.27% of the representative isolates showed positive amplification while about 12.72% were negative. As shown in [Table 1](#), the percent similarity of isolates to data in the NCBI gene bank ranged from 85% to 99%. In percent, a portion of 26% had 99% similarity while 85% had 94% similarity to *Salmonella*. The percent occurrence




Figure 1 Agarose gel (1%) electrophoresis showing the amplification of 16 S rDNA of representative presumptive bacteria isolates colonising the gut of chickens obtained from North-west province, South Africa.

Notes: Lane 1 to 29 shows the amplification of 16 S region of presumptive *Salmonella* DNA isolates while lanes 14 = no template (negative control), lane 13 = *Salmonella* Typhimurium ATCC 14028 (positive control).

Abbreviations: M, DNA marker (1kb); bp, base pairs.




Figure 2 Agarose gel (1%) electrophoresis showing the amplification of *invA* gene in representative presumptive *Salmonella* isolates in chickens obtained from Mafikeng, North-west province, South Africa.

Notes: Lane 1 = no template (negative control), Lane 2 = *Salmonella* Typhimurium ATCC 14028TM (positive control); Lane 6 = *Escherichia coli* ATCC 259622TM (negative control), lane 2 to 18 = positive amplification of *invA* gene at 286 bp, lane 6 = no amplification.

Abbreviations: M, DNA marker (100 bp); bp, base pairs.

of *Salmonella* in chicken is presented in Figure 3. The percent occurrence based on subspecies ranged from 2% to 61% (Supplementary material S2). The highest occurring subspecies belongs to the *Salmonella enterica* subsp. *enterica* (61%) while the least occurring serotype was *Salmonella* Salamae, *Salmonella* Weltevreden, *Salmonella* Chingola, *Salmonella* Houten, *Salmonella* Bareilly (2%). Based on source, *Salmonella* Typhimurium was highest in indigenous chickens (9.08%) followed by layers (3.63%), while in broilers, *Salmonella* Arizona was highest as described in Figure 3. *Salmonella* Salamae was not isolated in the indigenous and broiler chickens from the study site. Likewise, in layers, *Salmonella* Arizona and *Salmonella* Weltevreden were not isolated except in indigenous breeds and in broiler chickens. Autoagglutination was obtained in some *Salmonella bongori* and *Salmonella enterica* subspecies *enterica* isolates, hence the inability to determine the serotypes of these isolates. Non-specific agglutination as a product of loss of antigen expression could give pseudo-positive results as earlier reported by.^{23,24}

Based on the cluster algorithm of the Neighbour Joining method used, the percent evolutionary relatedness of *Salmonella* isolates is presented in Figure 4. The evolutionary distance of *Salmonella* isolates was 35.13655429. Most of the *Salmonella* isolates were found to evolve from the same ancestral origin with similarities higher than 70% and comparable to strains sourced from the gene bank. *Salmonella enterica* subsp. *enterica* (MG663457, MG663509, MG663461 and MG663502) were found to evolve from the same ancestor which we presumed to be *Salmonella* spp. However, a genetic evolution was observed in the isolates with a 72% homology compared to MG663500, MG663459,

and MG663456 having 99% homology to the genetic sequences of the parent's genome.

Also, *Salmonella bongori* (MG663487) exhibited a 100% concatenated homology with other *Salmonella* isolates. All the comparable sequences from gene bank, *Salmonella* spp. (KU641443), *Salmonella* Arizonae (CP006693) and *Salmonella bongori* (KR350635), showed relatedness and were comparable to sequences identified as *Salmonella bongori* (MG663486), MG663492 and *Salmonella* Blockley (MGG3495). Isolates *Salmonella enterica* subsp. *enterica* (MG663485, MG663489, MG663485 and MG663462) had 100% evolutionary relation to the parental genus. *Salmonella* Houten (MG663464) and *Salmonella* Heidelberg (MG663483) clustered together on the same cladograph. *Salmonella enterica* subsp. *enterica* (MG663464) had 100% homology to an out-group (*Salmonella enterica* subsp. *enterica* KY656601). However, *Salmonella enterica* subsp. *enterica* (MG663468) was similar to *Salmonella* Enteritidis (CP018655) and had a 94% homology to *Salmonella* Typhimurium (MG663465). *Salmonella* Typhimurium (MG663510, MH086979 and MG663473) had the same node showing that they both evolved from the same ancestor.

Table 2 presents the antibiotic sensitivity profile of *Salmonella* isolates from chickens in Mafikeng. About 56% of the total *Salmonella* isolates were resistant to ampicillin treatment, 18% had intermediate resistance while 26% were susceptible (Supplementary material S3). The ampicillin-resistant strains were found more in the indigenous chickens (81%) followed by broilers (36%) and lowest in the layers (27%). About 69% of all *Salmonella* isolates were resistant to oxy-tetracycline with 9% being intermediate-resistant. Higher occurrence of oxy-tetracycline

Table 1 Identity Of Presumptive *Salmonella* Isolates From Chickens In Mafikeng, South Africa

Isolate Number	Sample Source	Sequence	Assession Number	InvA	Serotype	Similarity (%)	Name of Organism
1	Broiler	Seq1	MG663456	+VE	AAG	98	<i>Salmonella enterica</i> subsp. <i>enterica</i>
2	Broiler	Seq2	MG663457	+VE	AAG	92	<i>Salmonella enterica</i> subsp. <i>enterica</i>
3	Broiler	Seq3	MG663458	+VE	AAG	97	<i>Salmonella enterica</i> subsp. <i>enterica</i>
4	Broiler	Seq4	MG663459	+VE	AG	92	<i>Salmonella enterica</i> ser. Weltevreden
5	Broiler	Seq5	MG663460	+VE	AG	92	<i>Salmonella enterica</i> ser. Chingola
6	Broiler	Seq6	MG663461	+VE	AG	92	<i>Salmonella enterica</i> ser. Arizonae
7	Broiler	Seq7	MG663462	+VE	AG	98	<i>Salmonella enterica</i> ser. Bovismorbificans
8	Layer	Seq8	MG663463	+VE	AAG	99	<i>Salmonella enterica</i> subsp. <i>enterica</i>
9	Layer	Seq9	MG663464	+VE	AAG	92	<i>Salmonella enterica</i> subsp. <i>enterica</i>
10	Layer	Seq10	MG663465	+VE	AG	97	<i>Salmonella enterica</i> ser. Typhimurium
11	Layer	Seq11	MG663466	+VE	AG	92	<i>Salmonella enterica</i> ser. Salamae
12	Layer	Seq12	MG663467	+VE	AG	99	<i>Salmonella enterica</i> ser. Houten
13	Layer	Seq13	MG663468	+VE	AAG	99	<i>Salmonella enterica</i> subsp. <i>enterica</i>
14	Indigenous	Seq14	MG663469	+VE	AAG	98	<i>Salmonella enterica</i> ser. Bareilly
15	Indigenous	Seq15	MG663470	+VE	AAG	99	<i>Salmonella enterica</i> subsp. <i>enterica</i>
16	Indigenous	Seq16	MG663471	+VE	AAG	98	<i>Salmonella enterica</i> subsp. <i>enterica</i>
17	Indigenous	Seq17	MG663472	+VE	AAG	92	<i>Salmonella enterica</i> subsp. <i>enterica</i>
18	Indigenous	Seq18	MG663473	+VE	AG	92	<i>Salmonella enterica</i> ser. Heidelberg
19	Indigenous	Seq19	MG663474	+VE	AG	92	<i>Salmonella enterica</i> ser. Arizonae
20	Indigenous	Seq20	MG663475	+VE	AAG	99	<i>Salmonella enterica</i> subsp. <i>enterica</i>
21	Indigenous	Seq21	MG663476	+VE	AG	99	<i>Salmonella enterica</i> ser. India
22	Indigenous	Seq22	MG663477	+VE	AG	97	<i>Salmonella enterica</i> ser. Crossness
23	Indigenous	Seq23	MG663478	+VE	AG	97	<i>Salmonella enterica</i> ser. Albany
24	Indigenous	Seq24	MG663479	+VE	AG	99	<i>Salmonella enterica</i> ser. Yovokome
25	Indigenous	Seq25	MG663480	+VE	AG	98	<i>Salmonella enterica</i> ser. Pullorum
26	Indigenous	Seq26	MG663481	+VE	AG	98	<i>Salmonella enterica</i> ser. Infantis
27	Broiler	Seq27	MG663482	+VE	AG	92	<i>Salmonella enterica</i> ser. Arizonae
28	Broiler	Seq28	MG663483	+VE	AG	99	<i>Salmonella enterica</i> ser. Heidelberg
29	Broiler	Seq29	MG663484	+VE	AAG	92	<i>Salmonella enterica</i> subsp. <i>enterica</i>
30	Broiler	Seq30	MG663485	+VE	AAG	92	<i>Salmonella enterica</i> subsp. <i>enterica</i>
31	Broiler	Seq31	MG663486	-VE	AAG	92	<i>Salmonella bongori</i>
32	Broiler	Seq32	MG663487	-VE	AAG	92	<i>Salmonella bongori</i>
33	Broiler	Seq33	MG663488	+VE	AG	92	<i>Salmonella enterica</i> ser. Arizonae
34	Layer	Seq34	MG663489	+VE	AAG	92	<i>Salmonella enterica</i> subsp. <i>enterica</i>
35	Layer	Seq35	MG663490	+VE	AG	92	<i>Salmonella enterica</i> ser. Wandsworth
36	Layer	Seq36	MG663491	+VE	AAG	92	<i>Salmonella enterica</i> subsp. <i>enterica</i>
37	Layer	Seq37	MG663492	-VE	AAG	92	<i>Salmonella bongori</i>
38	Layer	Seq38	MG663493	+VE	AG	92	<i>Salmonella enterica</i> ser. Kentucky
39	Layer	Seq39	MG663494	-VE	AAG	92	<i>Salmonella bongori</i>
40	Layer	Seq40	MG663495	+VE	AG	94	<i>Salmonella enterica</i> ser. Blockley
41	Layer	Seq41	MG663496	+VE	AG	98	<i>Salmonella enterica</i> ser. Newport
42	Layer	Seq42	MG663497	+VE	AG	98	<i>Salmonella enterica</i> ser. Typhimurium
43	Indigenous	Seq43	MG663498	-VE	AAG	98	<i>Salmonella bongori</i>
44	Indigenous	Seq44	MG663499	+VE	AAG	99	<i>Salmonella enterica</i> ser. Manchester
46	Indigenous	Seq46	MG663500	+VE	AAG	98	<i>Salmonella enterica</i> subsp. <i>enterica</i>
47	Indigenous	Seq47	MG663501	+VE	AG	99	<i>Salmonella enterica</i> ser. Typhimurium
48	Indigenous	Seq48	MG663502	+VE	AAG	92	<i>Salmonella enterica</i> subsp. <i>enterica</i>
49	Indigenous	Seq49	MG663503	+VE	AG	99	<i>Salmonella enterica</i> ser. Typhimurium
50	Indigenous	Seq50	MG663504	+VE	AG	85	<i>Salmonella enterica</i> ser. Typhimurium
51	Indigenous	Seq51	MG663505	+VE	AG	97	<i>Salmonella enterica</i> ser. Typhimurium

(Continued)

Table 1 (Continued).

Isolate Number	Sample Source	Sequence	Assession Number	InvA	Serotype	Similarity (%)	Name of Organism
52	Indigenous	Seq52	MG663506	+VE	AAG	99	<i>Salmonella enterica</i> ser. Koessen
53	Indigenous	Seq53	MG663507	+VE	AAG	98	<i>Salmonella bongori</i>
54	Indigenous	Seq54	MG663508	-VE	AG	99	<i>Salmonella enterica</i> ser. Blegdam
55	Indigenous	Seq55	MG663509	+VE	AAG	92	<i>Salmonella enterica</i> subsp. <i>enterica</i>
Control 1	Control 1		ATCC 14028 TM	+VE	AG		<i>Salmonella enterica</i> ser. Typhimurium
Control 2	Control 2		MG663511	+VE		99	<i>Escherichia coli</i> O157:H7

Notes: Lane 1 = control 1 = *Salmonella* Typhimurium (positive control); control 2 = *Escherichia coli* (negative control) was an environmental strain.

Abbreviations: +VE, positive amplification; -VE, negative amplification; AAG, auto-agglutination against antisera; AG, positive agglutination.

resistance was obtained in indigenous chickens (65%). More than 30% of the isolates were resistant to ciprofloxacin while about 20% had intermediate resistance with a distribution largest (46%) in the indigenous chickens. Ninety-five percent resistance to streptomycin was obtained in this study and was dominant in the layers. The percent resistance to trimethoprim/sulphamethoxazole ranged from 64% to 84% in the different samples investigated. When exposed to chloramphenicol, only a small proportion (8–20%) of isolates were resistant to this drug. Against erythromycin, an 100% resistance was observed and was found not to depend on sample source. Hence, the use of erythromycin in the treatment of *Salmonella*-borne infection should be avoided.

The multiple antibiotic resistance index and antibiotic-resistant phenotypes of *Salmonella* isolate are presented in Table 3. The MAR index ranged from 0.27 to 0.81 and was highest in *Salmonella* Weltevreden (AMP-OXT-STR-SXT-C30-ERY-NOR-KF-NAL), *Salmonella* Pullorum (AMP-OXT-CIP-STR-SXT-ERY-NOR-KF-NAL) and *Salmonella* Typhimurium (AMP-OXT-CIP-STR-SXT-ERY-NOR-KF-NAL) with resistance against nine different groups of antibiotics investigated. However, the MAR index of isolates was lowest in *Salmonella enterica* subsp. *enterica*, *Salmonella* Arizonae, *Salmonella* Albany and *Salmonella* Heidelberg strains having resistance to only three groups of the antibiotics studied.




Figure 3 Percentage occurrence of *Salmonella* isolates based on its source.




Figure 4 Neighbor joining method of phylogenetic tree based on partial 16S rDNA gene sequence, showing the phylogenetic relationships between *Salmonella* species and the most closely related strains from the genbank.

Notes: Numbers at the nodes indicate the ranks of bootstrap based on 1000 resampled data sets, and the cut-off points were placed at 70% for condensed tree. The scale bar indicates 0.5 base substitution per site. *Salmonella enterica* were set as the out-group. Sequences obtained in this study are denoted with a circle shape.

Salmonella bongori strains from the indigenous chickens had the highest multiple resistance index of 0.72 with a wide range of resistant phenotypes (AMP-OXT-CIP-STR-SXT-ERY-NOR-KF) while similar strains from broilers (0.3) had the lowest MAR index with resistance

phenotype patterns (OXT-STR-ERY-NAL). Also, a large portion of the *Salmonella* spp. and *Salmonella enterica subsp. enterica* had a hepta-multi-antibiotic-resistant patterns (AMP-OXT-CIP-STR-SXT-ERY-NOR) with MAR of 0.63 majorly from the indigenous chickens.

Table 2 Prevalence Of Antibiotic Resistance And Multi-Drug Resistance Of *Salmonella* Isolates From Chicken Sourced From Mafikeng, South Africa

Antibiotics	Resistance				Intermediate Resistance			
	Total n= 55 (%)	Broilers n =14 (%)	Layers n =15 (%)	Indigenous n = 26 (%)	Total n = 55 (%)	Broilers n = 14 (%)	Layers n = 15 (%)	Indigenous n = 26 (%)
AMP	31 (56)	5 (36)	4 (27)	21 (81)	10 (18)	4 (29)	5 (33)	1 (4)
OXT	38 (69)	12 (86)	9 (60)	17 (65)	9 (16)	1 (7)	4 (27)	4 (15)
CIP	17 (31)	3 (21)	5 (33)	10 (39)	11 (20)	5 (36)	3 (20)	2 (8)
STR	52 (95)	13 (93)	15 (100)	25 (96)	2 (4)	1 (7)	0 (0)	1 (4)
GCN	2 (4)	0 (0)	1 (7)	1 (4)	1 (2)	1 (7)	0 (0)	0 (0)
SXT	43 (78)	9 (64)	12 (80)	22 (85)	0 (0)	0 (0)	0 (0)	0 (0)
C30	7 (13)	2 (14)	3 (20)	2 (8)	8 (15)	4 (29)	2 (13)	2 (8)
ERY	55 (100)	14 (100)	15 (100)	26 (100)	0 (0)	0 (0)	0 (0)	0 (0)
NOR	27 (49)	7 (50)	6 (40)	14 (54)	5 (9)	1 (7)	0 (0)	4 (15)
KF	19 (36)	8 (57)	3 (20)	8 (31)	15 (27)	0 (0)	7 (47)	8 (31)
NAL	26 (47)	8 (57)	9 (60)	9 (35)	5 (9)	0 (0)	1 (7)	4 (15)
Number of MAR phenotypes								
0	0 (0)	0 (0)	0 (0)	0 (0)				
1	0 (0)	0 (0)	0 (0)	0 (0)				
2	1 (2)	1 (7)	0 (0)	0 (0)				
3	4 (7)	0 (0)	0 (0)	4 (15)				
4	13 (24)	4 (29)	4 (27)	5 (19)				
5	13 (24)	3 (21)	1 (4)	9 (35)				
6	4 (7)	2 (14)	0 (0)	2 (8)				
7	8 (15)	3 (21)	0 (0)	5 (19)				
8	9 (17)	1 (7)	0 (0)	8 (31)				
9	3 (6)	1 (7)	0 (0)	2 (8)				

Notes: Ampicillin (AMP) 10 µg; chloramphenicol (C30) 30 µg; nalidixic acid (NAL) 30 µg; streptomycin (STR) 10 µg; oxy-tetracycline (OXT) 30 µg; cephalothin (KF) 30 µg; erythromycin (ERY) 15 µg; sulphamethoxazole/trimethoprim (SXT) 22 µg; gentamycin (GCN) 10µg; ciprofloxacin (CIP) 10 µg; norfloxacin (NOR) 10 µg.

Abbreviation: MAR, multiple antibiotic resistance.

Salmonella Koessen, *Salmonella* India, *Salmonella* Crossness, *Salmonella* Yovokome (AMP-STR-SXT-ERY-KF) had penta-resistant phenotype patterns.

An octa-antibiotics resistance was obtained in *Salmonella* Houten, *Salmonella* Bovismorbificans, *Salmonella* Blegdam, *Salmonella* Typhimurium and *Salmonella bongori*. *Salmonella* Typhimurium isolated in this study were found to belong to the indigenous chickens only, with a MAR index ranging from 0.72 to 0.81. This is indicative of a high multi-antibiotic resistance profiles against the main streams of antibiotics often prescribed in the treatment of *Salmonella* infections in both humans and animals. All octa-antibiotic-resistant strains had resistance to nalidixic acid except *Salmonella* Blegdam and *Salmonella bongori*.

The relatedness and differences between the MAR-resistant strains of *Salmonella* are as shown in Figure 5. Three (3) major clusters (Clusters I, II and III) were

observed and were traced to the source of isolates as described in Table 5. A total of 18 isolates clustered in cluster I, while in cluster II (17) and cluster III (20).⁷ Cluster III recorded the highest distribution of *Salmonella* isolates. Table 4 presents the percentage distribution of *Salmonella* based on sample source and antibiotics resistance clustering patterns. The percentage distribution of *Salmonella* isolates based on antibiotics resistance clusters ranged from 15% to 73.3%. The distribution of resistant strains in cluster I ranged from 16.6% to 73.3% and the largest proportion were from indigenous chickens (11; 73.3%). In clusters I and III, respectively, isolates from layers (22.2%; 35%) and indigenous chickens (73.3%; 50%) had the highest relatedness in terms of antibiotic resistance patterns as opposed to isolates from broilers. However, there was no significant difference between isolates from layers, clustered in clusters II (4; 23%) and I (4; 22.2%) at $P \geq 0.05$. The

Table 3 Multiple Antibiotic Resistance Index And Phenotype Pattern Of *Salmonella* Isolates From Chickens In Mafikeng, South Africa

<i>Salmonella</i> Strains	Sample Source	No. Of Strains	Antibiotics Resistance Profiles	MAR Index
<i>Salmonella bongori</i>	Indigenous	1	AMP-CIP-STR-SXT-ERY	0.45
	Indigenous	1	AMP-OXT-CIP-STR-SXT-ERY-NOR-KF	0.72
	Broilers, layers	2	OXT-CIP-STR-ERY-NOR-NAL	0.54
	Indigenous	1	OXT-CIP-STR-SXT-ERY-NOR-NAL	0.63
	Broiler	1	OXT-STR-ERY-NAL	0.36
<i>Salmonella enterica subsp. enterica</i>	Indigenous	1	OXT-STR-ERY-NOR	0.36
	Broiler	1	OXT-STR-SXT-ERY	0.36
	Layer	1	OXT-STR-SXT-ERY-NAL	0.45
	Indigenous	1	OXT-STR-SXT-ERY-NOR	0.45
	Indigenous	1	AMP-OXT-CIP-STR-SXT-ERY-NOR	0.63
	Indigenous	1	AMP-OXT-CIP-STR-SXT-C30-ERY	0.63
	Broiler	1	AMP-SXT- ERY-KF	0.36
	Indigenous	1	AMP-SXT-ERY	0.27
	Indigenous, layer	2	AMP-STR-SXT-ERY	0.36
	Broiler	1	AMP-STR-SXT-ERY-KF	0.45
	Indigenous	1	AMP-STR-SXT-ERY-NOR	0.45
	Broiler	1	OXT-CIP-STR-SXT-ERY-NOR-NAL	0.63
	Broiler	1	OXT-STR-ERY-KF	0.36
	Indigenous	1	OXT-STR-ERY-NAL	0.36
	Layer	1	OXT-STR-ERY-NOR	0.36
	Indigenous	1	STR-SXT-C30-ERY-KF-NAL	0.54
<i>Salmonella enterica ser. Arizonae</i>	Indigenous	1	AMP-STR-ERY	0.27
	Broiler	1	OXT-CIP-STR-SXT-ERY-NOR-NAL	0.63
	Broiler	1	OXT-STR-ERY-NOR-KF-NAL	0.54
	Broiler	1	OXT-STR-SXT-ERY-KF	0.45
<i>Salmonella enterica ser. Typhimurium</i>	Indigenous	1	AMP-OXT-CIP-STR-SXT-ERY-NOR-KF	0.72
	Indigenous	1	AMP-OXT-CIP-STR-SXT-ERY-NOR-NAL	0.72
	Indigenous	1	AMP-OXT-STR-SXT-ERY-NOR-KF-NAL	0.72
	Indigenous	1	AMP-STR-GCN-SXT-C30-ERY-NOR-NAL	0.72
	Indigenous	1	AMP-OXT-CIP-STR-SXT-ERY-NOR-KF-NAL	0.81
	Indigenous	1	STR-SXT-ERY-KF	0.36
	Indigenous	1	STR-SXT-ERY-KF-NAL	0.45
<i>Salmonella enterica ser. Heidelberg</i>	Broiler	1	AMP-OXT-STR-SXT-C30-ERY-KF	0.63
	Indigenous	1	OXT-STR-ERY	0.27
<i>Salmonella enterica ser. Manchester</i>	Indigenous	1	AMP-CIP-STR-SXT-ERY-NOR-NAL	0.63
<i>Salmonella enterica ser. Weltevreden</i>	Broiler	1	AMP-OXT-STR-SXT-C30-ERY-NOR-KF-NAL	0.81
<i>Salmonella enterica ser. Salamae</i>	Indigenous	1	OXT-STR-SXT-ERY-NOR-NAL	0.54
<i>Salmonella enterica ser. Blegdam</i>	Indigenous	1	AMP-OXT-CIP-STR-SXT-ERY-NOR-KF	0.72
<i>Salmonella enterica ser. Pullorum</i>	Indigenous	1	AMP-OXT-CIP-STR-SXT-ERY-NOR-KF-NAL	0.81
<i>Salmonella enterica ser. Koessen</i>	Indigenous	1	AMP-OXT-STR-SXT-ERY	0.45
<i>Salmonella enterica ser. India</i>	Indigenous	1	AMP-OXT-STR-SXT-ERY	0.45
<i>Salmonella enterica ser. Crossness</i>	Indigenous	1	AMP-OXT-STR-SXT-ERY	0.45
<i>Salmonella enterica ser. Bovismorbificans</i>	Broiler	1	AMP-OXT-STR-SXT-ERY-NOR-KF-NAL	0.72

(Continued)

Table 3 (Continued).

Salmonella Strains	Sample Source	No. Of Strains	Antibiotics Resistance Profiles	MAR Index
<i>Salmonella enterica</i> ser. Infantis	Indigenous	1	AMP-OXT-STR-SXT-ERY-NOR-NAL	0.63
<i>Salmonella enterica</i> ser. Albany	Indigenous	1	AMP-STR-ERY	0.27
<i>Salmonella enterica</i> ser. Newport	Layer	1	AMP-STR-SXT-ERY	0.36
<i>Salmonella enterica</i> ser. Yovokome	Indigenous	1	AMP-STR-SXT-ERY-KF	0.45
<i>Salmonella enterica</i> ser. Wandsworth	Indigenous	1	OXT-CIP-STR-SXT-ERY-NOR-NAL	0.63
<i>Salmonella enterica</i> ser. Kentucky	Layer	1	OXT-STR-ERY-NAL	0.36
<i>Salmonella enterica</i> ser. Blockley	Indigenous	1	STR-SXT-ERY-KF	0.36
<i>Salmonella enterica</i> ser. Chingola	Broiler	1	OXT-STR-ERY-NOR-NAL	0.45
<i>Salmonella enterica</i> ser. Bareilly	Indigenous	1	OXT-STR-SXT-ERY-NAL	0.45
<i>Salmonella enterica</i> ser. Houten	Indigenous	1	AMP-OXT-CIP-STR-S300-ERY-NOR-NAL	0.72

Note: Ampicillin⁵³ 10 µg; chloramphenicol (C30) 30 µg; nalidixic acid (NAL) 30 µg; streptomycin (STR) 10 µg; oxy-tetracycline (OXT) 30 µg; cephalothin (KF) 30 µg; erythromycin (ERY) 15 µg; sulphamethoxazole/trimethoprim (SXT) 22 µg; gentamycin (GCN) 10µg; ciprofloxacin (CIP) 10 µg; norfloxacin (NOR) 10 µg.

Abbreviation: MAR, multi-antibiotic resistance index.

percent distribution of *Salmonella* isolates from indigenous chickens ranged from 29.4% to 73.3%. As shown in Table 5, there exists a positive correlation in the

antibiotic-resistance patterns of *Salmonella* isolates from layers and broilers while a negative correlation was obtained against indigenous chickens.




Figure 5 Dendrogram of antibiotic resistance profiles of *Salmonella* strains isolated from chickens in Mafikeng, South Africa, using cluster analysis.

Notes: Seq represents sequence numbering of *Salmonella* isolates from different types of chickens in Mafikeng. The tree was constructed using the Ward's method and Euclidean distances in the Statistica version 7 software (Statsoft US).

Abbreviation: Seq, sequences of *Salmonella* isolates from chickens.

Table 4 The Percent Distribution Of Resistant *Salmonella* Isolates Based On Sample Source And Antibiotic Resistant Clusters

Type of Chicken/ Source	Cluster I N = 18	Cluster II N = 17	Cluster III N = 20
Broiler	3 (16.6%)	8 (47.1%)	3 (15%)
Layers	4 (22.2%)	4 (23.5%)	7 (35%)
Indigenous	11 (73.3%)	5 (29.4%)	10 (50%)

Table 5 Pearson Correlations Between Percent Antibiotic Resistance Of *Salmonella* Isolates And Source

Antibiotic Resistance	Total Resistance (%)	Broilers (%)	Layers (%)	Indigenous (%)
Total resistance (%)	1			
Broilers (%)	0.925**	1		
Layers (%)	0.931**	0.874**	1	
Indigenous (%)	0.959**	0.800**	0.825**	1

Note: **Correlation is significant at the 0.01 level (2-tailed).

Discussion

The morphological characteristics observed in this study support the previous observation contained in the WHO Global Salm-Surv as described by Hendriksen et al.²⁵ The triple sugar iron test of the presumptive *Salmonella* isolates depicts their ability to utilize lactose, saccharose and dextrose sugars. A positive to indole test depicts the ability of isolates to utilize amino acid in the form of tryptophan to produce the enzyme indole. However, some isolates showed a positive reaction to urease and indole which contradicts the expected results stipulated in the Bergey's Manual of Determinative Bacteriology. Nevertheless, a similar variation has been reported by Shan et al.²⁶ However, the observation in this study might be due to a shift in the nutrient utilisation pattern of *Salmonella* as a result of ecological stress emanating from competition for food and other stress inducers. Hence, biochemical characteristics might not be adequate to effectively discriminate a microbial community, hence the need for the use of more reliable approaches such as the molecular techniques. The PCR discrimination method was effective in the discrimination of *Salmonella spp.* as opposed to the use of biochemical and morphological characteristics as obtained from this study. Therefore, the polymerase chain

reaction could present a rapid, sensitive and reliable method for pathogen detection.

The positive amplification of the *invA* genes in *Salmonella* isolates supports the findings of previous authors on the presence of invasive genes in *Salmonella*.²⁷ Demonstrated the presence and functionality of *invA*, B and C genes which are regions of high similarity in diverse *Salmonella* serovars except in *Salmonella arizona* in which *invD* gene regions were detected. The presence of *invA* genes has been reported in *Salmonella* isolates from broiler chickens in Iran,²⁸ poultry, pigs, humans and other food commodities in Brazil.²⁹ The use of *invA* genes has been regarded as the most reliable in *Salmonella* discrimination since many possess the *invA* gene within their genomes. Therefore, it is pertinent in the tracking of the pathogenesis of *Salmonella*-borne infections in animals and humans. The virulence of *Salmonella* in hosts has been linked to their ability to invade the epithelial tissues. On ingestion, *Salmonella* attaches itself to the intestinal mucosa lining contributing to a decrease in the pH of the gastrointestinal tract, thus causing an irritation. Invasive *Salmonella* species could deplete the mucosa layer by penetrating through the M cells overlying the Peyer's patches.³⁰ In some patients, this situation may progress to a systemic infection resulting from the invasion of the intestinal lymphoid follicles by *Salmonella* strains which presents clinical signs associated with drained mesenteric lymph nodes.

From the study, some isolates were found not to possess the *invA* gene, thus implying their inability to cause infections in hosts. However, the occurrence of the invasive *Salmonella* isolates among the chicken samples within the Mafikeng Community suggests that consumers and other stakeholders within the food and value chain might be at a risk of *Salmonella*-borne infections. This can hamper the safety and health of both veterinary and humans and the socioeconomic status of the people living in Mafikeng community, South Africa.

Salmonella species such as *Salmonella Bovismorbificans* was isolated from this study as opposed to the previous report of it been found only in humans.³¹ Many diverse *Salmonella* strains identified in this study have been implicated to possess the extended spectrum of the β -lactamases (ESBLs) enzymes coding for antibiotic-resistant genes and have been linked to salmonellosis in humans.³¹⁻³⁴ The isolation of these strains from chickens may have resulted from human-to-animal interaction along the available interfaces

such as contaminated feed, water, handling, infected hosts (rodents) and animal care personnel on the farms.

The dominance of *Salmonella enterica* species in chickens supports the findings of³⁵ who reported a high occurrence of *Salmonella* Typhimurium compared to the serotype *Salmonella* Enteritidis in a study conducted in the Democratic Republic of Congo. Also, the isolation of *Salmonella* Heidelberg, *Salmonella* Koessen, *Salmonella* Pullorum and *Salmonella* Gallinarum³⁶ has previously been isolated from poultry justifying that aves (poultry) are reservoirs of *Salmonella*.³⁴ *Salmonella* Koessens, *Salmonella* Pullorum and *Salmonella* Gallinarum have been linked to salmonellosis originating from eggs. These *Salmonella* serovars constitute a threat to food safety and are capable of causing human sicknesses. Variations in *Salmonella* serovars occurrences have been reported in different countries and said to be a function of geographical location.^{36–38} The typhoidal groups have been implicated as the most cause of ill health in the developing countries.^{39,40} However, the non-typhoidal *Salmonella* serovars (*Salmonella* Typhimurium and *Salmonella* Enteritidis) are regarded as the major causes of salmonellosis outbreak in developing countries like India, Iran and many sub-saharan Africa.⁴¹ Hence, the isolation of virulent strains of *Salmonella* in chickens which happens to form a major part of South African cuisines brings a concern to food security and safety of consumers.

The bootstrap values within the evolutionary trend were higher than 70%, which supports the previous report of Wayne et al⁴² on decision of a close relatedness between organisms. However, the disparity in homology between CP012144 and other isolates could be due to mutation and development of new traits as evolution proceeds. However, CP012144 was comparable to *Salmonella enterica* subsp. *enterica* (MG663470) isolated in this study confirming the sharing of the same ancestral origin. The development of new traits having a 99% homology in isolates MG663482, MG66347, and MG663480 was observed showing that they all evolved from the same ancestor. This corroborates the previous report that *Salmonella* strains evolved from two broad genus *Salmonella enterica* and *Salmonella bongori*.⁴³ The bootstrap clustering of MG663495 with *S. bongori* could be due to evolutionary traits that are not pronounced in the isolates. However, the high bootstrap values within the *Salmonella enterica* group are indicative of high genetic relatedness and reliability of traits developed which cannot easily disappear or wiped out overnight.⁴⁴ The bootstrap values of *Salmonella* isolates and control strains in this study were higher than 50% which showed a high level of

repetitive clustering within the isolates. This supports the findings of Soltis and Soltis⁴⁵ on the acceptable bootstrap value (100–70%) in the construction of phylogenies. The observation of 100% bootstrap values of isolates as shown in the phylogenetic tree showed that 100% level of repetition exists in the genome compared.

Ampicillin belongs to the group of aminopenicillins which are often administered in the treatment of diseases caused by the gram-negative pathogenic enteric bacteria. Gentamycin belongs to the group of aminoglycosides alongside streptomycin, but in this study, gentamycin showed effectiveness in the control of *Salmonella*. Gentamycin has been reported to have a higher sensitivity on *Salmonella* strains compared to other antibiotics used in previous studies.⁴⁶ This might be due to the fact that gentamycin does not fall within the most common antibiotics administered in the treatment of *Salmonella* caused infection. However, it must be noted that an uncontrolled use of these antibiotics could also lead to a build-up of resistance to these antibiotics. All isolated *Salmonella bongori* strains had resistance to ciprofloxacin and nalidixic acid which belong to the fluoroquinolones often regarded as the last resort in *Salmonella* infection treatment. Similar reports have been made on the isolation of fluoroquinolone-resistant *Salmonella* in Taiwan.⁴⁷ Nalidixic acid is a new generation of the fluoroquinolones often prescribed in the treatment of *Salmonella*-caused infections. This finding is concurrent with⁴⁸ reports on multiple antibiotics resistance of *Salmonella* isolates from poultry in India and Egypt, respectively.

Higher antibiotic resistance phenotype profile was observed in *Salmonella* Typhimurium as opposed to the previous report of its penta-antibiotic resistance phenotype profiles. The increased antibiotic resistance obtained in this study could be due to misuse of antibiotics, thus resulting in adaptation and change in the antibiotic sensitivity behaviour of this pathogen with the aim to survive stress condition within the eco-system. Isolation of multiple antibiotics resistance strains in indigenous chickens calls for concern as it is believed that this breed of chickens is not often administered antibiotics during ill health. Albeit, the occurrence of multi-drug resistance could be due to the effect of a possible lateral gene transfer within the ecological niche. With regard to sample source, the occurrence of *Salmonella* multiple-antibiotics resistance followed the order (layers ≤ broilers ≤ indigenous chickens).

The resort to the prolonged use of fluoroquinolones in the treatment of *Salmonella*-borne infections has led to many

cases of antimicrobial resistance globally.⁴⁹ This resistance has been explained to be caused by mutations of the gyrase DNA gene and change in the efflux pump which is a target for the fluoroquinolones.^{50,51} However, Lauderdale et al⁴⁷ have suggested the use of the extended spectrum of cephalosporins as the last resort in the treatment of *Salmonella* infections. Hence, there is the need to develop an effective therapeutic approach in the control of these evolving virulent *Salmonella* strains. Furthermore, a negative correlation exists between the antibiotic-resistant profiles of *Salmonella* isolates from broilers, indigenous and layer chickens. The Pearson partial correlation was significant at $p \leq 0.01$. A positive correlation shows closer similarities in the antibiotic-resistant patterns of different *Salmonella* isolates in the study area, while a negative correlation within the sample source is implicative of non-source-dependent profiles.

The high occurrence of the antibiotic-resistant *Salmonella* strains from indigenous chickens could be due to a pick-up of virulence determinants from the environment or through interaction hosts such as rodents and livestock whom they share feeding and drinking troughs. Also, the high percent distribution of antibiotic-resistant strains in the broilers as shown in the clustering profiles indicates that isolates do share the same antibiotic resistance histories. According to Forshell et al,⁵² the abuse and misuse of antibiotics is a major cause of increasing antibiotic resistance among microorganisms of public health significance such as *Salmonella*. During processing or dressing operations, care for poultry, situations of gastrointestinal content shedding could arise which could lead to spillages of gut content into the environment. Also, unhygienic practices on farms and processing industries could aid in the transport of these virulent strains to the public either through the release of untreated effluents into river channels or other water bodies. These water bodies form a major resource for livelihood among the rural dwellers.

Conclusion

From this study, it is reported that the similarities in the antibiotic-resistant patterns among isolates from broilers, layers and indigenous chickens reveal similarities in antibiotic exposure histories. It is therefore suggested that there is need to sensitize farmers to adhere to prescribed guidelines on the use of antibiotics. In addition, the implementation of good sanitation among farm workers as well as standard operating procedures in farms where animals are housed should be encouraged to curb the spread of multi-drug-resistant strains of *Salmonella* since the latter may pose a threat to public health. The detection

of large proportions of diverse multi-antibiotic-resistant *Salmonella* strains in chickens within Mafikeng community, especially in indigenous breed, indicates that these animals may pose a threat on food security and safety. Further studies on the antibiotic-resistant genes harboured by this pathogen could advance knowledge in the development of suitable antibiotics and other prophylaxis to curb *Salmonella*-caused infections.

Acknowledgments

Authors acknowledge the assistance received from Dr M. Ogunrinu and Dr L. Ngoma during this study. This work was supported by The World Academy of Science in collaboration with the Directorate of Science and Technology, National Research Foundation (Twas DST-NRF) of South Africa (grant number UID: 110854, 2018).

Disclosure

Mr Stephen Abiola Akinola reports grants from TWAS DST-NRF, outside the submitted work. The authors report no other conflicts of interest in this work.

References

1. Ulaya WD. *Determination of Virulence Factors in Salmonella Isolates of Human, Poultry and Dog Origin in Lusaka District, Zambia, in Department of Paraclinical Studies Lusaka.* The University of Zambia; 2013:98.
2. Acheson D, Hohmann EL. Nontyphoidal salmonellosis. *Clin Infect Dis.* 2001;32(2):263–269. doi:10.1086/318457
3. Gill CJ, Hamer DH. Foodborne illnesses. *Curr Treat Options Gastroenterol.* 2001;4(1):23. doi:10.1007/s11938-001-0044-0
4. Kemal J. A review on the public health importance of bovine salmonellosis. *Vet Sci Technol.* 2014;5(2):1.
5. Kagambèga A, Lienemann T, Aulu L, et al. Prevalence and characterization of *Salmonella enterica* from the feces of cattle, poultry, swine and hedgehogs in Burkina Faso and their comparison to human *Salmonella* isolates. *BMC Microbiol.* 2013;13(1):253. doi:10.1186/1471-2180-13-253
6. Drzewiecka D. Significance and roles of *Proteus* spp. bacteria in natural environments. *Microb Ecol.* 2016;72(4):741–758. doi:10.1007/s00248-015-0720-6
7. WHO. *World Health Statistics 2016: Monitoring Health for the SDGs Sustainable Development Goals.* WHO Press. 2016.
8. Zouboulis A, Loukidou M, Matis K. Biosorption of toxic metals from aqueous solutions by bacteria strains isolated from metal-polluted soils. *Process Biochem.* 2004;39(8):909–916. doi:10.1016/S0032-9592(03)00200-0
9. Ateba CN, Mochaiwa B. Use of *invA* gene specific PCR analysis for the detection of virulent *Salmonella* species in beef products in the north west province, South Africa. *J Food Nutr Res.* 2014;2(p):294–300. doi:10.12691/jfmr-2-6-5
10. Muyzer G, Teske A, Wirsén CO, Jannasch HW. Phylogenetic relationships of *Thiomicrospira* species and their identification in deep-sea hydrothermal vent samples by denaturing gradient gel electrophoresis of 16S rDNA fragments. *Arch Microbiol.* 1995;164:165–172. doi:10.1007/BF02529967

11. Sambrook J, Russel DW. *Molecular Cloning. A Laboratory Manual*. 3. ed. New York: Cold Spring Harbor Laboratory Press; 2001.
12. National Center for Biotechnology Information, U.S. National Library of Medicine, 8600 Rockville Pike, BethesdaMD, 20894, USA.
13. Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res*. 1994;22(22):4673–4680. doi:10.1093/nar/22.22.4673
14. Hall TA. *BioEdit: A User-Friendly Biological Sequence Alignment Editor and Analysis Program for Windows 95/98/NT*. In *Nucleic Acids Symposium Series*. London: Information Retrieval Ltd.; 1999: c1979–c2000.
15. Jukes TH, Cantor CR. Evolution of protein molecules. *Mamm Protein Metab*. 1969;3(21):132.
16. Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol*. 1987;4(4):406–425. doi:10.1093/oxfordjournals.molbev.a040454
17. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol*. 2013;30(12):2725–2729. doi:10.1093/molbev/mst197
18. Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics*. 2011;27(16):2194–2200. doi:10.1093/bioinformatics/btr381
19. Bauer AW, Kirby WM, Sherris JC, Turck M. Antibiotic susceptibility testing by a standardized single disk method. *Am J Clin Pathol*. 1966;45(4):493. doi:10.1093/ajcp/45.6_ts.764
20. Wayne PA. Clinical and laboratory standards institute. *Perform Stand Antimicrob Susceptibility Test*. 2007;M100-S17.
21. Rota C, Yangüela J, Blanco D, Carramiñana JJ, Ariño A, Herrera A. High prevalence of multiple resistance to antibiotics in 144 *Listeria* isolates from Spanish dairy and meat products. *J Food Prot*. 1996;59(9):938–943. doi:10.4315/0362-028X-59.9.938
22. Singh S, Yadav AS, Singh SM, Bharti P. Prevalence of *Salmonella* in chicken eggs collected from poultry farms and marketing channels and their antimicrobial resistance. *Food Res Int*. 2010;43:2027–2030. doi:10.1016/j.foodres.2010.06.001
23. Wattiau P, Weijers T, Andreoli P, et al. Evaluation of the Premi®Test *Salmonella*, a commercial low-density DNA microarray system intended for routine identification and typing of *Salmonella enterica*. *Int J Food Microbiol*. 2008;123(3):293–298. doi:10.1016/j.ijfoodmicro.2008.01.006
24. Schrader KN, Fernandez-Castro A, Cheung WKW, Crandall CM, Abbott SL. Evaluation of commercial antisera for *Salmonella* serotyping. *J Clin Microbiol*. 2008;46(2):685–688. doi:10.1128/JCM.01808-07
25. Hendriksen R, Jaap W, Van Bergen M. *Global Salm-Surv. A Global Salmonella Surveillance and Laboratory Support Project of the World Health Organization. Identification of Thermotolerant Campylobacter*. WHO; 2003.
26. Shan Y, Lai Y, Yan A. Metabolic reprogramming under microaerobic and anaerobic conditions in bacteria. *Subcell. Biochem*. 2012;64:159–179.
27. Galán JE, Curtiss R. Distribution of the *invA*, *-B*, *-C*, and *-D* genes of *Salmonella typhimurium* among other *Salmonella* serovars: *invA* mutants of *Salmonella typhi* are deficient for entry into mammalian cells. *Infect Immun*. 1991;59(9):2901–2908.
28. Salehi TZ, Mahzounieh M, Saeedzadeh A. Detection of *invA* gene in isolated *Salmonella* from broilers by PCR method. *Int J Poult Sci*. 2005;4(8):557–559. doi:10.3923/ijps.2005.557.559
29. Oliveira SDD, Rodenbusch CR, Michael GB, Cardoso MIR, Canal CW, Brandelli A. Detection of virulence genes in *Salmonella enteritidis* isolated from different sources. *Braz J Microbiol*. 2003;34:123–124. doi:10.1590/S1517-83822003000500042
30. Ohl ME, Miller SI. *Salmonella*: a model for bacterial pathogenesis. *Annu Rev Med*. 2001;52(1):259–274. doi:10.1146/annurev.med.52.1.259
31. Cavaco L, Hasman H, Xia S, Aarestrup FM. *qnrD*, a novel gene conferring transferable quinolone resistance in *Salmonella enterica* serovar Kentucky and Bovismorbificans strains of human origin. *Antimicrob Agents Chemother*. 2009;53(2):603–608. doi:10.1128/AAC.00997-08
32. Harakeh S, Yassine H, El-Fadel M. Antimicrobial-resistant patterns of *Escherichia coli* and *Salmonella* strains in the aquatic Lebanese environments. *Environ Pollut*. 2006;143(2):269–277. doi:10.1016/j.envpol.2005.11.027
33. Morpeth SC, Ramadhani HO, Crump JA. Invasive non-typhi *Salmonella* disease in Africa. *Clin Infect Dis*. 2009;49(4):606–611. doi:10.1086/603553
34. Qiao J, Zhang Q, Alali WQ, et al. Characterization of extended-spectrum β -lactamases (ESBLs)-producing *Salmonella* in retail raw chicken carcasses. *Int J Food Microbiol*. 2017;248:72–81. doi:10.1016/j.ijfoodmicro.2017.02.016
35. Phoba M-F, Boeck H, Ifeka BB, et al. Epidemic increase in *Salmonella* bloodstream infection in children, Bwamanda, the Democratic Republic of Congo. *Eur J Clin Microb Infect Dis*. 2014;33(1):79–87. doi:10.1007/s10096-013-1931-8
36. Shivaprasad H. Fowl typhoid and pullorum disease. *Revue Scientifique Et Technique-Office Int Des Epizooties*. 2000;19(2):405–416. doi:10.20506/rst.19.2.1222
37. Shilangale RP, Giannatale ED, Chimwamuombe PM, Kaaya GP. Prevalence and antimicrobial resistance pattern of *Salmonella* in animal feed produce in Namibia. *Vet. Ital*. 2012;48:125–132.
38. Foley S, Lynne A, Nayak R. *Salmonella* challenges: prevalence in swine and poultry and potential pathogenicity of such isolates 1 2. *J Anim Sci*. 2008;86(14 suppl):E149–E162. doi:10.2527/jas.2007-0464
39. Foley S, Lynne A. Food animal-associated *Salmonella* challenges: pathogenicity and antimicrobial resistance 1. *J Anim Sci*. 2008;86(14 suppl):E173–E187. doi:10.2527/jas.2007-0447
40. Santos RL, Zhang S, Tsoilis RM, Kingsley RA, Adams LG, Bäumlér AJ. Animal models of *Salmonella* infections: enteritis versus typhoid fever. *Microbes Infect*. 2001;3(14–15):1335–1344.
41. Gilks CF. Acute bacterial infections and HIV disease. *Br Med Bull*. 1998;54(2):383–393. doi:10.1093/oxfordjournals.bmb.a011695
42. Wayne L, Stackebrandt E, Kandler O, et al. Report of the ad hoc committee on reconciliation of approaches to bacterial systematics. *Int J Syst Evol Microbiol*. 1987;37(4):463–464. doi:10.1099/00207713-37-4-463
43. Tuchili L, Ulaya W, Kato Y, Kaneuchi C. Recent characterisation of *Salmonella* strains isolated from chickens in Zambia. *J Vet Med Sci*. 1996;58(1):77–78. doi:10.1292/jvms.58.77
44. Konstantinidis KT, Stackebrandt E. *Defining Taxonomic Ranks, in the Prokaryotes*. Springer; 2013:229–254.
45. Soltis PS, Soltis DE. Applying the bootstrap in phylogeny reconstruction. *Stat Sci*. 2003;256–267. doi:10.1214/ss/1063994980
46. Arora D, Kumar S, Singh D, Jindal N, Mahajan N. Isolation, characterization and antibiogram pattern of *Salmonella* from poultry in parts of Haryana, India. *Adv Anim Vet Sci*. 2013;1(5):161–163.
47. Lauderdale T-L, Aarestrup FM, Chen P-C, et al. Multidrug resistance among different serotypes of clinical *Salmonella* isolates in Taiwan. *Diagn Microbiol Infect Dis*. 2006;55(2):149–155. doi:10.1016/j.diagmicrobio.2006.01.002
48. Mahmoud MS, Esmail M, Khairy R, Mazher O. Detection of multi-drug resistant non-typhoid *Salmonella* isolates in cases of gastroenteritis in Egypt. *Br Microb Res J*. 2015;6(3):167. doi:10.9734/BMRJ
49. Threlfall EJ. Antimicrobial drug resistance in *Salmonella*: problems and perspectives in food-and water-borne infections. *FEMS Microbiol Rev*. 2002;26(2):141–148. doi:10.1111/j.1574-6976.2002.tb00606.x

50. Piddock LJ. Fluoroquinolone resistance in *Salmonella* serovars isolated from humans and food animals. *FEMS Microbiol Rev.* 2002;26(1):3–16. doi:10.1111/j.1574-6976.2002.tb00596.x
51. Yang B, Xi M, Cui S, et al. Mutations in gyrase and topoisomerase genes associated with fluoroquinolone resistance in *Salmonella* serovars from retail meats. *Food Res Int.* 2012;45(2):935–939. doi:10.1016/j.foodres.2011.01.031
52. Forshell LP, Wierup M. *Salmonella* contamination: a significant challenge to the global marketing of animal food products. *Rev Sci Tech off Int Epiz.* 2006;25(2):541–554.
53. Guariguata L, Whiting DR, Hambleton I, Beagley J, Linnenkamp U, Shaw JE. Global estimates of diabetes prevalence for 2013 and projections for 2035. *Diabetes Res Clin Pract.* 2014;103(2):137–149. doi:10.1016/j.diabres.2013.11.002

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>