

# Circ\_0058124 Upregulates MAPK1 Expression to Promote Proliferation, Metastasis and Metabolic Abilities in Thyroid Cancer Through Sponging miR-940

This article was published in the following Dove Press journal:  
*OncoTargets and Therapy*

Dezhong Sun<sup>1</sup>  
Li Chen<sup>2</sup>  
Huaiqing Lv<sup>1</sup>  
Yongli Gao<sup>3</sup>  
Xuelai Liu<sup>4</sup>  
Xiaoyan Zhang<sup>3</sup>

<sup>1</sup>Department of Otolaryngology, Linyi People's Hospital, Linyi, Shandong, People's Republic of China; <sup>2</sup>Department of Anesthesiology Operation, Linyi People's Hospital, Linyi, Shandong, People's Republic of China; <sup>3</sup>Department of Medicine Oncology, Linyi People's Hospital, Linyi, Shandong, People's Republic of China; <sup>4</sup>Department of Neurosurgery, Linyi Hospital of Traditional Chinese Medicine, Linyi, Shandong, People's Republic of China

**Background:** Thyroid cancer (TC) is an endocrine disease, and its progression is regulated by many factors, including circular RNAs (circRNAs). However, as a new circRNA, the role of circ\_0058124 in TC is worth further exploration.

**Methods:** The expression levels of circ\_0058124, microRNA-940 (miR-940) and mitogen-activated protein kinase 1 (MAPK1) were assessed by quantitative polymerase chain reaction (q-PCR). The circular characteristic of circ\_0058124 was identified by oligo (dT)<sub>18</sub> primers, Ribonuclease R (RNase R) and Actinomycin D (ActD), and its localization was determined by nuclear-cytoplasmic separation assay. Also, cell proliferation was detected by colony formation assay, and cell migration and invasion were assessed by transwell assay. Further, Seahorse XF Extracellular Flux Analyzer was used to measure the oxygen consumption rate (OCR) of cells. Besides, dual-luciferase reporter, RNA immunoprecipitation (RIP) and RNA pull-down assays were used to identify the mechanism of circ\_0058124. Western blot (WB) analysis was used to test the MAPK1 protein level. In addition, mice xenograft models were constructed to test the effect of circ\_0058124 on TC tumor growth in vivo.

**Results:** Circ\_0058124 was highly expressed in TC and is a stable cyclic transcript, mainly located in the cytoplasm. Circ\_0058124 knockdown suppressed proliferation, migration, invasion and metabolic abilities in TC cells. MiR-940 could be absorbed by circ\_0058124, and the inhibition effect of its overexpression on TC progression could be reversed by overexpressed-circ\_0058124. MAPK1 was a target of miR-940, and the suppression effect of its silencing on TC progression could be inverted by miR-940 inhibitor. Besides, MAPK1 expression was regulated by circ\_0058124 and miR-940. Interference of circ\_0058124 also reduced TC tumor growth in vivo.

**Conclusion:** Circ\_0058124 might play a carcinogenic role in TC progression by regulating the miR-940/MAPK1 axis, which might provide a new idea for the treatment of TC.

**Keywords:** TC, circ\_0058124, miR-940, MAPK1

Correspondence: Xiaoyan Zhang  
Department of Medicine Oncology, Linyi People's Hospital, No. 27, Jiefang Road East Section, Linyi, Shandong 276000, People's Republic of China  
Tel +86 539 8092120  
Email vgvlug@163.com

## Introduction

Thyroid cancer (TC) is a common thyroid malignant tumor, including papillary TC, follicular TC, anaplastic TC and medullary TC.<sup>1</sup> Papillary TC is a more common type of TC, which has a lower degree of malignancy and a better prognosis.<sup>2</sup> In recent years, the incidence of TC has increased significantly,

which is related to the region, race and gender.<sup>3</sup> Distant metastasis and local recurrence are indicators of poor prognosis in TC patients, which seriously affect the survival rate of patients.<sup>4,5</sup> Therefore, it is urgent to seek new biomarkers that can be used for early diagnosis.

Circular RNAs (circRNAs) are non-coding RNAs (ncRNAs), which have attracted more attention due to its unique covalently closed structure.<sup>6</sup> CircRNAs can be involved in the regulation of cancer progression and have potential as the cancer biomarkers.<sup>7</sup> For example, circ\_0044516 could promote proliferation and metastasis in prostate cancer, so it could function as a biomarker of prostate cancer.<sup>8</sup> Besides, hsa\_circ\_006848 could act as a novel biomarker of early gastric cancer.<sup>9</sup> In TC, the roles of circ\_0039411 and circ\_0025033 had been elucidated, both of which could promote the progression of TC.<sup>10,11</sup> Circ\_0058124 is a newly discovered circRNA differentially expressed in TC and has been proved to be associated with the poor prognosis of patients.<sup>12</sup> However, the role and mechanism of TC should be further explored.

MicroRNAs (miRNAs) are small ncRNAs with a length of 22 nucleotides, and their expression is closely related to cancer progression.<sup>13</sup> MiRNAs-targeted therapy technology has developed new strategies for cancer treatment.<sup>14</sup> With the deepening of the research on circRNAs mechanism, it has been found that circRNAs can act as complementary RNAs (ceRNAs) to participate in the regulation of target genes of miRNAs.<sup>15</sup> MiR-940 is widely under-expressed in many cancers and has been shown to be involved in the regulation of cancer progression, such as glioma cancer and breast cancer.<sup>16,17</sup> Studies have shown that miR-940 is lower expressed in TC tissues.<sup>18</sup> Nevertheless, the role of miR-940 in TC still remains unclear.

Mitogen-activated protein kinase 1 (MAPK1), a member of the MAP kinase family, is involved in the regulation of cell physiology and pathology.<sup>19</sup> Some studies indicate that MAPK1 is highly expressed in TC and participates in the regulation of TC progress by miRNA as a target gene.<sup>20,21</sup> Therefore, MAPK1 is a key gene that regulates the progression of TC.

The purpose of this study is to explore the role of circ\_0058124 in TC and explore its new mechanism of regulating the progression of TC, so as to provide a reference for circ\_0058124 to become a diagnostic biomarker of TC.

## Materials and Methods

### Patients and Tissue Samples

Our study was approved by the Ethics Committee of Linyi People's Hospital. 51 TC patients were recruited from Linyi People's Hospital and divided into I + II group (23) and III + IV group (28) according to their tumor stage. TC tissues and adjacent normal tissues of all patients were collected and stored at  $-80^{\circ}\text{C}$  until use. Patient consent was written informed consent. The study was performed in accordance with the Declaration of Helsinki.

### Cell Culture

TC cell lines (BCPAP, TPC-1, IHH-4 and HTH83) and human thyroid normal cell line (NTHY-OR13.1) were bought from Guandao Biological (Shanghai, China). All cells were cultured in Dulbecco's modified Eagle's medium (DMEM, Hyclone, Logan, Utah, USA) containing 10% fetal bovine serum (FBS; Hyclone), 100 U/mL penicillin and 100  $\mu\text{g}/\text{mL}$  streptomycin (Invitrogen, Carlsbad, CA, USA) at  $37^{\circ}\text{C}$  in an incubator with 5%  $\text{CO}_2$ .

### Quantitative Polymerase Chain Reaction (q-PCR)

Total RNA was extracted using Trizol reagent (Invitrogen). Complementary DNA (cDNA) was synthesized using the Transcriptor Universal cDNA Master (Roche, Basel, Switzerland). Q-PCR was performed using SYBR Green (Takara, Dalian, China). Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) and U6 were used as internal controls. The  $2^{-\Delta\Delta\text{Ct}}$  method was used to calculate the fold changes. All primers were listed as follows: circ\_0058124, F 5'-CCTTTGCAGAAGTCACCGGG-3', R 5'-CCCAGGAGACCACAAAGCTAC-3'; fibronectin1 (FN1), F 5'-TGATCACATGGACGCTGC-3', R 5'-GAGTCAAGCCGGACACAACG-3'; MAPK1, F 5'-CTGCTGCTCAACACCACCT-3', R 5'-GCCACATATTCTGTGTCAGGAACC-3'; GAPDH, F 5'-ACCACAGTCCATGCCATCAC-3', R 5'-TCACACCCTGTTGCTGTA-3'; miR-940, F 5'-GCATCGTTCCTTCAAGCCGATCT-3', R 5'-TGGGTGAGTCGTTCCGG-3'; U6, F 5'-GCAGGAGTCTTCACAGAGT-3', R 5'-TCTAGAGGAGAAGCTGGGGT-3'.

### CircRNA Identification and Localization

We used three experimental methods to identify the circular characteristic of circ\_0058124. Oligo (dT)<sub>18</sub> primers could only amplify RNA containing poly (A) tails, while random primers could amplify all RNA. Circ\_0058124 and mFN1

were amplified with random primers or oligo (dT)<sub>18</sub> primers, followed by q-PCR to confirm whether circ\_0058124 contained poly (A) tails. Ribonuclease R (RNase R; Duma, Shanghai, China) could only degrade linear RNA without affecting circRNA. The extracted RNA was treated with RNase R, followed by q-PCR to determine the circular characteristic of circ\_0058124. Also, the extracted RNA was treated with Actinomycin D (ActD; Amyjet, Wuhan, China) for 0 h, 6 h, 12 h, 18 h and 24 h, respectively, followed by q-PCR to verify the stability of circ\_0058124. Here, mRNA FN1 (mFN1) represented the linear RNA of circ\_0058124. For circRNA localization, the cytoplasmic and nuclear RNA of TPC-1 and HTH83 cells were isolated and extracted using Cytoplasmic & Nuclear RNA Purification Kit (Amyjet), and the circ\_0058124 expression in cytoplasmic and nuclear was measured by q-PCR. GAPDH and U6 were served as the cytoplasm control and nuclear control, respectively.

## Cell Transfection

Circ\_0058124 small interfering RNA (siRNA), overexpression plasmid and lentiviral short hairpin RNA (shRNA) (si-circ\_0058124, circ\_0058124 and sh-circ\_0058124) or their negative controls (si-NC, vector and sh-NC), miR-940 mimic and inhibitor (miR-940 and anti-miR-940) or their negative controls (miR-NC and anti-miR-NC), MAPK1 siRNA (si-MAPK1) and its negative control (si-NC) were synthesized by Ribobio (Guangzhou, China). All plasmid vectors were transfected into TPC-1 and HTH83 cells using Lipofectamine 3000 (Invitrogen).

## Colony Formation Assay

TPC-1 and HTH83 cells were seeded into 6-well plates. After transfection, cells were incubated for 2 weeks. Then, TPC-1 and HTH83 cells were fixed with methanol and stained with crystal violet to observe and count the number of colonies (> 50 cells).

## Transwell Assay

Cell migration and invasion assays were performed by transwell chambers (Corning Inc., Corning, NY, USA). TPC-1 and HTH83 cells were seeded into the upper chambers, which non-coated and pre-coated with Matrigel (BD Biosciences, San Jose, CA, USA) to detect migration and invasion, respectively. The upper chambers were filled with serum-free medium, and the lower chambers were filled with serum medium. After 24 h, cells on the lower chambers were fixed, stained and then counted the number of migrated and invaded cells.

## Cell Metabolic Ability Assay

This experiment was performed after cell transfection for 48 h. According to the manufacturer's agreement, oligomycin, mitochondrial uncoupling agent carbonyl cyanide 4-(trifluoromethoxy) phenylhydrazone (FCCP) and Antimycin/Rotenone (Seahorse Bioscience, Billerica, MA, USA) were successively added to TPC-1 and HTH83 cells, and the oxygen consumption rate (OCR) of cells was monitored by the Seahorse XF Extracellular Flux Analyzer (Seahorse Bioscience).

## Dual-Luciferase Reporter Assay

The sequences of circ\_0058124-WT/MUT or MAPK1-WT/MUT were inserted into the pGL3-control vectors (Promega, Madison, WI, USA). The above reporter vectors were co-transfected with miR-940 mimic or miR-NC into TPC-1 and HTH83 cells using Lipofectamine 3000 (Invitrogen). The luciferase activities of cells were measured by Dual-Lucy Assay Kit (Solarbio, Beijing, China).

## RNA Immunoprecipitation (RIP) Assay

RIP assay was performed using the Magna RIP RNA-Binding Protein immunoprecipitation kit (Millipore, Billerica, MA, USA). After transfection for 48 h, TPC-1 and HTH83 cells were lysed by RIP lysis buffer. Then, cell lysates were incubated with magnetic beads conjugated with argonaute2 antibody (anti-Ago2) or immunoglobulin G (IgG) antibody (anti-IgG). After purified, the enrichment of circ\_0058124 and miR-940 was tested by q-PCR.

## RNA Pull-Down Assay

Biotin-labeled miR-940 probe (Bio-miR-940) and negative control probe (Bio-NC) were purchased from Sangon Inc. (Shanghai, China). Bio-miR-940 or Bio-NC was incubated with magnetic beads at 4°C overnight. TPC-1 and HTH83 cells were lysed and collected the lysates. Then, cell lysates were incubated with magnetic beads mixture at room temperature for 1 h. After purified, the enrichment of circ\_0058124 was detected by q-PCR.

## Western Blot (WB) Analysis

TPC-1 and HTH83 cells were lysed with RIPA buffer (Beyotime, Shanghai, China). The same amount of protein was separated via sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) gel and transferred onto polyvinylidene fluoride (PVDF) membranes (Millipore). Then, the membranes were blocked with nonfat milk and incubated

with primary antibodies against MAPK1 (1:1,000, Amyjet) or GAPDH (1:2,000, Amyjet) at 4°C overnight. After incubated with a secondary antibody (1:2,000, Amyjet), the membranes were treated with enhanced chemiluminescence solution (Beyotime) to detect the protein signals.

## Mice Xenograft Models

Six-week-old BALB/c male nude mice were purchased from Sebia (Guangzhou, China). TPC-1 cells stably transfected with sh-circ\_0058124 or sh-NC were subcutaneously injected into nude mice. Tumor length and width were measured weekly to calculate tumor volume according to the formula: volume (mm<sup>3</sup>) = width<sup>2</sup> × length/2. Mice were killed after 5 weeks and tumor samples were taken for further study. Animal experiments were approved by the Animal Research Committee of Linyi People's Hospital and conducted in accordance with the animal welfare guidelines of Linyi People's Hospital.

## Statistical Analysis

All data were shown as the mean ± standard deviation. Student's *t*-test or one-way analysis of variance was used for statistical analysis in SPSS17.0 software (SPSS Inc., Chicago, IL, USA). *P* < 0.05 was considered to be statistically significant.

## Results


### Circ\_0058124 Was Upregulated in TC Tissues and Cells

We first measured the expression of circ\_0058124 in TC tissues. Compared with adjacent normal tissues,

circ\_0058124 expression was elevated in TC tissues (Figure 1A). Besides, in III and IV stages of TC tissues, the expression of circ\_0058124 was higher than that in I and II stages of TC tissues, indicating that circ\_0058124 expression was also related to TC tumor stage (Figure 1B). Meanwhile, we also found that the expression of circ\_0058124 was increased in four TC cell lines (especially TPC-1 and HTH83 cells) compared with that in NTHY-ORI3.1 cells (Figure 1C). These data suggested that circ\_0058124 might play an important role in TC.


### Identification and Validation of Circ\_0058124 in TC Cells

To confirm the circular characteristics of circ\_0058124, we used random primers or oligo (dT)<sub>18</sub> primers to perform q-PCR. The results showed that compared with random primers, the relative expression of circ\_0058124 was markedly decreased in TPC-1 and HTH83 cells when using the oligo (dT)<sub>18</sub> primers, while mFN1 was not (Figure 2A), indicating that circ\_0058124 had no poly-A tail. Besides, we detected the subcellular distribution of circ\_0058124 and found that circ\_0058124 was mainly distributed in the cytoplasm of TC cells (Figure 2B), suggesting that circ\_0058124 might be mainly involved in post-transcriptional regulation. Meanwhile, circ\_0058124 was resistant to RNase R, while linear RNA mFN1 could be digested by RNase R (Figure 2C), indicating that circ\_0058124 was circular. In addition, we used ActD to verify the stability of circ\_0058124 and found that circ\_0058124 was more stable than mFN1 in TPC-1 and HTH83 cells (Figure 2D). Hence, these results



**Figure 1** The expression of circ\_0058124 in tissues and cells.

**Notes:** (A) Circ\_0058124 expression in TC tissues (TC) and adjacent normal tissues (Normal) was detected by q-PCR. (B) Q-PCR was used to assess the expression of circ\_0058124 in the different stages of TC (I+II and III+IV). (C) The expression of circ\_0058124 in TC cell lines (BCPAP, TPC-1, IHH-4 and HTH83) and NTHY-ORI3.1 cells was measured by q-PCR. \**P* < 0.05.



**Figure 2** Identification and validation of circ\_0058124 in TC cells.

**Notes:** (A) The relative expression levels of circ\_0058124 and mFN1 in TPC-1 and HTH83 cells were analyzed by q-PCR after normalized with random primers and oligo (dT)<sub>18</sub> primers. (B) The expression levels of circ\_0058124, U6 and GAPDH in the nuclear and cytoplasmic of TPC-1 and HTH83 cells were detected by q-PCR after nuclear-cytoplasmic separation. (C) The relative expression levels of circ\_0058124 and mFN1 in TPC-1 and HTH83 cells were assessed by q-PCR after treatment with RNase R. (D) The relative expression levels of circ\_0058124 and mFN1 in TPC-1 and HTH83 cells were analyzed by q-PCR after treatment with ActD at the indicated time points. \**P* < 0.05.

demonstrated that circ\_0058124 is a circular and stable transcript in TC.


## Knockdown of Circ\_0058124 Hindered the Progression of TC Cells

To explore the role of circ\_0058124 on TC progression, we inhibited circ\_0058124 expression using si-circ\_0058124. The decrease of circ\_0058124 expression in TPC-1 and HTH83 cells indicated that the transfection of si-circ\_0058124 was successful (Figure 3A). Colony formation assay results showed that circ\_0058124 silencing reduced the number of colonies in TPC-1 and HTH83 cells (Figure 3B), indicating that its knockdown suppressed the proliferation ability of TPC-1 and HTH83 cells. Besides, transwell assay results revealed that the number of migrated and invaded TPC-1 and HTH83 cells were remarkably decreased in the si-circ\_0058124 group compared with that in the si-NC group (Figure 3C and D), suggesting that silenced-circ\_0058124 restrained the migration and invasion abilities of TPC-1 and HTH83 cells. In addition, we also tested the OCR of TPC-1 and HTH83 cells, and the results showed that after circ\_0058124 knockdown, the OCR of basal respiration and maximum respiration was significantly decreased in TPC-1 and HTH83 cells (Figure 3E and F), which

determined that the metabolic ability of cells was inhibited by circ\_0058124 silencing. All data revealed that circ\_0058124 played an important role in the progress of TC cells.

## Circ\_0058124 Could Sponge miR-940

We further investigated the complementary miRNAs paired with circ\_0058124. The Interactome tool was used for prediction, and the results showed that miR-940 had binding sites with circ\_0058124 (Figure 4A). To validate the binding ability between them, we constructed the WT-circ\_0058124 and MUT-circ\_0058124 reporter vectors. Dual-luciferase reporter assay demonstrated that miR-940 overexpression could markedly suppress the luciferase activity of WT-circ\_0058124, but not affect the luciferase activity of MUT-circ\_0058124 in TPC-1 and HTH83 cells (Figure 4B). Also, RIP assay results revealed that circ\_0058124 and miR-940 were remarkably enriched in anti-Ago2 compared with that in anti-IgG (Figure 4C). Besides, RNA pull-down assay results showed that the enrichment of circ\_0058124 was significantly increased in Bio-miR-940 (Figure 4D). These data confirmed that circ\_0058124 could absorb miR-940 in TPC-1 and HTH83 cells. Moreover, we also detected the expression of miR-940 in different tissues and cells and discovered




**Figure 3** Effects of circ\_0058124 on the progression of TC cells.

**Notes:** TPC-1 and HTH83 cells were transfected with si-circ\_0058124 or si-NC. (A) The expression of circ\_0058124 was detected by q-PCR to evaluate transfection efficiency. (B) Colony formation assay was used to measure the number of colonies in TPC-1 and HTH83 cells. (C and D) The number of migrated and invaded TPC-1 and HTH83 cells were determined by transwell assay. (E and F) The OCR of basal respiration and maximum respiration in TPC-1 and HTH83 cells was analyzed by Seahorse XF Extracellular Flux Analyzer. \* $P < 0.05$ .

the fact that miR-940 was lower expressed in TC (Figure 4E and F). In addition, we explored the effect of circ\_0058124 expression on miR-940 expression. A significant increase in circ\_0058124 expression indicated that circ\_0058124 transfection was successful

(Figure 4G). Q-PCR results revealed that miR-940 expression was promoted by circ\_0058124 knock-down and inhibited by circ\_0058124 overexpression (Figure 4H). All these data proved the binding relationship between circ\_0058124 and miR-940.




**Figure 4** Circ\_0058124 could sponge miR-940.

**Notes:** (A) The sequences of circ\_0058124 containing the miR-940 binding sites or mutant binding sites were shown. (B) Dual-luciferase reporter assay was used to detect the interaction between miR-940 and circ\_0058124 in TPC-1 and HTH83 cells. (C) The enrichment of circ\_0058124 and miR-940 in anti-Ago2 or anti-IgG was determined by the RIP assay. (D) RNA pull-down assay was used to assess the enrichment of circ\_0058124 in Bio-miR-940 or Bio-NC. (E) Q-PCR was performed to measure the expression of miR-940 in TC tissues (TC) and adjacent normal tissues (Normal). (F) The expression of miR-940 in TC cells (TPC-1 and HTH83) and NTHY-OR13.1 cells was determined by q-PCR. (G) Circ\_0058124 expression was determined by q-PCR to evaluate the transfection efficiency of circ\_0058124 overexpression plasmid in TPC-1 and HTH83 cells. (H) The expression of miR-940 was detected by q-PCR in TPC-1 and HTH83 cells to assess the effects of si-circ\_0058124 and circ\_0058124 overexpression plasmid on miR-940 expression. \* $P < 0.05$ .

## Overexpression of Circ\_0058124 Could Revert the Suppression Effect of miR-940 Mimic on the Progression of TC Cells

To confirm the effect of miR-940 on the progression of TC cells, we co-transfected miR-940 mimic and circ\_0058124 overexpression plasmid into TPC-1 and HTH83 cells. The detection results of miR-940 expression showed that miR-940 mimic could markedly improve miR-940 expression, while circ\_0058124 hindered its expression, indicating that the transfection efficiency of both was excellent (Figure 5A). Through colony formation assay, we found

that miR-940 overexpression could suppress the number of colonies in TPC-1 and HTH83 cells, while this inhibition effect could be recovered by circ\_0058124 overexpression (Figure 5B). Subsequently, transwell assay also showed that overexpressed circ\_0058124 could reverse the suppression effects of miR-940 overexpression on the migration and invasion of TPC-1 and HTH83 cells (Figure 5C and D). Besides, OCR detection assay results revealed that miR-940 indeed inhibited the metabolic ability of TPC-1 and HTH83 cells, while overexpressed circ\_0058124 inverted this suppression effect (Figure 5E and F). These



**Figure 5** Effects of miR-940 on the progression of TC cells.

**Notes:** TPC-1 and HTH83 cells were co-transfected with miR-940 mimic and circ\_0058124 overexpression plasmid. (A) Q-PCR was used to detect the expression of miR-940 to evaluate the transfection efficiency of miR-940 mimic and circ\_0058124 overexpression plasmid. (B) The number of colonies in TPC-1 and HTH83 cells was measured by colony formation assay. (C and D) Transwell assay was used to determine the number of migrated and invaded TPC-1 and HTH83 cells. (E and F) Seahorse XF Extracellular Flux Analyzer was used to assess the OCR of basal respiration and maximum respiration in TPC-1 and HTH83 cells. \* $P < 0.05$ .


results further confirmed that the function of miR-940 was regulated by circ\_0058124 in TC.

## miR-940 Could Target MAPK1

At the same time, we used the Targetscan tool to predict the downstream target gene of miR-940. As shown in Figure 6A, MAPK1 3'UTR had complementary binding sites with miR-940. Then, WT-MAPK1 and MUT-MAPK1 reporter vectors were constructed to perform the Dual-luciferase reporter assay. The results determined that miR-940 mimic markedly decreased the luciferase activity of WT-MAPK1 in TPC-1 and HTH83 cells, while had no effect on MUT-MAPK1 (Figure 6B). Meanwhile, we detected MAPK1 expression in TC tissues and adjacent

normal tissues and found that MAPK1 expression was increased in TC tissues (Figure 6C). Through WB analysis, we also discovered that the protein level of MAPK1 was upregulated in TC tissues and cells (Figure 6D and E). Besides, we investigated the effect of miR-940 expression on MAPK1 expression. The decrease of miR-940 expression indicated that the transfection efficiency of miR-940 inhibitor was excellent (Figure 6F). WB analysis results showed that miR-940 overexpression remarkably suppressed MAPK1 protein level in TPC-1 and HTH83 cells, while miR-940 inhibition markedly promoted its protein level (Figure 6G), suggesting that MAPK1 expression was regulated by miR-940. Therefore, these results suggested that MAPK1 was a target of miR-940.





**Figure 6** miR-940 could target MAPK1.

**Notes:** (A) The sequences of MAPK1 3'UTR containing the miR-940 binding sites or mutant binding sites were shown. (B) The interaction between miR-940 and MAPK1 in TPC-1 and HTH83 cells was assessed by dual-luciferase reporter assay. (C and D) The mRNA and protein expression levels of MAPK1 in TC tissues (TC) and adjacent normal tissues (Normal) were determined by q-PCR and WB analysis. (E) WB analysis was used to measure the MAPK1 protein level in TC cells (TPC-1 and HTH83) and NTHY-OR13.1 cells. (F) The expression of miR-940 was detected by q-PCR to evaluate the transfection efficiency of anti-miR-940. (G) The effects of miR-940 mimic and inhibitor on the protein level of MAPK1 in TPC-1 and HTH83 cells were measured by WB analysis. \* $P < 0.05$ .


## miR-940 Inhibitor Could Invert the Suppression Effect of MAPK1 Silencing on the Progression of TC Cells

To further confirm our conclusion, we co-transfected si-MAPK1 and anti-miR-940 into TPC-1 and HTH83 cells. We first measured the transfection efficiency of both and found that si-MAPK1 could hinder MAPK1 protein level and anti-miR-940 could recover its protein level, indicating that the transfection of both was successful (Figure 7A). Subsequently, we performed colony formation and transwell assays. All results revealed that MAPK1 silencing inhibited proliferation, migration and invasion in TPC-1 and HTH83 cells, while miR-940 inhibitor could reverse this inhibition effect and recover the normal function of TC cells (Figure 7B–D). Furthermore, OCR detection assay also showed that miR-940 inhibitor inverted the suppression effect of silenced-MAPK1 on the metabolic

ability of TPC-1 and HTH83 cells (Figure 7E and F). Therefore, these results confirmed that the role of MAPK1 in TC was associated with miR-940 expression.

## MAPK1 Expression Was Regulated by Circ\_0058124 and miR-940

To further verify the existence of the circ\_0058124/miR-940/MAPK1 axis, we explored the effect of circ\_0058124 expression on the MAPK1 level. We co-transfected circ\_0058124 overexpression and miR-940 mimic into TPC-1 and HTH83 cells and then performed WB analysis. The results revealed that circ\_0058124 overexpression could markedly improve the protein level of MAPK1, and miR-940 mimic could partially reverse this promotion effect, thus significantly suppressing the MAPK1 protein level (Figure 8). These data suggested that MAPK1 indeed was a downstream gene of the circ\_0058124/miR-940 axis.



**Figure 7** Effects of MAPK1 on the progression of TC cells.

**Notes:** TPC-1 and HTH83 cells were co-transfected with si-MAPK1 and anti-miR-940. **(A)** WB analysis was used to assess the protein level of MAPK1 to evaluate the transfection efficiency of si-MAPK1 and anti-miR-940. **(B)** Colony formation assay was performed to detect the number of colonies in TPC-1 and HTH83 cells. **(C and D)** Transwell assay was used to measure the number of migrated and invaded TPC-1 and HTH83 cells. **(E and F)** The OCR of basal respiration and maximum respiration in TPC-1 and HTH83 cells was determined by Seahorse XF Extracellular Flux Analyzer. \* $P < 0.05$ .


## Knockdown of Circ\_0058124 Reduced TC Tumor Growth Through Regulating miR-940 and MAPK1 Expression

To further confirm our conclusion, we constructed mice xenograft models to perform *in vivo* experiments. After 5 weeks of recording, we found that the tumor volume of mice showed significant inhibition in the sh-circ\_0058124 group compared with that in the sh-NC group (Figure 9A). Besides, the tumor weight was also markedly decreased in the sh-circ\_0058124 group (Figure 9B). Meanwhile, we detected circ\_0058124 expression to determine whether the knockdown of circ\_0058124 was successful and the results determined that circ\_0058124 expression was remarkably suppressed in the sh-circ\_0058124 group (Figure 9C). To confirm the regulation

of circ\_0058124 on miR-940 and MAPK1, we detected miR-940 and MAPK1 expression in tumor tissues. The results showed that in the sh-circ\_0058124 group, miR-940 expression was significantly increased (Figure 9D), and MAPK1 protein level was obviously decreased (Figure 9E). These data again confirmed that the regulation of circ\_0058124 on TC progression through the miR-940/MAPK1 axis.


## Discussion

The pathogenesis of TC is quite complex, including the imbalance of many hormones, so TC is also an endocrine disease.<sup>22</sup> In addition, insufficient or high iodine intake is also a risk factor for the development of TC.<sup>23</sup> At present, the incidence of TC is still increasing year by year.<sup>24</sup>



**Figure 8** Effects of circ\_0058124 and miR-940 on MAPK1 expression.

**Notes:** TPC-1 and HTH83 cells were co-transfected with circ\_0058124 overexpression plasmid and miR-940 mimic. The protein level of MAPK1 in TPC-1 and HTH83 cells was measured by WB analysis. \* $P < 0.05$ .



**Figure 9** Effects of circ\_0058124 on TC tumor growth in vivo.

**Notes:** (A) Tumor volume was calculated with length  $\times$  width<sup>2</sup>/2 method at the indicated time points. (B) Tumor weight was measured in mice. (C) The expression of circ\_0058124 was detected by q-PCR to evaluate the transfection efficiency of sh-circ\_0058124. (D) MiR-940 expression was tested by q-PCR in tumors. (E) The protein level of MAPK1 was determined by WB analysis in tumors. \* $P < 0.05$ .

Although the prognosis of patients is good, the metastasis and recurrence of cancer still greatly increase the mortality of patients.<sup>4,5</sup> Therefore, elucidating the mechanisms that affect TC progression can help develop new clinical treatment strategies. In our study, we found that circ\_0058124 was highly expressed in TC and was higher in late TC (III and IV stages) than in early TC (I and II stages), which

was consistent with the results of Yao et al.<sup>12</sup> Besides, we demonstrated that circ\_0058124 was a stable cyclic transcript without the poly-A tail and resistant to RNase R digestion. Circ\_0058124 was mainly located in the cytoplasm, unlike the nucleus-dominated linear mRNA.

The role of circRNAs in gene expression regulation has attracted increasing attention. Although a large number of

circRNAs have been identified, only a small number of circRNAs have been confirmed. Here, we selected a novel circRNA, circ\_0058124, for functional verification. We found that knockdown of circ\_0058124 inhibited proliferation, migration, invasion and metabolic abilities in TC cells, suggesting that circ\_0058124 expression was crucial to the normal biological function of TC cells. Also, circ\_0058124 silencing reduced TC tumor growth, which once again confirmed the importance of circ\_0058124 for TC progression. Therefore, circ\_0058124 had the potential to be a target for the diagnosis and treatment of TC.

In view of the mechanism of circRNA that could act as ceRNA, we explored the circRNA-miRNA-mRNA regulatory network of circ\_0058124. The results showed that circ\_0058124 could sponge miR-940 to promote MAPK1 expression. Previous studies had demonstrated that miR-940 inhibited glioma cancer progression by suppressing mitochondrial metabolism, and it could restrain the proliferation and migration of breast cancer.<sup>25,26</sup> MAPK1 has been proved to be involved in the regulation of tumor progression through mediating proliferation and metastasis.<sup>27,28</sup> In the StarBase database, we found that miR-940 expression was lower and MAPK1 expression was higher in TC. Here, we confirmed that miR-940 was markedly decreased and MAPK1 was increased in TC. The functional test suggested that circ\_0058124 overexpression obviously reversed the inhibition effects of miR-940 overexpression on the proliferation, migration, invasion and metabolic abilities of TC cells. Also, the suppression of MAPK1 silencing on TC cell progression could be inverted by miR-940 inhibitor. Meanwhile, miR-940 and MAPK1 expression were regulated by circ\_0058124 in vitro and in vivo. All these data confirmed the existence of circ\_0058124/miR-940/MAPK1 regulatory network in TC. As a new circRNA, the proposed mechanism of circ\_0058124 provided a reference for us to explore the mechanism of circ\_0058124 in other cancers further, and also enriched the research background of circ\_0058124.

As a biomarker of cancer, many circRNAs have been proved to be related to the clinical and pathological characteristics of cancer, which can provide theoretical support for the functional studies of circRNAs.<sup>29</sup> Therefore, the research on circ\_0058124 still needed us to enrich. Our study focused on the effect of circ\_0058124 on TC cell proliferation, metastasis and metabolic abilities, but the expression of related proteins was not detected, which was also the shortcoming of our experimental design, so further exploration was needed in the follow-up experiments.

## Conclusion

In summary, our results suggested that circ\_0058124 promoted TC progression through regulating the miR-940/MAPK1 axis. Therefore, targeted therapy based on circ\_0058124/miR-940/MAPK1 might be an effective therapeutic strategy for TC.

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Hugen N, Sloot YJE, Netea-Maier RT, et al. Divergent metastatic patterns between subtypes of thyroid carcinoma results from the nationwide dutch pathology registry. *J Clin Endocrinol Metab.* 2019. doi:10.1210/clinem/dgz078
- Young S, Harari A, Smooke-Praw S, et al. Effect of reoperation on outcomes in papillary thyroid cancer. *Surgery.* 2013;154(6):1354–1361; discussion 1361–1352. doi:10.1016/j.surg.2013.06.043
- Kitahara CM, Sosa JA. The changing incidence of thyroid cancer. *Nat Rev Endocrinol.* 2016;12(11):646–653. doi:10.1038/nrendo.2016.110
- Ringel MD. Metastatic dormancy and progression in thyroid cancer: targeting cells in the metastatic frontier. *Thyroid.* 2011;21(5):487–492. doi:10.1089/thy.2011.2121
- Byeon HK, Kim SB, Oh HS, et al. Clinical analysis of pediatric thyroid cancer: a single medical institution experience of 18 years. *Ann Otol Rhinol Laryngol.* 2019;128(12):1152–1157.
- Li X, Yang L, Chen LL. The biogenesis, functions, and challenges of circular RNAs. *Mol Cell.* 2018;71(3):428–442. doi:10.1016/j.molcel.2018.06.034
- Ng WL, Mohd Mohidin TB, Shukla K. Functional role of circular RNAs in cancer development and progression. *RNA Biol.* 2018;15(8):995–1005. doi:10.1080/15476286.2018.1486659
- Li T, Sun X, Chen L. Exosome circ\_0044516 promotes prostate cancer cell proliferation and metastasis as a potential biomarker. *J Cell Biochem.* 2019.
- Lu J, Zhang PY, Xie JW, et al. Circular RNA hsa\_circ\_0006848 related to ribosomal protein L6 acts as a novel biomarker for early gastric cancer. *Dis Markers.* 2019;2019:3863458. doi:10.1155/2019/3863458
- Pan Y, Xu T, Liu Y, et al. Upregulated circular RNA circ\_0025033 promotes papillary thyroid cancer cell proliferation and invasion via sponging miR-1231 and miR-1304. *Biochem Biophys Res Commun.* 2019;510(2):334–338. doi:10.1016/j.bbrc.2019.01.108
- Yang Y, Ding L, Li Y, et al. Hsa\_circ\_0039411 promotes tumorigenesis and progression of papillary thyroid cancer by miR-1179/ABCA9 and miR-1205/MTA1 signaling pathways. *J Cell Physiol.* 2019.
- Yao Y, Chen X, Yang H, et al. Hsa\_circ\_0058124 promotes papillary thyroid cancer tumorigenesis and invasiveness through the NOTCH3/GATAD2A axis. *J Exp Clin Cancer Res.* 2019;38(1):318. doi:10.1186/s13046-019-1321-x
- Kwan JY, Psarianos P, Bruce JP, et al. The complexity of microRNAs in human cancer. *J Radiat Res.* 2016;57(Suppl S1):i106–i111. doi:10.1093/jrr/rw009
- Tutar L, Tutar E, Ozgur A, et al. Therapeutic targeting of microRNAs in cancer: future perspectives. *Drug Dev Res.* 2015;76(7):382–388. doi:10.1002/ddr.v76.7
- Tay Y, Rinn J, Pandolfi PP. The multilayered complexity of ceRNA crosstalk and competition. *Nature.* 2014;505(7483):344–352. doi:10.1038/nature12986

16. Luo H, Xu R, Chen B, et al. MicroRNA-940 inhibits glioma cells proliferation and cell cycle progression by targeting CKS1. *Am J Transl Res*. 2019;11(8):4851–4865.
17. Liu W, Xu Y, Guan H, et al. Clinical potential of miR-940 as a diagnostic and prognostic biomarker in breast cancer patients. *Cancer Biomark*. 2018;22(3):487–493. doi:10.3233/CBM-171124
18. Hu J, Li C, Liu C, et al. Expressions of miRNAs in papillary thyroid carcinoma and their associations with the clinical characteristics of PTC. *Cancer Biomark*. 2017;18(1):87–94. doi:10.3233/CBM-161723
19. Sun Y, Liu WZ, Liu T, et al. Signaling pathway of MAPK/ERK in cell proliferation, differentiation, migration, senescence and apoptosis. *J Recept Signal Transduct Res*. 2015;35(6):600–604. doi:10.3109/10799893.2015.1030412
20. Wang J, Xiao T, Zhao M. MicroRNA-675 directly targets MAPK1 to suppress the oncogenicity of papillary thyroid cancer and is sponged by long non-coding RNA RMRP. *Onco Targets Ther*. 2019;12:73 07–7321. doi:10.2147/OTT.S213371
21. Wang J, Yang H, Si Y, et al. Iodine promotes tumorigenesis of thyroid cancer by suppressing Mir-422a and up-regulating MAPK1. *Cell Physiol Biochem*. 2017;43(4):1325–1336. doi:10.1159/000481844
22. Mughal BB, Demeneix BA. Endocrine disruptors: flame retardants and increased risk of thyroid cancer. *Nat Rev Endocrinol*. 2017;13 (11):627–628. doi:10.1038/nrendo.2017.123
23. Dal Maso L, Bosetti C, La Vecchia C, et al. Risk factors for thyroid cancer: an epidemiological review focused on nutritional factors. *Cancer Causes Control*. 2009;20(1):75–86. doi:10.1007/s10552-008-9219-5
24. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2018. *CA Cancer J Clin*. 2018;68(1):7–30. doi:10.3322/caac.21442
25. Xu T, Zhang K, Shi J, et al. MicroRNA-940 inhibits glioma progression by blocking mitochondrial folate metabolism through targeting of MTHFD2. *Am J Cancer Res*. 2019;9(2):250–269.
26. Hou L, Chen M, Yang H, et al. MiR-940 inhibited cell growth and migration in triple-negative breast cancer. *Med Sci Monit*. 2016; 22:3666–3672. doi:10.12659/MSM.897731
27. Li W, Liang J, Zhang Z, et al. MicroRNA-329-3p targets MAPK1 to suppress cell proliferation, migration and invasion in cervical cancer. *Oncol Rep*. 2017;37(5):2743–2750. doi:10.3892/or.2017.5555
28. Li XW, Tuergan M, Abulizi G. Expression of MAPK1 in cervical cancer and effect of MAPK1 gene silencing on epithelial-mesenchymal transition, invasion and metastasis. *Asian Pac J Trop Med*. 2015;8(11):93 7–943. doi:10.1016/j.apjtm.2015.10.004
29. Jiang Z, Shen L, Wang S, et al. Hsa\_circ\_0028502 and hsa\_circ\_0076251 are potential novel biomarkers for hepatocellular carcinoma. *Cancer Med*. 2019. doi:10.1002/cam4.2584

## OncoTargets and Therapy

Dovepress

### Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic

agents and protocols on patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/oncotargets-and-therapy-journal>