

RETRACTED ARTICLE: Pumpkin Oil–Based Nanostructured Lipid Carrier System for Antiulcer Effect in NSAID-Induced Gastric Ulcer Model in Rats

This article was published in the following Dove Press journal:
International Journal of Nanomedicine

Osama AA Ahmed ^{1,2}
Usama A Fahmy¹
Rana Bakhaidar¹
Mohamed A El-Moselhy^{3,4}
Mohamed A Alfaleh⁵
Al-Shaimaa F Ahmed ⁴
Asmaa SA Hammad ⁴
Hibah Aldawsari¹
Nabil A Alhakamy ^{1,6–8}

¹Advanced Drug Delivery Research Group, Faculty of Pharmacy, King Abdulaziz University, Jeddah 21589, Saudi Arabia; ²Department of Pharmaceutics and Industrial Pharmacy, Faculty of Pharmacy, Minia University, Minia 61519, Egypt; ³Department of Pharmacology, School of Pharmacy, Ibn Sina National College, Jeddah 22413, Saudi Arabia; ⁴Department of Pharmacology and Toxicology, Faculty of Pharmacy, Minia University, Minia 61519, Egypt; ⁵Department of Natural Products and Alternative Medicine, Faculty of Pharmacy, King Abdulaziz University, Jeddah, Saudi Arabia; ⁶Department of Pharmaceutics, Faculty of Pharmacy, King Abdulaziz University, Jeddah 21589, Saudi Arabia; ⁷Center of Excellence for Drug Research and Pharmaceutical Industries, King Abdulaziz University, Jeddah 21589, Saudi Arabia; ⁸King Fahd Medical Research Center, King Abdulaziz University, Jeddah 21589, Saudi Arabia

Correspondence: Osama AA Ahmed
Advanced Drug Delivery Research
Group, Faculty of Pharmacy, King
Abdulaziz University, Jeddah 21589, Saudi
Arabia
Tel +966 599120686
Email oaahmed@kau.edu.sa

Background: Peptic ulcer disease, a painful lesion of the gastric mucosa, is considered one of the most common gastrointestinal disorders. This study aims to investigate the formulation of pumpkin seed oil (PSO)-based nanostructured lipid carriers (NLCs) to utilize PSO as the liquid lipid component of NLCs and to achieve oil dispersion in the nano-range in the stomach.

Methods: Box–Behnken design was utilized to deduce the optimum formula with minimum particle size. The optimized PSO-NLCs formula was investigated for gastric ulcer protective effects in Wistar rats by evaluating ulcer index and determination of gastric mucosa oxidative stress parameters.

Results: PSO was successfully incorporated as the liquid lipid (LL) component of NLCs. The prepared optimum PSO-NLCs formula showed a size of 64.3 nm. Pretreatment of animals using the optimized PSO-NLCs formula showed significantly ($p < 0.001$) lower ulcer index compared to indomethacin alone group and significantly ($p < 0.05$) less mucosal lesions compared to the raw oil.

Conclusion: These results indicated great potential for future application of optimized PSO-NLCs formula for antiulcer effect in non-steroidal anti-inflammatory drug (NSAID)-induced gastric ulcer.

Keywords: natural products, gastric ulcer, pumpkin seed oil, nano-lipid carriers, optimization, Box–Behnken experimental design

Introduction

Peptic ulcer disease (PUD) is a common gastrointestinal disorder with 10% prevalence in the human society.¹ It is a disease related to damage caused by balance disturbance between aggressive and defense factors in the stomach. The aggressive factors include pepsin and stomach acid secretion, active free radicals and oxidants, leukotrienes, endothelins, in addition to exogenous factors such as alcohol intake and nonsteroidal anti-inflammatory drugs (NSAIDs). Contrastingly, gastric mucin, prostaglandins (PGs), bicarbonate, nitric oxide (NO), growth factors, and antioxidant enzymes or antioxidant peptides like glutathione (GSH) constitute the defensive factors. Nonetheless, the most commonly affected organs are the lesser curvature in the stomach and proximal duodenum, however, ulceration may also occur anywhere in the gastrointestinal tract (GIT) from pylorus to cardia.^{2,3}

Importantly, the prolonged use of NSAIDs is the second most common cause of PUD.⁴ NSAIDs used for anti-inflammatory, antipyretic, pain-relief, anti-platelet

aggregation, and anti-thrombogenesis indications.⁵ In particular, Indomethacin, a member of NSAIDs family, is widely used for the management of rheumatoid arthritis, several inflammatory diseases, and for its well-established cardiovascular protection properties; however, its contribution to gastric ulceration has been documented in literature.² The induced inhibition of the cyclooxygenase enzyme (COX-2) enzyme is responsible for indomethacin's anti-inflammatory effect. Nevertheless, when used to alleviate inflammation and pain, it is known to exert a severe damaging effect on epithelial cells of the digestive tract, which constitutes its serious side effect. It is believed that the pathogenesis of indomethacin-induced gastric ulceration occurs via its potential to block the activities of the COX-1 enzyme, the major protective factor of gastrointestinal system, and the subsequent deficiency of protective factors such as prostaglandin E2 (PGE2), the production and secretion of mucus and bicarbonate, decreased mucosal blood flow, platelet aggregation dysfunction, impairment of microvascular structures.^{6,7} In addition, indomethacin increases aggressive factors, eg, acid, and oxidant parameters. On the other hand, indomethacin reduces anti-oxidant parameters; altogether indomethacin previously indicated the effects lead to epithelial damage.^{6,7}

Numerous treatment modalities are presently available to prevent indomethacin-induced peptic ulceration and to promote healing of mucosal damage, for instance, histamine receptor antagonists (H2RAs), proton pump inhibitors, PGs analogues, and gastroprotective agents.⁵ A superior drug to prevent and treat gastric-related side effects caused by NSAIDs in general remains somewhat controversial in clinical practice. Besides, most of these drugs have been reported to produce severe adverse reactions and toxicity upon chronic usage.⁸ Hence, a search for less toxic drugs is highly required, particularly in cases when they are to be used for an extended period.

Amongst the novel compounds recently researched for alleviation of gastric ulcer is pumpkin seed oil (PSO). Research has been carried out to investigate the potential efficacy of the aforementioned drug as a potent anti-oxidant for management and protection against peptic ulcer; yet data with this regard remain scarce in literature.⁹ PSO is rich in mono- and polyunsaturated fatty acids, mainly oleic and linoleic acid (37–41.7%).¹⁰ In addition, PSO contains carotenoids, in high concentration, and sterols as stigmastatrienol, stigmastadienol, and spinasterol.^{10,11} Reports have shown the therapeutic effects

of PSO, primarily highlighting the antidiabetic, antibacterial, anti-oxidant and anti-inflammatory properties of the edible oil with the highest contribution to the anti-oxidant capability being related to the polar fraction of the oil, mainly tocopherols.^{12–15} The mechanism underlying the anti-oxidant activity involves the blockage of 5-alpha reductase enzyme action.^{16,17}

Nanostructured lipid carriers (NLCs), second-generation solid lipid nanoparticles (SLNs), are high-performance pharmaceutical nanocarrier systems developed to enhance water solubility, stability as well as oil compound bioavailability.¹⁸ Mainly intended for parenteral administration of anti-cancer therapeutics, SLNs introduced in 1991, are nanosized particulate carrier system prepared either with physiological lipids or phospholipids, forming a lipid matrix that is solid at physiological temperature, with a size range of 50 to 1000 nm, dispersed in water vehicle, or alternatively, in an aqueous surfactant solution.^{19–25} Unlike most polymeric microspherical and nanoparticulate carrier systems, the production of both lipid-based nano-formulations, SLNs and NLCs, eliminates the employment of potentially toxic organic solvents, which often lead to detrimental effect on certain drugs. Nevertheless, due to their lipophilic nature, they are primarily developed for the purpose of incorporating lipophilic active pharmaceutical ingredients. However, hydrophilic drugs are also incorporated yet to a lesser extent.^{26–28}

In fact, NLCs were developed in order to overcome the formulation-associated pitfalls of the SLNs.²⁸ They are formulated using physiological, non-irritating lipids, unlike those used in forming polymeric nanoparticles (NPs). With a slight modification to the SLNs, NLCs are prepared by incorporating bioactive liquid oil component into the lipid-based formulation. Particularly enhancing oral delivery of drugs, the notable advantage raised by NLCs is the ability to encapsulate extensive drug quantities by the formation of imperfect, less structured lipid matrices, for better encapsulation efficiency.^{27,29,30} Importantly, the imperfection is owed to liquid oil incorporation within the core matrix of solid lipid (SL).³¹ Further to the remarkable advantages imparted by SLNs and other novel drug delivery systems of nanoparticles, NLCs demonstrated further enhancement of stability, reduced expulsion of the encapsulated drug from the carrier during storage period due to the imperfection of crystalline lattice, properties not possessed by SLNs.^{31,32} Moreover, using SLNs and NCLs has minimized the fairly large number of shortcomings associated with liposomes and NPs, such as difficulty of upscaling, high-cost production process and materials, and potential toxicity.^{19,33–35}

Lately, the trend towards using NLCs as vehicles for oils is extremely promising.²⁶ The NLC as a liquid core demonstrated minimal toxicity. In addition, it has enhanced the in vivo performance and potentiated the immunosuppressive effects of the carried drug Tacrolimus, through inhibition of Interleukin-2 (IL-2) cytokine release. Furthermore, Muchow et al have developed a paste-like formulation of omega-3-loaded NLC in order to chemically stabilize the fatty acids.³⁶

This study aims to investigate the formulation of PSO-based NLCs in order to utilize PSO as the liquid lipid (LL) component of NLCs and to achieve oil dispersion in the nano-range in the stomach. Box–Behnken design was utilized to deduce the optimum formula. The optimized PSO-NLCs formula was investigated for gastric ulcer protective effects by evaluating ulcer index and determination of gastric mucosa oxidative stress parameters.

Materials and Methods

Materials

Pumpkin seed oil (PSO), d- α -tocopheryl polyethylene glycol 1000 succinate (TPGS) and Tween 80 were purchased from Sigma-Aldrich (St. Louis, MI, USA). Precirol[®] ATO was obtained as a kind gift from Gattefosse (Saint-Priest, France). Soybean L- α -phosphatidylcholine (soybean lecithin) was purchased from Lipoid (Ludwigshafen, Germany). The PSO oil has been used as supplied.

Formulation of PSO-NLCs

Precirol was utilized as the solid lipid (SL), PSO was utilized as the liquid lipid (LL) and Tween 80: TPGS: soybean lecithin (2:2 ratio) mixture was used as surfactant in the formulation of the NLCs. PSO-NLCs were formulated by high-shear homogenization and ultrasonication technique as previously reported.^{25,37,38} Briefly, lipid formula components, precirol (SL) and PSO (LL), were melted at 75 °C. Separately, aqueous surfactant mixture solution (Tween 80: TPGS: soybean lecithin) was heated to 75 °C and mixed with the melted lipids. The mixture was homogenized (T25 Ultra-Turrax (IKA[®] Werke GmbH & Co. KG, Staufen, Germany)) at 12,000 rpm for 60 seconds, in a water bath at 75 °C. The mixture was then subjected to probe-sonication for 5 minutes. The final volume was adjusted to 20 mL using distilled water. The formed nanoemulsion was then left to cool to form PSO-NLCs and stored at 20 °C.

Optimization of PSO-NLCs

The development of the Box–Behnken experimental design for PSO-NLCs formulation components was carried out, based on preliminary investigation, using the Statgraphics plus, version 4 (Statgraphics software) (Manugistic Inc., PA, USA). The selected factors were the concentrations of precirol as SL (X1); the concentration of oil PSO as LL (X2); and the sonication time (ST) (X3). The design response was PSO-NLCs size (nm). The design goal was set to minimize the size of the investigated response (PSO-NLCs). The design generated 15 formulations of various combination of the investigated factors that were prepared and described in the 'Formulation of PSO-NLCs' section.

PSO-NLCs Size Determination

PSO-NLCs size was determined utilizing Nano-ZS particle size analysis (Malvern Instrument, Worcestershire, UK). One hundred microliters of each PSO-NLCs formulation were 100 times diluted with distilled water that was passed through a 0.1- μ m membrane filter, vortexed for 1 minute and then measured.

Prediction and Preparation of Optimized PSO-NLCs Formulation

The data collected from PSO-NLCs formulations, proposed by the experimental design, were statistically analyzed utilizing the software (ANOVA and multiple response optimization). The proposed optimum formulation obtained (predicted formula) was practically prepared and compared to the predicted formula by the design for result validation.

Fourier-Transform Infrared Physicochemical Characterization of the Optimized PSO-NLCs

The optimized PSO-NLCs formula and formula components were assessed using Fourier-transform infrared (FTIR) analysis as previously described.³⁹ Briefly, FTIR spectra of PSO, precirol, TPGS, Tween 80, soybean lecithin and the prepared optimized PSO-NLCs formula were recorded over the wavelength range from 400 to 4000 cm⁻¹ using FTIR spectrophotometer (Nicolet IZ 10, Thermo Fisher Scientific, Waltham, MA, USA). Samples were directly applied to the FTIR spectrophotometer without treatment.

In vivo Evaluation of Optimized PSO-NLCs Formulation

Animals

Adult male Wistar rats (180–200 gm) were obtained from the animal house of King Fahd Medical Research Center, King Abdulaziz University. Animals were acclimatized for 1 week before the experiment. The study protocol was approved by the Faculty of Pharmacy Research Ethics Committee, King Abdulaziz University (Reference #: PH 122–41). Care and use of animals according to the EU Directive 2010/63/EU and DHEW publication NIH 80–23 was ensured. One day prior the induction of gastric ulcers, all rats were fasted in mesh-bottomed cages to minimize coprophagia with free access to water. The rats were then divided to four groups (8 animals each): 1) control group: non-treated rats with no induction of ulcer, 2) indomethacin group: in which the rats received 50 mg/kg of indomethacin, 3) PSO group: in which the rats received pure PSO 30 minutes before injection of indomethacin and finally 4) optimized PSO-NLCs formula group: in which the rats received optimized PSO-NLCs formula 30 minutes before injection of indomethacin. PSO was given at a dose of 100mg/kg and PSO-NLCs optimized formulation was given as an equivalent dose and was administered orally 30 minutes before the induction of ulcer. Gastric ulceration was induced by an intraperitoneal injection of indomethacin (50mg/kg). Four hours later, all rats were sacrificed by decapitation. Their stomachs were removed, opened along the greater curvature, and washed with ice-cold saline. Animal stomachs were scored for macroscopic gross mucosal lesions. Gastric mucosae were collected and stored at -80°C until used for estimation of oxidative stress parameters. Another set of stomachs from each group was immersed in 10% formalin for histopathological examination.

Gastric Mucosal Lesion Assessment

Mucosal lesions in all animal groups were quantified according to the previously described method by Szabo and Hollander.⁴⁰ Briefly, images were captured for pinned stomachs and areas of mucosal damage were measured using ImageJ software and then expressed as % of the total stomach surface area. For each group, mean ulcer score expressed as ulcer index (U.I) and the percentage of inhibition (preventive index) against indomethacin-induced ulcers was determined using the equation:

$$\text{Ulcer inhibition (\%)} = \left(\frac{\text{U.I. in indomethacin} - \text{U.I. in treated rats}}{\text{U. I. in indomethacin}} \right) \times 100 \quad (1)$$

Determination of Gastric Mucosa Oxidative Stress Parameters

Gastric mucosal tissues were homogenized (0.1 g/mL) using phosphate buffer saline (ice-cold) using then centrifuged for 20 minutes at 4°C . The following parameters were calculated from the aspirated supernatant:

- Malondialdehyde (MDA), a measure of lipid peroxidation, was determined according to the method of Uchiyama and Mihara.⁴¹
- Nitric oxide (NO) was assayed colorimetrically using Griess reagent.⁴²
- Catalase activity was determined using a commercially available kit (Biodiagnostic, Egypt), according to the method of Fossati, Principe.⁴³

Results and Discussion

Formulation and Optimization of PSO-NLCs

NLCs composed of SL and LL. The inclusion of LL in NLCs, different from SL nanoparticles, aims to reduce crystallinity and increase the fluidity of the matrix with reduced packaging density.^{44–46} This leads to improved storage life when compared with SL nanoparticles.⁴⁴ The important criteria for efficacy and efficiency of nanoparticles (drug release, biodistribution and cellular uptake) are particle size and size distribution.^{46,47} Table 1 shows the PSO-NLCs size variabilities for the prepared formulations. The results revealed that the size ranged from 65.0 to 284.0 nm for formulations F13 and F14, respectively. The polydispersity index for the prepared PSO-NLCs formulations was in the range of 0.2–0.5 that shows acceptable unimodal size distribution. Two-way ANOVA analysis showed a significant antagonistic effect of the SL (X1) and ST (X3) percentages on the PSO-NLCs size with p -values of 0.00001 and 0.0001, respectively (Table 2, Figure 1). In addition, the quadratic term of X3 showed a significant synergistic effect on PSO-NLCs size with a p -value of 0.0313. The equation of PSO-NLCs size prediction according to correlation with the factors is shown in Equation (1).

$$\begin{aligned} \text{PSO - NLCs size (nm)} = & 491.65 - 162.5X_1 - 17.32X_2 \\ & - 24.44X_3 - 242.59X_1^2 \\ & - 11.11X_1X_2 - 5.83X_1X_3 \\ & - 20.37X_2^2 + 17.5X_2X_3 \\ & + 2.26X_3^3 \end{aligned}$$

Table 1 Experimental Runs and the Observed Globule Sizes (Observed and Fitted Values)

PSO NLCs Formula No.	Factors (X1–X3)			Response	
	SL (%)	LL (%)	ST (min)	Globule Size (nm)	
				Observed	Fitted
1	0.75	0.25	3.0	173.0	172.7
2	0.90	0.10	3.0	86.0	82.5
3	0.75	0.25	3.0	174.0	172.7
4	0.60	0.40	3.0	248.0	251.5
5	0.75	0.10	1.0	205.0	205.8
6	0.90	0.40	3.0	92.0	87
7	0.75	0.25	3.0	171.0	172.7
8	0.75	0.40	1.0	198.0	200.3
9	0.75	0.40	5.0	168.0	167.3
10	0.60	0.10	3.0	241.0	246.0
11	0.75	0.10	5.0	154.0	151.8
12	0.90	0.25	1.0	115.0	117.8
13	0.90	0.25	5.0	65.0	70.8
14	0.60	0.25	1.0	284.0	278.3
15	0.60	0.25	5.0	241.0	238.3

Abbreviations: PSO, pumpkin seed oil; NLCs, nanostructured lipid carriers; X1, concentrations of precirol as solid lipid; X2, the concentration of oil PSO as liquid lipid; X3, the sonication time; SL, solid lipid; LL, liquid lipid; ST, sonication time.

Table 2 Statistical Analysis of Variance (ANOVA) of the PSO-NLCs Size

Factor	PSO-NLCs Size	
	Estimate	p-value
X1	-164.0	0.0001*
X2	5.0	0.2815
X3	-43.5	0.0001*
X1X1	-10.17	0.1335
X1X2	0.5	0.9353
X1X3	-3.5	0.5763
X2X2	-0.9137	0.8864
X2X3	1.3	0.1331
X3X3	8.083	0.0313*
R ²	99.18%	
Adj R ²	99.18%	
SEE	5.86	
MAE	2.89	
PRESS	12.99	

Note: *Significant effect of factors on PSO-NLCs size at p < 0.05.
Abbreviations: PSO, pumpkin seed oil; NLCs, nanostructured lipid carriers; X1, concentrations of precirol as solid lipid; X2, the concentration of oil PSO as liquid lipid; X3, the sonication time; R², coefficient of determination; Adj R², adjusted coefficient of determination; SEE, standard error of estimate; MAE, mean absolute error; PRESS, predicted residual error sum of squares.

The results indicated that increasing SL (precirol) percent (content) in the formulation showed a reduction in the produced PSO-NLCs size. This has been observed in

Standardized Pareto Chart for SIZE




Figure 1 Standardized Pareto chart showing the significance of X1, X2 and X3 and their combined effects on PSO-NLCs size.
Abbreviations: X1, concentrations of precirol as solid lipid; X2, the concentration of oil PSO as liquid lipid; X3, the sonication time.

formulation number, 2, 6, 12 and 13. The reduction in PSO-NLCs size with increased SL content is attributed to the formation of more dense rigid crystalline structure of the formed nanoparticles.⁴⁶ In addition, the inverse relationship between ST (X3) and PSO-NLCs size is attributed to the ability of ultrasound sonication force to breakdown the coarse emulsion droplets to smaller nano-range sizes. Consequently, increased ST provides more energy to breakdown emulsion droplets to smaller sizes.⁴⁸

On the other hand, the results of pareto chart, **Figure 1**, showed a direct relationship, although non-significant at the specified concentration range, between LL and PSO-NLCs size. Previous reports revealed the direct relationship between LL and PSO-NLCs size.^{45,46,48,49} The




Figure 2 3D response surface plots showing the effects of X1, X2 and X3 on the investigated PSO-NLCs size.
Abbreviations: SL, solid lipid; LL, liquid lipid; ST, sonication time.

Table 3 Optimum Levels for PSO-NLCs Factors and Predicted, Actual and Residual Values for PSO-NLCs Size

Factor	Optimum Level	Low Level	High Level
X1	0.9	0.6	0.9
X2	0.1	0.1	0.4
X3	5.0	1.0	5.0
Response	Prediction	Actual	Residual
PSO-NLCs size	62.9 nm	64.3 nm	1.4
Desirability constraint		Minimize the particle size	

Abbreviations: PSO, pumpkin seed oil; NLCs, nanostructured lipid carriers; X1, concentrations of precirol as solid lipid; X2, the concentration of oil PSO as liquid lipid; X3, the sonication time.

rational for this relation either unknown⁴⁸ or attributed to the inability of surfactant to cover the melted lipid droplets' surface when the LL-to-SL ratio was more than

50%.⁴⁹ Pareto chart and response surface plot revealed the relationship between the investigated factors (X1–X3) and PSO-NLCs size (Figures 1 and 2).

Validation of the PSO-NLCs Optimized Formula

The obtained data from the 15 formulations generated by the experimental design were analyzed with ANOVA. The Box–Behnken design predicted the optimum formulation that was practically prepared and evaluated and compared with the predicted results generated by the design (Table 3). The prepared optimum formula showed a size of 64.3 nm that was compared with the predicted value (62.9 nm) of PSO-NLCs size generated by the design (Table 3). The optimized PSO-NLCs formulation was utilized in the in vivo evaluation




Figure 3 Size distribution of optimized PSO-NLCs. **Abbreviation:** PSO-NLCs, pumpkin seed oil nanostructured lipid carriers.




Figure 4 FTIR spectra of PSO, Tween 80, precirol, TPGS, soybean lecithin and the optimized PSO-NLCs formula. **Abbreviations:** FTIR, Fourier-transform infra-red; TPGS, d- α -Tocopheryl polyethylene glycol 1000 succinate; PSO-NLCs, pumpkin seed oil nanostructured lipid carriers.




Figure 5 Bar graphs showing the effect of indomethacin, PSO and PSO-NLCs formula on ulcer index (**A**) and preventive index (**B**). Representative photos of the stomachs from the four different groups (**C**).

Notes: Data are presented as mean \pm S.E.M. *Significantly different from indomethacin at $p < 0.05$; **significantly different from indomethacin at $p < 0.01$; ***significantly different from indomethacin at $p < 0.001$; #significantly different from PSO at $p < 0.01$.

Abbreviation: Indo, Indomethacin.

studies. Size distribution of the optimized formula is shown in Figure 4 that revealed unimodal narrow size distribution.

FTIR Physicochemical Characterization of the Optimized PSO-NLCs

Figure 4 shows the FTIR spectra of optimized PSO-NLCs and its individual formula components. The main PSO IR peaks were $3300:3500\text{ cm}^{-1}$ as weak broad peaks that refer to the OH and COOH groups. The indicated PSO OH and COOH did not interfere with the characteristic peak region (around 3000 cm^{-1}) of other formula components. The results indicated no change in the characteristic functional group peaks of individual components when formulated in

the optimized PSO-NLCs. FTIR is a useful tool for the evaluation of possible formula components interaction. Incompatibility among formula components could be predicted by changes in the characteristic peaks of the functional groups of each component of the optimized formula.

In vivo Evaluation of Optimized PSO-NLCs Formulation

Effect of PSO and PSO-NLCs Formula on Indomethacin-Induced Gastric Lesions

As shown in Figure 5, indomethacin resulted in the development of ulcer lesions, which were quantified as ulcer index of 6.2 ± 0.6 (Figure 5A). Pretreatment using both PSO and




Figure 6 Bar graphs showing the effect of indomethacin, PSO and PSO-NLCs formula on mucosal MDA (A), mucosal catalase (B) and mucosal nitrites (C). **Notes:** Data are presented as mean \pm SEM. **Significantly different from indomethacin at $p < 0.01$. ***Significantly different from indomethacin at $p < 0.001$. #Significantly different from PSO at $p < 0.05$. **Abbreviation:** Indo, indomethacin.

optimized PSO-NLCs formula resulted in a significantly ($p < 0.01$ and 0.001 , respectively) lower UI compared to indomethacin (Figure 5B). The effect of the formula was more pronounced showing significantly ($p < 0.05$) less mucosal lesions compared to the raw oil. Representative photos of the stomachs from the four different groups are shown in Figure 5C.

Effect of PSO and PSO-NLCs Formula on Gastric Mucosal Oxidative Stress Parameters

Lipid peroxidation, catalase activity and total nitrite levels were evaluated in the gastric mucosal tissues. As shown

in Figure 6, the results for the indomethacin group indicated that gastric MDA and total nitrite levels were elevated compared to control that indicated increased oxidative stress. Similarly, the activity of catalase was higher than the control rat group ($p < 0.001$) reflecting a compensatory increase in antioxidative parameters to counteract the elevated reactive oxygen species generation. PSO and PSO-NLCs formula administration had protective effects against these alterations showing significantly lower MDA, NO and catalase activity (Figure 6A–C).




Figure 7 Representative photomicrographs of H&E stained stomachs of: **(A)** control; **(B)** indomethacin-treated rats (ulcer model) showed damage and loss of epithelial layer and gastric pits and increased mucosal thickness with distorted gastric glands with inflammatory cells infiltration of the submucosa; **(C)** PSO + indomethacin showed a mild damage and loss of epithelial layer with slight decreased in mucosal thickness and dilation of gastric glands; **(D)** PSO-NLCs formula + indomethacin showed marvelous amelioration of epithelial layer and gastric pits with normal thickness of mucosa (magnification = 200 \times). H&E stain.

Histopathological Examination of Stomach Section (PSO versus PSO-NLCs Formula)

Figure 7 shows the results of histopathological examination of H&E stained stomach sections showing normal structure with no evidence of inflammation or ulceration in control rats (Figure 7A). Sections from indomethacin-treated groups show features of acute gastritis in the form of foveolar hyperplasia, edema, hyperemia and focal necrosis of foveolar cells. The lamina propria shows signs of neutrophilic infiltration (Figure 7B). No pathologic lesions could be detected in muscularis propria. Sections from PSO-treated rats showed pits of normal gastric mucosal glands with no ulceration in which one of them is found to have focal foveolar necrosis, mild edema and hyperemia in the lamina propria with submucosal area of congestion and hyperemia and no abnormalities in muscularis propria (Figure 7C). The stomach of optimized PSO-NLCs formula-treated rats (Figure 7D) shows normal gastric mucosal glands with foveolar arrangement and of normal length. No inflammation or infiltrates in lamina propria could be detected.

Gastric ulcer occurs when there is an imbalance between certain aggressive factors and defensive endogenous factors.

There is therefore a great need for healthy, economic and effective antiulcer agents. Natural products have emerged as a source of compounds with potential antiulcer activity.^{50,51} Previous report by our group investigated the PSO solubilizing ability of ibuprofen in self-nanoemulsifying drug delivery system for improved solubility and as protection factor from peptic ulcer induced by the enhanced solubility of ibuprofen.⁵² These promising results were taken a step further to prove the efficacy of PSO in ulcer protection through formulation into NLCs with improved stability and efficacy characteristics when compared with the self-nanoemulsifying drug delivery system. Optimized PSO-NLCs showed improved efficacy in protection of anti-inflammatory drug-induced ulcer. The protection is attributed to PSO components (polyunsaturated fatty acids, tocopherol and sterols). In addition, PSO has been reported for wound healing characteristics.^{14,53} The optimized formula could have the ability to re-epithelialize the internal tissues as a result of tocopherol content of formula (from PSO and TPGS) that has scavenger activity of peroxy, hydroxyl, and superoxide radicals with the ability to heal the internal ulceration.¹² In addition, soybean lecithin content of the

surfactant mixture in the optimized PSO-NLCs formulation offers gastric mucosal barrier.⁵³

Conclusion

In this study, PSO was successfully incorporated as the LL component of NLCs. Box–Behnken experimental design for PSO-NLCs formulation components was carried out to achieve oil dispersion in the smallest formulation size in the stomach. Pretreatment using the optimized PSO-NLCs formula showed lower UI compared to indomethacin and less mucosal lesions compared to the raw oil. These results indicated great potential for future application of optimized PSO-NLCs formula for antiulcer effect in NSAID-induced gastric ulcer.

Acknowledgments

This project was funded by the Deanship of Scientific Research (DSR) at King Abdulaziz University, Jeddah, under grant no. RG-2–166–40. The authors, therefore, acknowledge with thanks the DSR for technical and financial support.

Disclosure

The authors report no conflicts of interest in this work.

References

- Zapata-Colindres JC, Zepeda-Gómez S, Montañez-Cruz A, Ballesteros E, de Jesús Villalobos J, Valdovinos-Andrade E. The association of *Helicobacter pylori* infection and non-steroidal anti-inflammatory drugs in peptic ulcer disease. *Can J Gastroenterol*. 2006;20(12):277–280. doi:10.1155/2006/175217
- Fazalda A, Quraisiah A, Nur Azhara MF. Antiulcer effect of honey in nonsteroidal anti-inflammatory drugs induced gastric ulcer model in rats: a systematic review. *Evidence-Based Complement Altern Med*. 2018;2018:7515692. doi:10.1155/2018/7515692
- Takeuchi K. Pathogenesis of NSAID-induced gastric damage: importance of cyclooxygenase inhibition and gastric hypermotility. *World J Gastroenterol*. 2016;18(18):2147–2160. doi:10.3748/wjg.v18.i18.2147
- Shahin NN, Abdelkader MM. A novel role of irbesartan in gastroprotection against indomethacin-induced gastric injury in rats: targeting DDAH, NO, MA and EGFR/ERK signaling. *Sci Rep*. 2018;8(1):1–12. doi:10.1038/s41598-018-22727-6
- Fang YF, Xu WL, Wang L, et al. Effect of hydrotalcite on indomethacin-induced gastric injury in rats. *Biomed Res Int*. 2019;2019:1–9. doi:10.1155/2019/4605748
- Wallace JL, McKnight W, Reuter BK, Vergnolle N. NSAID-induced gastric damage in rats: requirement for inhibition of both cyclooxygenase 1 and 2. *Gastroenterology*. 2000;119(3):706–714. doi:10.1053/gast.2000.16510
- Wright JM. The double-edged sword of COX-2 selective NSAIDs. *CMAJ*. 2002;167(10):1131–1137.
- Zainol S, Basri M, Basri HB, et al. Formulation optimization of a palm-based nanoemulsion system containing levodopa. *Int J Mol Sci*. 2012;13(10):13049–13064. doi:10.3390/ijms131013049
- Scally B, Emberson JR, Spata E, et al. Effects of gastroprotectant drugs for the prevention and treatment of peptic ulcer disease and its complications: a meta-analysis of randomised trials. *Lancet Gastroenterol Hepatol*. 2018;3(4):231–241. doi:10.1016/S2468-1253(18)30037-2
- Montesano D, Blasi F, Simonetti MS, Santini A, Cossignani L. Chemical and nutritional characterization of seed oil from *Cucurbita maxima* L. (Var. *Berrettina*) pumpkin. *Foods*. 2018;7(3):30. doi:10.3390/foods7030030
- Siano F, Straccia MC, Paolucci M, Fasulo G, Boscaino F, Volpe MG. Physico-chemical properties and fatty acid composition of pomegranate, cherry and pumpkin seed oils. *J Sci Food Agric*. 2016;96(5):1730–1735. doi:10.1002/jsfa.7279
- Farzaei MH, Shams-Ardekani MR, Abbasabadi Z, Rahimi R. Scientific evaluation of edible fruits and spices used for the treatment of peptic ulcer in traditional Iranian medicine. *IRN Gastroenterol*. 2013;2013:1–12. doi:10.1155/2013/136922
- Adams GG, Imran S, Wang S, et al. The hypoglycaemic effect of pumpkins as anti-diabetic and functional medicines. *Food Res Int*. 2011;44(4):862–867. doi:10.1016/j.foodres.2011.03.011
- Bardaa S, Ben Halima N, Alami F, et al. Oil from pumpkin (*Cucurbita pepo* L.) seeds: evaluation of its functional properties on wound healing in rats. *Lipids Health Dis*. 2016;15(1). doi:10.1186/s12944-016-0237-0
- Caili F, Huan S, Jianhong L. A review of pharmacological activities and utilization technologies of pumpkin. *Plant Foods Hum Nutr*. 2006;61(2):73–80. doi:10.1007/s11130-006-0016-6
- Fruhwieler GO, Wenzl M, El-Toukhy R, Wagner FS, Hermetter A. Fluorescence screening of antioxidant capacity in pumpkin seed oils and other natural oils. *Eur J Lipid Sci Technol*. 2003;105(6):266–274. doi:10.1002/ejlt.200390055
- Stevenson DG, Filer FJ, Wang L, Jane JL, Wang T, Inglett GE. Oil and phospholipid content and composition of pumpkin seed oil in 12 cultivars. *J Agric Food Chem*. 2007;55:4005–4013. doi:10.1021/jf06979
- Zhu QY, Guissi F, Yang RY, et al. Preparation of deep sea fish oil-based nanostructured lipid carriers with enhanced cellular uptake. *J Nanosci Nanotechnol*. 2015;15(12):9539–9547. doi:10.1166/jnn.2015.10880
- Wissing SA, Kayser O, Müller RH. Solid lipid nanoparticles for parenteral drug delivery. *Adv Drug Deliv Rev*. 2004;56(9):1257–1272. doi:10.1016/j.addr.2003.12.002
- Malam Y, Loizidou M, Seifalian AM. Liposomes and nanoparticles: nanosized vehicles for drug delivery in cancer. *Trends Pharmacol Sci*. 2009;30(11):592–599. doi:10.1016/j.tips.2009.08.004
- Müller RH, Mäder K, Gohla S. Solid lipid nanoparticles (SLN) for controlled drug delivery - a review of the state of the art. *Eur J Pharm Biopharm*. 2000;50(1):161–177. doi:10.1016/S0939-6411(00)00087-4
- Hao J, Fang X, Zhou Y, et al. Development and optimization of solid lipid nanoparticle formulation for ophthalmic delivery of chloramphenicol using a Box–Behnken design. *Int J Nanomedicine*. 2011;6:683–692. doi:10.2147/ijn.s17386
- Almeida AJ, Souto E. Solid lipid nanoparticles as a drug delivery system for peptides and proteins. *Adv Drug Deliv Rev*. 2007;59(6):478–490. doi:10.1016/j.addr.2007.04.007
- Mukherjee S, Ray S, Thakur RS. Solid lipid nanoparticles: a modern formulation approach in drug delivery system. *Indian J Pharm Sci*. 2009;71(4):349–358. doi:10.4103/0250-474X.57282
- Akhoond Zardini A, Mohebbi M, Farhoosh R, Bolurian S. Production and characterization of nanostructured lipid carriers and solid lipid nanoparticles containing lycopene for food fortification. *J Food Sci Technol*. 2018;55(1):287–298. doi:10.1007/s13197-017-2937-5
- Khan SS, Ganguli M, Aditya A, Khan SS, Baboota S, Ali J. Improved in vivo performance and immunomodulatory effect of novel Omega-3 fatty acid based Tacrolimus nanostructured lipid carrier. *J Drug Deliv Sci Technol*. 2019;52:138–149. doi:10.1016/j.jddst.2019.04.019

27. Gaba B, Fazil M, Ali A, Baboota S, Sahni JK, Ali J. Nanostructured lipid (NLCs) carriers as a bioavailability enhancement tool for oral administration. *Drug Deliv.* 2015;22(6):691–700. doi:10.3109/10717544.2014.898110
28. Joshi M, Patravale V. Nanostructured lipid carrier (NLC) based gel of celecoxib. *Int J Pharm.* 2008;346(1–2):124–132. doi:10.1016/j.ijpharm.2007.05.060
29. Tiwari R, Pathak K. Nanostructured lipid carrier versus solid lipid nanoparticles of simvastatin: comparative analysis of characteristics, pharmacokinetics and tissue uptake. *Int J Pharm.* 2011;415(1–2):232–243. doi:10.1016/j.ijpharm.2011.05.044
30. Naseri N, Valizadeh H, Zakeri-Milani P. Solid lipid nanoparticles and nanostructured lipid carriers: structure preparation and application. *Adv Pharm Bull.* 2015;5(3):305–313. doi:10.15171/apb.2015.043
31. Lee YH, Chang SH, Tsai YF, Fang JY, Hwang TL. Oleic acid-loaded nanostructured lipid carrier inhibit neutrophil activities in the presence of albumin and alleviates skin inflammation. *Int J Nanomedicine.* 2019;14:6539–6553. doi:10.2147/IJN.S208489
32. Khosa A, Reddi S, Saha RN. Nanostructured lipid carriers for site-specific drug delivery. *Biomed Pharmacother.* 2018;103:598–613. doi:10.1016/j.biopha.2018.04.055
33. Aboutaleb E, Atyabi F, Khoshayand MR, et al. Improved brain delivery of vincristine using dextran sulfate complex solid lipid nanoparticles: optimization and in vivo evaluation. *J Biomed Mater Res - Part A.* 2014;102(7):2125–2136. doi:10.1002/jbm.a.34890
34. Serini S, Cassano R, Corsetto PA, Rizzo AM, Calviello G, Trombino S. Omega-3 PUFA loaded in resveratrol-based solid lipid nanoparticles: physicochemical properties and antineoplastic activities in human colorectal cancer cells in vitro. *Int J Mol Sci.* 2018;19(2):586. doi:10.3390/ijms19020586
35. Zhang C, Gu C, Peng F, et al. Preparation and optimization of triptolide-loaded solid lipid nanoparticles for oral delivery with reduced gastric irritation. *Molecules.* 2013;18(11):13340–13356. doi:10.3390/molecules181113340
36. Muchow M, Schmitz EI, Despatova N, Maincent P, Müller RH. Omega-3 fatty acids-loaded lipid nanoparticles for patient-compliant oral bioavailability enhancement. *Pharmazie.* 2009;64(8):497–504. doi:10.1016/j.ph.2009.09.084
37. Beloqui A, Del Pozo-Rodríguez A, Isla J, Rodríguez-Galán A, Solinís MÁ. Nanostructured lipid carriers as oral delivery systems for poorly soluble drugs. *J Drug Deliv Sci Technol.* 2017;42:144–154. doi:10.1016/j.jds.2017.06.013
38. Weber S, Zimmer A, Pardeike L. Solid lipid nanoparticles (SLN) and nanostructured lipid carriers (NLC) for pulmonary application: a review of the state of the art. *Eur J Pharm Biopharm.* 2014;86(1):7–22. doi:10.1016/j.ejpb.2013.08.013
39. Ahmed OAA. Development and single dose clinical pharmacokinetics investigation of the whey protein assisted alpha lipoic acid nanoencapsulation. *Jordanian Sci Res.* 2018;8(1):15802. doi:10.1038/s41598-018-3423-8
40. Szabos, Hollander D. Pathways of gastrointestinal protection and repair: mechanisms of action of sucralfate. *Am J Med.* 1989;86(6):23–31. doi:10.1016/0002-9343(89)90153-8
41. Mihara M, Uchiyama M. Determination of malonaldehyde precursor in tissues by thiobarbituric acid test. *Anal Biochem.* 1978;86(1):271–278. doi:10.1016/0003-2697(78)90342-1
42. Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JS, Tannenbaum SR. Analysis of nitrate, nitrite, and [15N]nitrate in biological fluids. *Anal Biochem.* 1982;126(1):131–138. doi:10.1016/0003-2697(82)90118-X
43. Fossati P, Prencipe L, Berti G. Use of 3,5-dichloro-2-hydroxybenzenesulfonic acid/4-aminophenazone chromogenic system in direct enzymic assay of uric acid in serum and urine. *Clin Chem.* 1980;26(2):227–231. doi:10.1093/clinchem/26.2.227
44. Muller RH, Keck CM. Challenges and solutions for the delivery of biotech drugs - a review of drug nanocrystal technology and lipid nanoparticles. *J Biotechnol.* 2004;113:151–170. doi:10.1016/j.jbiotec.2004.06.007
45. Pinheiro M, Ribeiro R, Vieira A, Andrade T, Reis S. Design of a nanostructured lipid carrier intended to improve the treatment of tuberculosis. *Drug Des Devel Ind Pharm.* 2016;10:246–2475. doi:10.2147/DDDT.S104395
46. Bahari LAS, Hamisheh B, H. The impact of variables on particle size of solid lipid nanoparticles and nanostructured lipid carriers; A comparative literature review. *Adv Pharm Bull.* 2016;6(2):143–151. doi:10.15171/apb.2016.021
47. El-Say KM, Ahmed OAA, Mohamed AI, Safo MK, Omar AM. Zein-alpha lipoic acid-loaded nanoparticles to enhance the oral bioavailability of dapoxetine: optimization and clinical pharmacokinetic evaluation. *Int J Nanomedicine.* 2019;14:7461–7473. doi:10.2147/IJN.S24611
48. Das S, Ng WK, Tan RBH. Are nanostructured lipid carriers (NLCs) better than solid lipid nanoparticles (SLNs): development, characterizations and comparative evaluations of clotrimazole-loaded SLNs and NLCs. *Eur J Pharm Sci.* 2012;47(1):139–151. doi:10.1016/j.ejps.2011.05.010
49. Song A, Zhang X, Li Y, Mao X, Han F. Effect of liquid-to-solid lipid transition on characterizations of flurbiprofen-loaded solid lipid nanoparticles (SLNs) and nanostructured lipid carriers (NLCs) for transdermal administration. *Drug Dev Ind Pharm.* 2016;42(8):1308–1314. doi:10.3109/03639045.2015.1132226
50. Khémiri I, Bitri L. Effectiveness of *Opuntia ficus indica* L. inermis seed oil in the protection and the healing of experimentally induced gastric mucosa ulcer. *Oxid Med Cell Longev.* 2019;2019:1–17. doi:10.1155/2019/1568720
51. Tovey FI. Role of dietary phospholipids and phytosterols in protection against peptic ulceration as shown by experiments on rats. *World J Gastroenterol.* 2015;21(5):1377–1384. doi:10.3748/wjg.v21.i5.1377
52. Sindi AM, Hosny KM. Preparation and evaluation of protective effect of pumpkin seed oil based self nanoemulsifying oral delivery system against ibuprofen-induced peptic ulcer. *J Drug Deliv Sci Technol.* 2019;52:415–420. doi:10.1016/j.jddst.2019.05.009
53. Tovey FI, Bardhan KD, Hobsley M. Dietary phospholipids and sterols protective against peptic ulceration. *Phytother Res.* 2013;27(9):1265–1269. doi:10.1002/ptr.4865