

Higher Plasma APOC-III Was Associated with a Slower Reduction of β -Amyloid Levels in Cerebrospinal Fluid Among Older Individuals Without Dementia

This article was published in the following Dove Press journal:
Neuropsychiatric Disease and Treatment

Xiaoyan Zhang

On behalf of the
Alzheimer's Disease
Neuroimaging Initiative

Department of Child Healthcare, The
Second Affiliated Hospital and Yuying
Children's Hospital of Wenzhou Medical
University, Wenzhou, People's Republic
of China

Purpose: Although emerging evidence has suggested that apolipoprotein C-III (APOC-III) is involved in the pathogenesis of Alzheimer's disease (AD), the association of APOC-III with longitudinal changes in cerebrospinal fluid (CSF) AD pathologies (β -amyloid (A β 42) and tau proteins) is not clear. In the present study, we aimed to examine whether plasma APOC-III levels are associated with longitudinal changes in CSF A β 42, total-tau (t-tau), and phosphorylated-tau (p-tau) levels among older individuals without dementia.

Patients and Methods: Linear mixed models were fitted with plasma APOC-III used as a predictor for longitudinal changes in CSF AD biomarkers over a 7-year period. Data were obtained from the Alzheimer's Disease Neuroimaging Initiative database, and 195 older individuals without dementia (47 subjects with normal cognition (NC) and 148 subjects with mild cognitive impairment (MCI)) with baseline plasma APOC-III measurements were included.

Results: Among older individuals without dementia, we found that the tertiles of plasma APOC-III were associated with changes in CSF A β 42, but not t-tau or p-tau. Specifically, the CSF A β 42 reduction for individuals in the highest plasma APOC-III tertile was significantly slower compared with those in the middle tertile, whereas no other pairwise difference was found to be statistically significant.

Conclusion: Among older individuals without dementia, higher plasma APOC-III levels were associated with slower declines in CSF A β 42.

Keywords: Alzheimer's disease, apolipoprotein C-III, beta-amyloid, tau proteins, longitudinal study

Introduction

Emerging evidence has suggested that apolipoprotein C-III (APOC-III) is involved in the pathogenesis of Alzheimer's disease (AD).¹⁻⁹ For example, plasma APOC-III levels were found to be significantly reduced in patients with AD.⁶ In cross-sectional studies, higher APOC-III levels in plasma were correlated with better cognitive performance.^{3,6} Additionally, a recent longitudinal study found that higher APOC-III levels in CSF were associated with a slower cognitive decline in individuals with mild cognitive impairment (MCI),¹ further supporting that APOC-III may be neuroprotective. A previous study suggested that this beneficial effect of APOC-III on cognition may be because APOC-III is an A β -binding

Correspondence: Xiaoyan Zhang
The Second Affiliated Hospital and Yuying
Children's Hospital of Wenzhou Medical
University, 109 Xueyuan West Road,
Wenzhou 325027, Zhejiang Province,
People's Republic of China
Email 18815001229@163.com

protein that can promote A β efflux and indirectly reduce the accumulation of A β in brain.⁶ However, to our knowledge, no studies have attempted to examine the longitudinal associations of APOC-III levels in plasma with changes in CSF AD pathologies (including A β 42, total tau (t-tau) and phosphorylated tau (p-tau)) in older individuals without dementia.

In the cross-sectional analysis, baseline plasma APOC-III levels were analyzed in relation to baseline CSF A β 42, t-tau and p-tau levels in older individuals without dementia. Further, the longitudinal analysis was conducted with baseline plasma APOC-III used as a predictor for changes in CSF A β 42, t-tau and p-tau levels over a 7-year period.

Patients and Methods

Alzheimer's Disease Neuroimaging Initiative (ADNI)

This longitudinal study used data extracted from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu; 3 October 2018). The ADNI study was initiated in 2003 with the primary goal of examining whether a variety of markers, such as neuropsychological assessments, serial MRI, PET, and other fluid biomarkers, could be combined to predict the progression of MCI and early AD. Each participant of the ADNI study provided written informed consent, and each ADNI site obtained local institutional review board approval.

Participants

Patients inclusion criteria have been previously described elsewhere,¹⁰ and can be found at the ADNI website (adni.loni.usc.edu). In brief, participants with NC had a Mini-Mental State Examination (MMSE)¹¹ score ≥ 24 , and a Clinical Dementia Rating (CDR)¹² score of 0. Participants with MCI had an MMSE ≥ 24 , a CDR of 0.5, objective memory deficits as measured by delayed recall scores of the Wechsler Memory Scale Logical memory II, preserved activities of daily living, and an absence of dementia.

In this longitudinal study, we included subjects who met criteria for NC and MCI and had baseline plasma APOC-III samples and follow-up quantifications of CSF A β 42, t-tau, and p-tau levels. At baseline, there was a total of 195 non-demented older individuals, including 47 participants with NC and 148 participants with MC. As shown in Table 1, annual levels of CSF A β 42, t-tau, and p-tau were examined for up to 7 years.

Determination of APOC-III in Plasma

Plasma APOC-III levels were measured using xMAP multiplex panel (MyriadRBM),⁷ details of which can be found at the ADNI website (<http://adni.loni.ucla.edu>). The file "Biomarkers Consortium ADNI plasma QC Multiplex data" was downloaded from the ADNI website (3 October 2018). To better approximate a normal distribution, the APOC-III analyte was natural log transformed. Study subjects were divided into tertile groups according to APOC-III levels. Plasma levels of APOC-III were [1.79, 2.09], [2.09, 2.22], and [2.22, 2.61] $\mu\text{g/mL}$ in the lowest, middle and highest tertiles, respectively.

Determination of CSF A β 42, t-tau, and p-tau

CSF levels of A β 42, t-tau, and p-tau were determined with the multiplex xMAP Luminex platform,¹³ details of which can be found at the ADNI website (adni.loni.usc.edu). The file "UPENNBBIOMK_MASTER" was extracted from the ADNI dataset in October 2018. In the present study, yearly CSF A β 42, t-tau, and p-tau levels were examined for up to 7 years (Table 1).

Table 1 Demographic and Clinical Data by Tertiles of Plasma APOC-III Levels

Clinical Variables	Low (n = 65)	Middle (n = 65)	High (n = 65)	P value
Age, y	75.2 \pm 6.63	75 \pm 6.65	74.7 \pm 7.49	0.9
Education, y	16.3 \pm 2.92	16.1 \pm 2.82	15.3 \pm 2.96	0.1
Female, n (%)	18 (27.7)	23 (35.4)	28 (43.1)	0.186
APOE4, n (%)	26 (40)	26 (40)	31 (47.7)	0.59
MMSE scores	27.4 \pm 1.86	27.5 \pm 1.55	27.3 \pm 2.15	0.97
CSF A β 42, pg/mL	188 \pm 57.2	177 \pm 57.2	181 \pm 62.2	0.6
CSF t-tau, pg/mL	84.6 \pm 42.6	103 \pm 57.2	97.5 \pm 48.1	0.09
CSF p-tau, pg/mL	31.1 \pm 13.8	33.6 \pm 16.7	32.9 \pm 16	0.8
Plasma APOC-III, $\mu\text{g/mL}$	2 (0.08)	2.15 (0.04) ^a	2.32 (0.08) ^{b,c}	<0.001
Follow-Up Visits, n Subjects				
Baseline	65	65	65	
1 y	65	65	65	
2 y	19	24	19	
3 y	16	19	15	
4 y	12	11	18	
5 y	6	7	8	
6 y	2	3	5	
7 y	0	2	0	

Notes: Comparison between tertile 1 and tertile 2 is marked behind "tertile 2", ^ap < 0.05. Comparison between tertile 1 and tertile 3 is marked behind "tertile 3", ^bp < 0.05. Comparison between tertile 2 and tertile 3 is marked behind "tertile 3", ^cp < 0.05.

Abbreviations: MMSE, Mini-Mental State Examination; APOC-III, apolipoprotein C-III; A β 42, β -amyloid 42; t-tau, total tau; p-tau, phosphorylated tau

Statistical Analyses

First, we applied Kruskal–Wallis tests and χ^2 tests to assess differences in demographic and clinical variables across the tertiles of APOC-III levels. Second, to examine the cross-sectional relationships between APOC-III and CSF AD biomarkers, Kruskal–Wallis tests were conducted. Third, to evaluate the association of baseline plasma APOC-III with changes in CSF AD biomarkers over time, linear mixed models were fitted for each CSF AD biomarker (CSF A β 42, t-tau, and p-tau). All these models were adjusted for age, sex, education and APOE4 genotype. Further, each model included a random intercept for each subject. All statistical analyses were conducted using R software (V.3.6.0).

Results

Demographic and Clinical Variables

The demographic and clinical variables of the study participants by tertiles of plasma APOC-III are demonstrated in Table 1. We did not find significant differences in age or educational attainment across tertiles. Characteristics (sex, percentage of subjects with the APOE4 genotype or MMSE¹¹ scores) did not differ by APOC-III tertiles (Table 1). In addition, we did not find a significant difference in plasma APOC-III levels between males and

females ($t = -1.68$, $p = 0.095$). The numbers of subjects present at each follow-up visit are also listed in Table 1.

Associations of Plasma APOC-III Levels with CSF AD Pathologies

To examine the cross-sectional associations of tertiles of plasma APOC-III with CSF AD pathologies, Kruskal–Wallis tests were performed. However, no significant association of tertiles of plasma APOC-III with CSF AD pathologies was observed (all $p > 0.05$; Table 1, Figure 1).

Longitudinal Change Models

First, to evaluate the association of baseline plasma APOC-III levels (categorized as tertiles) with changes in CSF AD biomarkers over time, linear mixed models were fitted. We found that tertiles of plasma APOC-III were associated with changes in CSF A β 42, but not t-tau or p-tau (Table 2 and Figure 2). In addition, a plot of individual CSF A β 42 concentrations over time across the tertiles has also been displayed (Figure S1). Specifically, as shown in Table 3 and Figure 2, the CSF A β 42 reduction for individuals in the highest APOC-III tertile was significantly slower compared with those in the middle (the middle tertile of APOC-III – the highest tertile of APOC-




Figure 1 Associations of tertiles of plasma APOC-III with CSF AD biomarkers among older individuals without dementia. No significant associations of tertiles of plasma APOC-III with CSF AD pathologies were observed (all $p > 0.05$).

Abbreviations: APOC-III, apolipoprotein C-III; A β 42, β -amyloid 42; t-tau, total tau; p-tau, phosphorylated tau.

Table 2 Summary of Linear Mixed Models Examining the Associations of Plasma APOC-III with Changes in CSF AD Pathologies

Dependent Variable: CSF A β 42			
Predictors	Estimate	SE	P value
Middle APOC-III \times time	-0.2453	0.9750	0.8013
High APOC-III \times time	1.9053	0.9568	0.0464
Dependent Variable: CSF t-tau			
Predictors	Estimate	SE	P value
Middle APOC-III \times time	0.6904	1.2410	0.5780
High APOC-III \times time	-0.5342	1.2334	0.6649
Dependent Variable: CSF p-tau			
Predictors	Estimate	SE	P value
Middle APOC-III \times time	0.1225	0.8038	0.8788
High APOC-III \times time	0.4625	0.7889	0.5577

Note: The bold value indicates a statistically significant difference with a p-value < 0.05. Estimates are unstandardized values, representing the amount of change in each CSF AD biomarker per year. All linear mixed models were adjusted for age, education, sex and APOE4 genotype.

Abbreviation: APOC-III, apolipoprotein C-III; A β 42, β -amyloid 42; t-tau, total tau; p-tau, phosphorylated tau.

III: estimate = -2.151, SE = 0.909, p = 0.0484), whereas no other pairwise difference was found to be statistically significant (all p > 0.05, Table 3) after correcting for multiple comparisons using Tukey method.

In addition, we examined the association of plasma APOC-III with changes in CSF biomarkers from baseline to 1-year follow-up due to the fact that this approach may

seem to have the least “survival” bias (Please see supplementary information: Table S1 and Figure S2).

Discussion

In the present study, we investigated the association of APOC-III levels in plasma with changes in CSF AD biomarkers over time among older individuals without dementia. We found that the tertiles of plasma APOC-III were associated with changes in CSF A β 42, but not t-tau or p-tau. Specifically, the CSF A β 42 reduction for individuals in the highest plasma APOC-III tertile was significantly slower compared with those in the middle tertile, whereas no other pairwise difference was found to be statistically significant.

In terms of the biology of APOC-III, it is primarily expressed in the liver and intestine and has an important role in lipid metabolism.¹⁴ An increasing amount of evidence has suggested that APOC-III might also be involved in the pathogenesis of AD. For instance, the APOC-III 3017G allele was found to be associated with decreased risk of AD among individuals without the presence of the APOE4 genotype.⁹ In cross-sectional studies, levels of APOC-III in plasma were reduced in patients with AD⁶ and were positively correlated with cognitive performance.^{3,6} Additionally, one recent prospective and longitudinal study showed that higher CSF APOC-III levels were associated with a slower cognitive decline in individuals with MCI.¹ However, the mechanisms underlying this potentially neuroprotective effect of APOC-III on cognition




Figure 2 Associations of baseline plasma APOC-III levels with changes in CSF AD pathologies. The tertiles of plasma APOC-III were associated with changes in CSF A β 42 (A), but not t-tau (B) or p-tau (C). Specifically, the CSF A β 42 reduction for individuals in the highest APOC-III tertile was significantly slower compared with those in the middle (the middle tertile of APOC-III – the highest tertile of APOC-III: estimate = -2.151, SE = 0.909, p = 0.0484), whereas no other pairwise difference was found to be statistically significant (all p > 0.05).

Abbreviations: APOC-III, apolipoprotein C-III; A β 42, β -amyloid 42; t-tau, total tau; p-tau, phosphorylated tau.

Table 3 Multiple Comparisons Across the Plasma APOC-III Groups

Contrast	CSF A β 42 Levels		
	Estimate	SE	P value
Low APOC-III vs Middle APOC-III	0.245	0.979	0.9660
Low APOC-III vs High APOC-III	-1.905	0.961	0.1177
Middle APOC-III vs High APOC-III	-2.151	0.909	0.0484

Note: The bold value indicates a statistically significant difference with a p-value <0.05. Tukey method was used for multiple comparison correction. Estimates are unstandardized values, representing the amount of change in CSF A β 42 per year.

Abbreviation: APOC-III, apolipoprotein C-III; A β 42, β -amyloid 42.

remain elusive. One approach for examining the mechanism by which APOC-III affects cognitive decline is to investigate the association of APOC-III with longitudinal reductions in CSF A β 42. In the present study, we provided in vivo evidence that higher CSF APOC-III levels may decelerate reduction in CSF A β 42 in non-demented older adults, supporting the notion that APOC-III may be a potential agent that can slow disease progression in the early stage of AD. However, the precise mechanism is not clear. Levels of A β 42 in CSF demonstrated a substantial reduction in patients with AD as reported in previous studies,^{15–19} probably resulted from aggregation and accumulation of A β 42 in brain or a failure of A β 42 clearance which contributes to decreased amount of A β 42 proteins transport to CSF.²⁰ APOC-III, an amyloid-binding protein, may promote A β efflux and indirectly reduce the deposition of A β in the brain parenchyma.⁶ However, further preclinical studies are needed to clarify the precise mechanisms underlying the relationship between APOC-III and amyloid pathology. Taken together, our data supported the notion that APOC-III may be a major pathogenic factor of AD after A β and tau.

Several limitations of this longitudinal study should be noted. First, the findings of this study should be interpreted with caution because of a potential selection bias. Further studies, especially population-based studies, should be conducted to replicate our results. In addition, the observational nature of our study limits our ability to differentiate whether increased APOC-III leads to, results from, or is just correlated with changes in amyloid pathology.

In conclusion, among older individuals without dementia, higher plasma APOC-III levels were associated with slower declines in CSF A β 42. It should be explored in further studies if modification of APOC-III level slows disease progression.

Acknowledgments

Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of Southern California.

Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Disclosure

The author reports no conflict of interest in this work.

References

1. Wang Q, Zhou W, Zhang J. Higher Apolipoprotein C-III levels in cerebrospinal fluid are Associated with slower cognitive decline in mild cognitive impairment. *J Alzheimers Dis.* 2019;67(3):961–969. doi:10.3233/JAD-181096

2. Koch M, DeKosky ST, Fitzpatrick AL, et al. Apolipoproteins and Alzheimer's pathophysiology. *Alzheimer Dement*. 2018.
3. Muenchhoff J, Song F, Poljak A, et al. Plasma apolipoproteins and physical and cognitive health in very old individuals. *Neurobiol Aging*. 2017;55:49–60. doi:10.1016/j.neurobiolaging.2017.02.017
4. Das M, Gursky O. Amyloid-forming properties of human apolipoproteins: sequence analyses and structural insights. *Adv Exp Med Biol*. 2015;855(undefined):175–211.
5. Lin Q, Cao Y, Gao J. Decreased expression of the APOA1-APOC3-APOA4 gene cluster is associated with risk of Alzheimer's disease. *Drug Des Devel Ther*. 2015;9:5421–5431. doi:10.2147/DDDT.S89279
6. Shih YH, Tsai KJ, Lee CW, et al. Apolipoprotein C-III is an amyloid-beta-binding protein and an early marker for Alzheimer's disease. *J Alzheimers Dis*. 2014;41(3):855–865. doi:10.3233/JAD-140111
7. Mattsson N, Insel P, Nosheny R, et al. Effects of cerebrospinal fluid proteins on brain atrophy rates in cognitively healthy older adults. *Neurobiol Aging*. 2014;35(3):614–622. doi:10.1016/j.neurobiolaging.2013.08.027
8. Song F, Poljak A, Crawford J, et al. Plasma apolipoprotein levels are associated with cognitive status and decline in a community cohort of older individuals. *PLoS One*. 2012;7(6):e34078. doi:10.1371/journal.pone.0034078
9. Sun Y, Shi J, Zhang S, et al. The APOC3 SstI polymorphism is weakly associated with sporadic Alzheimer's disease in a Chinese population. *Neurosci Lett*. 2005;380(3):219–222. doi:10.1016/j.neulet.2005.01.038
10. Aisen PS, Petersen RC, Donohue MC, et al. Clinical core of the Alzheimer's disease neuroimaging initiative: progress and plans. *Alzheimer Dement*. 2010;6(3):239–246. doi:10.1016/j.jalz.2010.03.006
11. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975;12(3):189–198. doi:10.1016/0022-3956(75)90026-6
12. Morris JC. The Clinical Dementia Rating (CDR): current version and scoring rules. *Neurology*. 1993;43(11):2412–2414. doi:10.1212/WNL.43.11.2412-a
13. Shaw LM, Vanderstichele H, Knapiak-Czajka M, et al. Cerebrospinal fluid biomarker signature in Alzheimer's disease neuroimaging initiative subjects. *Ann Neurol*. 2009;65(4):403–413. doi:10.1002/ana.21610
14. Norata GD, Tsimikas S, Pirillo A, Catapano AL. Apolipoprotein C-III: from pathophysiology to pharmacology. *Trends Pharmacol Sci*. 2015;36(10):675–687. doi:10.1016/j.tips.2015.07.001
15. Motter R, Vigo-Pelfrey C, Kholodenko D, et al. Reduction of beta-amyloid peptide42 in the cerebrospinal fluid of patients with Alzheimer's disease. *Ann Neurol*. 1995;38(4):643–648. doi:10.1002/ana.410380413
16. Leinenbach A, Pannee J, Dulffer T, et al. Mass spectrometry-based candidate reference measurement procedure for quantification of amyloid-beta in cerebrospinal fluid. *Clin Chem*. 2014;60(7):987–994. doi:10.1373/clinchem.2013.220392
17. Olsson A, Vanderstichele H, Andreassen N, et al. Simultaneous measurement of beta-amyloid(1-42), total tau, and phosphorylated tau (Thr181) in cerebrospinal fluid by the xMAP technology. *Clin Chem*. 2005;51(2):336–345. doi:10.1373/clinchem.2004.039347
18. Wiltfang J, Esselmann H, Bibl M, et al. Highly conserved and disease-specific patterns of carboxyterminally truncated Abeta peptides 1-37/38/39 in addition to 1-40/42 in Alzheimer's disease and in patients with chronic neuroinflammation. *J Neurochem*. 2002;81(3):481–496. doi:10.1046/j.1471-4159.2002.00818.x
19. Andreassen N, Hesse C, Davidsson P, et al. Cerebrospinal fluid beta-amyloid(1-42) in Alzheimer disease: differences between early- and late-onset Alzheimer disease and stability during the course of disease. *Arch Neurol*. 1999;56(6):673–680. doi:10.1001/archneur.56.6.673
20. Blennow K, Mattsson N, Scholl M, Hansson O, Zetterberg H. Amyloid biomarkers in Alzheimer's disease. *Trends Pharmacol Sci*. 2015;36(5):297–309. doi:10.1016/j.tips.2015.03.002

Neuropsychiatric Disease and Treatment

Dovepress

Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS, and

is the official journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>