

The Combination of circEPSTII and MIF Offers Diagnostic Value for Endometrial Cancer

Zhili Cui¹, Liyuan Zhou¹, Xin An², Wenli Liu¹, Jingxia Li¹, Yueping Zhang³, Wei Zhang⁴

¹Department of Gynecology, Affiliated Hospital of Hebei University of Engineering, Handan, Hebei, 056002, People's Republic of China; ²Department of Pathology, Handan First Hospital, Handan, Hebei, 056000, People's Republic of China; ³Shexian Maternal and Child Health Hospital, Shexian, Hebei, 056004, People's Republic of China; ⁴Department of Gynecology, Handan Traditional Chinese Medicine Hospital, Handan, Hebei, 056001, People's Republic of China

Correspondence: Zhili Cui, Department of Gynecology, Affiliated Hospital of Hebei University of Engineering, No. 81 Congtai Road, Congtai District, Handan, Hebei, 056002, People's Republic of China, Tel + 86-3103962266, Email cuizhili72@yeah.net

Background: Circular RNAs (circRNAs) exhibit unique patterns of expression and high levels of stability in patient plasma samples such that they represent ideal non-invasive biomarkers that can be leveraged to detect a wide array of diseases including endometrial cancer (EC). This study was designed to identify circRNAs with potential diagnostic utility in serum samples from EC patients while also evaluating the utility of macrophage migration inhibitory factor (MIF) as a biomarker when screening for this form of cancer in the clinic.

Methods: Levels of circEPSTII and MIF were assessed in the plasma of EC patients and healthy subjects (n=186 each) through qPCR and ELISAs. The diagnostic utility of these biomarkers was assessed with receiver operating characteristic curve (ROC) analyses.

Results: Relative to healthy subjects, EC patient serum contained significantly elevated circEPSTII and MIF. An association was noted between circEPSTII expression in stages, histologic grade, and residual tumor. ROC curves confirmed that serum circEPSTII levels distinguished between controls and patients with EC with an Area of 0.835 and serum MIF levels distinguished between controls and patients with EC with an Area of 0.6646. When instead diagnosing patients based on the combination of MIF and circEPSTII, the Area further rose to 0.8604.

Conclusion: Assessing the combination of circEPSTII and MIF may be a viable approach to reliably diagnosing EC.

Keywords: circEPSTII, MIF, diagnostic biomarker, endometrial cancer

Introduction

Endometrial cancer (EC) is the third most prevalent malignancy among females worldwide,¹ yet no reliable approaches to detecting EC in its early stage currently exist for women at average risk who are free of any symptoms.² Under most established guidelines, initial detection strategies most commonly include endometrial biopsy or transvaginal ultrasonography, but both of these approaches are subject to limitations. Diagnostic curettage is also commonly employed when seeking to definitively identify this tumor type, but this procedure is inherently invasive and requires the scraping of tissue specimens without clear visibility such that missed diagnoses may occur, while also exposing patients to the potential for endometrial bleeding of varying severity levels. EC tumors can also invade the muscle layer, contributing to metastatic progression and adverse patient outcomes.³ This procedure is also highly specialized such that it can only be conducted by experienced gynecologists. As such, validated biomarkers that can more reliably aid in the diagnosis of EC are urgently needed in order to improve the prognosis of this form of cancer.

Circular RNAs (circRNAs) are a unique subset of transcripts that are extremely stable and resistant to degradation owing to their structural characteristics, defined by a closed covalent loop of RNA.⁴⁻⁶ Particular circRNAs have increasingly been shown to regulate a wide array of cancers and other diseases through their ability to modulate biological processes such as differentiation, glycolytic metabolism, survival, proliferation, and migration.^{7,8} These findings have spurred growing interest

in the application of circRNAs as therapeutic targets or diagnostic biomarkers in a range of pathological contexts.^{9,10} Widespread circRNA dysregulation has recently been reported in EC, with certain circRNAs serving as important regulators of disease-related processes.^{11,12} Of note, circEPSTI1 can reportedly enhance certain oncogenic activities in cervical cancer,¹³ oral squamous cell carcinoma,¹⁴ ovarian cancer,¹⁵ non-small cell lung cancer,¹⁶ and breast cancer.¹⁷ How circEPSTI1 functions in EC, however, remains poorly understood. One recent study also determined that macrophage migration inhibitory factor (MIF) levels are closely associated with EC patient prognostic outcomes.¹⁸ While the bulk of prior MIF research has centered on inflammatory diseases,¹⁹ an increasing number of studies suggest that it may function as a key factor that links inflammation to oncogenesis owing to the importance of inflammatory activity and tissue remodeling in the context of cancer development.²⁰

Here, circEPSTI1 and MIF levels were compared between samples of serum obtained from patients with EC and healthy subjects, while also assessing these expression levels across a range of tumor stages. To more fully assess the link between these biomarkers and the biology of EC, the associations between circEPSTI1 and MIF levels and the clinicopathological characteristics of EC patients were additionally examined. Receiver operating characteristic (ROC) curves were also constructed to test their diagnostic performance. Overall, these findings provided strong evidence for the importance of circEPSTI1 and MIF in the pathogenesis of EC, while also suggesting that they can be leveraged to help diagnose this form of cancer.

Materials and Methods

Sample Collection

This study received approval from the Ethics Committee of the Affiliated Hospital of Hebei University of Engineering, and was performed in accordance with the principles of the Helsinki Declaration. Serum samples were obtained from patients with EC and healthy individuals (n=186 each) between May 2021 and December 2022. The informed consent were gained from the patients and their families. Blood samples were collected in RNase-free tubes for RNA isolation, which was performed with 30 minutes after collection. Patient clinicopathological and molecular characteristics were additionally recorded.

qPCR

RNA was isolated from 500 μ L of serum per patient with an RNA Isolation Kit (Vazyme Biotech), after which cDNA was synthesized with a Prime Script RT reagent Kit (Takara). Then, qPCR reactions were conducted with SYBR Green (Takara) by 7500 Fast Real-Time PCR System, utilizing the GAPDH for the normalization of expression with the $2^{-\Delta\Delta Ct}$ method. The primers used in this study were as follows: circEPSTI1, Forward (F), AAGCTGAAGAAGCTGAACTC, Reverse (R), CCGGTTTATATTTGGTGCTATC. GAPDH, F, GGAGCGAGATCCCTCCAAAAT, R, GGCTGTTGTC ATACTTCTCATGG. The qPCR conditions used in this study were as follows: Initial 1 step at 95°C for 30 sec, followed by 40 cycles at 95°C for 5 sec and at 60°C for 34 sec.

ELISAs

MIF levels in appropriate samples were analyzed with an ELISA kit (R&D Systems) based on provided directions, with all samples being analyzed in triplicate.

Statistical Analysis

GraphPad Prism 8.0 (GraphPad Software) was used to analyze all data, which were compared with Student's t-tests or one-way ANOVAs. Survival analyses were conducted as per the Kaplan–Meier method. $P < 0.05$ served as the cut-off when defining statistical significance.

Results

Participant Characteristics

For this study, plasma samples from 186 EC patients and an equal number of normal control individuals were analyzed. The characteristics of these patients are summarized in [Table 1](#), and no differences in age were detected between these two groups of study subjects ($P > 0.05$).

Table I Clinicopathological Characteristics of EC Patients and Controls

Characteristics	EC (%)	Healthy Controls (%)
Age		
<55	39 (20.97)	60 (32.26)
≥55	147 (79.03)	126 (67.74)
Histological type		
Endometrioid	154 (82.80)	
Mixed	10 (5.38)	
Serous	22 (11.82)	
Stage		
I	120 (64.52)	
II	19 (10.22)	
III	32 (17.20)	
IV	15 (8.06)	
Diabetes		
No	102 (54.84)	
Yes	46 (24.73)	
NA	38 (20.43)	
Hypertension		
No	65 (34.95)	
Yes	98 (52.69)	
NA	23 (12.37)	
Histologic grade		
G1	35 (18.82)	
G2	56 (30.11)	
G3	88 (47.31)	
High grade	7 (3.76)	
Menopause status		
Indeterminate	11 (5.91)	
Peri-Menopause	13 (6.99)	
Post-menopause	129 (69.35)	
Pre-menopause	18 (9.68)	
NA	15 (8.07)	
Residual tumor		

(Continued)

Table 1 (Continued).

Characteristics	EC (%)	Healthy Controls (%)
R0	114 (61.29)	
R1	13 (6.99)	
R2	9 (4.83)	
Rx	23 (12.37)	
NA	27 (14.52)	

CircEPSTII Levels in Serum Samples are Correlated with the Clinicopathological Characteristics of Patients with EC

To confirm the identity of circEPSTII, the genomic loci of circEPSTII and the Sanger sequence of junction site of circEPSTII were detected by Sanger sequencing (Figure 1A). EC patient serum samples were found to exhibit higher levels of circEPSTII when analyzed via qPCR (Figure 1B). The EC patients were allocated to two groups according to the median circEPSTII level (n=93/group). Under these conditions, high circEPSTII expression in the serum was significantly related to the presence of stages (p=0.031), histologic grade (p=0.029), and residual tumor (p=0.022), as shown in Table 2. No significant relationships were seen with age (p=0.652), histological type (p=0.352), diabetes (p=0.186), hypertension (p=0.854), or menopause status (p=0.309), but patients expressing high circEPSTII did exhibit a reduction in both progression-free survival (PFS, Figure 2A) and overall survival (OS, Figure 2B) when compared to those patients exhibiting low expression levels. Consistently, univariate analyses detected a close relationship between the levels of this circRNA, stage, histologic grade and residual tumor, as shown in Tables 3 and 4. These associations remained significant when performing multivariate analyses.

Serum MIF Levels Over Values for the Diagnosis of EC

There is prior evidence for an association between MIF concentrations and the prognosis of individuals with EC.¹⁸ Here, the mean respective concentrations of MIF in the serum of EC patients and healthy subjects were 4.825




Figure 1 Patients with EC exhibit elevated serum levels of circEPSTII. **(A)** The genomic loci of circEPSTII and the Sanger sequence of junction site of circEPSTII. **(B)** Serum samples from EC patients and healthy subjects were assessed to compare serum circEPSTII expression, which was found to be elevated in individuals diagnosed with EC. ***p<0.001.

Table 2 Relationships Between circEPSTII Expression Levels and Patient Clinicopathological Characteristics

Clinical Characteristics	Number	circEPSTII Expression		P value
		Low	High	
Age				0.652
<55	39	18	21	
≥55	147	75	72	
Histological type				0.352
Endometrioid	154	77	77	
Mixed	10	6	4	
Serous	22	10	12	
Stage				0.031
I	120	72	48	
II	19	8	11	
III	32	9	23	
IV	15	4	11	
Diabetes				0.186
No	102	59	43	
Yes	46	19	27	
Hypertension				0.854
No	65	32	33	
Yes	98	42	56	
Histologic grade				0.029
G1	35	25	10	
G2	56	36	20	
G3	88	31	57	
High grade	7	1	6	
Menopause status				0.309
Indeterminate	11	6	5	
Peri-Menopause	13	6	7	
Post-menopause	129	63	66	
Pre-menopause	18	10	8	
Residual tumor				0.022
R0	114	70	44	
R1	13	8	5	
R2	9	3	6	
Rx	23	7	16	




Figure 2 Serum levels of circEPSTII are related to the clinicopathological characteristics of patients with EC. Log rank tests were utilized to assess the PFS (A) and OS (B) of patients exhibiting low or high circEPSTII expression.

± 1.05 ng/mL and 5.871 ± 3.37 ng/mL ($P < 0.0001$) (Figure 3A). As revealed by constructed ROC curves, levels of serum MIF were capable of reliably distinguishing between patients with EC and healthy individuals, with an Area of 0.6646 (95% CI; 0.6086–0.7205) (Figure 3B).

Table 3 Univariate and Multivariate Cox Analyses of PFS-Related Factors in EC Patients

Variables	Univariate Analysis		p value	Multivariate Analysis		p value
	HR	95% CI		HR	95% CI	
Age (Years)	0.521	0.251–0.985	0.351	–	–	–
Histological type	0.481	0.218–0.854	0.311	–	–	–
Stage	1.625	0.529–2.519	0.034	1.458	0.352–2.221	0.035
Diabetes	0.428	0.365–1.258	0.481	–	–	–
Hypertension	0.558	0.118–0.849	0.352	–	–	–
Histologic grade	2.015	0.511–2.719	0.025	1.982	0.658–2.419	0.028
Menopause status	0.522	0.185–0.971	0.246	–	–	–
Residual tumor	1.952	1.152–2.254	0.019	1.845	0.954–2.197	0.021
circEPSTII expression	2.512	1.025–3.251	0.011	2.197	0.981–2.971	0.015

Table 4 Univariate and Multivariate Cox Analyses of OS-Related Factors in EC Patients

Variables	Univariate Analysis		p value	Multivariate Analysis		p value
	HR	95% CI		HR	95% CI	
Age (Years)	0.452	0.341–0.881	0.391	–	–	–
Histological type	0.512	0.271–0.919	0.291	–	–	–
Stage	1.522	0.418–2.025	0.037	1.441	0.519–1.957	0.041
Diabetes	0.537	0.415–1.192	0.405	–	–	–
Hypertension	0.395	0.251–0.719	0.334	–	–	–
Histologic grade	1.952	0.397–2.518	0.026	1.852	0.418–2.228	0.03
Menopause status	0.494	0.251–0.819	0.319	–	–	–
Residual tumor	1.825	1.002–2.295	0.025	1.719	1.128–2.341	0.027
circEPSTII expression	2.417	1.102–3.025	0.014	2.027	1.249–2.843	0.018




Figure 3 Evaluation of the expression and diagnostic value of MIF in patients with EC. **(A)** Serum MIF concentrations were elevated in EC patients as compared to healthy subjects. **(B)** Serum MIF concentrations were able to distinguish between EC patients and control subjects in ROC curve analyses. *** $P < 0.001$.




Figure 4 Serum circEPSTII offers diagnostic value for the detection of EC. **(A)** Serum levels of circEPSTII were able to effectively distinguish between EC patients and controls when constructing an ROC curve. **(B)** An ROC curve analysis was conducted for the combination of circEPSTII and MIF to more effectively distinguish between healthy subjects and individuals with EC.

EC Can Be Effectively Diagnosed Based on Serum circEPSTII

To similarly assess the diagnostic performance of the serum levels of circEPSTII, ROC curves were again constructed, effectively allowing for differentiation between samples from individuals with EC and healthy subjects. At the optimal cut-off of 3.9496, the Area for the ROC curve was 0.8351 (95% CI: 0.7941–0.8761) (Figure 4A). CircEPSTII may thus outperform MIF with respect to its sensitivity when leveraged as a biomarker to diagnose EC. These analyses were further expanded by examining the diagnostic performance of a combination of serum circEPSTII and MIF as a means of detecting EC. The combined AUC when using these two biomarkers was 0.8604 (95% CI: 0.8217–0.8991) (Figure 4B). Predictive probability levels above this cut-off were indicative of EC positivity.

Discussion

EC is among the most frequently diagnosed malignancies among women. While many patients are diagnosed while the disease is in its early stages, the overall survival of individuals with recurrent disease or who are only diagnosed when the

disease is relatively advanced is substantially shorter. Analyses of biomarkers in the peripheral blood thus hold great promise as a noninvasive and cost-effective means of aiding in EC diagnosis.

Many studies have emphasized the value of circRNAs in the diagnosis of a variety of cancers as these molecules are both highly stable and abundant, in addition to playing important roles in key cancer-associated processes.^{21,22} The stability of circRNAs in patient serum and plasma remains consistent even with freezing and thawing or shifts in pH,^{23,24} making them optimal minimally invasive targets for clinical assessment. Serum Circ-FAF1/Circ-ELP3 offers great utility as a biomarker when applied in the context of diagnosing breast cancer,²⁵ while hsa_circ_0000437 can promote gastric cancer pathogenesis and metastasis to the lymph nodes.²⁶ Levels of hsa_circ_0000702 in patient serum have been proposed to be an effective gastric cancer-related biomarker.²⁷ Although identified only recently, circEPSTI1 has been noted to support tumorigenesis and progression in cervical cancer,¹³ oral squamous cell carcinoma,¹⁴ ovarian cancer,¹⁵ non-small cell lung cancer,¹⁶ and breast cancer.¹⁷ The functions and diagnostic performance of circEPSTI1 in EC, however, have not been previously documented.

This study is the only one we are aware of examining the relevance of MIF in EC patients. MIF has also been tied to angiogenic activity and the induction of tumor progression and growth in various cancers, although not in EC. In patients with malignant ascites, for example, Hagemann et al noted strong MIF expression, while ovarian cancer cell-derived MIF was capable of stimulating cytokine, chemokine, and pro-angiogenic factor production, contributing to angiogenic activity and vascularization.²⁸ Nishihira et al further highlighted a close association between MIF and both angiogenesis and tumor growth through studies in which they used antisense MIF to treat murine colon cancer cells.²⁹ These results support the use of MIF as a biomarker for diagnosing EC.

Here, serum circEPSTI1 levels in EC patients were evaluated to gauge their diagnostic value. These analyses revealed that higher circulating serum levels of this circRNA were related to EC patient stages, histologic grade, and residual tumor. This suggests that circEPSTI1 may play a role in the development or malignant progression of this form of cancer. The OS and PFS of patients expressing higher circEPSTI1 levels were worse than those of patients in which this circRNA was not highly abundant, further underscoring its potential application in the context of patient management. The ROC curves showed that serum circEPSTI1 levels could reliably differentiate between patients with EC and healthy serum samples at high levels of sensitivity and specificity better than those for MIF alone. When combining serum circEPSTI1 and MIF analyses together, EC could be accurately diagnosed with even greater specificity and sensitivity, indicating that this combined biomarker panel may be a robust strategy for EC diagnosis in the clinic.

Conclusion

In conclusion, EC patients exhibit pronounced circEPSTI1 upregulation that may be closely tied to oncogenic activity such that efforts to measure circulating levels of this circRNA can aid in the more effective and timely diagnosis of this disease. The combination of circEPSTI1 with a range of other clinical biomarkers that are part of standard analyses may further enhance its diagnostic performance. While these results are subject to some limitations, they nonetheless underscore important opportunities for research focused on assessing the clinical utility of circEPSTI1 in combination with MIF when seeking to screen for EC.

Funding

This study is supported by Handan City Science and Technology Research and Development Plan Project (No. 19422083029ZC).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Neacsu A, Marcu ML, Stanica CD, et al. Clinical and morphological correlations in early diagnosis of endometrial cancer. *Rom J Morphol Embryol.* 2018;59(2):527–531.
2. Tarney CM, Wang G, Bateman NW, et al. Biomarker panel for early detection of endometrial cancer in the prostate, lung, colorectal, and ovarian cancer screening trial. *Am J Obstet Gynecol.* 2019;221(5):472 e471–472 e410. doi:10.1016/j.ajog.2019.06.005

3. Gottwald L, Kubiak R, Pasz-Walczak G, et al. The value of progesterone and estrogen receptors expression in tissue microarray method in prognosis of patients with endometrioid endometrial cancer. *Ginekol Pol.* 2013;84(2):95–101. doi:10.17772/gp/1602
4. Panda AC, Grammatikakis I, Munk R, Gorospe M, Abdelmohsen K. Emerging roles and context of circular RNAs. *Wiley Interdiscip Rev RNA.* 2017;8(2). doi:10.1002/wrna.1386
5. Muller S, Appel B. In vitro circularization of RNA. *RNA Biol.* 2017;14(8):1018–1027. doi:10.1080/15476286.2016.1239009
6. Ebbesen KK, Kjems J, Hansen TB. Circular RNAs: identification, biogenesis and function. *Biochim Biophys Acta.* 2016;1859(1):163–168. doi:10.1016/j.bbagr.2015.07.007
7. Chen RX, Liu HL, Yang LL, et al. Circular RNA circRNA_0000285 promotes cervical cancer development by regulating FUS. *Eur Rev Med Pharmacol Sci.* 2019;23(20):8771–8778.
8. Chen Y, Wang J, Wang C, Liu M, Zou Q. Deep learning models for disease-associated circRNA prediction: a review. *Brief Bioinform.* 2022;23(6). doi:10.1093/bib/bbac364
9. Zhang HD, Jiang LH, Sun DW, Hou JC, Ji ZL. CircRNA: a novel type of biomarker for cancer. *Breast Cancer.* 2018;25(1):1–7. doi:10.1007/s12282-017-0793-9
10. Jiang B, Zhang J, Sun X, et al. Circulating exosomal hsa_circRNA_0039480 is highly expressed in gestational diabetes mellitus and may be served as a biomarker for early diagnosis of GDM. *J Transl Med.* 2022;20(1):5. doi:10.1186/s12967-021-03195-5
11. Liu Y, Chen S, Zong ZH, Guan X, Zhao Y. CircRNA WHSC1 targets the miR-646/NPM1 pathway to promote the development of endometrial cancer. *J Cell Mol Med.* 2020;24(12):6898–6907. doi:10.1111/jcmm.15346
12. Zhang J, Chen S, Wei S, et al. CircRAPGEF5 interacts with RBFOX2 to confer ferroptosis resistance by modulating alternative splicing of TFRC in endometrial cancer. *Redox Biol.* 2022;57:102493. doi:10.1016/j.redox.2022.102493
13. Wu P, Li C, Ye DM, et al. Circular RNA circEPSTI1 accelerates cervical cancer progression via miR-375/409-3P/515-5p-SLC7A11 axis. *Aging.* 2021;13(3):4663–4673. doi:10.18632/aging.202518
14. Wang J, Jiang C, Li N, et al. The circEPSTI1/mir-942-5p/LTBP2 axis regulates the progression of OSCC in the background of OSF via EMT and the PI3K/Akt/mTOR pathway. *Cell Death Dis.* 2020;11(8):682. doi:10.1038/s41419-020-02851-w
15. Xie J, Wang S, Li G, et al. circEPSTI1 regulates ovarian cancer progression via decoying miR-942. *J Cell Mol Med.* 2019;23(5):3597–3602. doi:10.1111/jcmm.14260
16. Xie Y, Wang L, Yang D. CircEPSTI1 promotes the progression of non-small cell lung cancer through miR-145/HMGB3 Axis. *Cancer Manag Res.* 2020;12:6827–6836. doi:10.2147/CMAR.S252893
17. Zhang Y, Tan D, Xie Y, et al. CircEPSTI1 Promotes the Proliferation of HER2-Positive Breast Cancer Cells via circEPSTI1/miR-145/ERBB3 Axis. *J Oncol.* 2022;2022:1028851. doi:10.1155/2022/1028851
18. Xiao W, Dong X, Zhao H, et al. Expression of MIF and c-erbB-2 in endometrial cancer. *Mol Med Rep.* 2016;13(5):3828–3834. doi:10.3892/mmr.2016.4992
19. Mizue Y, Ghani S, Leng L, et al. Role for macrophage migration inhibitory factor in asthma. *Proc Natl Acad Sci U S A.* 2005;102(40):14410–14415. doi:10.1073/pnas.0507189102
20. Bucala R, Donnelly SC. Macrophage migration inhibitory factor: a probable link between inflammation and cancer. *Immunity.* 2007;26(3):281–285. doi:10.1016/j.immuni.2007.03.005
21. Ma Y, Li Z, Ma D, Guo J, Sun W. Hsa_circ_0003195 as a biomarker for diagnosis and prognosis of gastric cancer. *Int J Clin Oncol.* 2022;27(2):354–361. doi:10.1007/s10147-021-02073-w
22. Lei B, Tian Z, Fan W, Ni B. Circular RNA: a novel biomarker and therapeutic target for human cancers. *Int J Med Sci.* 2019;16(2):292–301. doi:10.7150/ijms.28047
23. Veá A, Llorente-Cortes V, de Gonzalo-Calvo D. Circular RNAs in Blood. *Adv Exp Med Biol.* 2018;1087:119–130.
24. Guo S, Hu C, Zhai X, Sun D. Circular RNA 0006602 in plasma exosomes: a new potential diagnostic biomarker for hepatocellular carcinoma. *Am J Transl Res.* 2021;13(6):6001–6015.
25. Omid-Shafaat R, Moayeri H, Rahimi K, et al. Serum Circ-FAF1/Circ-ELP3: a novel potential biomarker for breast cancer diagnosis. *J Clin Lab Anal.* 2021;35(11):e24008. doi:10.1002/jcla.24008
26. Shen X, Kong S, Ma S, et al. Hsa_circ_0000437 promotes pathogenesis of gastric cancer and lymph node metastasis. *Oncogene.* 2022;41(42):4724–4735. doi:10.1038/s41388-022-02449-w
27. Yuan W, Fang R, Mao C, Chen H, Tai B, Cong H. Serum circular RNA hsa_circ_0000702 as a novel biomarker for diagnosis of gastric cancer. *J Clin Lab Anal.* 2023;37(3):e24842. doi:10.1002/jcla.24842
28. Hagemann T, Robinson SC, Thompson RG, Charles K, Kulbe H, Balkwill FR. Ovarian cancer cell-derived migration inhibitory factor enhances tumor growth, progression, and angiogenesis. *Mol Cancer Ther.* 2007;6(7):1993–2002. doi:10.1158/1535-7163.MCT-07-0118
29. Nishihira J, Ishibashi T, Fukushima T, Sun B, Sato Y, Todo S. Macrophage migration inhibitory factor (MIF): its potential role in tumor growth and tumor-associated angiogenesis. *Ann N Y Acad Sci.* 2003;995(1):171–182. doi:10.1111/j.1749-6632.2003.tb03220.x

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies across all disease areas. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.