

Lower creatinine as a marker of malnutrition and lower muscle mass in hemodialysis patients

Abdulmecit Yildiz¹
Fatih Tufan²

¹Department of Nephrology, Uludag University School of Medicine, Bursa, ²Department of Geriatrics, Istanbul Faculty of Medicine, Istanbul University, Istanbul, Turkey

Dear editor

We read the recently published and well-designed study of Lee et al¹ that suggests that bioimpedance analysis (BIA) gives relevant information about hydration status and malnutrition in hemodialysis patients. The authors recommend that utilization of BIA routinely in hemodialysis patients would be rational. We would like to make a few comments about their study.

In their study, elderly subjects had significantly lower creatinine levels. Recent studies indicate that lower creatinine levels in patients undergoing hemodialysis are associated with lower muscle mass, malnutrition, and mortality.² We have also observed lower creatinine levels and higher urea/creatinine ratios in elderly hemodialysis patients compared with younger ones and in those with poorer nutritional status compared with those with normal nutritional status.³ Thus, we recommend that the authors report on the association between creatinine, handgrip strength, BIA measurements, and malnutrition inflammation score in their study that may provide clinically relevant and practical information.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Lee JE, Jo IY, Lee SM, et al. Comparison of hydration and nutritional status between young and elderly hemodialysis patients through bioimpedance analysis. *Clinical Interventions in Aging*. 2015;10:1327–1334.
2. Park J, Mehrotra R, Rhee CM, et al. Serum creatinine level, a surrogate of muscle mass, predicts mortality in peritoneal dialysis patients. *Nephrol Dial Transplant*. 2013;28(8):2146–2155.
3. Tufan F, Yildiz A, Dogan I, Yildiz D, Sevinir S. Urea to creatinine ratio: a forgotten marker of poor nutritional state in patients undergoing hemodialysis treatment. *Aging Male*. 2015;18(1):49–53.

Correspondence: Fatih Tufan
Department of Geriatrics, Istanbul
Faculty of Medicine, Istanbul University,
Fatih, Istanbul, PO Box 34093, Turkey
Email fatihtufan@gmail.com

Authors' reply

Jung Eun Lee^{1,2}

In Young Jo³

Song Mi Lee³

Woo Jeong Kim³

Hoon Young Choi^{2,4}

Sung Kyu Ha⁴

Hyung Jong Kim⁵

Hyeong Cheon Park^{2,4}

¹Department of Internal Medicine, Yongin Severance Hospital, Yonsei University College of Medicine, ²Severance Institute for Vascular and Metabolic Research, Yonsei University College of Medicine, ³Department of Nutrition Services, Gangnam Severance Hospital, ⁴Department of Internal Medicine, Gangnam Severance Hospital, Yonsei University College of Medicine, Seoul, ⁵Department of Internal Medicine, CHA Bundang Medical Center, CHA University, Seongnam, Korea

Correspondence: Hyeong Cheon Park
Department of Internal Medicine, Gangnam Severance Hospital,
Yonsei University College of Medicine, 211 Eonju-ro,
Gangnam-gu, Seoul, 135-720, Korea
Tel +82 2 2019 3310
Fax +82 2 3463 3882
Email amp97@yuhs.ac

Hyung Jong Kim
Department of Internal Medicine, Bundang CHA Medical Center,
CHA University, 59 Yatap-ro, Bundang-gu, Seongnam-si,
Gyeonggi-do, 463-712, Korea
Tel +82 31 780 5895
Fax +82 31 780 5981
Email khj04@cha.ac.kr

Dear editor

We would like to thank Drs Yildiz and Tufan for their thoughtful comments on our recent study.¹ Malnutrition is common in elderly hemodialysis (HD) patients and is associated with increased morbidity and mortality.² Therefore, proper assessment of nutritional status and its management is essential for elderly HD patients. A panel of tests that assess

body composition as well as body protein stores are recommended. The serum creatinine (Cr) level before dialysis is a strong predictor of low muscle mass and poor outcome.³ Our study also demonstrated that elderly HD patients have lower serum Cr levels similar to the study by Tufan et al.⁴ Univariate analysis showed serum Cr level was significantly associated with malnutrition-inflammation score ($\beta=-0.289$, $P=0.008$). However, on multivariate analysis, serum Cr level lost its significance ($\beta=-0.015$, $P=0.899$). The serum Cr level has been used as a biochemical nutritional marker along with serum albumin or prealbumin. However, the observed serum Cr is to some extent dependent on the intake of animal proteins and study population differences may also contribute to discrepancies. Measuring serum Cr level is cheap compared to other biochemical nutritional markers and is routinely measured in HD patients. However, serum Cr measurements may not give the full picture of nutritional status and lack significance compared to bioimpedance analysis measurement especially in elderly HD patients.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Lee JE, Jo IY, Lee SM, et al. Comparison of hydration and nutritional status between young and elderly hemodialysis patients through bioimpedance analysis. *Clinical Interventions in Aging*. 2015;10:1327–1334.
2. Wolfson M. Nutrition in elderly dialysis patients. *Semin Dial*. 2002;15(2):113–115.
3. Heimbürger O, Qureshi AR, Blaner WS, Berglund L, Stenvinkel P. Hand-grip muscle strength, lean body mass, and plasma proteins as markers of nutritional status in patients with chronic renal failure close to start of dialysis therapy. *Am J Kidney Dis*. 2000;36(6):1213–1225.
4. Tufan F, Yildiz A, Dogan I, Yildiz D, Sevinir S. Urea to creatinine ratio: a forgotten marker of poor nutritional state in patients undergoing hemodialysis treatment. *Aging Male*. 2015;18(1):49–53.

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Clinical Interventions in Aging 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Clinical Interventions in Aging editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Clinical Interventions in Aging

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine,

CAS, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/clinical-interventions-in-aging-journal>

Dovepress