

# Chronic pain and geriatric syndromes in community-dwelling patients aged $\geq 65$ years

Orly Liberman<sup>1</sup>  
Tamar Freud<sup>2</sup>  
Roni Peleg<sup>2,3</sup>  
Ariela Keren<sup>1</sup>  
Yan Press<sup>2,4,5</sup>

<sup>1</sup>Nursing Department, Recanati School for Community Health Professions, Ben-Gurion University of the Negev, Beer-Sheva, Israel; <sup>2</sup>Department of Family Medicine, Sial Research Center for Family Medicine and Primary Care, Faculty of Health Sciences, Ben-Gurion University of the Negev, Beer-Sheva, Israel; <sup>3</sup>Pain Clinic, Clalit Health Services, Beer-Sheva, Israel; <sup>4</sup>Yasski Clinic, Comprehensive Geriatric Assessment Unit, Clalit Health Services, Beer-Sheva, Israel; <sup>5</sup>Unit for Community Geriatrics, Division of Health in the Community, Ben-Gurion University of the Negev, Beer-Sheva, Israel

**Background:** In growing elderly populations, there is a heavy burden of comorbidity and a high rate of geriatric syndromes (GS) including chronic pain.

**Purpose:** To assess the prevalence of chronic pain among individuals aged  $\geq 65$  years in the Southern District of Israel and to evaluate associations between chronic pain and other GS.

**Methods:** A telephone interview was conducted on a sample of older adults who live in the community. The interview included the Brief Pain Inventory and a questionnaire on common geriatric problems.

**Results:** Of 419 elderly individuals who agreed to be interviewed 232 (55.2%) suffered from chronic pain. Of those who reported chronic pain, 136 participants (68.6%) noted that they had very severe or unbearable pain. There were statistically significant associations between the pain itself and decline in patient's functional status, increased falls, reduced mood, and cognitive decline.

**Conclusion:** The results of this study show that chronic pain is very common in older adults and that it is associated with other GS. There is a need to increase awareness of chronic pain in older adults and to emphasize the important role that it plays in their care.

**Keywords:** pain, impact of pain, older adults, geriatric syndromes, community dwelling

## Introduction

GS<sup>1,2</sup> are clinical conditions in older adults that do not fit into discrete disease categories.<sup>3</sup> Conditions such as delirium, pressure ulcers, falls, and incontinence are “widely known”,<sup>3,4</sup> but other conditions, such as visual and hearing problems, elder abuse, malnutrition, sleep problems, and even dizziness and syncope too have been included under the heading of GS.<sup>5</sup>

Pain is a very common problem among adults. The prevalence of chronic pain in the adult population ranges from 20% to 50%, depending on the study population, the sampling method, the interview method, the definition of “chronicity” and the definition of the site of pain.<sup>6–10</sup>

Many studies have investigated associations between pain and specific GS, such as functional limitations,<sup>6,7,9,11–15</sup> affective disorders,<sup>6,10,14,16–26</sup> cognitive decline,<sup>27–30</sup> and falls.<sup>12</sup> However, a literature review on the association between pain and GS and their mutual effect on outcomes, such as mortality, functional decline, falls, etc., did not yield a clear-cut picture. Lohman et al<sup>31</sup> found that the inclusion of persistent pain as an additional criterion for frailty led to a potentially better prediction of incident adverse outcomes. Andrews et al<sup>32</sup> looked at the cross-sectional relationship between

Correspondence: Yan Press  
Unit for Comprehensive Geriatric Assessment, Clalit Health Services, Yasski Clinic, 24 King David Street, Beer-Sheva, Israel  
Tel +972 8 640 7738  
Fax +972 8 640 7795  
Email yanp@bgu.ac.il

pain and basic functional limitations and their effect on ADL disability after a 10-year follow-up. They found that pain itself is not an independent risk factor. They linked this finding to the possibility that pain and functional limitation, such as other GS, have common underlying risk factors and mechanisms. In one of the important studies that investigated the prevalence and co-occurrence of GS among community-dwelling people aged  $\geq 75$  years, the authors cited the fact that they did not assess the co-occurrence of pain GS as one of the study limitations.<sup>33</sup> It should be noted that many of the studies that looked at the effect of GS on negative outcomes did not take pain into account.<sup>34–36</sup>

In the present study, we assessed associations between chronic pain and GS among individuals aged  $\geq 65$  years. We hypothesized that GS are very common in our study population of older individuals with chronic pain.

## Methods

### Study population

This was a cross-sectional telephone survey of Hebrew-speaking men and women aged  $\geq 65$  years living in the community in Beer-Sheba and its surrounding area who were insured by the Clalit Healthcare Services. The Clalit Healthcare Service is the largest health care service in Israel and insures over two-thirds of the Israeli population.

### Study design and sampling process

We conducted a random sampling from a roster of 1103 patients in clinics in the Jewish sector,  $\geq 65$  years, drawn from the Clalit Healthcare Services systematic databases. As part of their training course, nursing students from the Recanati School for Community Health Professions of the Faculty of Health Sciences of Ben-Gurion University of the Negev have a research seminar on the subject of pain in older adults. They also have three communication courses on how to interview patients through different media including by telephone and then have to pass a simulated test. Before the study interviews were conducted, the 51 students, who took part in the research seminar, underwent training by one of the investigators (TF) on how to conduct the study interview by telephone, how to address the participants, and how to fill in the questionnaire.

Each completed questionnaire was examined by two of the investigators (AK and OL) and those that were not completed correctly were not used in the study.

Interviewers contacted these patients and asked them to consent to participate in the study. Those who met the study criteria and agreed verbally to participate underwent a

comprehensive telephone interview. At the beginning of the interview the patients were asked whether they had suffered from persistent or intermittent pain over the previous three months? Only those patients who answered this question in the affirmative underwent a full interview with the study questionnaire.

Patients who refused to participate, who did not speak Hebrew, who were not able to participate due to hearing or comprehension impairment, who resided in old age settings, who were hospitalized, or who were away from Israel were excluded from the study.

### Study instruments

The patients underwent a structured personal telephone interview. The questionnaire included questions on chronic diseases including chronic pain, sociodemographic data, and the following formal questionnaires:

- To measure pain intensity, we used a five-point verbal descriptor scale, which has been shown to have low error rates and good face convergent and criterion validity.<sup>37</sup> Patients were asked to grade their pain in terms of the following response options: no pain, slight pain, moderate pain, severe pain, and unbearable pain.
- A 10-point verbal numeric scale, which was found to have high correlation with a 10-point visual analog scale,<sup>38</sup> has also been used to measure pain intensity. Patients were asked to grade their average pain during the previous 3 months on a scale from 0 (no pain) to 10 (unbearable pain).
- The BPI<sup>39</sup> has been used to measure pain interference. We extracted questions from this questionnaire related to the site of pain and its effect on the patient's daily activity: overall activity, mood, walking capacity, amount and quality of sleep at night, relationships with others, need for bedrest, and routine work. The questionnaire was translated to and validated in Hebrew from previous research.<sup>40</sup>
- A GS questionnaire including dichotomous questions (yes/no) on BADL ("Do you dress alone?", "Do you wash alone?"), IADL ("Do you shop on your own?", "Do you prepare your own meals?"), questions on falls ("Did you fall over the previous year? If yes, how many times?"), a question on weight loss ("Did you have an unplanned loss of weight of 5 kg or more over the last six months?"), a question on cognitive status ("Do you have memory problems?"), a question on mood ("Have you mostly felt depressed or sad over the last two months?"),

and a question on polypharmacy (“Do you take six or more drugs on a regular basis?”). These questions were taken from a computerized screening questionnaire that is routinely filled in by nurses in the Clalit Healthcare Services clinics while screening patients with geriatric problems.<sup>41</sup>

## Sample size calculation

The Southern District has 52,000 patients aged 65 years or above. Assuming that 25% suffer from chronic pain,<sup>6</sup> the required sample size to assess the rate of patients with chronic pain with an accuracy of 10% would be 75 patients. It also has 27,000 elderly patients aged 75 years or above. Assuming that 50% of these have chronic pain,<sup>10</sup> the required sample size to assess the rate of chronic pain with accuracy of 10% would be 96 patients. To cover both of these conditions with accuracy of 10% and under the assumption that we would need 96 patients 75 years or above and that this group represents about 50% of the patients aged  $\geq 65$  years (27000/52000), we would need 210 patients. To compensate for potential mistakes and missing values, the final sample size was 232 patients.

## Statistical analyses

A descriptive analysis was performed with means and standard deviations for continuous variables and frequencies for categorical variables. The association between severity of pain and its effect on daily living with the presence of various GS was tested by Student’s *t*-test. The association between the number of GS and the severity of pain and its effect on daily living were analyzed by analysis of variance. Logistic regression models were used to find associations among chronic pain and GS. Statistical significance was determined at  $P < 0.05$ .

The study was approved by the Helsinki Committee of the Meir Medical Center (approval #095/2014k).

## Results

Nine hundred and twenty-two patients, of the original random sample of 1103 individuals insured by the Clalit Healthcare Services, were located for an interview. Of these, 223 refused to be interviewed, 255 were not capable of being interviewed (77 due to hearing or other communication problems and the other 178 because they did not speak Hebrew), and 25 died before telephone contact with them. In all, 420 patients could be interviewed and consented to participate. Of these, 188 (44.8%) reported that they did not suffer from chronic pain, so the interview with them was discontinued. Chronic

pain over the previous 3 months at least was reported by 232 patients (55.2%), and they underwent the full interview (Figure 1). The mean age was  $73.7 \pm 6.5$  years and 89 (38.4%) were men. The sociodemographic characteristics of the study participants are shown in Table 1.

## The characteristics of the pain and its effects

One hundred eighty-two (78.4%) reported a single pain site, 40 (17.2%) reported two sites, and 10 (4.4%) reported three or more sites. The three most common pain sites were the back (37.5%), the knee (26.7%), and the limbs (22.8%).

Based on the five-point verbal descriptor scale, 136 of 232 (58.6%) graded their pain as severe or unbearable. The mean pain severity on the 10-point numeric rating scale was  $7.1 \pm 2.1$ . Based on the BPI measure of the effect of pain on overall activity (pain interference), the mean score was  $4.9 \pm 2.4$  on the scale from 0 to 10 (Table 2). Only 41.4% of the patients with chronic pain were treated with analgesics.


## The prevalence of GS

The data on the prevalence of GS is presented in Table 3. One hundred ninety-eight (85.4%) reported any type of GS. The most common problem was dependency at grocery shopping, with 51.7% reporting this as a problem. The least common problem was dependency at dressing as only 15.5% of the participants were not capable of dressing themselves.

Table 4 shows the association between the presence of a GS and the severity of pain and its effect on overall activity. There was no statistically significant association between the severity of pain and falls over the previous year. A statistically significant association was found with the severity of pain and its effect on overall activity for all the other GS.

The severity of pain ranged from a mean of  $6.3 \pm 2.4$  among participants without GS to  $6.7 \pm 2.1$  for those with 1–2 GS and  $7.7 \pm 1.9$  for those with three or more GS ( $P < 0.0001$ ). Similarly, the mean score for the effect of pain on overall activity (pain interference) ranged from  $3.1 \pm 2.1$  for participants without GS to  $6.2 \pm 1.9$  among those with three or more GS ( $P < 0.0001$ ).

To check the effect of pain on the spectrum of GS, we built logistic regression models with the following dependent variables: functional state (any dependency in IADL, any dependency in BADL, and any dependency in ADL), depressive mood, and memory loss (Table 5). The only variable that predicted pain intensity was any dependency in BADL (odds ratio = 1.41, 95% confidence interval: 1.15–1.71,  $P = 0.001$ ).



**Figure 1** Recruitment of study patients

**Note:** <sup>a</sup>Could not be interviewed: do not hear, do not communicate, nursing home, hospitalized, abroad, other problems.

## Discussion

In a telephone survey of patients  $\geq 65$  years living in the community, we found that more than half of the participants reported pain that lasted for at least 3 months. We also found that the severity of chronic pain and its interference with overall activity were of significance in a large number of patients with GS (Table 4). The prevalence of pain in our results is similar to those of Raftery et al<sup>10</sup> who reported that about 50% of patients in the  $\geq 65$  years age group living in the community reported pain for 3 or more months. Our results are higher than those of other studies.<sup>6,8,9,42</sup> These differences stem from lack of uniformity among these studies in terms of age, setting, and definition of chronic pain. Thus, for

example, Pereira et al<sup>9</sup> found a rate of 30% among elderly patients who were hospitalized for rehabilitation, while Johannes et al<sup>8</sup> reported a rate of 38.5% among patients in this age group living in the community. In our sample, 94% of patients with chronic pain reported at least a moderate severity of pain and almost 60% complained of severe or very severe pain that had a negative effect on their daily activity. However, only 41.4% participants with chronic pain took analgesics. Similar results have been reported in the past<sup>28,43</sup> and they may be related to problems in the diagnosis of pain<sup>44,45</sup> on the part of both the patients and the physicians and poor patient–doctor concordance in evaluation of the pain.<sup>12</sup> These findings are of great concern since, according

**Table 1** Descriptive characteristics of the study population (N = 232)

	N	Percentage
Gender		
Male	89	38.4
Female	142	61.2
Total	231	(mis = 1)
Age (years)		
Mean $\pm$ SD	73.7 $\pm$ 6.5	
Range	65–101	
Total	232	
Place of birth		
Israel	19	8.2
Asia/North Africa	123	53.0
Eastern Europe/former USSR	69	29.7
Western Europe/North and South America	19	8.2
Total	230	(mis = 2)
Marital status		
Single	0	0.0
Married	142	61.2
Divorced	14	6.0
Widowed	74	31.9
Separated	2	0.9
Total	232	
Education (years)		
Mean $\pm$ SD	10.95 $\pm$ 4.29	
Range	0–25	
Total	211	(mis = 11)
Employment		
Salaried	10	4.3
Partially salaried	3	1.3
Self-employed	10	4.3
Retired	184	79.3
Housewife	12	5.2
Unemployed	6	2.6
Total	225	(mis = 7)
Religious identity		
Agnostic	72	31.0
Religious	37	15.9
Traditional	123	53.0
Total	232	

**Abbreviation:** mis, missing data regarding some participants.

to the results of a recently published study,<sup>7</sup> direct intervention in the treatment of joint pain can, theoretically, reduce functional impairment.

The findings of the present study on the association between chronic pain and GS is supported by the results of many previous studies, which also reported associations between chronic pain and functional impairment,<sup>6,7,9,11–15</sup> falls,<sup>12</sup> cognitive problems,<sup>27–30</sup> and impaired mood.<sup>6,10,14,16–26</sup>

The overlap between pain and cognition would appear to be the fact that pain has a clear cognitive element that

**Table 2** Pain characteristics and impact (N = 232)

	N	Percentage
Number of pain sites		
1	182	78.4
2	40	17.2
3	8	3.4
4+	2	0.9
Total	232	
Pain site (more than one is possible)		
Back	87	37.5
Knee	62	26.7
Limbs	53	22.8
Shoulder	32	13.8
Head	21	9.1
Other	21	9.1
Chest	10	4.3
Abdomen	9	3.9
Total	232	
Pain level—five-point verbal descriptor scale		
No pain	0	0
Mild pain	14	6.0
Moderate pain	82	35.3
Severe pain	95	40.9
Unbearable pain	41	17.7
Total	232	
Pain intensity—10-point verbal numeric rating Scale (0 = no pain, 10 = unbearable pain)		
Mean $\pm$ SD	7.1 $\pm$ 2.1	
Range	1–10	
Total	230	(mis = 2)
Impact of pain on daily activity (0 = no impact, 10 = very high impact)		
General activity	5.7 $\pm$ 2.9	
Mood	5.1 $\pm$ 3.4	
Ambulation	5.3 $\pm$ 3.1	
Work	4.8 $\pm$ 3.3	
Relationships	2.7 $\pm$ 3.1	
Sleep	5.3 $\pm$ 3.4	
Pleasure	5.3 $\pm$ 3.3	
Pain interference		
Mean $\pm$ SD	4.9 $\pm$ 2.4	
Range	0–10	
Total	223	(mis = 9)

**Abbreviation:** mis, missing data regarding some participants.

necessitates recall of past experience and decision making,<sup>29</sup> since these two processes (chronic pain and cognition) utilize the element of attention.<sup>46</sup> In the review by Moriarty et al,<sup>29</sup> the authors raised an important question as to the association between the treatment administered for chronic pain and cognitive impairment: in some cases, this treatment can cause cognitive impairment, while in others, not only does analgesic therapy not cause cognitive impairment but it might even improve it. In fact, cognitive impairment can lead to the failure of appropriate treatment for chronic pain.<sup>28</sup>

**Table 3** Rate of geriatric syndromes (N=232)

	N	Percentage
Independent with grocery shopping		
Yes	112	48.3
No	120	51.7
Total	232	
Independent with preparing meals		
Yes	133	57.3
No	99	42.7
Total	232	
Independent with bathing		
Yes	185	79.7
No	47	20.3
Total	232	
Independent with dressing		
Yes	196	84.5
No	36	15.5
Total	232	
Any dependency in ADL <sup>a</sup>		
Yes	200	86.2
No	32	13.8
Total	232	
Falls in the past year		
Yes	84	36.8
No	144	63.2
Total	228	(mis = 4)
If yes, number of falls		
Mean ± SD	2.4 ± 1.8	
Range	1–10	
Total	79	(mis = 5)
Unplanned loss of weight of 5 kg or more over the last 6 months		
Yes	35	15.8
No	186	84.2
Total	221	(mis = 11)
Memory loss		
Yes	103	45.4
No	124	54.6
Total	227	(mis = 5)
Depressed or sad in the past 2 months		
Yes	97	43.1
No	128	56.9
Total	225	(mis = 7)
Six or more regular medications		
Yes	104	45.8
No	123	54.2
Total	227	(mis = 5)
Geriatric syndromes <sup>b</sup>		
0	34	14.7
1–2	90	38.8
3–6	108	46.6
Total	232	

**Notes:** <sup>a</sup>Any dependence on ADL—if the answer to one of the ADL questions (dressing, washing, shopping, or preparing food) was positive. <sup>b</sup>If the answer to one of the ADL questions was positive, ADL was considered to be impaired and none of the other questions in this category were taken into account for the calculation of the number of geriatric syndromes.

**Abbreviations:** ADL, activities of daily living; mis, missing data regarding some participants.

As noted, the association between chronic pain and affective changes has been investigated extensively in the past. Most of the longitudinal studies showed that the relationship between depression and chronic pain is bidirectional, where chronic pain can cause depression and depression can lead to development or exacerbation of chronic pain.<sup>18</sup>

The combination of pain and depression was found to be related to a negative health cognitive bias that makes patients more exposed to and more focused on their pain.<sup>47</sup> Depression can lead patients to a negative and pessimistic perception of the future and can have a negative effect on the patient's capacity to cope with pain.<sup>48</sup> In neurobiologic terms, the main noradrenergic and serotonergic nuclei in the central nervous system are responsible for the chronicity of pain and development of depression.<sup>49,50</sup>

The finding in the present study of a high rate of chronic pain that is associated with a high prevalence of GS in the elderly population highlights the need for a comprehensive plan to actively identify elderly patients with these problems.

As noted above, the Clalit Healthcare Services has a computerized system for initial screening of the elderly for the identification of GS,<sup>41</sup> but this system does not relate at all to the issue of pain. It should be upgraded to include this condition.

Many studies assessed associations between GS and pain, but each of these reported on only one or two GS.<sup>6,7,9–30</sup> In contrast, studies that assessed multiple, rather than specific, GS usually did not assess the co-occurrence of pain.<sup>33,51</sup> In the present study, we assessed the co-occurrence of chronic pain with multiple GS.

In the univariate analysis, we found a high rate of co-occurrence of pain in all the GS that were evaluated. In the logistic regression analysis, we found that chronic pain, as an independent variable, along with more advanced age and the presence of depression, predicts dependency in BADL. Although it is not possible, using a cross-sectional research design, to determine whether chronic pain predicts dependency in BADL or whether the association is bidirectional, our finding does strengthen the assumption of Andrews et al, on common underlying risk factors and mechanisms between pain and the other GS.<sup>32</sup>

Our study has some advantages. It was conducted in a population of elderly individuals living in the community who were selected at random, so it is reasonable to assume that the results can be generalized to other elderly populations.

**Table 4** Association between the number of geriatric syndromes, the intensity of pain, and the effect of pain on overall activity

	N	Pain intensity <sup>a</sup>				Pain interference <sup>b</sup>			
		Mean	SD	n	P	Mean	SD	n	P
Any dependency in ADL <sup>c</sup>									
Yes	145	7.4	2.1	144	0.004	5.7	2.2	139	<0.0001
No	87	6.6	2.2	86		3.5	2.3	84	
Falls over the previous year									
Yes	84	7.4	2.1	83	0.051	5.7	2.1	81	<0.0001
No	144	6.9	2.2	143		4.3	2.5	139	
Unplanned loss of weight of 5 kg or more over the last 6 months									
Yes	35	7.9	1.9	34	0.016	5.7	2.3	33	0.014
No	186	6.9	2.1	185		4.6	2.4	180	
Memory loss									
Yes	103	7.5	2.1	101	0.009	5.5	2.4	100	<0.0001
No	124	6.7	2.2	124		4.3	2.4	119	
Depressed or sad in the past 2 months									
Yes	97	7.5	2.1	95	0.005	6.2	2.1	91	<0.0001
No	128	6.7	2.1	128		3.8	2.2	126	
Six or more regular medications									
Yes	104	7.6	1.9	103	0.001	5.9	2.1	101	<0.0001
No	123	6.7	2.2	122		3.9	2.3	118	
Number of geriatric syndromes									
0	34	6.3	2.4	34	<0.0001	3.1	2.1	33	<0.0001
1–2	90	6.7	2.1	90		3.9	2.3	85	
3–6	108	7.7	1.9	106		6.2	1.9	105	

**Notes:** <sup>a</sup>10-point verbal numeric scale (0 = no pain, 10 = unbearable pain); <sup>b</sup>Effect of pain on overall activity (0 = usual activity, 10 = no activity); <sup>c</sup>Unable to perform at least one of the following activities independently: dressing, washing, preparing a meal, shopping.

**Abbreviation:** ADL, activities of daily living.

However, it also has several limitations. First, the evaluation was conducted by telephone interview and not face to face, and the data received from the patient, such as medication or cognitive state, were not confirmed vis-a-vis data from the medical records. The large number of interviewers in this study is also a potential limitation. Even though all the students who conducted the interviews underwent intensive training, the large number of interviewers (51) made it impossible to ensure the quality of all the interviews.

In the present study we cannot rule out the possibility of systematic biases, such as selection or information bias. Because of the study design, our data were based on subjective reporting of pain without any objective assessment, so we cannot confirm that the patients included in the study actually suffered from chronic pain as they reported. Therefore, we cannot rule out information bias. The study was conducted on a random sample of patients registered in the Clalit Healthcare Services, which provides medical care for two-thirds of Israel's aged  $\geq 65$  years population. This reduces the risk of selection bias but does not totally eliminate the possibility that proper randomization was not achieved.

Another significant limitation is the less than optimal planning of the study. Of the 419 patients who agreed to

participate in and completed the interview, 188 (44.8%) did not report any chronic pain so they were not asked to answer questions on co-morbid conditions or GS. In addition, 223 of the 922 elderly patients (24.2%) refused to be interviewed and another 255 (27.7%) could not be interviewed because they did not speak Hebrew ( $N = 178$ ) or because communication with them was impossible ( $N = 77$ ), usually due to impaired hearing. We cannot rule out the possibility that the prevalence of GS among those who did not participate may have been different from those who did participate.

Furthermore, over 25% of the study participants reported cognitive decline or a depressed mood, both of which could have affected their self-assessment of chronic pain. The patients were not asked about specific drug therapy in the interview, except indirectly through the questions of polypharmacy and chronic co-morbidity. Clearly, medications and medical history are important information in a study on pain, but these areas were not covered in the study questionnaire out of concern that the questionnaire would become too long and difficult to complete. The absence of this important information is an additional limitation of the study.

Another weakness is that the presence of GS was determined by means of a single dichotomous question and not by

**Table 5** Relationship between pain level and GS (IADL, BADL, any dependency in ADL, depression, and memory problems) by logistic regression analysis

Model	Variable	OR	95% CI		P-value
			Lower	Upper	
Any dependency in IADL <sup>a</sup>	Age	1.138	1.076	1.203	0.000
	Gender (male)	1.330	0.683	2.589	0.402
	Memory problems	1.386	0.712	2.699	0.337
	Depression	5.028	2.466	10.252	0.000
	Pain level (0–10)	1.129	0.967	1.317	0.124
Any dependency in BADL <sup>b</sup>	Age	1.099	1.041	1.160	0.001
	Gender (male)	0.609	0.284	1.307	0.203
	Memory problems	0.737	0.343	1.584	0.434
	Depression	6.330	2.924	13.700	0.000
	Pain level (0–10)	1.405	1.154	1.711	0.001
Any dependency in ADL <sup>c</sup>	Age	1.139	1.076	1.205	0.000
	Gender (male)	1.421	0.721	2.802	0.310
	Memory problems	1.349	0.684	2.660	0.388
	Depression	6.024	2.877	12.611	0.000
	Pain level (0–10)	1.157	0.989	1.353	0.069
Depressed or sad in the past 2 months	Age	1.007	0.960	1.057	0.771
	Gender (male)	0.620	0.324	1.189	0.150
	Memory problems	3.253	1.752	6.041	0.000
	Depression	5.969	2.863	12.447	0.000
	Pain level (0–10)	1.076	0.924	1.253	0.346
Memory loss	Age	0.992	0.948	1.039	0.742
	Gender (male)	0.770	0.423	1.402	0.393
	Memory problems	3.248	1.749	6.031	0.000
	Depression	1.396	0.708	2.755	0.336
	Pain level (0–10)	1.114	0.968	1.281	0.132

**Notes:** <sup>a</sup>Any dependency in IADL—if the answer to one of the IADL (instrumental activities of daily living) questions (shopping or preparing food) was positive. <sup>b</sup>Any dependency in BADL—if the answer to one of the BADL questions (dressing, washing, shopping, or preparing food) was positive. <sup>c</sup>Any dependency in ADL—if the answer to one of the ADL questions (dressing, washing, shopping, or preparing food) was positive.

**Abbreviations:** ADL, activities of daily living; BADL, basic activities of daily living; BPI, Brief Pain Inventory; CI, confidence interval; GS, geriatric syndromes; IADL, instrumental activities of daily living; OR, odds ratio

a more comprehensive assessment, for example, a full functional assessment or a formal cognitive assessment. Thus, the true rate of GS in the community may be different from the data that we presented here. Furthermore, we used the standard screening questionnaire of Clalit Healthcare Services, which did not include well-recognized GS, such as delirium, pressure sores, and incontinence, making the drawing of general conclusions more problematic.

Because of the cross-sectional design of the study, we can only discuss association, not causality, between chronic pain and GS.

### Conclusion

In a random study of individuals aged ≥65 years living in the community, we found a high rate of chronic pain and GS as well as an association between them. There is a need for further studies to investigate these issues in depth, using

accepted geriatric instruments for the determination of GS. This will facilitate more optimal care for the elderly population.

### Abbreviations

ADL, activities of daily living; BADL, basic activities of daily living; BPI, Brief Pain Inventory; GS, geriatric syndromes; IADL, instrumental activities of daily living.

### Acknowledgments

We would like to thank the nursing students of the Recanati School for Community Health Professions of Ben-Gurion University of the Negev for conducting the study interviews. This study was partially supported by a grant from the Recanati School for Community Health Professions, Faculty of Health Sciences, Ben-Gurion University of the Negev, Israel.



## Author contributions

Orly Liberman designed the study, supervised the data collection, and wrote the article. Tamar Freud was responsible for the statistical design of the study and for carrying out the statistical analysis. Roni Peleg designed the study and assisted with writing the article. Ariela Keren supervised the data collection and assisted with writing the article. Yan Press designed the study and wrote the article. All authors contributed toward data analysis, drafting and critically revising the paper, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

## Disclosure

Orly Liberman and Ariela Keren are employed at the Recanati School for Community Health Professions. The authors report no other conflicts of interest in this work.

## References

- Weber P, Meluzinova H, Matejovska-Kubesova H, et al. Geriatric giants--contemporary occurrence in 12,210 in-patients. *Bratisl Med J*. 2015;116(7):408–416.
- Noguchi N, Blyth FM, Waite LM, et al. Prevalence of the geriatric syndromes and frailty in older men living in the community: The Concord Health and Ageing in Men Project. *Australas J Ageing*. 2016;35(4):255–261.
- Inouye SK, Studenski S, Tinetti ME, Kuchel GA. Geriatric syndromes: clinical, research, and policy implications of a core geriatric concept. *J Am Geriatr Soc*. 2007;55(5):780–791.
- Tinetti ME, Inouye SK, Gill TM, Doucette JT. Shared risk factors for falls, incontinence, and functional dependence. Unifying the approach to geriatric syndromes. *J Am Med Assoc*. 1995;273(17):1348–1353.
- Senn N, Monod S. Development of a comprehensive approach for the early diagnosis of geriatric syndromes in general practice. *Front Med*. 2015;2:78.
- Hairi NN, Cumming RG, Blyth FM, Naganathan V. Chronic pain, impact of pain and pain severity with physical disability in older people--is there a gender difference? *Maturitas*. 2013;74(1):68–73.
- Henchoz Y, Bula C, Guessous I, et al. Chronic symptoms in a representative sample of community-dwelling older people: a cross-sectional study in Switzerland. *BMJ Open*. 2017;7(1):e014485.
- Johannes CB, Le TK, Zhou X, Johnston JA, Dworkin RH. The prevalence of chronic pain in United States adults: results of an Internet-based survey. *Journal Pain*. 2010;11(11):1230–1239.
- Pereira LS, Sherrington C, Ferreira ML, et al. Self-reported chronic pain is associated with physical performance in older people leaving aged care rehabilitation. *Clin Interv Aging*. 2014;9:259–265.
- Rafty MN, Sarma K, Murphy AW, De la Harpe D, Normand C, McGuire BE. Chronic pain in the Republic of Ireland--community prevalence, psychosocial profile and predictors of pain-related disability: results from the Prevalence, Impact and Cost of Chronic Pain (PRIME) study, part 1. *Pain*. 2011;152(5):1096–1103.
- Covinsky KE, Lindquist K, Dunlop DD, Yelin E. Pain, functional limitations, and aging. *J Am Geriatr Soc*. 2009;57(9):1556–1561.
- Kruschinski C, Wiese B, Dierks ML, Hummers-Pradier E, Schneider N, Junius-Walker U. A geriatric assessment in general practice: prevalence, location, impact and doctor-patient perceptions of pain. *BMC Fam Pract*. 2016;17:8.
- Scudds RJ, Ostbye T. Pain and pain-related interference with function in older Canadians: the Canadian Study of Health and Aging. *Disabil Rehabil*. 2001;23(15):654–664.
- Shega JW, Tiedt AD, Grant K, Dale W. Pain measurement in the National Social Life, Health, and Aging Project: presence, intensity, and location. *J Gerontol B Psychol Sci Soc Sci*. 2014;69(Suppl 2):S191–S197.
- Carmaci C, Iliffe S, Kharicha K, et al. Health risk appraisal in older people 3: prevalence, impact, and context of pain and their implications for GPs. *Br J Gen Pract*. 2007;57(541):630–635.
- Aguera-Ortiz L, Failde I, Cervilla JA, Mico JA. Unexplained pain complaints and depression in older people in primary care. *J Nutr Health Aging*. 2013;17(6):574–577.
- Biddulph JP, Iliffe S, Kharicha K, et al. Risk factors for depressed mood amongst a community dwelling older age population in England: cross-sectional survey data from the PRO-AGE study. *BMC Geriatr*. 2014;14:5.
- Kroenke K, Wu J, Bair MJ, Krebs EE, Damush TM, Tu W. Reciprocal relationship between pain and depression: a 12-month longitudinal analysis in primary care. *J Pain*. 2011;12(9):964–973.
- Leong IY, Farrell MJ, Helme RD, Gibson SJ. The relationship between medical comorbidity and self-rated pain, mood disturbance, and function in older people with chronic pain. *J Gerontol A Biol Sci Med Sci*. 2007;62(5):550–555.
- Lerman SF, Rudich Z, Brill S, Shalev H, Shahar G. Longitudinal associations between depression, anxiety, pain, and pain-related disability in chronic pain patients. *Psychosom Med*. 2015;77(3):333–341.
- Mallen CD, Peat G. Screening older people with musculoskeletal pain for depressive symptoms in primary care. *Br J Gen Pract*. 2008;58(555):688–693.
- Meyer T, Cooper J, Raspe H. Disabling low back pain and depressive symptoms in the community-dwelling elderly: a prospective study. *Spine*. 2007;32(21):2380–2386.
- Mossey JM, Gallagher RM. The longitudinal occurrence and impact of comorbid chronic pain and chronic depression over two years in continuing care retirement community residents. *Pain Med*. 2004;5(4):335–348.
- Parmelee PA, Harralson TL, McPherron JA, Schumacher HR. The structure of affective symptomatology in older adults with osteoarthritis. *Int J Geriatr Psychiatry*. 2013;28(4):393–401.
- Salazar A, Duenas M, Mico JA, et al. Undiagnosed mood disorders and sleep disturbances in primary care patients with chronic musculoskeletal pain. *Pain Med*. 2013;14(9):1416–1425.
- Shipton E, Ponnampereuma D, Wells E, Trewin B. Demographic characteristics, psychosocial measures, and pain in a sample of patients with persistent pain referred to a new zealand tertiary pain medicine center. *Pain Med*. 2013;14(7):1101–1107.
- Karp JF, Reynolds CF 3rd, Butters MA, et al. The relationship between pain and mental flexibility in older adult pain clinic patients. *Pain Med*. 2006;7(5):444–452.
- Landi F, Onder G, Cesari M, et al. Pain management in frail, community-living elderly patients. *Arch Intern Med*. 2001;161(22):2721–2724.
- Moriarty O, McGuire BE, Finn DP. The effect of pain on cognitive function: a review of clinical and preclinical research. *Prog Neurobiol*. 2011;93(3):385–404.
- Weiner DK, Rudy TE, Morrow L, Slaboda J, Lieber S. The relationship between pain, neuropsychological performance, and physical function in community-dwelling older adults with chronic low back pain. *Pain Med*. 2006;7(1):60–70.
- Lohman MC, Whiteman KL, Greenberg RL, Bruce ML. Incorporating persistent pain in phenotypic frailty measurement and prediction of adverse health outcomes. *J Gerontol A Biol Sci Med Sci*. 2017;72(2):216–222.
- Andrews JS, Cenzer IS, Yelin E, Covinsky KE. Pain as a risk factor for disability or death. *J Am Geriatr Soc*. 2013;61(4):583–589.
- Tabue-Teguio M, Grasset L, Avila-Funes JA, et al. Prevalence and co-occurrence of geriatric syndromes in people aged 75 years and older in France: results from the Bordeaux Three-city Study. *J Gerontol A Biol Sci Med Sci*. 2017;73(1):109–116.
- Anpalahan M, Gibson SJ. Geriatric syndromes as predictors of adverse outcomes of hospitalization. *Intern Med J*. 2008;38(1):16–23.
- Huang CC, Lee JD, Yang DC, Shih HI, Sun CY, Chang CM. Associations between geriatric syndromes and mortality in community-dwelling elderly: results of a national longitudinal study in Taiwan. *J Am Med Dir Assoc*. 2017;18(3):246–251.

36. Kane RL, Shamliyan T, Talley K, Pacala J. The association between geriatric syndromes and survival. *J Am Geriatr Soc.* 2012;60(5): 896–904.
37. Gagliese L, Weizblit N, Ellis W, Chan VW. The measurement of post-operative pain: a comparison of intensity scales in younger and older surgical patients. *Pain.* 2005;117(3):412–420.
38. Holdgate A, Asha S, Craig J, Thompson J. Comparison of a verbal numeric rating scale with the visual analogue scale for the measurement of acute pain. *Emerg Med.* 2003;15(5-6):441–446.
39. Cleeland CS, Ryan KM. Pain assessment: global use of the Brief Pain Inventory. *Ann Acad Med Singapore.* 1994;23(2):129–138.
40. Neville A, Peleg R, Singer Y, Sherf M, Shvartzman P. Chronic pain: a population-based study. *Isr Med Assoc J.* 2008;10(10):676–680.
41. Press Y, Hazzan R, Clarfield A, Dwolatzky T. A semistructured computerized screening interview for the assessment of older patients in the primary care setting *Int J Disabil Hum Dev* 2009;3:259–266.
42. Blyth FM, March LM, Brnabic AJ, Jorm LR, Williamson M, Cousins MJ. Chronic pain in Australia: a prevalence study. *Pain.* 2001;89(2–3): 127–134.
43. Pitkala KH, Strandberg TE, Tilvis RS. Management of nonmalignant pain in home-dwelling older people: a population-based survey. *J Am Geriatr Soc.* 2002;50(11):1861–1865.
44. Gloth FM 3rd. Pain management in older adults: prevention and treatment. *J Am Geriatr Soc.* 2001;49(2):188–199.
45. Von Roenn JH, Cleeland CS, Gonin R, Hatfield AK, Pandya KJ. Physician attitudes and practice in cancer pain management. A survey from the Eastern Cooperative Oncology Group. *Ann Intern Med.* 1993;119(2):121–126.
46. Oosterman JM, de Vries K, Dijkerman HC, de Haan EH, Scherder EJ. Exploring the relationship between cognition and self-reported pain in residents of homes for the elderly. *Int Psychogeriatr.* 2009;21(1): 157–163.
47. Rusu AC, Pincus T, Morley S. Depressed pain patients differ from other depressed groups: examination of cognitive content in a sentence completion task. *Pain.* 2012;153(9):1898–1904.
48. Von Korff M, Simon G. The relationship between pain and depression. *Br J Psychiatry Suppl.* 1996(30):101–108.
49. Alba-Delgado C, Borges G, Sanchez-Blazquez P, et al. The function of alpha-2-adrenoceptors in the rat locus coeruleus is preserved in the chronic constriction injury model of neuropathic pain. *Psychopharmacology.* 2012;221(1):53–65.
50. Alba-Delgado C, Mico JA, Sanchez-Blazquez P, Berrococo E. Analgesic antidepressants promote the responsiveness of locus coeruleus neurons to noxious stimulation: implications for neuropathic pain. *Pain.* 2012;153(7):1438–1449.
51. Lakhan P, Jones M, Wilson A, Courtney M, Hirdes J, Gray LC. A prospective cohort study of geriatric syndromes among older medical patients admitted to acute care hospitals. *J Am Geriatr Soc.* 2011;59(11): 2001–2008.

## Journal of Pain Research

### Publish your work in this journal

The Journal of Pain Research is an international, peer reviewed, open access, online journal that welcomes laboratory and clinical findings in the fields of pain research and the prevention and management of pain. Original research, reviews, symposium reports, hypothesis formation and commentaries are all considered for publication.

Submit your manuscript here: <https://www.dovepress.com/journal-of-pain-research-journal>

Dovepress

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.