

Indole-core-based novel antibacterial agent targeting FtsZ

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Wenchang Yuan,^{1,*}
Zhiwu Yu,^{2,*} Weiqi Song,^{3,*}
Yanan Li,⁴ Zhiyuan Fang,¹
Baizhen Zhu,¹ Xiaomei Li,¹
Hao Wang,⁵ Wei Hong,⁶
Ning Sun^{1,7}

¹The Fifth Affiliated Hospital of Guangzhou Medical University, Guangzhou 510700, People's Republic of China; ²Division of Laboratory Science, Affiliated Cancer Hospital & Institute of Guangzhou Medical University, Guangzhou 510095, People's Republic of China; ³School of Public Health, Guangzhou Medical University, Guangzhou 511436, People's Republic of China; ⁴Department of Pharmacy, The Fifth Affiliated Hospital of Sun Yat-sen University, Zhuhai 519000, People's Republic of China; ⁵School of Pharmacy, Ningxia Medical University, Yinchuan 750004, People's Republic of China; ⁶School of Chemistry and Chemical Engineering, North Minzu University, Yinchuan 750021, People's Republic of China; ⁷State Key Laboratory of Chemical Biology and Drug Discovery, and Department of Applied Biology and Chemical Technology, The Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong, People's Republic of China

*These authors contributed equally to this work

Correspondence: Ning Sun
The Fifth Affiliated Hospital of Guangzhou Medical University, 621 Gangwan Road, Huangpu District, Guangzhou 510700, People's Republic of China
Email ning.sun@connect.polyu.hk

Wei Hong
School of Chemistry and Chemical Engineering, North Minzu University, 204 Wenchang North Street, Xixia District, Yinchuan 750021, People's Republic of China
Email hongwei136@hotmail.com

Background: The prevalence of drug-resistant bacterial infections urges the development of new antibacterial agents that possess a mechanism of action different from traditional antibiotics. FtsZ has been recognized as a key functional protein in bacterial cell division and it is currently believed to be a potential target for the development of novel antibacterial agents.

Purpose: The primary aim of the study is to screen out an inhibitor targeting at FtsZ and followed to investigate its antibacterial activity and mode of action.

Methods: Cell-based cell division inhibitory screening assay, antimicrobial susceptibility test, minimum bactericidal concentration assay, time-killing curve determination, FtsZ polymerization assay, GTPase activity assay, and molecular modeling were performed in the present study.

Results: The screening study from a small library consisting of benzimidazole and indole derivatives discovered a compound (CZ74) with an indole-core structure. The compound exhibited strong cell division inhibitory effect. In addition, CZ74 shows high antibacterial potency against a number of tested Gram-positive bacteria, such as methicillin-resistant *Staphylococcus aureus* and vancomycin-resistant *Enterococcus*. The minimum inhibitory concentration values obtained were within the range of 2–4 µg/mL. The results of biological study revealed that CZ74 at 2 µg/mL is able to disrupt FtsZ polymerization and inhibit GTPase activity and cell division. From molecular modeling study, CZ74 is found possibly binding into the interdomain cleft of FtsZ protein and then leads to inhibitory effects.

Conclusion: This indole-cored molecule CZ74 could be a potential lead compound and could be further developed as a new generation of antibacterial agents targeting FtsZ to combat against multidrug-resistant bacteria.

Keywords: indole-core-based derivative, antibacterial, cell division, FtsZ, antibiotic resistant

Introduction

Over the past few decades, the use of antibiotics has greatly improved public health worldwide. However, due to the improper use of antibiotics, bacteria develop drug resistance rapidly to most antibiotics. This may imply that the battle against bacterial infections is endless.^{1,2} Nowadays, the issue of antimicrobial resistance is critical and is a global problem. For example, two typical representative bacteria, methicillin-resistant *Staphylococcus aureus* (MRSA) and vancomycin-resistant *Enterococcus faecium* (VREF), are resistant to the clinically used antibiotics such as methicillin and vancomycin.^{3,4} Considering the rapid emergence of drug-resistant bacteria and limited clinically used antibiotic treatment options, to develop

alternative strategies to fight against bacterial infections is therefore urgently needed.

With respect to the recent progress on new drug discovery, bacterial division could be a new area for the development of new generation of antibiotics. Among the proteins in the division, FtsZ has been recognized as a key factor.^{5,6} During bacterial cell division, FtsZ assembled into a circumferential ring (Z-ring) at the division site. The process involves GTP binding and hydrolysis followed by stabilization and anchoring onto the inner surface of the cytoplasmic membrane via FtsZ-binding proteins. When Z-ring is ready, it recruits a series of other cell division proteins that ultimately trigger cytokinesis.⁷⁻⁹ The high conservation and functional importance of FtsZ make it as a potential target for the development of new antibacterial agents.¹⁰ Recently, a few FtsZ-targeting compounds have been reported to inhibit bacterial cell division through interfering FtsZ activity.¹⁰⁻¹⁶ However, no FtsZ-targeting small molecule is found in clinical trial stage, meaning that further advancement on the molecular scaffold design to achieve higher inhibition potency is required.

Benzimidazoles and indoles are natural compounds and their aromatic heterocyclic structures are widely used for pharmaceutical synthesis and modification. Many

benzimidazole or indole core-based derivatives have good activity against various microbials, such as fungi and bacteria.^{17,18} Previous studies revealed that tri-substituted benzimidazole derivatives and indole derivatives (Figure 1) can inhibit effectively the growth of bacterial strains such as *M. tuberculosis*, *S. aureus*, and *B. subtilis* via disrupting FtsZ activity.¹⁹⁻²² In this study, we screened a small library of compounds containing 100 benzimidazole or indole core-based derivatives through a cell-based screening assay and successfully identified an indole core-based compound (CZ74, (E)-2-((1-(Phenylsulfonyl)-5-(4-(trifluoromethyl)phenyl)-1H-indol-3-yl)methylene) hydrazine-1-carboximide hydrochloride) (Figure 1). This small molecule was found possessing strong cell division inhibitory effect. In addition, we further investigated its antibacterial activity against several bacterial strains and the antibacterial mechanism.

Materials and methods

Cell-based cell division inhibitory screening assay

To search for compounds possessing strong cell inhibitory effects, a screening assay as described previously was performed.²³ In brief, *B. subtilis* 168 (ATCC 23857) cells




Figure 1 Structures of benzimidazole and indole derivatives.

were grown in starvation medium containing 1% hydrolyzed casein overnight. The culture was then diluted to OD₆₀₀ of ~0.01 in starvation medium containing 3% hydrolyzed casein and incubated at 37°C. The stationary phase culture was diluted to OD₆₀₀ of ~0.06, and 10 µL aliquots were transferred to 96-well microtiter plates containing different concentrations of the screening compounds in 100 µL volumes of medium. After incubation for approximately 5 hrs at 37°C, cell morphology was observed by phase-contrast light microscopy. Berberine, a natural compound has been reported to inhibit *B. subtilis* cell division at 64 µg/mL,²⁴ was used as a positive control in the assay. Each sample was repeated three times. The compound inhibited the *B. subtilis* cell division at the concentration <64 µg/mL would be selected to further investigation.

Antimicrobial susceptibility test

All the strains used in this test were purchased from American Type Culture Collection (ATCC, USA). And all of them are clinically relevant pathogens except *B. subtilis*. The minimum inhibitory concentrations (MICs) of CZ74 against the tested strains were determined following procedures described in the Clinical and Laboratory Standards Institute (CLSI) guidelines.²⁵ *S. aureus* strains were cultured in cation-adjusted Mueller Hinton broth (CA-MHB), *E. faecalis* and *E. faecium* strains were cultured in Brain Heart Infusion broth (BHI), the other strains were cultured in Mueller Hinton broth (MHB) in the assays. The MIC was defined as the lowest compound concentration at which prevents visible growth of bacterium. Methicillin and berberine were applied as a reference in this assay. Each test was repeated three times.

Minimum bactericidal concentration (MBC) assay

This assay was used to detect the MBC value of an antibacterial agent, and to further confirm whether the compound is bactericidal or not. The test performed using the protocol described above. The MBC was determined by placing 10 µL of culture volume from the MIC assay onto CA-MHB agar plate for *S. aureus* strains, MHB agar plate for *B. subtilis*, and BHI agar plate for *E. faecium*. Then, the plates were incubated at 37°C, and the colony formation was examined after 24 hrs. MBC was recorded as the lowest concentration at which bacterial colonies were not observed.²⁶ Each test was repeated three times.

Time-killing curve determination

This assay was used to detect the kinetics of antibacterial activity of CZ74. An overnight culture of *S. aureus* ATCC 25923 and *B. subtilis* 168 was diluted to 10⁶ CFU/mL approximately. Different concentrations of CZ74 were added to bacterial culture. The cultures were then incubated at 37°C 250 rpm. After that, 100 µL of the culture were taken for serial dilutions and then spread on CA-MHB or MHB agar plates at different time intervals (0, 2, 4, 6, 8, 24 hrs). The plates were incubated at 37°C for 24 hrs and then the cell counts (CFU/mL) were recorded. Each test was repeated three times.

FtsZ polymerization assay

To investigate the possible antibacterial mechanism of CZ74, *S. aureus* FtsZ was prepared according to the previous study.²⁴ Some biochemical assays were also performed to confirm the effect of CZ74 on the FtsZ. Firstly, the light scattering assay and transmission electron microscopy (TEM) analysis were performed by following a protocol from the literature²⁷ to investigate the effect of CZ74 on the FtsZ polymerization. In the light scattering assay, FtsZ (6 µM) was incubated with different concentrations of CZ74 at room temperature for 10 mins. The light scattering signal was immediately detected by a thermostatically (37°C) controlled fluorescence spectrometer after additions of 20 mM KCl, 5 mM MgCl₂, 1 mM GTP. One percentage DMSO and 5 µg/mL methicillin were tested as vehicle and negative control in the assay. In the TEM analysis, 12 µM FtsZ was incubated with one percentage DMSO or 4 µg/mL of CZ74 in 50 mM buffer for 10 mins at room temperature. Then, 5 mM MgCl₂, 50 mM KCl, and 1 mM GTP were added to the mixtures. The samples were then incubated in water bath at 37°C. After 15 mins, 10 µL of the sample was transferred to a Formvar carbon-coated copper grid and subjected to 0.5% phosphotungstic acid for negative staining. And the air-dried sample was observed with a transmission electron microscope.²⁷

GTPase activity assay

The GTPase activity assay was conducted by following the previously described procedures in the investigation of CZ74 on the GTPase activity of FtsZ.¹⁶ In short, 3.5 µM FtsZ was incubated with different concentrations of CZ74 (0, 1, 2, 4, 8, and 16 µg/mL) at room temperature in 20 mM Tris buffer. After 10 mins, 5 mM of MgCl₂, 200 mM

of KCl, and 500 mM GTP were added. The samples were then incubated at 37°C for 30 mins. After that, the reactions were quenched by adding 100 mL of Cytophos reagent for 10 mins. Inorganic phosphate was quantified with a microplate reader at 650 nm.

Molecular modeling

Discovery Studio (DS) 2016 was used to perform the molecular modeling study. The X-ray crystal structure of SaFtsZ was downloaded from the PDB database (PDB entry: 4DXD).²⁸ FtsZ and the ligand CZ74 were prepared for this study using the own tools in DS. And then, the CDocker protocol was used to predict the binding poses and the top-scoring poses were visually inspected.

Preparation of compound CZ74

The preparation and characterization of CZ74 can be referred to supplementary material and Figure S1.

Results

Cell inhibitory effect of CZ74 against *B. subtilis*

The cell-based screening assay identified three potential compounds (Table S1), which inhibited the growth of *B. subtilis* at the concentration equal to or below 64 µg/mL. However, only CZ74 possesses bacterial cell division inhibitory effect. Compared to the control experiments, CZ74 elongated the cell length of *B. subtilis* obviously at the concentration of 2 µg/mL (Figure 2). The untreated *B. subtilis* cells have typical short rod morphology with cell lengths from 5 to 15 µm (Figure 2A). After incubation with CZ74 at 2 µg/mL, the cells were found longer than 50 µm (Figure 2B). This phenomenon was similar to the *B. subtilis* cells treated with 64 µg/mL berberine (Figure S2). These results are also consistent with the reported FtsZ inhibitors, such as benzamide and pyrimidine derivatives,^{13,29,30} and




Figure 2 Cell division inhibitory effect of CZ74. Cells of *B. subtilis* 168 were grown in the absence (A) and presence of 2 µg/mL of CZ74 (B). Scale bar=15 µm.

suggest that CZ74 may disrupt FtsZ function and lead to the abnormal cell division.

Antibacterial activity of CZ74

To investigate the antibacterial activity of CZ74, the compound was examined against several bacterial strains including some drug-resistant strains. The results showed that CZ74 inhibited all the tested Gram-positive strains effectively. The MIC values are found in the range of 2–4 µg/mL. For instance, the growth of *B. subtilis* 168, *S. aureus* ATCC 25923, and *S. epidermidis* ATCC 12228 were inhibited at 2 µg/mL. The MIC value is comparable to that of methicillin and is much lower than that of berberine. Furthermore, the MIC values of CZ74 against the MRSA (ATCC BAA-41, 33591, 33592, and 43300) is as low as 2 µg/mL. Compared with methicillin and berberine, the antibacterial activity of CZ74 against MRSA is much more potent (Table 1). The compound was also evaluated against 50 clinical isolated MRSA. The results also show that CZ74 has a potent antibacterial activity with the MIC value in the range from 2 to 4 µg/mL (Table S2). CZ74 also inhibited the growth of *E. faecalis* and *E. faecium* effectively with the MIC value of 4 µg/mL (Table 1). Since the MICs of vancomycin against vancomycin-resistant *E. faecalis* and *E. faecium* (VREs) are higher than 96 µg/mL, CZ74 possesses higher antibacterial activity against VREs than that of vancomycin.²⁴

Table 1 Antibacterial activity of CZ74 against a panel of bacterial strains

Organism	MIC (µg/mL)		
	CZ74	MET	BER
<i>B. subtilis</i> 168	2	<1	96
<i>S. aureus</i> ATCC 25923	2	1	192
<i>S. aureus</i> ATCC 29247 ^a	2	4	192
<i>S. aureus</i> ATCC BAA-41 ^b	2	1024	192
<i>S. aureus</i> ATCC 33591 ^b	2	1024	192
<i>S. aureus</i> ATCC 33592 ^b	2	512	192
<i>S. aureus</i> ATCC 43300 ^b	2	512	192
<i>S. epidermidis</i> ATCC 12228	2	0.75	192
<i>E. faecalis</i> ATCC 29212	4	4	>192
<i>E. faecalis</i> ATCC 51575 ^c	4	4	>192
<i>E. faecium</i> ATCC 49624	4	4	>192
<i>E. faecium</i> ATCC 700221 ^c	4	4	>192
<i>E. coli</i> ATCC 25922	>64	4	>192
<i>P. aeruginosa</i> ATCC BAA-2108	>64	>1024	>192
<i>K. pneumoniae</i> ATCC BAA-1144	>64	>1024	>192

Notes: ^aAmpicillin-resistant strain; ^bMRSA; ^cVancomycin-resistant strains.

Abbreviations: MET, methicillin; BER, berberine.

However, in the antibacterial effect against Gram-negative strains such as *P. aeruginosa*, *K. pneumoniae*, and *E. coli*, the compound at 64 $\mu\text{g/mL}$ did not show any antibacterial effect. The antibacterial property of CZ74 observed is similar to those of reported FtsZ-targeting compounds such as benzamide derivatives.^{30,31}

Bactericidal test of CZ74 against the selected bacterial strains

After obtained the antibacterial activity of CZ74, it is important to further investigate whether the antibacterial activity is bactericidal or bacteriostatic. In this study, the MBC values of CZ74 against *B. subtilis* 168, *S. aureus* ATCC 25923, MRSA (ATCC 43300), and VREF (ATCC 700221) were determined (Table 2). According to the CLSI standards, an MBC/MIC ratio of 1–2 is considered indicative of bactericidal behavior.²⁵ The results show that all the MBC/MIC ratios obtained are equal to or lower than 2, indicating bactericidal activity of CZ74 against the tested bacterial species.

Table 2 The MBC of CZ74 against drug-resistant bacterial strains ($\mu\text{g/mL}$)

Bacteria	MBC	MIC	MBC/MIC
<i>B. subtilis</i> 168	4	2	2
<i>S. aureus</i> ATCC 25923	4	2	2
<i>S. aureus</i> ATCC 43300	2	2	1
<i>E. faecium</i> ATCC 700221	8	4	2

Kinetics of the bactericidal activity of CZ74

After the bactericidal profile of CZ74 was confirmed, the time-killing curve determinations were performed to investigate the kinetics of killing of CZ74 against *S. aureus* ATCC 25923 and *B. subtilis* 168. As shown in Figure 3A, *S. aureus* grew rapidly in the first 6 hrs without CZ74. After 6 hrs incubation, the growth of *S. aureus* went into the stationary phase. On the other hand, the concentration of CZ74 at 2 \times MIC and 4 \times MIC can completely inhibit the growth of *S. aureus* after incubation overnight. At the concentration of 1 \times MIC, the compound also reduces the viable counts of *S. aureus* below 10^3 CFU mL^{-1} after incubation for 24 hrs. The growth inhibition with different concentration gradients of CZ74 demonstrated that the rate of killing *S. aureus* was dose-dependent. In addition, a similar kinetic was found when *B. subtilis* incubated with CZ74 for 24 hrs (Figure 3B).

Effect of CZ74 on the FtsZ polymerization

In order to understand the mechanism of antibacterial activity of CZ74, the effect of CZ74 on the dynamic polymerization of FtsZ was investigated by using a light scattering assay. Figure 4A shows the time-dependent polymerization profile of FtsZ in the absence and in the presence of CZ74 at a concentration range from 1 to 4 $\mu\text{g/mL}$. The results revealed that CZ74 stimulated FtsZ polymerization in a concentration-dependent profile. The similar phenomena can also be observed in other reported




Figure 3 Time-killing curves of CZ74. (A) Samples of a growing culture of *S. aureus* ATCC 25923 were incubated with concentrations of CZ74 equivalent to 1 \times (red), 2 \times (green), or 4 \times (blue) the MIC. (B) Samples of a growing culture of *B. Subtilis* 168 were incubated with concentrations of CZ74 equivalent to 1 \times (red), 2 \times (green), or 4 \times (blue) the MIC. Vehicle (1% DMSO; black) was included.




Figure 4 Time-dependent polymerization profiles of FtsZ in the absence and presence of CZ74 at a concentration range from 1 to 4 μg/mL (A). Electron micrographs of FtsZ polymers in the absence (B) and in the presence (C) of 4 μg/mL of CZ74. Scale bar = 1000 nm.

FtsZ-targeting compounds.^{10,11} Methicillin was also tested under the same conditions as a negative control. As expected, it does not affect FtsZ polymerization. Apart from the light scattering assay, we also used TEM to visualize the effect of CZ74 on the FtsZ polymerization. The results showed that both the size of the FtsZ polymers and the bundles of the FtsZ protofilaments were dramatically increased when using 2 μg/mL CZ74 (Figure 4B and C).

Effect of CZ74 on the GTPase activity of FtsZ

The rate of GTP hydrolysis is correlated with the dynamic polymerization of FtsZ.^{8,9} To investigate whether the antibacterial mechanism of CZ74 is related to the GTPase inhibitory effect, the effect of this compound on the GTPase activity of the FtsZ was determined. The results revealed that the compound inhibited the GTPase activity of FtsZ in a dose-dependent profile. In the assay, 1, 2, 4, 8, and 16 μg/mL of CZ74 inhibited the GTPase activity by 15%, 30%, 45%, 60%, and 62%, respectively (Figure 5). The result indicates that the inhibitory effect of CZ74 on GTPase activity may probably lead to the disruption of FtsZ polymerization.

Predicted the binding mode of CZ74

Molecular modeling study was also conducted to predict the binding sites and the interaction pose of CZ74 in FtsZ. The optimal docking pose suggests that the compound probably binds to a hydrophobic and narrow cleft constituted by the T7-loop, H7-helix, and a four-stranded β-sheet (Figure 6A). The possible interactions between CZ74 and the amino acids of FtsZ were shown in a 2D ligand interaction diagram (Figure 6B). The guanidyl group of CZ74 could interact




Figure 5 Inhibition of GTPase activity of FtsZ by different concentrations of CZ74.




Figure 6 Predicted binding mode of CZ74 bound to FtsZ. **(A)** CZ74 bound to the C-terminal interdomain cleft of FtsZ; **(B)** predicted interaction between CZ74 and amino acids of FtsZ.

with Val203, Asn208, and Leu209 of FtsZ protein through four conventional hydrogen bonds. Several favorable hydrophobic interactions between CZ74 and the side chains of Asp199, Val203, Met226, Thr265, Val297, and Ile311 were predicted in the binding model. In addition, the trifluoromethyl group of CZ74 was predicted participating in one carbon-hydrogen bond with the side chain of Gly193 and three halogen bonds with Gly193, Gly227, and Val310. Moreover, a lot of amino acids in this pocket (such as Gly196, Val307, Thr309, etc.) could interact with CZ74 via van der Waals force (data not shown).

Discussion

Indoles are a class of important heterocycles possessing diverse biological activities, such as anticancer, antibacterial, and antiviral.^{32–34} In this study, we found that an indole derivative CZ74 possesses strong antibacterial activity against the tested Gram-positive bacteria, including MRSA and VRE. The MIC values obtained are in the range of 2–4 µg/mL, which is similar to or slightly better than certain reported inhibitors targeting at FtsZ. In particular, berberine derivative inhibited the growth of MRSA with a MIC value of 2 µg/mL and VREF with a MIC value of 8 µg/mL.²⁴ PC 190723 can effectively against MRSA at the concentration of 1 µg/mL.³⁰ However, CZ74 has not any antibacterial activity against the tested Gram-negative

strain at the concentration of 64 µg/mL, likely owing to the intrinsically lower permeability of the Gram-negative cell envelope.³⁵

In the biological assays, CZ74 was found to stimulate FtsZ polymerization with GTPase inhibitory effect. The similar biochemical results were also found in some FtsZ inhibitors, such as PC190723 and quinoline derivative.^{30,36} FtsZ polymerization is found to be regulated by GTPase activity of FtsZ⁸ and the increasing of FtsZ polymers may probably contribute from the conformational change to the high-affinity state, which enables FtsZ assembly.³⁷ In addition, same as those reported FtsZ-targeting compounds,¹⁰ the strong inhibitory effect of CZ74 on the bacterial cell division was observed. Recently, our group has reported some FtsZ-targeting compounds with strong antibacterial activity, such as berberine derivatives, pyrimidine derivatives, and quinolinium derivatives.^{15,16,24,27,29} Unfortunately, the low solubility of these compounds hindered their further investigation in animal study. We expected that the polar guanidyl group of CZ74 could overcome the solubility issue and make it possible for in vivo study. In addition, this hydrophilic guanidyl group was predicted to interact with FtsZ through hydrogen bonds (Figure 6).

In summary, an indole-core-based compound was demonstrated as a potential antibacterial agent targeting

FtsZ. The compound exhibits good antibacterial activity against several clinically representative Gram-positive strains including some drug-resistant bacterial strains, such as MRSA and VRE. From the biochemical evaluations with the compound, the mode of action was probed. It was suggested that the compound could possibly bind into the interdomain cleft of FtsZ and take functions in disrupting the GTPase activity, stimulating FtsZ polymerization, and eventually inhibiting bacterial cell division to cause cell death. Therefore, this indolecored small molecule is suitable for further investigation on their biological activities, particularly the antibacterial mechanism.

Acknowledgments

We acknowledge the support from the National Natural Science Foundation of China (81703333 and 81660588), the Science and Technology Program of Guangzhou Municipal Health Commission (20192A011018), and the Medical Scientific Research Foundation of Guangdong Province (A2017588).

Disclosure

The authors declare no conflicts of interest in this work.

References

- Zhang SX, Zhou YM, Tian LG, et al. Antibiotic resistance and molecular characterization of diarrheagenic *Escherichia coli* and nontyphoidal *Salmonella* strains isolated from infections in Southwest China. *Infect Dis Poverty*. 2018;7(1):53. doi:10.1186/s40249-018-0427-2
- Wright GD. Antibiotics: a new hope. *Chem Biol*. 2012;19(1):3–10. doi:10.1016/j.chembiol.2011.10.019
- Li Y, Yang L, Fu J, Yan M, Chen D, Zhang L. Microbial pathogenicity and virulence mediated by integrons on Gram-positive microorganisms. *Microb Pathog*. 2017;111:481–486. doi:10.1016/j.micpath.2017.09.035
- Gould IM, David MZ, Esposito S, et al. New insights into methicillin-resistant *Staphylococcus aureus* (MRSA) pathogenesis, treatment and resistance. *Int J Antimicrob Agents*. 2012;39(2):96–104.
- Adams DW, Errington J. Bacterial cell division: assembly, maintenance and disassembly of the Z ring. *Nat Rev Microbiol*. 2009;7(9):642–653.
- Lock RL, Harry EJ. Cell-division inhibitors: new insights for future antibiotics. *Nat Rev Drug Discov*. 2008;7(4):324–338.
- Roseboom W, Nazir MG, Meiresonne NY, et al. Mapping the contact sites of the *Escherichia coli* division-initiating proteins FtsZ and ZapA by BAMG cross-linking and site-directed mutagenesis. *Int J Mol Sci*. 2018;19(10):2928.
- Yang X, Lyu Z, Miguel A, McQuillen R, Huang KC, Xiao J. GTPase activity-coupled treadmilling of the bacterial tubulin FtsZ organizes septal cell wall synthesis. *Science*. 2017;355(6326):744–747.
- Bisson-Filho AW, Hsu YP, Squyres GR, et al. Treadmilling by FtsZ filaments drives peptidoglycan synthesis and bacterial cell division. *Science*. 2017;355(6326):739–743.
- Hurley KA, Santos TM, Nepomuceno GM, Huynh V, Shaw JT, Weibel DB. Targeting the bacterial division protein FtsZ. *J Med Chem*. 2016;59(15):6975–6998. doi:10.1021/acs.jmedchem.5b01098
- Haranahalli K, Tong S, Ojima I. Recent advances in the discovery and development of antibacterial agents targeting the cell-division protein FtsZ. *Bioorg Med Chem*. 2016;24(24):6354–6369. doi:10.1016/j.bmc.2016.05.003
- Foss MH, Eun Y-J, Weibel DB. Chemical–biological studies of subcellular organization in bacteria. *Biochemistry-US*. 2011;50(36):7719–7734. doi:10.1021/bi200940d
- Fang Z, Zheng S, Chan KF, et al. Design, synthesis and antibacterial evaluation of 2,4-disubstituted-6-thiophenyl-pyrimidines. *Eur J Med Chem*. 2019;161:141–153. doi:10.1016/j.ejmech.2018.10.039
- Bi F, Song D, Zhang N, et al. Design, synthesis and structure-based optimization of novel isoxazole-containing benzamide derivatives as FtsZ modulators. *Eur J Med Chem*. 2018;159:90–103. doi:10.1016/j.ejmech.2018.09.053
- Sun N, Du RL, Zheng YY, et al. Antibacterial activity of 3-methylbenzo[d]thiazol-methylquinolinium derivatives and study of their action mechanism. *J Enzyme Inhib Med Chem*. 2018;33(1):879–889. doi:10.1080/14756366.2018.1465055
- Sun N, Du RL, Zheng YY, et al. Antibacterial activity of N-methylbenzofuro[3,2-b]quinoline and N-methylbenzoindolo[3,2-b]quinoline derivatives and study of their mode of action. *Eur J Med Chem*. 2017;135:1–11. doi:10.1016/j.ejmech.2017.04.018
- Zhao F, Liu N, Zhan P, Jiang X, Liu X. Discovery of HCV NS5B thumb site I inhibitors: core-refining from benzimidazole to indole scaffold. *Eur J Med Chem*. 2015;94:218–228. doi:10.1016/j.ejmech.2015.03.012
- Zhang MZ, Mulholland N, Beattie D, et al. Synthesis and antifungal activity of 3-(1,3,4-oxadiazol-5-yl)-indoles and 3-(1,3,4-oxadiazol-5-yl)methyl-indoles. *Eur J Med Chem*. 2013;63:22–32. doi:10.1016/j.ejmech.2013.01.038
- Kumar K, Awasthi D, Lee SY, et al. Novel trisubstituted benzimidazoles, targeting Mtb FtsZ, as a new class of antitubercular agents. *J Med Chem*. 2011;54(1):374–381. doi:10.1021/jm1012006
- Margalit DN, Romberg L, Mets RB, et al. Targeting cell division: small-molecule inhibitors of FtsZ GTPase perturb cytotubulin ring assembly and induce bacterial lethality. *Proc Natl Acad Sci U S A*. 2004;101(32):11821–11826. doi:10.1073/pnas.0404439101
- Jennings LD, Foreman KW, Rush TS 3rd, et al. Combinatorial synthesis of substituted 3-(2-indolyl)piperidines and 2-phenyl indoles as inhibitors of ZipA-FtsZ interaction. *Bioorg Med Chem*. 2004;12(19):5115–5131. doi:10.1016/j.bmc.2004.07.031
- Sun N, Zheng YY, Du RL, et al. New application of tiplaxtinin as an effective FtsZ-targeting chemotype for an antimicrobial study. *Medchemcomm*. 2017;8(10):1909–1913. doi:10.1039/c7md00387k
- Stokes NR, Sievers J, Barker S, et al. Novel inhibitors of bacterial cytokinesis identified by a cell-based antibiotic screening assay. *J Biol Chem*. 2005;280(48):39709–39715. doi:10.1074/jbc.M506741200
- Sun N, Chan F-Y, Lu Y-J, et al. Rational design of berberine-based FtsZ inhibitors with broad-spectrum antibacterial activity. *PLoS One*. 2014;9(5):e97514. doi:10.1371/journal.pone.0097514
- Wikler MA, Hindler JF, Cookerill FR, Patel JB, Bush K, Powell M. *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically*. Wayne, PA, USA: Clinical and Laboratory Standards Institute. Vol. 29; 2009:M07–A08.
- Zheng Z, Tharmalingam N, Liu Q, et al. Synergistic efficacy of aedes aegypti antimicrobial peptide cecropin A2 and tetracycline against *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother*. 2017;61:7. doi:10.1128/AAC.00686-17
- Sun N, Lu YJ, Chan FY, et al. A thiazole orange derivative targeting the bacterial protein FtsZ shows potent antibacterial activity. *Front Microbiol*. 2017;8:855. doi:10.3389/fmicb.2017.00855
- Tan CM, Therien AG, Lu J, et al. Restoring methicillin-resistant *Staphylococcus aureus* susceptibility to beta-lactam antibiotics. *Sci Transl Med*. 2012;4(126):126ra135. doi:10.1126/scitranslmed.3003592

29. Chan KF, Sun N, Yan SC, et al. Efficient synthesis of amine-linked 2,4,6-trisubstituted pyrimidines as a new class of bacterial FtsZ inhibitors. *ACS Omega*. 2017;2(10):7281–7292. doi:10.1021/acsomega.7b00701
30. Haydon DJ, Stokes NR, Ure R, et al. An inhibitor of FtsZ with potent and selective anti-staphylococcal activity. *Science*. 2008;321(5896):1673–1675. doi:10.1126/science.1159961
31. Haydon DJ, Bennett JM, Brown D, et al. Creating an antibacterial with in vivo efficacy: synthesis and characterization of potent inhibitors of the bacterial cell division protein FtsZ with improved pharmaceutical properties. *J Med Chem*. 2010;53(10):3927–3936. doi:10.1021/jm9016366
32. Kaur M, Singh M, Chadha N, Silakari O. Oxindole: A chemical prism carrying plethora of therapeutic benefits. *Eur J Med Chem*. 2016;123:858–894. doi:10.1016/j.ejmech.2016.08.011
33. Yang YT, Zhu JF, Liao GC, Xu HJ, Yu B. The development of biologically important spirooxindoles as new antimicrobial agents. *Curr Med Chem*. 2018;25(19):2233–2244.
34. Zhang MZ, Chen Q, Yang GF. A review on recent developments of indole-containing antiviral agents. *Eur J Med Chem*. 2015;89:421–441.
35. Keffer JL, Huecas S, Hammill JT, Wipf P, Andreu JM, Bewley CA. Chrysopaentins are competitive inhibitors of FtsZ and inhibit Z-ring formation in live bacteria. *Bioorg Med Chem*. 2013;21(18):5673–5678.
36. Fang Z, Ban L, Li Y, et al. A quinoline-based FtsZ inhibitor for the study of antimicrobial activity and synergistic effects with beta-lactam antibiotics. *J Pharmacol Sci*. 2018;137(3):283–289.
37. Elsen NL, Lu J, Parthasarathy G, et al. Mechanism of action of the cell-division inhibitor PC190723: modulation of FtsZ assembly cooperativity. *J Am Chem Soc*. 2012;134(30):12342–12345.

Supplementary material

Part I: Preparation and characterization of CZ74

The preparation of CZ74 was performed using a procedure as previously described (*The Journal of Antibiotics*. 2017;70:832–844).¹

Characterization:

¹H-NMR (400 MHz, DMSO): δ 12.05 (s, 1H, NH), 8.57 (s, 1H, Ar-H), 8.49 (d, $J=1.5$, 1H, Ar-H), 8.43 (s,

1H, CH), 8.02–8.13 (m, 3H, Ar-H), 7.95 (d, $J=8.1$, 2H, Ar-H), 7.70–7.85 (m, 7H, Ar-H and NH), 7.64 (t, $J=7.8$, 2H, Ar-H); ES-MS 486.1 (M + H)⁺; HRMS calcd for C₂₁H₁₉F₃N₅O₂S + 486.1212, found 486.1201.







Figure S1 ¹H-NMR spectrum of CZ74.

Part 2

Table S1 Compounds possess antibacterial activity against *B. subtilis* generated from cell-based screening assay

Structure of compound	Concentration (inhibit the growth of <i>B. subtilis</i>)	Concentration (inhibit bacterial cell division)
 <p>Cmpd 1</p>	1 µg/mL	No effect on bacterial cell division
 <p>Cmpd 2</p>	64 µg/mL	No effect on bacterial cell division
 <p>Cmpd 3 (CZ74)</p>	2 µg/mL	Inhibit bacterial cell division at 2 µg/mL

Part 3



Figure S2 Cell inhibitory effect of 64 µg/mL berberine on *B. subtilis* cells. Scale bar=15 µm.

Part 4

Table S2 Antibacterial activity of CZ74 against clinical isolated MRSA

MRSA ^a	MIC µg/mL	
	CZ74	Vancomycin
1428	2	0.5
8162	2	1
7917	4	1
3123	2	1
7247	4	1
7717	4	1
2671	2	1
3170	2	2
4127	2	1
3096	2	1
7971	2	1
7807	2	2
2491	4	2
2931	2	1
7863	4	1
7943	4	2
7947	4	1
1610	4	1
7813	2	1
5426	2	1
6334	2	1
7864	4	1
7972	4	1
1220	2	0.5
6515	2	1
4023	2	2
7960	4	2
1651	2	2
8179	2	1
8071	2	1
4955	2	1
5395	2	1
7844	4	1
7916	4	1
2158	2	1
7682	4	1
5239	2	1
2350	2	1
7664	2	0.5
7536	2	1
7587	4	2
1708	2	1
5035	2	1
4548	2	2
1947	2	2

(Continued)

Table S2 (Continued).

MRSA ^a	MIC µg/mL	
	CZ74	Vancomycin
5056	2	2
8078	2	2
7434	4	1
7449	4	1
7183	4	2

Notes: ^aAll the strains are resistant to methicillin (MIC>64 µg/mL).

Reference

1. Hong W, Li J, Chang Z, et al Synthesis and biological evaluation of indole core-based derivatives with potent antibacterial activity against resistant bacterial pathogens. *The Journal of Antibiotics*. 2017;70(7):832.

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>