

# Deficient Regulatory Innate Lymphoid Cells and Differential Expression of miRNAs in Acute Myeloid Leukemia Quantified by Next Generation Sequence

This article was published in the following Dove Press journal:  
*Cancer Management and Research*

Jifeng Yu   
Yingmei Li  
Yue Pan   
Yu Liu  
Haizhou Xing  
Xinsheng Xie  
Dingming Wan  
Zhongxing Jiang

Department of Hematology, The First  
Affiliated Hospital of Zhengzhou  
University, Zhengzhou 450052, People's  
Republic of China

**Background:** A new regulatory subpopulation of ILCs, IL1Creg has been identified in mouse and human intestines. IL1Cregs share characteristics with both innate lymphoid cells and regulatory cells; however, the significance of CD45<sup>+</sup>Lin<sup>-</sup>CD127<sup>+</sup>IL-10<sup>+</sup> IL1Cregs in patients with AML remains unclear. Intriguingly, IL1Cregs constitutively express id2, id3, sox4, tgfr1, tgfr2, il2rb and il2rg, but the significance of miRNAs associated with these genes has yet to be explored. In this study, we evaluate IL1Creg frequency, IL1Creg gene-associated miRNA quantification, and its significance in patients with AML and normal donors.

**Methods:** Using 4 color combinations of surface and intracellular antibody staining, the CD45<sup>+</sup>Lin<sup>-</sup>CD127<sup>+</sup>IL-10<sup>+</sup> IL1Cregs from 12 normal donors and 42 patients newly diagnosed with AML were measured by flow cytometry. Plasma samples and bone marrow cells from 6 normal donors and 9 patients with AML were studied by next-generation sequence miRNAs quantification.

**Results:** Our results showed that the frequency of IL1Cregs was 0.8924±1.3791% in bone marrow (BM) cells from normal donors and 0.2434±0.5344% in BM cells from AML patients. The frequency of IL1Creg cells in AML patients was significantly lower than that in normal donors (P<0.01). Furthermore, the frequency of the CD45<sup>+</sup>Lin<sup>-</sup>CD127<sup>+</sup>IL-10<sup>-</sup> subset was 4.0869±6.7701% and 0.2769±0.2526% from normal donors and AML patients, respectively. There was a statistically significant difference of CD45<sup>+</sup>Lin<sup>-</sup>CD127<sup>+</sup>IL-10<sup>-</sup> cells between normal donors and AML patients (p<0.01). miRNA detection results showed 376 miRNAs from plasma and 182 miRNAs from BM cell samples with expression levels with a statistically significant difference between AML patients and normal donors (both Q and P-value < 0.001). Analysis of miRNAs from IL1Cregs associated genes including id2, id3, sox4, tgfr1, tgfr2, il2rb, and il3rg, from normal donors and AML patients demonstrated 34 miRNA from plasma samples and 14 miRNA segments from BM cell samples with a statistically significant difference between AML patients and normal donors (both Q and P-value <0.001). Among them, 4 miRNAs (hsa-miR-193b-3p, hsa-miR-1270, hsa-miR-210-3p, and hsa-miR-486-3p) were detected in both plasma and BM cell samples.

**Conclusion:** Our study enumerated IL1Cregs, then measured miRNAs from those IL1Cregs in AML samples for the first time. The results demonstrated the deficiency of IL1Creg and differential expression of miRNAs in patients with AML.

**Keywords:** regulatory innate lymphoid cells, IL1Cregs, flow cytometry, FCM, miRNAs, next generation sequence, acute myeloid leukemia, AML

Correspondence: Jifeng Yu; Zhongxing  
Jiang  
Department of Hematology, The First  
Affiliated Hospital of Zhengzhou  
University, 1 East Jianshe Road,  
Zhengzhou 450052, People's Republic of  
China  
Tel +86-371-6629 5132  
Email yujifengz@163.com;  
jiangzx@zzu.edu.cn

## Introduction

Innate lymphoid cell (ILC) is a kind of innate immune lymphocyte discovered in recent years. Following the traditional classification method of T cells, it has

been classified into ILC1, ILC2, and ILC3 subsets: group 1 comprises natural killer (NK) cells and ILC1s, group 2 comprises ILC2, and group 3 comprises ILC3.<sup>1</sup> For clarity, the term ILCs will only refer to ILC1, ILC2, and ILC3 not to conventional NK cells. ILCs have been described to have a lymphoid morphology but lack rearranged antigen receptors and can secrete various cytokines.<sup>2</sup> Although ILCs express CD127, the cell surface receptors of most hematopoietic lineages (lin) are absent.<sup>3</sup> ILC1s are described as CD45<sup>+</sup>Lin<sup>-</sup>CD127<sup>+</sup>CD161<sup>+</sup>CD117<sup>-</sup>CRTH2<sup>-</sup>,<sup>4</sup> ILC2s as CD45<sup>+</sup>Lin<sup>-</sup>CD127<sup>+</sup>CD161<sup>+</sup>CD117<sup>+</sup>CRTH2<sup>+</sup>,<sup>5</sup> ILC3s as CD45<sup>+</sup>Lin<sup>-</sup>CD127<sup>+</sup>CD161<sup>+</sup>CD117<sup>+</sup>CRTH2<sup>-</sup>,<sup>6,7</sup> and NK cells as CD45<sup>+</sup>Lin<sup>-</sup>CD161<sup>+</sup>CD94<sup>+</sup>CD56<sup>+</sup>.<sup>8</sup> Human ILCs have been identified in various tissues including peripheral blood<sup>9</sup> and cord blood.<sup>10</sup> Recently, a flow cytometric Optimized Multicolor Immunofluorescence Panel (OMIP) for characterization of human innate lymphoid cells from neonatal and peripheral blood has been suggested.<sup>11</sup>

In 2017, Wang et al discovered a sub-group of ILC cells with phenotype of Lin<sup>-</sup>CD45<sup>+</sup>CD127<sup>+</sup>IL-10<sup>+</sup> by flow cytometry in the intestines of mice and humans and named them regulatory innate lymphoid cells (ILCregs).<sup>12,13</sup> While the morphology of these cells has similar characteristics of lymphocytes, they do not express CD4 and FoxP3, and are therefore not Tregs.<sup>14</sup> These cells express ILC markers such as CD25 (IL-2Ralpha) and CD90 (Thy1), and over-express IL-2R gamma and SCA-1, but they do not express ILC1 markers (NK1.1, NKp46), ILC2 markers (ST2 and KLRG1), ILC3 markers (NKp46, CD4, and RORγt) and white blood cell markers.<sup>15,16</sup> Infusion of ILCregs can prevent congenital colitis, alleviate inflammatory bowel disease, save Rag1<sup>-/-</sup>IL10<sup>-/-</sup> mice, and amplified in DSS-stimulated models. Flow cytometry assay showed that the highest proportion of the cells in the lamina propria of small intestine was 13.1% and 15.7%, in mice and human, respectively. Immunofluorescence and immunohistochemistry methods were also used to verify the presence of these cells in human and mouse intestines.<sup>12,13</sup>

Since newly confirmed ILCregs may exist in both mouse and human intestines and bone marrow, the function and mechanism of ILCregs in patients of acute myeloid leukemia are still unclear and no relevant research reports have been reported yet. Therefore, based on the knowledge of ILCregs and AML, we hypothesize that ILCregs may be deficient in the patients of AML. In this study, we explore the expression levels of the

CD45<sup>+</sup>Lin<sup>-</sup>CD127<sup>+</sup>IL-10<sup>+</sup> ILCregs population and its CD45<sup>+</sup>Lin<sup>-</sup>CD127<sup>+</sup>IL-10<sup>-</sup> subset population, evaluating the clinical significance between the normal donors and patients with AML by using the 4 color monoclonal antibody combination detected by flow cytometry.

miRNAs are 19–22 nucleotides long non-coding RNAs which regulate the expression of genes by sequence-specific binding to mRNA to either promote or block its translation.<sup>17</sup> This is a powerful level of epigenetic control for gene expression that can influence the phenotype of a cell.<sup>18</sup> In recent years, regulation of miRNAs is extensively studied for their role in biological processes as well as in the development and progression of various human diseases, miRNAs hold huge potential for diagnostic and prognostic biomarkers as well as predictors of drug response.<sup>19</sup> Insights into the roles of miRNAs in development and disease, particularly in cancer have made miRNAs attractive tools and targets for novel therapeutic approaches.<sup>20</sup> Some studies have shown miRNAs play a role in blocking differentiation of leukemic cells and promoting their unchecked cell division.<sup>21,22</sup>

To further explore the importance of miRNA in patients of AML and the relationship of miRNA with the frequency of ILCregs, we used the next-generation sequence (NGS) method to measure the miRNA from both plasma and BM mono-nuclear cell samples. We compared the expression levels of miRNAs from AML patients to normal controls to find a statistically significant increase or decrease in expression levels of approximately 3280 miRNA sequences. Analysis of miRNAs in ILCreg-associated genes yields further information regarding statistically significant changes in expression.

## Methods

### Patients with Acute Myeloid Leukemia

The bone marrow samples were obtained from 42 patients newly diagnosed with acute myeloid leukemia prior to the induction chemotherapy were obtained following written informed consent from patients and/or their legal guardians at the First Affiliated Hospital of Zhengzhou University from September 2018 to August 2019. Study protocols were approved by Ethics Committee of The First Affiliated Hospital of Zhengzhou University (Zhengzhou, Henan, China) and based on the ethical principles for medical research involving human subjects of the Helsinki Declaration. Diagnosis was done according to the standard diagnosis with MICM criteria based on the 2016 WHO AML Classification<sup>23</sup>

and the 2017 NCCN Clinical Practice Guidelines.<sup>24</sup> All clinical data regarding patients are summarized in Table 1.

## Control Group

In the control group, 12 bone marrow samples from 6 male and 6 female normal donors with the age of 20–55 years old were obtained following written informed consent from healthy volunteers and tested side by side with some of the AML patients' samples.

## Regulatory Innate Lymphoid Cell Immunophenotyping by Multicolor Flow Cytometry

Freshly obtained samples up to 4 hrs from the collection were processed for flow cytometry assay. Bone marrow samples were collected and cultured in media containing 1:1000 BD GolgiPlug (BD Biosciences, USA) for 4 hr at 37°C. Cells were harvested and stained with surface markers CD45-FITC (BD Biosciences, USA, Cat # 555482), Lineage-APC (BD Biosciences, USA, Cat# 562722), CD127-PE-Cy7 (BD Biosciences, USA, Cat # 560822) for 15 mins. Red blood cells were lysed with lysing buffer (BD Biosciences, USA, Cat # 555899). Cells were then fixed and permeabilized by BD Cytofix/Cytoperm buffer set (BD Biosciences, USA, Cat# 555028) after surface marker staining, followed by application of anti-IL-10-PE (BD Biosciences, USA, Cat#554498) antibody (JES5-9D7) staining intracellular IL-10 before analysing through flow cytometry. The readout was done with Beckman Coulter Gallios 4-laser and 10-color model flow cytometer (Beckman Coulter, USA). Data analysis was done by using FlowJo software (BD Biosciences, USA).

## Gating Strategy

Single-cell populations with SSC vs FSC were selected and were further gated on the CD45<sup>+</sup>Lin<sup>-</sup> population.

**Table 1** Clinical Characteristics of AML Patients


Total Patients=42	Mean±SD	Range (%)
Age (year-old)	48	16-74
Gender Males (n, %)	20	47.62%
WBC (×10 <sup>9</sup> /L)	30.48±48.82	0.32–230.00
Hb (g/L)	82.01±16.65	39.00–123.00
Plate (×10 <sup>9</sup> /L)	45.17±42.93	5.00–210.00
Blast in BM	53.10±26.78	12.80–96.40
Complex karyotype (%)	8/42	19.1
Fusion Genes (%)	9/42	21.4
Gene mutations (%)	36/42	85.7

Furthermore, gating of CD127<sup>+</sup>IL-10<sup>+</sup> and CD127<sup>+</sup>IL-10<sup>-</sup> sub-population was done by different fluorescence-conjugated antibodies combinations. Regulatory innate lymphoid cells were identified as CD45<sup>+</sup>Lin<sup>-</sup>CD127<sup>+</sup>IL-10<sup>+</sup> population. At least a total of 1 million cells were collected for analysis of ILCCreg cells. A representative example of ILCCreg subpopulation is shown in Figure 1.

## Samples Preparation for RNA Extraction & miRNA Library Construction

Fresh bone marrow samples from 10 AML patients and 6 normal donors were obtained for miRNA testing. Plasma and mono-nuclear cells were collected with Ficoll-Hypaque centrifugation method. Mono-nuclear cells were collected and washed once with PBS. Plasma and mono-nuclear cell samples were put into RNA preparation tubes and stored at -80°C for batch processing. The RNA samples were centrifuged at 12,000×g for 5mins at 4°C. The supernatant was transferred to a new 2.0mL tube which was added 0.3mL of Chloroform/isoamyl alcohol (24:1) per 1.5mL of Trizol reagent. After the mix was centrifuged at 12,000×g for 10mins at 4°C, the aqueous phase was transferred to a new 1.5mL tube which added equal volume of supernatant of isopropyl alcohol. The mix was centrifuged at 12,000×g for 20mins at 4°C and then removed the supernatant. After washed with 1mL 75% ethanol, the RNA pellet was air-dried in the biosafety cabinet and then dissolved by add 25µL~100µL of DEPC-treated water. Subsequently, total RNA was qualified and quantified using a Nano Drop and Agilent 2100 Bioanalyzer (Thermo Fisher Scientific, MA, USA).

Library was prepared with 1µg total RNA for each sample. Total RNA was purified by electrophoretic separation on a 15% urea denaturing polyacrylamide gel electrophoresis (PAGE) gel and small RNA regions corresponding to the 18–30 nt bands in the marker lane (14–30 ssRNA Ladder Marker, TAKARA) were excised and recovered. Then, the 18–30 nt small RNAs were ligated to adenylated 3' adapters annealed to unique barcodes, followed by the ligation of 5' adapters. The adapter-ligated small RNAs were subsequently transcribed into cDNA by SuperScript II Reverse Transcriptase (Invitrogen, USA) and then several rounds of PCR amplification with PCR Primer Cocktail and PCR Mix were performed to enrich the cDNA fragments. The PCR products were selected by agarose gel electrophoresis with target fragments 110~130 bp, and then purified by QIAquick Gel Extraction Kit (QIAGEN, Valencia, CA). The yield and



**Figure 1** ILCregs from normal donors and AML patients.

**Notes:** The different expression patterns of ILCregs from normal donors (right panel) and AML patients (left panel) are shown in Figure 1. Single-cell populations with SSC vs FSC were selected and were further gated on the CD45+Lin- population. Furthermore, gating of CD127+IL-10+ and CD127+IL-10- sub-population was done by different fluorescence-conjugated antibodies' combinations. Regulatory innate lymphoid cells were identified as CD45+Lin-CD127+IL-10+ population. At least a total of 1 million cells were collected for analysis of ILCreg cells.

quality of the RNA samples was assessed using the Agilent Bioanalyzer prior to library construction using the Illumina TruSeq Small RNA Sample Prep protocol (Illumina; San Diego, California). Multiplexed samples of RNA that exceed quality control metrics (RIN > 6.0) were run on an Illumina NextSeq500 instrument at a targeted depth of 10 million reads per sample. After filtering and trimming of index and adapter sequences, whole-genome alignment of the miR FASTQ reads was performed using the Homo sapiens/hg21 reference genome in the SHRiMPS aligner included in the miRNAs analysis application available in BaseSpace (Illumina), as well as the sRNA Toolbox application suite.

## Statistical Analysis

Clinical data, laboratory findings from the patients were collected in the spreadsheet constructed in Microsoft Excel software for Windows 10 (Microsoft). Data were analysed using IBM SPSS Statistics software version 17.0 (IBM, USA). For intra-group comparison, Student's *t*-test was used. Significance level was  $p < 0.05$ .

The analysis of the RNA-seq data was performed following the pipeline available from the limma packages in the Bioconductor project. Log2counts per million (logCPM) transformation was applied before normalization and linear model fitting. Empirical Bayes moderation

was carried out to obtain robust estimates of gene-wise variability and the final *p* values from the linear model with appropriately designed contrasts were adjusted by the Benjamin-Hochberg procedure for a targeted false discover rate of 0.05. Volcano plots and box plots were used to graphically examine the differences between groups.

## Results

### Patient Cohort: Clinical Characteristics

A total of 42 newly diagnosed AML patients, including 20 males and 22 females, were included in this study. The clinical manifestations of these patients were summarized (Table 1). The median age at diagnosis was 48 years (range 16–72 years). 12 normal donors including 6 males and 6 females with the age of 20–55 years old were used for controls.

### ILCregs Detection by Flow Cytometry

The percentage of individual subpopulations of ILCregs in the bone marrow from patients with acute myeloid leukemia was determined. Using the 4 color monoclonal antibody combination, we measured the ILCregs defined as CD45<sup>+</sup>Lin<sup>-</sup>CD127<sup>+</sup>IL-10<sup>+</sup> in the BM from both normal donor and AML patients. The different expression patterns



of ILCregs from normal donors (right panel) and AML patients (left panel) are shown in Figure 1.

Comparison of ILCreg numbers in the bone marrow between normal donors and patients with acute myeloid leukemia are shown as box plots below (Figure 2). The average frequencies of ILCregs were  $0.8924 \pm 1.3791\%$  and  $0.2434 \pm 0.5344\%$  for normal donors and patients with AML, respectively. In comparison with the normal donors, the frequency of ILCregs cells in AML patients was significantly lower ( $P < 0.01$ ). Interestingly, the average frequency of  $CD45^+Lin^-CD127^+IL-10^-$  subset was  $4.0869 \pm 6.7701\%$  and  $0.2769 \pm 0.2526\%$  for normal donors and AML patients, respectively. In comparison with the normal donors, the frequency of  $CD45^+Lin^-CD127^+IL-10^-$  cells in AML patients was also significantly lower ( $P < 0.01$ ).


## miRNAs Sequenced by NGS

Furthermore, we sequenced miRNAs from both plasma and BM cells samples from 9 patients with newly diagnosed AML and 6 normal donors by using the next-generation sequence method. We obtained sequence for 3280 miRNAs and averaged the amount of each miRNA from 9 AML patients and 6 controls. Levels of miRNA expression were compared with a *t*-test while the P-values and Q values were calculated for each miRNA. Different miRNA patterns were found from both BM cell samples and plasma samples between the AML patients and normal donors. There were 62 miRNAs up-regulated and 120 miRNAs down-regulated from the AML BM cell samples and 103 miRNAs up-regulated and 273

miRNAs down-regulated from the plasma samples from the box plots shown below (Figure 3). The pathway-related expression patterns were also analyzed and shown below (Figure 4).


In our analysis of plasma samples from 9 AML patients with 6 controls, there were 376 miRNAs with expression levels that showed a statistically significant difference (both Q value and P-value  $< 0.001$ ). Meanwhile, in the BM cell samples from AML patients and normal controls, there were 182 miRNAs with expression levels that showed a statistically significant difference (both Q value and P-value  $< 0.001$ ). This was also revealed in a heatmap and cluster analysis of differentially expressed miRNAs of AML patients and normal donors (data not shown). The fold change analysis shows the miRNA differential signature between the AML patients BM cell samples and plasma samples. The color pattern describes the level of expression: red shows miRNAs with down-regulated expression, black shows miRNAs with unchanged expression among the samples, and green shows those miRNAs with up-regulated expression. The size of the branches represents the Euclidian distance of the average expression of the miRNAs.

In Figure 5–7, miRNA of BM cell samples from 9 AML patients versus 6 control is shown as below. Log<sub>2</sub> Fold Change is shown on the x-axis and  $-\log_{10}$  p-value is shown on the y-axis. Points in bold font indicate miRNAs with statistically significant log fold change and adjusted p-value, green color indicated down-regulation and red color indicated up-regulation. Figure 5 shows the volcano plot for the BM cell samples of patients versus controls with



**Figure 2** Boxplots showing positive percentages of ILCregs and CD127+IL-10- cells with a statistically significant difference between control and AML patient bone marrow samples.

**Notes:** In comparison with the normal donors, the frequency of  $CD45^+Lin^-CD127^+IL-10^-$  ILCregs and  $CD45^+Lin^-CD127^+IL-10^-$  cells in AML patients was significantly lower ( $P < 0.01$ ).



**Figure 3** Differential miRNA expression in AML patients BM cells versus plasma.

**Notes:** Up-regulated and down-regulated genes from the AML BM cell and plasma samples comparing to the normal donors as the box plots shown below (Figure 3 box plots).

1053 miRNAs having statistically significant change in expression, 56 down-regulated and 24 up-regulated. Similarly, Figure 6 shows the difference of plasma samples from AML patients and normal donors with 135 statistically significant changes in miRNA expression, including 16 down-regulated and 18 up-regulated. Figure 7 shows the volcano plot for the difference of plasma samples and BM cell samples from AML patients with 454 statistically significant changes in miRNA expression, including 162 down-regulated and 9 up-regulated. Clearly, the subset analysis identified a greater number of miRNAs with statistically significant changes in expression in the AML patients.

We then performed subset analysis of our data on the miRNAs with the ILCreg associated *id2*, *id3*, *sox4*, *tgfbr1*, *tgfbr2*, *il2rb* and *il2rg* genes. Results showed 14 miRNAs with a statistically significant difference in expression level (both Q value and P-value < 0.001) from BM cell samples between AML patients and normal donors (Table 2). Meanwhile, there were 34 miRNAs with a statistically significant difference (both Q value and P-value < 0.001). Figure 9 shows greatest increase from BM cell samples when comparing AML patients to controls.


34 miRNA from plasma samples and 14 from BM cell samples that showed a statistically significant difference (both Q value and P-value < 0.001) between AML patients and normal donors are shown in Table 3. The # indicated miRNAs were segments found in both plasma and BM cell

samples. miRNAs that have not previously been associated with AML are indicated with asterisks. Figure 8 showed greatest increase and decrease from plasma samples when comparing AML patients to controls.

Furthermore, we identified each individual ILCregs-associated genes miRNAs from both plasma and BM cell samples that showed a statistical significance (Table 4). Among these ILCregs associated genes include *id2*, *id3*, *sox4*, *tgfbr1*, *tgfbr2*, *il2rb* and *il2rg* which all comprise miRNAs that had been found in either plasma or BM cell samples or in both with the exception of the *il2rb* gene. For *id2* gene, only *hsa-miR-210-3p* miRNA showed a statistically significant difference (both Q value and P-value < 0.001) from BM cell samples between AML patients and normal donors. Consequently, we identified 13 miRNAs related to *id2* gene that showed a statistically significant difference (both Q value and P-value < 0.001) from plasma samples between AML patients and normal donors.

For *id3* gene-related miRNAs, only *hsa-miR-3194-3p* miRNA that showed a statistically significant difference (both Q value and P-value < 0.001) from BM cell samples between AML patients and normal donors. Meanwhile, only *hsa-miR-6751-5p* and *hsa-miR-4722-5p* showed a statistically significant difference (both Q value and P-value < 0.001) from plasma samples between AML patients and normal donors.

For *il2rg* gene-related miRNAs, only *hsa-miR-23a-5p* and *hsa-miR-486-3p* miRNAs showed a statistically




**Figure 4** Kegg pathway-related expression patterns were analyzed and shown in AML patients.

significant difference (both Q value and P-value < 0.001) from BM cell samples between AML patients and normal donors. Meanwhile, only hsa-miR-491-5p and hsa-miR-486-3p that showed a statistically significant difference (both Q value and P-value < 0.001) from plasma samples between AML patients and normal donors.


For sox4 gene-related miRNAs, four miRNAs hsa-miR-193b-3p, hsa-miR-128-2-5p, hsa-miR-1270, hsa-

miR-210-5p miRNAs showed a statistically significant difference (both Q value and P-value < 0.001) from BM cell samples between AML patients and normal donors. Meanwhile, there were 13 miRNAs that showed a statistically significant difference (both Q value and P-value < 0.001) from plasma samples between AML patients and normal donors.


For tgfbr1 gene related miRNAs, four miRNAs hsa-miR-24-3p, hsa-miR-5195-3p, hsa-miR-584-3p, hsa-miR-6722-3p



**Figure 5** Volcano plot: BM cell samples from AML patients vs normal donors.  
**Notes:** BM cell samples from AML patients vs normal donors. Log<sub>2</sub> Fold Change is shown on the x-axis and  $-\log_{10}$  p-value is shown on the y-axis. Points in bold font indicate miRNAs with statistically significant log fold change and adjusted p-value, green color indicated down-regulation and red color indicated up-regulation.



**Figure 7** Volcano plot: Plasma samples vs BM cell samples from AML patients.  
**Notes:** Difference of plasma samples and BM cell samples from AML patients. Log<sub>2</sub> Fold Change is shown on the x-axis and  $-\log_{10}$  p-value is shown on the y-axis. Points in bold font indicate miRNAs with statistically significant log fold change and adjusted p-value, green color indicated down-regulation and red color indicated up-regulation.



**Figure 6** Volcano plot: Plasma samples from AML patients vs normal donors.  
**Notes:** Log<sub>2</sub> Fold Change is shown on the x-axis and  $-\log_{10}$  p-value is shown on the y-axis. Points in bold font indicate miRNAs with statistically significant log fold change and adjusted p-value, green color indicated down-regulation and red color indicated up-regulation.

miRNAs showed a statistically significant difference (both Q value and P-value < 0.001) from BM cell samples between AML patients and normal donors. Meanwhile, five hsa-miR-101-3p, hsa-miR-1275, hsa-miR-3200-3p, hsa-miR-3187-3p, hsa-miR-4722-5p miRNAs that showed a statistically significant difference (both Q value and P-value < 0.001) from plasma samples between AML patients and normal donors.

For *tgfrb2* gene-related miRNAs, hsa-miR-3151-5p and hsa-miR-34c-5p miRNAs showed a statistically significant difference (both Q value and P-value < 0.001) from BM cell samples between AML patients and normal donors. Meanwhile, hsa-miR-432-5p, hsa-miR-342-5p, hsa-miR-370-3p, hsa-miR-3074-5p miRNAs that showed

a statistically significant difference (both Q value and P-value < 0.001) from plasma samples between AML patients and normal donors. No *IL2rb*-related miRNAs were found in our test.

## Discussion

Acute myeloid leukemia (AML) is a clonal neoplastic disease with the proliferation of leukemic myeloid hematopoietic precursor cells and impaired production of normal hematopoiesis.<sup>25–27</sup> Research results have confirmed that AML patients have immune system impairment. Regulatory T cells are the most important part of the immune system, which have numerically and functionally deficiencies in AML patients.<sup>28–31</sup> These defects are reported to suppress the proliferation and function of T helper cells.<sup>32–34</sup> ILC is a kind of innate immune lymphocyte discovered in recent years.<sup>35</sup> ILCregs cells, one of its subpopulation, had been discovered in both human and mouse intestines.<sup>12,13</sup> The newly discovered ILCregs share the characteristics of innate immune cells and Tregs while playing an important role in the pathogenesis of intestinal inflammatory diseases and other related immune regulation. The function and mechanism of ILCregs in AML are still unclear. Studies have shown that ILCreg cell populations continuously express *Tgfr1*, *Tgfr2*, *Il2rb*, and *Il2rg*, suggesting that ILCregs could respond to TGF-beta and IL-2 signals. IL-10 and TGF-beta 1 are also highly expressed in human ILCregs.<sup>12,13</sup> Although



**Table 2** Comparison of BM Cell Sample miRNA Expression Levels Between AML Patients and Normal Donors


Gene ID	AML2_BMC Expression	ND_BMC Expression	AML2_BMC Average Read Count	ND_BMC Average Read Count	Log2 (ND_BMC/ AML2_BMC)	P value (AML2_BMC- Vs-ND_BMC)	Q value (AML2_BMC- Vs-ND_BMC)
hsa-miR-24-3p	17,801.222	38,942.5	23,090.778	60,768	1.11232	0	0
hsa-miR-370-3p	8	17.167	8	17.333	0.8318	1.40E-04	5.10E-04
hsa-miR-3074-5p	5310.222	10,925.667	5901	12,285.5	0.77425	0	0
hsa-miR-342-5p	1158.111	2391	1189.667	2472.167	0.77154	0	0
hsa-miR-3200-3p	7.667	16	8	16	0.71632	0.00133	0.00412
hsa-miR-128-1-5p	1.333	2.833	1.444	2.833	0.68831	0.19214	0.33203
hsa-miR-3154	53.778	76	54	76.5	0.21882	0.01943	0.04804
hsa-miR-1275	916.111	1287.333	937.556	1318.167	0.20788	0	0
hsa-miR-1271-5p	233.333	311.167	236.889	317.333	0.13811	0.00231	0.00682
hsa-miR-1307-3p	1155.111	1505.667	1195.667	1555.333	0.09573	0	1.00E-05
hsa-miR-18a-3p	251	318.333	261.556	330.167	0.0524	0.23142	0.38122
hsa-miR-212-3p	50.333	61	51.333	61.833	-0.01519	0.87927	0.93717
hsa-miR-339-3p	487.111	583.167	499.556	593.333	-0.03548	0.27004	0.4214
hsa-miR-3187-3p	77.333	89.833	78.889	91.5	-0.06973	0.39184	0.55886
hsa-miR-138-5p	24.889	38.5	36.222	40	-0.14055	0.24797	0.39494
hsa-miR-432-5p	19.889	21.333	20	21.5	-0.17934	0.27658	0.43017
hsa-miR-149-5p	22.667	26.5	25.667	27.333	-0.19291	0.18593	0.32473
hsa-miR-378e	4.778	5	5	5	-0.28368	0.39806	0.56686
hsa-miR-25-5p	159.556	149	164.889	151.833	-0.40268	0	0
hsa-miR-324-3p	676.333	504.833	693.778	515.333	-0.71265	0	0
hsa-miR-324-5p	549.556	356.667	561.667	361.167	-0.92073	0	0
hsa-miR-1270	18.222	9.833	18.667	10	-1.18414	0	0
hsa-miR-193b-3p	630.778	248.833	651.333	251.167	-1.65843	0	0
hsa-miR-210-3p	111.222	34.667	113.111	35.5	-1.95553	0	0
hsa-miR-34c-5p	59.333	18.333	128.889	19	-3.04574	0	0

ILCregs had been studied in the intestines of mice and humans, the frequency and expression patterns in patients with AML had not been explored yet. By using the combination of surface markers of CD45, Lin, CD127, and IL-10 for intracellular staining followed by flow cytometry analysis, we were able to detect ILCregs in BM from both normal donors and patients with AML. The frequency of the CD45<sup>+</sup>Lin<sup>-</sup>CD127<sup>+</sup>IL-10<sup>+</sup> ILCregs subset populations was significantly decreased in the patients with AML comparing to normal donors. Our results have shown for the first time the enumeration and deficiency of ILCregs in patients with AML.


CD127 had been used by many researchers as the major specific surface marker to distinguish Tregs from normal T cells. Research results have shown that CD25<sup>+</sup>CD127<sup>hi</sup> T cells play a role in maintaining non-inflammatory environment by deviating immune responses away from pro-inflammatory Th1 response and encourage an anti-inflammatory Th2-type response.<sup>28</sup> A variety of studies have also shown the

deficiencies of the regulatory T cells in AML patients.<sup>32,34,36,37</sup> CD45<sup>+</sup>Lin<sup>-</sup>CD127<sup>+</sup>IL-10<sup>-</sup> cell population had not been explored in patients with AML and its function remains unknown. Our results show that in comparison with the normal donors, the frequency of CD45<sup>+</sup>Lin<sup>-</sup>CD127<sup>+</sup>IL-10<sup>-</sup> cells in AML patients was significantly decreased for BM cells population (P<0.01).

ILCreg cell populations have continuously expressed Tgfb1, Tgfb2, Il2rb, and Il2rg,<sup>12,13</sup> but the miRNAs related to these genes have not yet been explored. We chose to sequence all miRNAs that were obtained in both plasma and BM cell samples from patients with newly diagnosed AML and controls. We quantified expression of 3280 miRNAs from each patient and control and performed statistical analyses to determine which miRNAs were increased or decreased in AML patients. Interestingly, our results demonstrate that several ILCreg miRNAs have been found from both plasma and BM cell samples with either up-regulated or down-regulated variants. This suggested that these miRNAs may play an important role in the development and function of ILCregs. Future studies in ILCreg regulation may give



**Figure 8** Box plots showing greatest increase or greatest decrease of miRNAs from plasma samples when comparing AML to controls. **Notes:** Box plots showing greatest increase (upper panel) or greatest decrease (lower panel) of miRNAs from AML patient plasma samples when comparing to controls. Y-axis showed average measurement of relative expression level of miRNAs from AML patients and normal donors.



**Figure 9** Box plots showing greatest increase from AML patient BM cell samples when comparing to controls. **Notes:** Box plots showing greatest increase from AML patient BM cell samples when comparing to controls. Y-axis showed average measurement of relative expression level of miRNAs from AML patients and normal donors.

additional information about the role of ILCregs in the pathogenesis of AML.

Some of the studies have demonstrated that certain miRNAs play an important role in certain malignant diseases such as AML with NPM1+/FLT3+ mutations.<sup>38</sup> One of the miRNAs associated with NPM1+/FLT3+ AML, miR-10a-5p showed the most statistically significant adjusted p-value in the yield as well as the highest fold change. This miRNA had been described in patients with NPM1 mutations. It's noted that high

expression levels are associated with good response to induction chemotherapy.<sup>39</sup> More recent studies demonstrate a role for miR-10a/b in regulating the proliferation and differentiation of HL-60 leukemic cells in vitro.<sup>40</sup> Other literature demonstrate that high expression of miR-338 is associated with poor prognosis in acute myeloid leukemia undergoing chemotherapy.<sup>41</sup> However, our results show that certain ILCreg genes targeted miRNAs had been found in both plasma and BM cells samples either up-regulated or down-regulated.

**Table 3** A Total of 34 miRNA from Plasma Samples and 14 from BM Cell Samples That Showed a Statistically Significant Difference

Gene ID	Log2 FC(ND_XJ/AML2_XJ)	ILCregs Related Genes	Gene ID	Log2 (ND_BMC/AML2_BMC)	ILCregs Related Genes
hsa-miR-1271-5p*	9.34877	d2	hsa-miR-486-3p#	1.53652	il2rg
hsa-miR-3200-3p	8.85439	tgfbr1	hsa-miR-23a-5p	1.44084	il2rg
hsa-miR-193b-3p*#	8.55935	sox4	hsa-miR-584-3p	1.36018	tgfbr1
hsa-miR-432-5p	8.36976	tgfbr2	hsa-miR-24-3p*	1.11232	tgfbr1
hsa-miR-3154*	6.94874	id2, sox4	hsa-miR-3194-3p	1.07301	id3
hsa-miR-18a-3p*	6.14571	id2	hsa-miR-3151-5p	-1.11375	tgfbr2
hsa-miR-1298-5p	5.75325	id2	hsa-miR-1270*#	-1.18414	sox4
hsa-miR-324-3p	4.59904	sox4	hsa-miR-210-5p*	-1.24805	sox4
hsa-miR-1270*#	3.3365	sox4	hsa-miR-193b-3p*#	-1.65843	sox4
hsa-miR-370-3p*	2.59359	id2, tgfbr2	hsa-miR-210-3p*#	-1.95553	id2
hsa-miR-3074-5p*	2.1621	tgfbr2	hsa-miR-34c-5p*	-3.04574	tgfbr2
hsa-miR-138-5p*	2.0674	id2	hsa-miR-6722-3p	-4.50607	tgfbr1
hsa-miR-212-3p*#	1.79612	sox4	hsa-miR-128-2-5p	-4.72552	sox4
hsa-miR-210-3p*	-1.53215	id2	hsa-miR-5195-3p	-5.28368	tgfbr1
hsa-miR-1275	-1.6481	id2, sox4, tgfbr1			
hsa-miR-3187-3p*	-1.68836	tgfbr1			
hsa-miR-491-5p*	-3.06053	il2rg			
hsa-miR-342-5p*	-3.0985	tgfbr2			
hsa-miR-486-3p#	-3.17156	il2rg			
hsa-miR-378d*	-3.24956	sox4			
hsa-miR-101-3p*	-3.45966	tgfbr1			
hsa-miR-1268b	-4.92447	sox4			
hsa-miR-4722-5p	-4.961	id3, tgfbr1			
hsa-miR-1268a	-5.92447	sox4			
hsa-miR-3909*	-5.961	id2			
hsa-miR-181a-2-3p*	-5.97892	sox4			
hsa-miR-339-3p*	-7.23575	sox4			
hsa-miR-193b-5p*	-7.41166	id2			
hsa-miR-1908-5p	-7.59423	id2			
hsa-miR-146b-3p*	-8.42697	id2			
hsa-miR-128-1-5p	-9.25745	sox4			
hsa-miR-6751-5p	-9.86823	id3			
hsa-miR-25-5p	-9.9698	sox4			
hsa-miR-4747-5p	-13.08938	id2			

**Notes:** \*miRNAs that have not previously been associated with AML are indicated with asterisks. #miRNAs that have been found in both plasma and BM cell samples.

These miRNAs may play an important role in the ILCregs's immuno-modulation.

MiR-181 family has been shown by others to be consistently increased in AML patients.<sup>42-44</sup> There are numerous possible target mRNAs for the miR-181 family leading to its interest as a therapeutic target. However, our results did not show the significant difference between AML patients and normal donors from both plasma and BM cell samples. In addition, there are no reported associations between miRNA-181 family and ILCreg associated genes. Another report has shown that miR-199b is consistently decreased in AML,<sup>45</sup> but our results did not show the expression difference between AML patients and normal donors. There was another report of increased expression of miR-223-3p in AML patients;<sup>46</sup>

however, our results did not show significant difference between AML patients and normal donors from both plasma and BM cell samples.

Subset analysis of our data on the miRNAs with the ILCregs associated id2, id3, sox4, tgfbr1, tgfbr2, il2rb and il2rg genes showed that there were 34 miRNA from plasma samples and 14 miRNAs from BM cells samples with expression levels that showed a statistically significant difference between AML patients and normal donors. The first identification of each individual ILCreg miRNA segments with a significant difference from AML patients will be helpful for further research on the ILCregs in AML molecular pathogenesis. However, our data are only a beginning to further studies for large sampling and

**Table 4** ILCregs Associated Genes Related miRNA Expression Patterns from Plasma and BM Cell Samples

Genes	Samples	Gene Associated miRNAs
SOX	BM cell samples	hsa-miR-193b-3p; hsa-miR-128-2-5p; hsa-miR-1270; hsa-miR-210-5p
	Plasma samples	hsa-miR-324-3p; hsa-miR-1268b; hsa-miR-128-1-5p; hsa-miR-193b-3p; hsa-miR-181a-2-3p; hsa-miR-1275; hsa-miR-3154; hsa-miR-1268a; hsa-miR-1270; hsa-miR-339-3p; hsa-miR-378d; hsa-miR-25-5p; hsa-miR-212-3p
ID2	BM cell samples	hsa-miR-210-3p
	Plasma samples	hsa-miR-193b-5p; hsa-miR-3909; hsa-miR-138-5p; hsa-miR-4747-5p; hsa-miR-1275; hsa-miR-1298-5p; hsa-miR-1908-5p; hsa-miR-3154; hsa-miR-18a-3p; hsa-miR-1271-5p; hsa-miR-370-3p; hsa-miR-146b-3p; hsa-miR-210-3p
ID3	BM cell samples	hsa-miR-3194-3p
	Plasma samples	hsa-miR-6751-5p; hsa-miR-4722-5p
IL2RG	BM cell samples	hsa-miR-23a-5p; hsa-miR-486-3p
	Plasma samples	hsa-miR-491-5p; hsa-miR-486-3p
TGFBRI	BM cell samples	hsa-miR-24-3p; hsa-miR-5195-3p; hsa-miR-584-3p; hsa-miR-6722-3p
	Plasma samples	hsa-miR-101-3p; hsa-miR-1275; hsa-miR-3200-3p; hsa-miR-3187-3p; hsa-miR-4722-5p
TGFB2	BM cell samples	hsa-miR-3151-5p; hsa-miR-34c-5p
	Plasma samples	hsa-miR-432-5p; hsa-miR-342-5p; hsa-miR-370-3p; hsa-miR-3074-5p

observation in similar AML patients but at different time points, such as at diagnosis, after complete remission, etc.

## Conclusion

Our study enumerates ILCregs, then measures miRNAs from those ILCregs in AML samples. The ILCregs and miRNA deficiencies are detected in patients with AML. Our results demonstrate the deficiency of ILCreg and differential expression of miRNAs in patients with AML.

## Ethics Approval and Consent to Participate

Informed consent was obtained from all patients and the protocol was approved by the Ethics Committee of the First Affiliated Hospital of Zhengzhou University.

## Consent for Publication

All authors contributed to data analysis, drafting or revising the article, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

## Data Sharing Statement

Data and material will be available upon corresponding author approval. All data sets analyzed for this study are included in the manuscript and the additional files.

## Author Contributions

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

## Funding

This study was funded by the Key Scientific Research Project of Henan Provincial Education Department (20A320062 and 19A320046); National Natural Science Foundation of China (U1804192); and Special Talents Project Fund of the First Affiliated Hospital of Zhengzhou University, Zhengzhou, China. The funding bodies did not participate in study design, in data collection, analysis, and interpretation, and in writing the manuscript.

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Spits H, Artis D, Colonna M, et al. Innate lymphoid cells -a proposal for uniform nomenclature. *Nat Rev Immunol.* 2013;13(2):145-149. doi:10.1038/nri3365

2. Spits H, Cupedo T. Innate lymphoid cells: emerging insights in development, lineage relationships, and function. *Annu Rev Immunol.* 2012;30(1):647–675. doi:10.1146/annurev-immunol-020711-075053
3. Walker JA, Barlow JL, McKenzie ANJ. Innate lymphoid cells -how did we miss them? *Nat Rev Immunol.* 2013;13(2):75–87. doi:10.1038/nri3349
4. Bernink JH, Peters CP, Munneke M, et al. Human type 1 innate lymphoid cells accumulate in inflamed mucosal tissues. *Nat Immunol.* 2013;14(3):221–229. doi:10.1038/ni.2534
5. Mjösberg JM, Trifari S, Crellin NK, et al. Human IL-25- and IL-33-responsive type 2 innate lymphoid cells are defined by expression of CRTH2 and CD161. *Nat Immunol.* 2011;12:1055–1062. doi:10.1038/ni.2104
6. Hoorweg K, Peters CP, Cornelissen F, et al. Functional differences between human NKp44(-) and NKp44(+) RORC(+) innate lymphoid cells. *Front Immunol.* 2012;3:72. doi:10.3389/fimmu.2012.00072
7. Tang Q, Ahn Y-O, Southern P, Blazar BR, Miller JS, Verneris MR. Development of IL-22-producing NK lineage cells from umbilical cord blood hematopoietic stem cells in the absence of secondary lymphoid tissue. *Blood.* 2011;117(15):4052–4055. doi:10.1182/blood-2010-09-303081
8. Borrego F, Masilamani M, Kabat J, Sanni TB, Coligan JE. The cell biology of the human natural killer cell CD94/NKG2A inhibitory receptor. *Mol Immunol.* 2005;42(4):485–488. doi:10.1016/j.molimm.2004.07.031
9. Montaldo E, Vacca P, Vitale C, et al. Human innate lymphoid cells. *Immunol Lett.* 2016;179(Supplement C):2–8. doi:10.1016/j.imlet.2016.01.007
10. Lim AI, Li Y, Lopez-Lastra S, et al. Systemic human ILC precursors provide a substrate for tissue ILC differentiation. *Cell.* 2017;168(6):1086–100.e10. doi:10.1016/j.cell.2017.02.021
11. Bianca Bennstein S, Riccarda Manser A, Weinhold S, Scherschlich N, Uhrberg M. OMIP-055: characterization of human innate lymphoid cells from neonatal and peripheral blood. *Cytometry A.* 2019. doi:10.1002/cyto.a.23741
12. Wang S, Xia P, Chen Y, et al. Regulatory innate lymphoid cells control innate intestinal inflammation. *Cell.* 2017;171(1):201–216. doi:10.1016/j.cell.2017.07.027
13. Zeng B, Shi S, Liu J, Xing F. Commentary: regulatory innate lymphoid cells control innate intestinal inflammation. *Front Immunol.* 2018;9:1522. doi:10.3389/fimmu.2018.01522
14. Sakaguchi S, Yamaguchi T, Nomura T, Ono M. Regulatory T cells and immune tolerance. *Cell.* 2008;133(5):775–787. doi:10.1016/j.cell.2008.05.009
15. Gury-BenAri M, Thaïss CA, Serafini N, et al. The spectrum and regulatory landscape of intestinal innate lymphoid cells are shaped by the microbiome. *Cell.* 2016;166(5):1231–1246.e13. doi:10.1016/j.cell.2016.07.043.
16. Robinette ML, Fuchs A, Cortez VS, et al. Transcriptional programs define molecular characteristics of innate lymphoid cell classes and subsets. *Nat Immunol.* 2015;16(3):306–317. doi:10.1038/ni.3094.
17. Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell.* 2004;116(2):281–297. doi:10.1016/S0092-8674(04)00045-5
18. Baltimore D, Boldin MP, O'connell RM, Rao DS, Taganov KD. MicroRNAs: new regulators of immune cell development and function. *Nat Immunol.* 2008;9(8):839–845. doi:10.1038/ni.f.209
19. Vishnoi A, Rani S. MiRNA biogenesis and regulation of diseases: an overview. *Methods Mol Biol.* 2017;1509:1–10.
20. Rupaimoole R, Slack FJ. MicroRNA therapeutics: towards a new era for the management of cancer and other diseases. *Nat Rev Drug Discov.* 2017;16(3):203–222. doi:10.1038/nrd.2016.246.
21. Wallace JA, O'Connell RM. MicroRNAs and acute myeloid leukemia: therapeutic implications and emerging concepts. *Blood.* 2017;130(11):1290–1301. doi:10.1182/blood-2016-10-697698
22. Trino S, Lamorte D, Caivano A, et al. MicroRNAs as new biomarkers for diagnosis and prognosis, and as potential therapeutic targets in acute myeloid leukemia. *Int J Mol Sci.* 2018;19(2):460. doi:10.3390/ijms19020460
23. Arber DA, Orazi A, Hasserjian R, et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood.* 2016;127(20):2391–2405. doi:10.1182/blood-2016-03-643544.
24. O'Donnell MR, Tallman MS, Abboud CN, et al. Acute myeloid leukemia, version 3.2017, NCCN clinical practice guidelines in oncology. *J Natl Compr Canc Netw.* 2017;15(7):926–957. doi:10.6004/jcnccn.2017.0116
25. Steensma DP, Bejar R, Jaiswal S, et al. Clonal hematopoiesis of indeterminate potential and its distinction from myelodysplastic syndromes. *Blood.* 2015;126(1):9–16. doi:10.1182/blood-2015-03-631747
26. Bowman RL, Busque L, Levine RL. Clonal hematopoiesis and evolution to hematopoietic malignancies. *Cell Stem Cell.* 2018;22(2):157–170. doi:10.1016/j.stem.2018.01.011
27. Young AL, Challen GA, Birmann BM, Druley TE. Clonal haematopoiesis harbouring AML-associated mutations is ubiquitous in healthy adults. *Nat Commun.* 2016;7:12484. doi:10.1038/ncomms12484
28. Wang M, Zhang C, Tian T, et al. Increased regulatory T cells in peripheral blood of acute myeloid leukemia patients rely on tumor necrosis factor (TNF)- $\alpha$ -TNF receptor-2 pathway. *Front Immunol.* 2018;9:1274. doi:10.3389/fimmu.2018.01274.
29. Sander FE, Nilsson M, Rydström A, et al. Role of regulatory T cells in acute myeloid leukemia patients undergoing relapse-preventive immunotherapy. *Cancer Immunol Immunother.* 2017;66(11):1473–1484. doi:10.1007/s00262-017-2040-9.
30. Yang W, Xu Y. Clinical significance of treg cell frequency in acute myeloid leukemia. *Int J Hematol.* 2013;98(5):558–562. doi:10.1007/s12185-013-1436-3.
31. Han Y, Dong Y, Yang Q, et al. Acute myeloid leukemia cells express ICOS ligand to promote the expansion of regulatory T cells. *Front Immunol.* 2018;9:2227. doi:10.3389/fimmu.2018.02227.
32. Williams P, Basu S, Garcia-Manero G, et al. The distribution of T-cell subsets and the expression of immune checkpoint receptors and ligands in patients with newly diagnosed and relapsed acute myeloid leukemia. *Cancer.* 2018;125(9):1470–1481. doi:10.1002/cncr.31896
33. Knaus HA, Berglund S, Hackl H, et al. Signatures of CD8 $\beta$  T cell dysfunction in AML patients and their reversibility with response to chemotherapy. *JCI Insight.* 2018;3:pil: 120974. doi:10.1172/jci.insight.120974
34. Ustun C, Miller JS, Munn DH, Weisdorf DJ, Blazar BR. Regulatory T cells in acute myelogenous leukemia: is it time for immunomodulation? *Blood.* 2011;118(19):5084–5095. doi:10.1182/blood-2011-07-365817
35. Bjorklund AK, Forkel M, Picelli S, et al. The heterogeneity of human CD127+ innate lymphoid cells revealed by single-cell RNA sequencing. *Nat Immunol.* 2016;17(4):451–460. doi:10.1038/ni.3368
36. Szczepanski MJ, Szajnik M, Czysowska M, et al. Increased frequency and suppression by regulatory T cells in patients with acute myelogenous leukemia. *Clin Cancer Res.* 2009;15(10):3325–3332. doi:10.1158/1078-0432.CCR-08-3010
37. Shenghui Z, Yixiang H, Jianbo W, et al. Elevated frequencies of CD4(+) CD25(+) CD127lo regulatory T cells is associated to poor prognosis in patients with acute myeloid leukemia. *Int J Cancer.* 2011;129(6):1373–1381. doi:10.1002/ijc.25791
38. Pandita A, Ramadas P, Poudel A, et al. Differential expression of miRNAs in acute myeloid leukemia quantified by Nextgen sequencing of whole blood samples. *PLoS One.* 2019;14(3):e0213078. doi:10.1371/journal.pone.0213078
39. Havelange V, Ranganathan P, Geyer S, et al. Implications of the miR-10 family in chemotherapy response of NPM1-mutated AML. *Blood.* 2014;123(15):2412–2415. doi:10.1182/blood-2013-10-532374.



40. Bi L, Sun L, Jin Z, Zhang S, Shen Z. MicroRNA-10a/b are regulators of myeloid differentiation and acute myeloid leukemia. *Oncol Lett.* 2018;15(4):5611–5619. doi:10.3892/ol.2018.8050.
41. Fu L, Qi J, Gao X, et al. High expression of miR-338 is associated with poor prognosis in acute myeloid leukemia undergoing chemotherapy. *J Cell Physiol.* 2019:1–9. doi:10.1002/jcp.28676
42. Marcucci G, Mrozek K, Radmacher MD, Garzon R, Bloomfield CD. The prognostic and functional role of microRNAs in acute myeloid leukemia. *Blood.* 2011;117(4):1121–1129. doi:10.1182/blood-2010-09-191312.
43. Weng H, Lal K, Yang FF, Chen J. The pathological role and prognostic impact of miR-181 in acute myeloid leukemia. *Cancer Genetics.* 2015;208(5):225–229. doi:10.1016/j.cancergen.2014.12.006.
44. Su R, Lin H, Zhang X, et al. MiR-181 family: regulators of myeloid differentiation and acute myeloid leukemia as well as potential therapeutic targets. *Oncogene.* 2015;34(25):3226–3239. doi:10.1038/onc.2014.274.
45. Favreau AJ, McGlauffin RE, Duarte CW, Sathyanarayana P. miR-199b, a novel tumor suppressor miRNA in acute myeloid leukemia with prognostic implications. *Experimental Hematology & Oncology.* 2015;5(1):4. doi:10.1186/s40164-016-0033-6
46. Trissal MC, DeMoya RA, Schmidt AP, Link DC. MicroRNA-223 regulates granulopoiesis but is not required for HSC maintenance in mice. *PLoS One.* 2015;10(3):e0119304. doi:10.1371/journal.pone.0119304.

## Cancer Management and Research

Dovepress

### Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient.

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/cancer-management-and-research-journal>