

# *Candida auris*: From Multidrug Resistance to Pan-Resistant Strains

This article was published in the following Dove Press journal:  
*Infection and Drug Resistance*

Mulneh Ademe<sup>1</sup>  
Friehiwot Girma<sup>2</sup>

<sup>1</sup>Department of Microbiology, Immunology and Parasitology, College of Health Sciences, Addis Ababa University, Addis Ababa, Ethiopia; <sup>2</sup>Department of Pediatrics and Child Health Nursing, School of Health Sciences, College of Medicine and Health Sciences, Bahir Dar University, Bahir Dar, Ethiopia

**Abstract:** *Candida auris* is an emerging multidrug-resistant fungus that is rapidly spreading worldwide. Currently, *C. auris* cases have been reported globally from >30 countries. Most reported infections involve critically ill patients in hospitals, mainly in intensive care unit settings. Infection with *C. auris* is associated with high mortality rates, and it is often resistant to multiple classes of antifungal drugs. Despite the rapid global spread, it is difficult to predict the actual burden of the infection as the standard laboratory methods fail to correctly identify the fungi. Longer stays in healthcare facilities, use of tracheostomies and percutaneous endoscopic gastrostomy tubes, ventilators in clinical care units and mobile equipment in healthcare settings are shown as major risk factors of *C. auris* infection. Due to its propensity to cause outbreaks and its antifungal resistance, *C. auris* poses a risk for patients in healthcare facilities. The emergence of pan-resistant *C. auris* strains in some areas is an alarming signal for the disease with limited treatment options, high mortality rates, and the ability of the pathogen to spread easily in healthcare settings. In this regard, susceptibility testing on clinical isolates, mainly for patients treated with echinocandins, is needed. Increasing awareness about *C. auris* infection and advancing the diagnostic methods are also essential for early detection and control of the deadly fungal infection.

**Keywords:** *Candida auris*, multidrug resistance, pandrug resistance, risk factors, challenges

## Background

*C. auris* is one of the few species of the genus *Candida* which cause candidiasis in humans. It is one of the emerging fungus that can cause invasive infections. As it was detected in the external ear canal of the patient, it was named as *Candida auris*. Auris is the Latin word for ear. Currently, *C. auris* has emerged globally as a multidrug-resistant nosocomial pathogen and it is considered a major threat to healthcare settings. Worldwide reports of *C. auris* have considerably increased within a decade.<sup>1,2</sup> The aim of this review is, therefore, to describe the current status of *C. auris*, mainly focusing on the current challenges in clinical and laboratory diagnosis, emergence of pan-drug resistance and possible ways of preventing and controlling the spread of the infection. For this purpose, a systematic search of bibliographical databases was done using the search engines: Google Scholar, Google search, Scopus and PubMed Central. The following key words (phrases) were used in the search engine including but not limited to *C. auris*, mechanisms of drug resistance, risk factors, diagnosis, treatment, prevention and control. Peer-reviewed research articles, reviews and short communications by international organizations were included. The search was restricted to English language and duplicates were removed.

Correspondence: Mulneh Ademe  
Department of Microbiology, Immunology  
and Parasitology, College of Health  
Sciences, Addis Ababa University, P.O.  
Box 9086, Addis Ababa, Ethiopia  
Email mulneh.ademe@aau.edu.et



## Risk Factors of *C. auris* Infection

Usually, longer stays in certain types of post-acute care facilities such as intensive care units (ICUs) is considered as a major risk factor of *C. auris* infection. For instance, the first and only *C. auris* candidemia case in Kuwait was a patient in the ICU with chronic renal failure.<sup>17</sup> In India, more than two-thirds of ICUs (19 out of 27 ICUs) were found to have *C. auris* isolates.<sup>18</sup> In Spain, an average length of stay of 25 days in healthcare institutions was associated with developing *C. auris* infection.<sup>19</sup> This might be attributed to the nature of disease transmission which mostly involves exposure to contaminated facilities in healthcare institutions.<sup>7</sup> *C. auris* colonizes both biotic (skin and other body sites) and abiotic surfaces. Notably, its ability to adhere to surfaces and plastic materials (eg, catheters), biofilm formation, and salt tolerance facilitate the acquisition of the fungus in healthcare settings such as ICUs.<sup>20</sup> The majority of *C. auris* infected patients have had a recent exposure to an indwelling device or have undergone some invasive procedures.<sup>21</sup> Infections have been observed several days to weeks after hospitalization in susceptible patients, suggesting exogenous sources of infection.

In agreement with this, patients with tracheostomies and percutaneous endoscopic gastrostomy (PEG) tubes are more likely to acquire *C. auris* infection.<sup>22</sup> Ventilators in clinical care units were reported to spread *C. auris* infection. In this regard, the prevalence of *C. auris* in nursing home units with ventilator beds was 7.7% compared to regular nursing homes without ventilator (0.7%).<sup>23</sup> Mobile equipment's in healthcare settings has also been implicated in the transmission of *C. auris* infection.<sup>23</sup> Furthermore, the incidence of *C. auris* increases among patients with primary or acquired altered immune response, therapeutic management of broad-spectrum antibiotics, transplantation, patients with different comorbidities, and other conditions requiring immunosuppressive agents.<sup>24,25</sup>

*C. auris* has been isolated from patients of both sexes and of all age groups. Though the reported *C. auris* isolates were mostly isolated from males,<sup>13</sup> no reason has been provided yet for the sexual differences in terms of frequency of *C. auris* infections.

## Current Challenges of *C. auris* Infection

### Non-Specific Clinical Presentations

*C. auris* can cause severe invasive infections. However, clinical conditions associated with *C. auris* are non-specific

and it is often difficult to differentiate between other types of systemic infections.<sup>10</sup> Clinical conditions associated with most of the reported *C. auris* cases include bloodstream infections, urinary tract infection, otitis, surgical wound infections, skin abscesses, myocarditis, meningitis, bone infections and wound infections.<sup>18,26</sup> According to the CDC report,<sup>8</sup> fever and chills that do not improve after antibiotic treatment for a suspected bacterial infection are usually considered among the common symptoms of invasive *C. auris* infection. However, none of the aforementioned clinical conditions provide definitive diagnosis, and laboratory investigation should be considered for confirmation.

## Misidentification by Conventional Diagnostic Tests

Early detection of *C. auris* infections has been shown to be beneficial as earlier initiation of appropriate antifungal therapy saved many lives.<sup>27</sup> However, identification of *C. auris* remained challenging because most phenotypic methods misidentify *C. auris*. Until the sequence analysis correctly identified isolates as *C. auris*, they were initially misidentified as other *Candida* species.<sup>5</sup> So far, culture and microscopy characteristics have been used to differentiate different *Candida* species. In view of this, *C. auris* cells are ellipsoid in shape, an ovoid to elongate budding yeast, which seldom forms rudimentary pseudohyphae and typically appears as pink. *C. auris* also grows as yeast and forms smooth, shiny, whitish-gray, viscous colonies on growth media and it has high tolerance for salinity and heat.<sup>10,28</sup> However, these features may not provide definitive diagnosis for *C. auris* as there are other *Candida* species, such as *C. haemulonii*, with the most similar phenotypic characteristics.<sup>29</sup>

It was confirmed by molecular techniques that conventional diagnostic tests using biochemical typing often misidentify *C. auris*.<sup>30</sup> *C. auris* isolates have been misidentified as a range of other *Candida* species with phenotypic and biochemical methods (Table 1), including API 20C, Vitek 2 (bioMérieux), Phoenix (BD), and MicroScan.<sup>28,31,32</sup> As recommended by CDC (Table 1), correct identification of *C. auris* can be done using accepted diagnostic methods such as matrix-assisted laser desorption ionization–time of flight (MALDI-TOF) platforms.<sup>33</sup> Moreover, molecular identification techniques like polymerase chain reaction, sequencing, and amplified fragment length polymorphism fingerprinting will detect *C. auris* and differentiate it

**Table 1** Identification of *Candida auris* by Different Diagnostic Methods

Tests Which Misidentify <i>C. auris</i> as Other <i>Candida</i> Species <sup>1,28,30-32,55</sup>	Tests Which Correctly Identify <i>C. auris</i> <sup>33,34</sup>
API 20CAUX	Bruker Biotyper MALDI-TOF (RUO libraries (v 2014[5627] and CA system library (v claim4))
API Candida	bioMerieux VITEK MS MALDI-TOF (ROU library (Saramis v 4.14))
Phoenix (BD Diagnostics)	VITEK 2 YST (Software v 8.01)
Vitek	Polymerase chain reaction (PCR)
MicroScan (Beckman Coulter)	Amplified fragment length polymorphism fingerprinting
Vitek MS (bioMerieux)	

from closely related species such as *C. haemulonii*, *C. duobushaemulonii*, or *C. lusitanae*.<sup>34</sup>

It has to be noted that isolations from non-sterile body sites such as lungs, urinary tract, skin and soft tissue, and genital apparatus may represent colonization rather than infections.<sup>26,35</sup> Hence, the presence of signs and symptoms of *C. auris* infections should be considered to differentiate simple colonization from infection.<sup>36</sup> Certainly, it is important to identify *C. auris* even from a non-sterile body site because colonization poses the risk of transmission, which requires implementation of infection control precautions.<sup>10</sup>

## Multidrug Resistance and Emergence of Pan-Resistant Strains

*C. auris* is characterized by high rate of antifungal resistance with reduced susceptibility to azoles, polyenes, and echinocandins.<sup>37</sup> In vitro, more than 90% of *C. auris* isolates have shown resistance to fluconazole.<sup>38</sup> *C. auris* resistance to voriconazole and amphotericin B was shown to be 3–73% and 13–35%, respectively.<sup>21,27</sup> In the US, more than 99% of the *C. auris* isolates were shown to be resistant to fluconazole, nearly two-thirds were resistant to amphotericin B, and roughly 4% were resistant to echinocandins.<sup>39</sup> A systematic review and meta-analysis from Sekyere et al also reported that most of *C. auris* isolates are resistant to fluconazole (44.29%), followed

by amphotericin B (15.46%), voriconazole (12.67%), caspofungin (3.48%) and flucytosine (1.95%).<sup>13</sup> Thus, higher resistance to fluconazole in a *Candida* non-albicans species has become one of the distinguishing characteristics indicative of a potential *C. auris* infection.<sup>40</sup>

Indeed, echinocandins are the first-line therapy for *C. auris* infection. As synergistic interactions have a better efficacy, a combination therapy of an echinocandin and liposomal amphotericin B is prescribed in cases of unresponsiveness to echinocandins.<sup>41–43</sup> However, most recently in US, three cases of pan-resistant *C. auris* with resistance to all three classes of commonly prescribed antifungal drugs have been reported.<sup>39</sup> In the same report, it was mentioned that a total of 801 patients with *C. auris* were identified in New York as of June 28, 2019. Among these patients, three of them developed resistance to all antifungal medications, including echinocandins.<sup>39</sup> In fact, echinocandin-resistant *C. auris* isolates have been previously described<sup>44</sup>; however, the isolates were susceptible to azoles. Chowdhary et al also reported 14 (4%) pan-azole-resistant *C. auris* cases<sup>9</sup>; however, this data was limited to only micafungin and anidulafungin. Although the overall numbers of pan-resistant cases reported so far are few, it is an alarming signal for the disease with limited treatment options, high mortality rates, and the ability of the pathogen to spread easily in healthcare settings.

Data regarding the molecular mechanisms of resistance of *C. auris* to antifungal agents are scarce, and the precise mechanism of resistance in isolates is not well known. As shown by some studies (Table 2), *C. auris* escapes from the microbicidal effect of all the classes of anti-fungal agents through different mechanisms, including but not

**Table 2** Mechanism of Antifungal Resistance by *C. auris*

Antifungal	Resistance Mechanism	References
Polyenes	Mutations, 5 SNPs in different genome loci	[28,56]
Azoles	Point mutations in the lanosterol 14 $\alpha$ -demethylase (ERG11) gene	[21,24,57]
	ERG 11 up-regulation due to mutation in Upc2 TF	
	Up-regulation of efflux pump genes	
Nucleoside analogs	Amino acid substitution (F211) in FURI	[58]
Echinocandin	Mutation in FKS1/2 gene	[10]



limited to mutations in ERG3 and ERG11 genes, up-regulation of efflux pump genes and single-nucleotide polymorphisms (SNPs) in different genome loci.<sup>13</sup> The South Asia clade is shown to exhibit increased antifungal resistance compared to other clades of *C. auris*.<sup>9</sup> Likewise, the reported U.S. pan-resistant cases emerged in New York where the South Asia clade (clade 1) predominates.<sup>45</sup>

## Increased Risk of Mortality

Invasive infections of *C. auris* are fatal unless early detection and treatment is initiated. The ability of *C. auris* to develop resistance to commonly used antifungal agents is responsible for its high rates of mortality.<sup>37</sup> In fact, the reported mortality rates due to *C. auris* vary. The 30-day mortality rate of *C. auris* in Colombia was 35.2%.<sup>46</sup> Mortality rates in Asia, Far East, and the United States is estimated to be more than 50% for those with invasive infections.<sup>5,47,48</sup> In Spain, a thirty-day mortality rate was reported to be 41.4%.<sup>19</sup> In general, the crude in-hospital mortality rate for *C. auris* candidemia is estimated to range from 30% to 72%.<sup>10</sup>

## Awareness About the Infection

*C. auris* is a newly emerging fungus and has subsequently been associated with invasive infections and outbreaks in healthcare settings worldwide. Lack of awareness about this drug-resistant *Candida* could lead to unnoticed transmission and outbreaks in healthcare settings.<sup>40</sup> In this regard, the healthcare workers should be aware of this fatal invasive infection and need to adapt the laboratory testing strategies and implement enhanced control measures early enough to prevent healthcare outbreaks.<sup>49</sup> So far, there is no concrete evidence on the current state of knowledge of the public and healthcare workers regarding *C. auris* infection. CDC itself claimed that CDC fungal experts had never heard and received a report describing a *Candida* infection resistant to all antifungal medications before 2016. It was after hearing the news that CDC sounded the alarm in the US about *C. auris*.<sup>50</sup> Currently, only few countries worldwide (about 30 countries) have reported *C. auris*,<sup>8</sup> suggesting that the infection is either undetected or unreported in the majority of countries. However, there is no much information whether the disease does not actually exist in those countries where *C. auris* is not reported yet. In this regard, the awareness of health workers about the whole picture of *C. auris* infection including the diagnostic challenges could not be underestimated as most of the routine diagnostic methods miss the fungus and the infection may continue to be unreported.

## Prevention and Control

*C. auris* strains are characterized by their ability to form a biofilm structure on biotic and abiotic surfaces.<sup>20</sup> Patients are often indefinitely colonized by *C. auris* primarily on skin, nares and other body sites. Indeed, there is no currently known decolonization strategy and colonization may persist for months leading to invasive infection and transmission to others.<sup>51</sup> *C. auris* also persists in the environment for months, and persistent environmental contamination, contaminated medical equipment and other fomites are believed to play a role in nosocomial *C. auris* transmission.<sup>52</sup> In view of this, rigorous attention to environmental cleaning is important for preventing transmission within a healthcare facility. Indeed, common disinfectants such as quaternary ammonia compounds do not work.<sup>8</sup> But, surface disinfectants such as chlorine and hydrogen peroxide were shown to have good efficacy against *C. auris*.<sup>53,54</sup>

*C. auris* was made nationally notifiable at the 2018 Council for State and Territorial Epidemiologists (CSTE) Annual Conference.<sup>8</sup> Hence, detected cases have to be reported to local or state health departments. Patients with confirmed infections or suspected to have *C. auris* infections should be kept in separate wards under strict contact precautions.<sup>8</sup> Patients or healthcare workers coming in close contact with infected persons should also be placed under strict contact precautions until they are proven to be negative for confirmatory diagnostic tests. Moreover, it is recommended to thoroughly disinfect the wards of patients found to be colonized or infected with *C. auris*. Cleaning and disinfection of shared medical equipment is encouraged.<sup>6</sup>

## Conclusion

Infections by *C. auris* are progressively emerging in hospitals and ICU settings. *C. auris* with high mortality rates, multi-drug resistance, environmental resilience, difficulty in microbiological identification and horizontal transmission has become an issue in clinical practice. Most reported infections of *C. auris* involved critically ill patients. The emergence of pan resistance is an alarming signal and controlling the spread of the resistance strain is needed. Because of the potential for the development of resistance, patients on antifungal treatment for *C. auris* should be monitored closely for clinical improvement. Susceptibility testing should also be conducted, especially in patients treated with echinocandins. Moreover, increasing awareness about the infection and

advancing the diagnostic methods is essential to control *C. auris* infection.

## Abbreviations

AFLP, amplified fragment length polymorphism; ICU, intensive care unit; MALDI-TOF, matrix-assisted laser desorption/ionization–time of flight mass spectrometry; RUO, research use only.

## Acknowledgments

First and foremost, we would like to thank God for giving us the courage and strength. Our sincere gratitude also goes to our families and friends for their support.

## Author Contributions

Both authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

## Disclosure

The authors declare no competing interests.

## References

- Spivak ES, Hanson KE. *Candida auris*: an emerging fungal pathogen. *J Clin Microbiol*. 2018;56(2):1–10.
- Satoh K, Makimura K, Hasumi Y, Nishiyama Y, Uchida K, Yamaguchi H. *Candida auris* sp. nov., a novel ascomycetous yeast isolated from the external ear canal of an inpatient in a Japanese hospital. *Microbiol Immunol*. 2009;53:41–44. doi:10.1111/j.1348-0421.2008.00083.x
- Chowdhary A, Sharma C, Meis JF. *Candida auris*: a rapidly emerging cause of hospital-acquired multidrug-resistant fungal infections globally. *PLoS Pathog*. 2017;13:e1006290. doi:10.1371/journal.ppat.1006290
- Chowdhary A, Sharma C, Duggal S, et al. New clonal strain of *Candida auris*, Delhi, India. *Emerg Infect Dis*. 2013;19(10):1670–1673.
- Lee WG, Shin JH, Uh Y, et al. First three reported cases of nosocomial fungemia caused by *Candida auris*. *J Clin Microbiol*. 2011;49(9):3139–3142. doi:10.1128/JCM.00319-11
- Schelenz S, Hagen F, Rhodes JL, et al. First hospital outbreak of the globally emerging *Candida auris* in a European hospital. *Antimicrob Resist Infect Control*. 2016;5(35):1–7. doi:10.1186/s13756-016-0132-5
- Tsay S, Welsh RM, Adams EH, et al. Notes from the field: ongoing transmission of *Candida auris* in health care facilities—United States, June 2016–May 2017. *MMWR Morb Mortal Wkly Rep*. 2017;66:514–515. doi:10.15585/mmwr.mm6619a7
- CDC. *Candida auris*: a drug-resistant germ that spreads in healthcare facilities. Available from: <https://www.cdc.gov/fungal/candida-auris/tracking-c-auris.html>. Accessed January 15, 2020.
- Chowdhary A, Prakash A, Sharma C, et al. A multicentre study of antifungal susceptibility patterns among 350 *Candida auris* isolates (2009–17) in India: role of the ERG11 and FKS1 genes in azole and echinocandin resistance. *J Antimicrob Chemother*. 2018;73:891–899. doi:10.1093/jac/dkx480
- Cortegiani A, Misseri G, Fasciana T, Giammanco A, Giarratano A, Chowdhary A. Epidemiology, clinical characteristics, resistance, and treatment of infections by *Candida auris*. *J Intensive Care*. 2018;6(69):1–13. doi:10.1186/s40560-018-0342-4
- Sharma C, Kumar N, Pandey R, Meis JF, Chowdhary A. Whole genome sequencing of emerging multidrug resistant *Candida auris* isolates in India demonstrates low genetic variation. *New Microbes New Infect*. 2016;13:77–82. doi:10.1016/j.nmni.2016.07.003
- Lockhart SR, Berkow EL, Chow N, Welsh RM. *Candida auris* for the clinical microbiology laboratory: not your grandfather's *Candida* species. *Clin Microbiol News*. 2017a;39:99–103. doi:10.1016/j.clinmicnews.2017.06.003
- Sekyere JO. *Candida auris*: A systematic review and meta-analysis of current updates on an emerging multidrug-resistant pathogen. *Microbiolgyopen*. 2018;7(4):1–29.
- Chow NA, de Groot T, Badali H, Abastabar M, Chiller TM, Meis JF. Potential Fifth Clade of *Candida auris*, Iran, 2018. *Emerg Infect Dis*. 2019;25(9):1780–1781. doi:10.3201/eid2509.190686
- Abastabar M, Haghani I, Ahangarkani F, Rezai MS, Armaki MT, Roodgari, et al. *Candida auris* otomycosis in Iran and review of recent literature. *Mycoses*. 2019;62:101–105.
- Chow NA, Gade L, Tsay SV, Forsberg K, Greenko JA, Southwick KL. Multiple introductions and subsequent transmission of multidrug-resistant *Candida auris* in the USA: a molecular epidemiological survey. *Lancet Infect Dis*. 2018;18(12):1377–1384. doi:10.1016/S1473-3099(18)30597-8
- Emara M, Ahmad S, Khan Z, et al. *Candida auris* candidemia in Kuwait, 2014. *Emerg Infect Dis*. 2015;21:1091–1092. doi:10.3201/eid2106.150270
- Rudramurthy SM, Chakrabarti A, Paul RA, et al. *Candida auris* candidaemia in Indian ICUs: analysis of risk factors. *J Antimicrob Chemother*. 2017;72:1794–1801. doi:10.1093/jac/dkx034
- Ruiz-Gaitán A, Moret AM, Tásias-Pitarch M, Alexandre-López AI, Martínez-Morel H, Calabuig E. An outbreak due to *Candida auris* with prolonged colonisation and candidaemia in a tertiary care European hospital. *Mycoses*. 2018;61(7):498–505. doi:10.1111/myc.12781
- Larkin E, Hager C, Chandra J, et al. The emerging pathogen *Candida auris*: growth phenotype, virulence factors, activity of antifungals, and effect of SCY-078, a novel glucan synthesis inhibitor, on growth morphology and biofilm formation. *Antimicrob Agents Chemother*. 2017;61(5):1–13. doi:10.1128/AAC.02396-16
- Lockhart SR, Etienne KA, Vallabhaneni S, Farooqi J, Chowdhary A, Govender NP. Simultaneous emergence of multidrug-resistant *Candida auris* on 3 continents confirmed by whole-genome sequencing and epidemiological analyses. *Clin Infect Dis*. 2017b;64(2):134–140. doi:10.1093/cid/ciw691
- Forsberg K, Woodworth K, Walters M, et al. Brendan Jackson1, Tom Chiller *Candida auris*: the recent emergence of a multidrug-resistant fungal pathogen. *Med Mycol*. 2019;57:1–12. doi:10.1093/mmy/myy054
- Adams E, Quinn M, Tsay S, et al. *Candida auris* in healthcare facilities, New York, USA, 2013–2017. *Emerg Infect Dis*. 2018;24(10):1816–1824. doi:10.3201/eid2410.180649
- Ben-Ami R, Berman J, Novikov A. Multidrug-resistant *Candida haemulonii* and *C. auris*, Tel Aviv, Israel. *Emerg Infect Dis*. 2017;23(1):195–203. doi:10.3201/eid2302.161486
- Azar MM, Turbett SE, Fishman JA, Pierce VM. Donor-derived transmission of *Candida auris* during lung transplantation. *Clin Infect Dis*. 2017;65:1040–1042. doi:10.1093/cid/cix460
- Chowdhary A, Voss A, Meis JF. Multidrug-resistant *Candida auris*: “new kid on the block” in hospital-associated infections? *J Hosp Infect*. 2016;94:209–212. doi:10.1016/j.jhin.2016.08.004
- Chowdhary A, Anil Kumar V, Sharma C, et al. Multidrug-resistant endemic clonal strain of *Candida auris* in India. *Eur J Clin Microbiol Infect Dis*. 2014;33:919–926. doi:10.1007/s10096-013-2027-1

28. Kathuria S, Singh PK, Sharma C, et al. Multidrug-resistant *Candida auris* misidentified as *Candida haemulonii*: characterization by matrix-assisted laser desorption ionization-time of flight mass spectrometry and DNA sequencing and its antifungal susceptibility profile variability by Vitek 2, CLSI broth microdilution, and Etest method. *J Clin Microbiol*. 2015;53:1823–1830. doi:10.1128/JCM.00367-15
29. Borman AM, Szekely A, Johnson EM. Comparative pathogenicity of United Kingdom isolates of the emerging pathogen *Candida auris* and other key pathogenic *Candida* species. *mSphere*. 2016;1(4):1–8. doi:10.1128/mSphere.00189-16
30. Jeffery-Smith A, Taori SK, Schelenz S, et al. *Candida auris*: a review of the literature. *Clin Microbiol Rev*. 2018;31:1–18.
31. Wattal C, Oberoi JK, Goel N, Raveendran R, Khanna S. Matrix-assisted laser desorption ionization time of flight mass spectrometry (MALDI-TOF MS) for rapid identification of micro-organisms in the routine clinical microbiology laboratory. *Eur J Clin Microbiol Infect Dis*. 2017;36:807–812. doi:10.1007/s10096-016-2864-9
32. Kim TH, Kweon OJ, Kim HR, Lee MK. Identification of uncommon *Candida* species using commercial identification system. *J Microbiol Biotechnol*. 2016;26:2206–2213. doi:10.4014/jmb.1609.09012
33. CDC. Algorithm to identify *Candida auris* based on phenotypic laboratory method and initial species identification. Available from: <https://www.cdc.gov/fungal/.../candidiasis/pdf/Testing-algorithm-by-Method-temp.pdf>. Accessed June 13, 2019.
34. Kordalewska M, Zhao Y, Lockhart SR, Chowdhary A, Berrio I, Perlin DS. Rapid and accurate molecular identification of the emerging multidrug-resistant pathogen *Candida auris*. *J Clin Microbiol*. 2017;55:2445–2452. doi:10.1128/JCM.00630-17
35. Kumar D, Banerjee T, Pratap CB, Tilak R. Itraconazole-resistant *Candida auris* with phospholipase, proteinase and hemolysin activity from a case of vulvovaginitis. *J Infect Dev Ctries*. 2015;9:435–437. doi:10.3855/jidc.4582
36. Pappas PG, Kauffman CA, Andes DR, et al. Clinical practice guideline for the management of candidiasis: 2016 update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2016;62:e1–50. doi:10.1093/cid/civ1194
37. Navalkele BD, Revankar S, Chandrasekar P. *Candida auris*: a worrisome, globally emerging pathogen. *Expert Rev Anti Infect Ther*. 2017;15:819–827. doi:10.1080/14787210.2017.1364992
38. Alfouzan W, Dhar R, Albarrag A, Al-Abdely H. The emerging pathogen *Candida auris*: a focus on the Middle-Eastern countries. *J Infect Public Heal*. 2019;12(4):451–459. doi:10.1016/j.jiph.2019.03.009
39. Ostrowsky B, Greenko J, Adams E, et al. *Candida auris* isolates resistant to three classes of antifungal medications—New York, 2019. A CDC report, Available from: <https://www.cdc.gov/mmwr/volumes/69/wr/mm6901a2.htm>. Accessed January 13, 2020.
40. ECDC (The European Centre for Disease Prevention and Control). Rapid risk assessment: *candida auris* in healthcare settings – europe. Available from: <https://www.ecdc.europa.eu/en/publications-data/rapid-risk-assessment-candida-auris-healthcare-settings-europe>. Accessed January 9, 2020.
41. Guarner J. Buruli ulcer. Review of a neglected skin mycobacterial disease. *J Clin Microbiol*. 2018;56(4):1–8.
42. Singhala T, Kumara A, Boradea P, Shaha S, Soman R. Successful treatment of *C. auris* shunt infection with intraventricular caspofungin. *Med Mycol Case Rep*. 2018;22:35–37. doi:10.1016/j.mmc.2018.08.005
43. Fakhim H, Vaezi A, Dannaoui E, et al. Comparative virulence of *Candida auris* with *Candida haemulonii*, *Candida glabrata* and *Candida albicans* in a murine model. *Mycoses*. 2018;61:377–382. doi:10.1111/myc.12754
44. Biagi MJ, Wiederhold NP, Gibas C, et al. Development of high-level echinocandin resistance in a patient with recurrent *Candida auris* candidemia secondary to chronic candiduria. *Open Forum Infect Dis*. 2019;6:1–5. doi:10.1093/ofid/ofz262
45. Leach L, Zhu Y, Chaturvedi S. Development and validation of a real-time PCR assay for rapid detection of *Candida auris* from surveillance samples. *J Clin Microbiol*. 2018;56(2):1–7.
46. Morales-Lopez SE, Parra-Giraldo CM, Ceballos-Garzon A, et al. Invasive infections with multidrug-resistant yeast *Candida auris*, Colombia. *Emerg Infect Dis*. 2017;23:162–164. doi:10.3201/eid2301.161497
47. Vallabhaneni S, Kallen A, Tsay S, et al. Investigation of the first seven reported cases of *Candida auris*, a globally emerging invasive, multidrug-resistant fungus—United States, May 2013–August 2016. *Am J Transplant*. 2016;17(1):296–299. doi:10.1111/ajt.14121
48. Sarma S, Kumar N, Sharma S, et al. Candidemia caused by amphotericin B and fluconazole resistant *Candida auris*. *Indian J Med Microbiol*. 2013;31:90–91. doi:10.4103/0255-0857.108746
49. Public Health Ontario (Ontario Agency for Health Protection and Promotion), Provincial Infectious Diseases Advisory Committee. *Interim Guide for Infection Prevention and Control of Candida Auris*. Toronto, ON: Queen’s Printer for Ontario; 2019:1–16.
50. CDC. Antibiotic resistance threats report. Available from: [www.cdc.gov/DrugResistance/Biggest-Threats.html](http://www.cdc.gov/DrugResistance/Biggest-Threats.html). Accessed August 1, 2019.
51. Caceres DH, Forsberg K, Welsh RM, et al. *Candida auris*: a review of recommendations for detection and control in healthcare settings. *J Fungi*. 2019;5(11):1–11. doi:10.3390/jof5040111
52. Shackleton J, Schelenz S, Rochon M, Hall A, Ryan L, Cervera-Jackson R. The impact of environmental decontamination in a *Candida auris* outbreak. *J Hosp Infect*. 2016;94(1):S88–9.
53. Abdolrasouli A, Armstrong-James D, Ryan L, Schelenz S. In vitro efficacy of disinfectants utilised for skin decolonisation and environmental decontamination during a hospital outbreak with *Candida auris*. *Mycoses*. 2017;60:758–763. doi:10.1111/myc.12699
54. Cadnum JL, Shaikh AA, Piedrahita CT, et al. Effectiveness of disinfectants against *Candida auris* and other *Candida* species. *Infect Control Hosp Epidemiol*. 2017;38:1240–1243. doi:10.1017/ice.2017.162
55. Lamoth F, Kontoyiannis DP. The *Candida auris* alert: facts and perspectives. *J Infect Dis*. 2018;217:516–520. doi:10.1093/infdis/jix597
56. Escandón P, Chow NA, Caceres DH, et al. Molecular epidemiology of *Candida auris* in colombia reveals a highly related, countrywide colonization with regional patterns in amphotericin B resistance. *Clin Infect Dis*. 2019;68:15–21. doi:10.1093/cid/ciy411
57. Prakash A, Sharma C, Singh A, Kumar Singh P, Kumar A, Hagen F. Evidence of genotypic diversity among *Candida auris* Isolates by multilocus sequence typing, matrix-assisted laser desorption ionization time-of-flight mass spectrometry and amplified fragment length polymorphism. *Clin Microbiol Infect*. 2016;22(3):277.e1-277.e9. doi:10.1016/j.cmi.2015.10.022
58. Rhodes J, Abdolrasouli A, Farrer RA, et al. Genomic epidemiology of the UK outbreak of the emerging human fungal pathogen *Candida auris*. *Emerg Microbes Infect*. 2018;7(43):1–12. doi:10.1038/s41426-017-0002-0

## Infection and Drug Resistance

Dovepress

### Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>