

Screening of potential key ferroptosis-related genes in Chronic Obstructive Pulmonary Disease

Yumeng Cao ^{1,*}, Huaqin Pan ^{2,*}, Yanwei Yang ³, Jingrun Zhou ¹, Guqin Zhang ¹

¹Department of Respiratory and Critical Care Medicine, Zhongnan Hospital of Wuhan University, Wuhan, Hubei, 430071, People's Republic of China; ²Transplantation Intensive Care Unit, Transplant Center of Wuhan University, Hubei Key Laboratory of Medical Technology on Transplantation, Wuhan, Hubei, 430071, People's Republic of China; ³Department of Critical Care Medicine, Zhongnan Hospital of Wuhan University, Wuhan, 430071, People's Republic of China

*These authors contributed equally to this work

Correspondence: Guqin Zhang, Department of Respiratory and Critical Care Medicine, Zhongnan Hospital of Wuhan University, Wuhan, Hubei, 430071, People's Republic of China, Email zgqxiaoxiao@126.com

Purpose: Ferroptosis plays essential roles in the development of COPD. We aim to identify the potential ferroptosis-related genes of COPD through bioinformatics analysis.

Methods: The RNA expression profile dataset GSE148004 was obtained from the GEO database. The ferroptosis-related genes were obtained from the FerrDb database. The potential differentially expressed ferroptosis-related genes of COPD were screened by R software. Then, protein-protein interactions (PPI), correlation analysis, gene-ontology (GO) enrichment analysis, and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis were applied for the differentially expressed ferroptosis-related genes. Finally, hub gene-microRNA(miRNA), hug gene-transcription factor interaction networks were constructed by miRTarBase v8.0 and JASPAR respectively, and hub gene drugs were predicted by the Enrichr database.

Results: A total of 41 differentially expressed ferroptosis-related genes (22 up-regulated genes and 19 down-regulated genes) were identified between 7 COPD patients and 9 healthy controls. The PPI results demonstrated that these ferroptosis-related genes interacted with each other. The GO and KEGG enrichment analyses of differentially expressed ferroptosis-related genes indicated several enriched terms related to ferroptosis, central carbon metabolism in cancer, and the HIF-1 signaling pathway. The crucial miRNAs and drugs associated with the top genes were identified.

Conclusion: We identified 41 potential ferroptosis-related genes in COPD through bioinformatics analysis. HIF1A, PPARG, and KRAS may affect the development of COPD by regulating ferroptosis. These results may expand our understanding of COPD and might be useful in the treatment of COPD.

Keywords: ferroptosis, COPD, bioinformatics analysis, gene expression omnibus dataset

Introduction

Chronic obstructive pulmonary disease (COPD) is a common respiratory system disease that often results in distal bronchiectasis, accompanied by tracheal wall degradation, with a high incidence rate and high mortality characteristics.¹⁻³ Characterized by persistent respiratory symptoms and progressive airflow blockage.³ Cigarette smoke (CS) exposure can cause an elevation in reactive oxygen species (ROS), leading to oxidative stress. This can further disrupt iron metabolism, resulting in inflammation, cellular aging, and cell death.⁴⁻⁸ Currently, COPD has become the third leading cause of death in the world. Due to the aging population and high smoking rates, the global burden of COPD is increasing. By 2030, the overall prevalence and mortality rate of COPD will increase by 8%.^{9,10} At present, there is no effective targeted drug for clinical use, so there is an urgent need to explore promising drug targets.

Ferroptosis, first proposed by Brent R. Stockwell of Columbia University in 2012, is a programmed cell death induced by lipid peroxidation through an iron dependent pathway with unique morphological and biological characteristics.^{11,12} Ferroptosis is characterized by increased ROS, iron dependent lipid peroxidation, and plasma membrane damage.¹³

Numerous studies have indicated the crucial role of ferroptosis in the development of acute lung injury,¹⁴ asthma,¹⁵ COPD,^{16–18} Alzheimer's disease¹⁹, and renal failure.¹⁵ Prolonged inhalation of particulate matter found in smoke, which includes iron particles, can contribute to the accumulation of ROS in the epithelial lining of the airways. This accumulation can cause damage to the plasma membrane, leading to the release of damage-related molecular patterns (DAMPs). Consequently, iron homeostasis is altered, resulting in excessive oxidative stress, inflammation, and eventually iron death. These processes can exacerbate the progression of emphysema and airway narrowing in individuals with COPD.^{7,20} Glutathione peroxidase 4 (GPX4) is an enzyme that plays a significant role in regulating abnormal amino acid metabolism during iron death. It functions as a crucial mediator in clearing lipid free radicals and acts as a vital protective metabolite in preventing iron death. Upon exposure to smoke, GPX4 plays a negative regulatory role in preventing the accumulation of unstable iron and reducing lipid peroxidation in lung epithelial cells. Additionally, it aids in mitigating non-apoptotic cell death, thereby reducing oxidative stress and lung injury associated with COPD.^{21,22} Nuclear Factor E2 Related Factor 2 (Nrf2), is an important protein transcription factor that plays a crucial role in intracellular oxidative stress response and antioxidant defense. When cells are subjected to oxidative stress conditions, Nrf2 gradually migrates to the nucleus, binds to the promoter of the target gene, and promotes the production of a series of antioxidant enzymes, such as GPX, which can neutralize harmful oxygen radicals and eliminate toxic compounds, thereby protecting cells from oxidative damage.²³ However, the ferroptosis-related genes of COPD remain largely unknown and require further exploration. Exploring and revealing the potential ferroptosis-related genes of COPD will provide us with potential biomarkers for the treatment of COPD.

We analyzed the dataset GSE148004 in the GEO database to explore the differentially expressed ferroptosis related genes in COPD. Then, differentially expressed ferroptosis related genes were analyzed using protein-protein interaction (PPI), correlation analysis, gene ontology (GO) enrichment analysis, and Kyoto Encyclopedia of Genes and Genomics (KEGG) pathway enrichment analysis.

Materials and Methods

Microarray Dataset Collection and Data Process

Microarray datasets were screened from GEO (<http://www.ncbi.nlm.nih.gov/geo/>). The search keywords were “COPD” and “macrophage” with the following searching strategies: ((“COPD”[MeSH Terms] OR COPD[All Fields]) AND (“macrophage”[MeSH Terms] OR macrophage[All Fields])) AND “Homo sapiens”[porgn] AND (“gse”[Filter] AND “Expression profiling by array”[Filter]). Finally, GSE148004 (as training set) was obtained. GSE148004 contained 16 whole sputum-derived alveolar macrophage RNA samples from 7 patients with COPD and 9 normal samples.

We transformed the probe into gene symbol in dataset based on the platform's annotation file, when there were multiple probes mapped to the same gene symbol; the mean value of probes was selected as the gene expression value. Differentially expressed genes (DEGs) between COPD and healthy were analyzed via the “limma package” in R software, with the following cutoff for adjustment: p value < 0.05 and FC (fold changes) > 1.0.

Screening Ferroptosis-Related Differentially Expressed Genes

A total of 369 ferroptosis-related genes were performed from the FerrDb Database (<http://www.zhounan.org/ferrdb>).²⁴ The intersection of DEGs and ferroptosis-related genes was visualized by the Venn plot.

Protein–Protein Interaction(PPI) Network Analysis of Ferroptosis-Related DEGs

The protein–protein interaction (PPI) network analysis of ferroptosis-related DEGs was conducted using the STRING online website (<https://string-db.org/>).²⁵ This creature is limited to “human”. Set a minimum valid constraint score of 0.4. The results of STRING were then imported into Cytoscape (version 3.9.1),²⁶ and the key subnetworks were extracted using the cytoHubba plugin. Then, correlation analysis of ferroptosis-related DEGs was identified using Spearman correlation in the R software “corrplot” package. Finally, the three genes with the highest scores (maximum correlation criterion, degree algorithm) in the subnetworks were selected as hub genes.

Functional Enrichment Analysis of Ferroptosis-Related DEGs

We used the “heatmap” and “ggplot2” packages of R to draw the thermal and volcanic maps. And we used the “clusterProfiler” package of R to perform the Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) enrichment analyses of ferroptosis-related DEGs. GO analysis included three categories, biological process (BP), cellular component (CC), and molecular function (MF), which was important in the exploration of biological functions. KEGG analysis was used to explore potential pathways.

Construction of Gene-miRNA and TF-Gene Network

In this study, TF-gene regulatory network and gene-miRNA interaction network was established by the NetworkAnalyst platform (<https://www.networkanalyst.ca/>).²⁷ The parameters used are as follows: Specify organism: H. sapiens (human); Set ID type: Official Gene Symbol; Gene miRNA interaction database: miRTarBase v8.0 database; and TF-gene interaction database: JASPAR database. The analysis results of NetworkAnalyst were then imported into Cytoscape software for further visualization.

Potential Drug Prediction

Potential drugs targeting hub genes were predicted by the Enrichr platform.²⁸ The access of the DSigDB database was acquired using Enrichr (<https://amp.pharm.mssm.edu/Enrichr/>).²⁹

Statistical Analysis

The statistical analyses were performed using R software (version 4.2.2).

Results

Ferroptosis-Related DEGs in COPD

First, the GSE148004 dataset was downloaded from GEO. We next analyzed the expression of 2238 DEGs in 7 COPD patients and 9 healthy individuals, which were presented using a volcanic map (Figure 1A); Then, we downloaded 369 ferroptosis related genes by using the FerrDd databases. A total of 41 ferroptosis-related genes were presented using the Wayne diagram (Figure 1B), including 19 down-regulated genes and 22 up-regulated genes (Table 1); Following the analysis of the GSE148004 dataset with R software, the 41 differentially expressed ferroptosis-related genes between COPD and normal groups were presented in heatmap (Figure 1C).

PPI Network and Correlation Analysis

To determine the interactions among ferroptosis-related DEGs, we performed PPI analysis. The results demonstrated that these ferroptosis-related genes interacted with each other (Figure 2A). and a visual analysis was performed using Cytoscape (Figure 2B). In order to explore the expression correlation of these ferroptosis-related DEGs, a correlation analysis was conducted (Figure 2C).

GO and KEGG Enrichment Analysis

We used R software to conduct GO and KEGG enrichment analysis to analyze the potential biological functions of these differentially expressed ferroptosis-related genes. The findings revealed that the most significant GO enriched terms in the inflammatory response, iron metabolism, and cellular response to hypoxia after trauma (biological process); mitochondrial outer membrane, organ outer membrane, outer membrane (cellular component); heme binding, tetrapyrrole binding, and L-amino acid transmembrane transporter activities regulate heme binding (molecular function). The outcomes were depicted using bubble, spin, and circle diagrams (Figure 3A–C). In KEGG enrichment analysis, differentially expressed ferroptosis-related genes are mainly involved in the process of central carbon metabolism in cancer, ferroptosis, and HIF-1 signaling pathways, and the results are presented using bubble charts (Figure 3D).




Figure 1 The differentially expressed ferroptosis-related genes (DEGs) in COPD and healthy samples. **(A)** Volcano plot of GSE148004. The cutoff logFC (fold change) was $-1/1$, and P-value is 0.05. Red dots indicate upregulated genes; blue dots indicate downregulated genes; gray dots indicate non-statistically significant genes. **(B)** Venn diagram of the ferroptosis-related DEGs. **(C)** Heatmap of the 41 differentially expressed ferroptosis-related genes in COPD and healthy samples. **Abbreviations:** COPD, chronic obstructive pulmonary disease; DEGs, differentially expressed genes.

Table 1 The 41 Differentially Expressed Ferroptosis-Related Genes in COPD Samples Compared to Healthy Samples

Gene Symbol	log2FC	P value	Adj. P value	Changes
BID	1.025055	0.000685	0.00401	Up
CYP4F8	1.582676	6.46E-05	0.000978	Up
ELOVL5	1.076668	0.000227	0.001975	Up
FADS2	1.062637	0.023175	0.057396	Up
HIF1A	1.518513	5.71E-05	0.000923	Up
IDO1	1.70048	0.009171	0.027269	Up
KDM6B	1.873145	0.000282	0.002257	Up
KLF2	1.781937	1.10E-05	0.000507	Up
KRAS	1.95944	2.46E-05	0.000649	Up
MAP3K14	1.344475	0.000122	0.001344	Up
MDM4	1.711914	0.000432	0.002921	Up
NDRG1	1.491937	1.38E-05	0.000538	Up
NR1D1	1.442638	0.001458	0.006775	Up
PTEN	1.352104	0.0003	0.002319	Up
SAT1	1.293103	2.84E-06	0.000334	Up
SLC38A1	1.360358	5.84E-07	0.000202	Up
SLC7A11	3.383393	2.11E-05	0.000616	Up
TBK1	1.198555	9.85E-06	0.00048	Up
TIMPI	1.649049	0.000193	0.001761	Up
TNFAIP3	2.372131	0.000121	0.001335	Up
ULK1	1.141252	0.00024	0.002043	Up
ZEB1	2.370521	3.66E-06	0.000355	Up
ACO1	-1.94494	5.69E-05	0.000923	Down
ACSF2	-1.39019	0.000159	0.001563	Down
ALOX5	-1.14873	0.000114	0.001294	Down
AQP3	-1.00798	0.010273	0.029818	Down
ATP5MC3	-1.17978	6.97E-05	0.001017	Down
EPAS1	-1.58164	8.34E-05	0.001099	Down
G6PD	-1.03218	0.001617	0.007283	Down
GSTZ1	-1.26395	3.10E-07	0.000174	Down
HDDC3	-1.06104	0.000245	0.002074	Down
HMOX1	-1.12132	8.39E-07	0.000234	Down
IDH1	-1.28995	0.000126	0.001361	Down
MMD	-1.58256	3.80E-05	0.00077	Down
MYB	-1.751	6.44E-05	0.000978	Down
PEBP1	-1.19204	0.000539	0.003388	Down
PEX12	-1.10679	3.38E-05	0.000724	Down
PPARG	-1.63734	0.000863	0.004718	Down
SLC1A5	-1.10518	0.001279	0.006189	Down
TFRC	-1.88876	9.10E-05	0.001147	Down
WVTR1	-1.11093	0.000832	0.004605	Down

Predictive Results of miRNA and TF Targeting the Hub Genes

Through the Network analyst, the gene-miRNA networks of hub genes HIF1A, RRARG, and KRAS were predicted and shown in [Supplementary Table 1](#). Based on the prediction results, a 163 node gene and miRNA co-expression network was constructed by Cytoscape and was visualized ([Figure 4A](#)), similarly, a 24 node gene and TF co-expression network was constructed by Cytoscape and was visualized ([Figure 4B](#)), which showed that has-mir-27a-3p could regulate HIF1A, PPARG, and KRAS expression; and the gene-TF network of hug genes HIF1A, PPARG, and KRAS was constructed, which identified GATA binding protein 2 (GATA2), NFKB transcription factor family member 1 (NFKB1), and




Figure 2 Protein–protein interactions (PPI) analysis the 41 differentially expressed ferroptosis-related genes. **(A)** The PPI among 41 differentially expressed ferroptosis-related genes. **(B)** The PPI network constructed with cytoscape. In the figure, nodes in crimson color represent 3 hub genes and yellow and Orange represent other common genes. **(C)** Spearman correlation analysis of the 41 differentially expressed ferroptosis-related genes. **Abbreviation:** PPI, Protein–protein interactions.

transcription factor p65 (RELA) as modulators of PPARG and HIF1A; Histone H4 Transcription Factor (HINFP) and Forkhead box(FOXC1) could regulate PPARG and KRAS.

Drug Prediction Targeting the Hub Genes

We use the DSigDB database to predict drugs associated with hub genes that may be used to treat COPD by regulating ferroptosis. Finally, a total of 380 targeted drugs were predicted, and the top 10 drugs predicted based on comprehensive scores are listed in Table 2. The top five drugs are simvastatin, N-acetyl-L-cysteine, actinomycin, diclofenac sodium, and phorbol-1-myristic acid-12-acetate (PMA).

Discussion

COPD is a major global health problem, and its development is influenced by smoking, inflammation, airway remodeling, and other factors. More and more evidence suggests that ferroptosis may be related to the pathogenesis of COPD. Smoking can lead to oxidative stress in lung cells, leading to ferroptosis in the lung epithelium. By analyzing different




Figure 3 The enrichment of GO terms and KEGG pathways based on 41 ferroptosis-related DEGs in COPD. **(A)** GO enrichment analysis of the ferroptosis-related DEGs, including biological process (BP), cell composition (CC), and molecular function (MF). **(B)** GO analysis of the top 5 differentially expressed ferroptosis-related genes. **(C)** GO analysis of the top 10 differentially expressed ferroptosis-related genes. **(D)** Kyoto Encyclopedia of Genes and Genomes (KEGG) analysis of 41 differentially expressed ferroptosis-related genes.

Abbreviations: GO, Gene Ontology; BP, biological process; CC, cellular component; MF, molecular function; KEGG, Kyoto Encyclopedia of Genes and Genomes.

ferroptosis-related differential genes in COPD patients and healthy controls, this study has identified 41 ferroptosis-related DEGs in COPD, and explored the diagnostic value of three pivotal genes, HIF1A, PPARG, and KRAS in COPD. Then, the Network analyst database is used to predict the miRNA and TF of the hub gene, and the DSigDB database is used to predict the potential target drugs of the hub gene. These drugs can play an important role in improving the severity of COPD through the ferroptosis mechanism of macrophages.

Hypoxia inducible factor-1 (HIF-1) is an important transcriptional regulator of cellular responses to hypoxia, oxidants, and inflammation.³⁰ HIF1A is the gene of hypoxia inducible factor 1 α (HIF1 α) that plays important transcriptional regulatory roles in oxygen regulation, iron death, inflammation, and immune responses. Firstly, HIF1 α is over-expressed in the lungs of COPD patients which can lead to adaptive changes in lung cells, such as angiogenesis, erythropoiesis, and cell proliferation, to adapt to hypoxic environments.^{31–33} Other studies suggest that hydrogen sulfide can inhibit CS induced PHD2/HIF1 α / MAPK signaling activation reduces the secretion of IL-6, and TNF- α , reduces oxidative stress, ROS loading, cells damage, alters iron homeostasis, thereby alleviating emphysema and improving lung function.^{31,34} Secondly, under hypoxic conditions, HIF1 α play a key role in iron metabolism by regulating the expression of iron-related proteins, such as DMT1, FPN1, Dcytb, and transferrin receptor (TfR), which may lead to abnormal accumulation of intracellular iron and the production of excessive oxygen free radicals, thereby exacerbating the progression of COPD.^{35,36}

Peroxisome proliferator-activated receptors (PPARs) are members of the steroid hormone receptor superfamily and are fatty acid sensors that regulate various aspects of lipid metabolism.³⁷ Currently, there are three subtypes of PPARs, including α , β/δ and γ . PPAR α promotes oxidation of fatty acids in the liver,³⁸ PPAR δ is an important regulator of lipid




Figure 4 Predictive networks of microRNAs (miRNAs), transcription factors (TFs). **(A)** Target gene-miRNA network. **(B)** Target gene-TF network. **Abbreviation:** TFs, transcription factors.

oxidation in skeletal muscle,³⁹ PPAR γ has been found in lung tissue and alveolar macrophages (AMs),⁴⁰ which could combined with NF κ B to inhibit the expression levels of inflammatory mediators, such as TNF- α and CC chemokine ligand(CCL)5, which overall reduced granulocyte aggregation, and lessened the degree of inflammation in the lungs of COPD patients.⁴¹ In addition, hydrogen sulfide can improve CS induced emphysema and airway inflammation by activating the Nrf2 and PPAR γ signaling pathways, maintaining iron metabolism and lipid metabolism balance.⁴²

Table 2 Top 10 Drugs with Higher Scores

Rank	Term	Odds Ratio	Combined Score	Genes
1	Simvastatin CTD 00007319	59,088	742,688.9	PPARG;KRAS;HIF1A
2	N-Acetyl-L-cysteine CTD 00005305	59,055	735,956.3	PPARG;KRAS;HIF1A
3	Cycloheximide CTD 00005731	59,028	730,615.4	PPARG;KRAS;HIF1A
4	Diclofenac CTD 00005804	58,719	678,032	PPARG;KRAS;HIF1A
5	Phorbol 12-myristate 13-acetate CTD 00006852	58,551	654,397.8	PPARG;KRAS;HIF1A
6	Curcumin CTD 00000663	58,416	637,246.8	PPARG;KRAS;HIF1A
7	5-Fluorouracil CTD 00005987	56,394	475,831	PPARG;KRAS;HIF1A
8	Genistein CTD 00007324	56,307	471,066.5	PPARG;KRAS;HIF1A
9	Bisphenol A CTD 00000312	56,214	466,090.8	PPARG;KRAS;HIF1A
10	Arsenenous acid CTD 00000922	56,151	462,786.2	PPARG;KRAS;HIF1A

Kirsten rat sarcoma virus oncogene (KRAS) is the most frequently mutated member of the Ras family in human tumors, usually associated with pancreatic cancer and lung cancer,⁴³ and one of the most important regulators of cell proliferation, differentiation and survival.⁴⁴ COPD is an independent risk factor for lung cancer. KRAS can also affect COPD and iron death through several pathways, but its specific mechanism is still unclear. Moghadamet used non-classifiable *Haemophilus influenzae* (NTHi) to induce COPD-like airway inflammation to study the effect of KRAS mutation expression in airway epithelium on induced lung cancer, and found that it has a promoting effect on lung cancer.⁴⁵ Another study showed that the high expression of HIF1 α is also sufficient to enhance KRAS-induced lung neoplasia, while promoting tumor growth, increasing tumor vessel survival, and cell proliferation.⁴⁶

Two drugs with the highest scores selected by us through DsigDB are currently in clinical use. Studies have shown that Simvastatin enhances the anti-tumor effects of bevacizumab against lung adenocarcinoma A549 Cells by abating the HIF-1 α -Wnt/ β -Catenin signaling pathway.^{47,48} However, research on simvastatin targeting PPARG and KRAS for the treatment of COPD is relatively scarce. Research has shown that the antioxidant N-acetylcysteine (NAC) has been identified as a novel inhibitor of the hypoxic response pathway. It can serve as a precursor to glutathione (GSH), increasing GSH content, clearing reactive oxygen species, and reducing the ferrous ions required for the oxidation of proline (PHDs), this weakens the stability of HIF-1 α under hypoxic conditions.⁴⁹ NAC has been widely used in the treatment of COPD. Additionally, other studies have revealed that the imbalance between T helper (Th)17 cells and regulatory T (Treg) cells plays a vital role in the development and outcome of COPD.^{50,51} NAC can downregulate the expression of HIF-1 α , thus regulating the balance of Th1/Treg and reducing the release of the inflammatory cytokine IL-17, thereby improving lung function in COPD patients.⁵² Currently, clinical drugs for treating COPD mainly focus on the research direction of HIF1A, but there is relatively little research on PPARG and KRAS. Therefore, future research directions can focus on treatment strategies for PPARG and KRAS to gain a more comprehensive understanding and treatment of COPD.

We also evaluated the biological functions and pathways of ferroptosis-related DEGs, indicating that the HIF-1 signaling pathway is involved in the occurrence and development of COPD. However, this study has some limitations. First, the number of samples chosen is relatively small, and a larger sample size is required. Second, due to time and funding constraints, this study only employs bioinformatics methods. Changes in the mRNA or protein levels of DEGs can be detected as a supplement to this study using RT-PCR or Western blotting, and clinical samples or animal experiments can also be supplemented.

Conclusions

In the present study, 41 potential ferroptosis-related genes of COPD were identified through bioinformatics analysis. The Network analyst and Enrichr platform was used to predict the miRNAs and TF and drugs associated with the hub genes, indicating that these drugs may affect the development of COPD by regulating ferroptosis in macrophages. These results expand our understanding of COPD and may contribute to the treatment of COPD.

Abbreviations

COPD, chronic obstructive pulmonary disease; GEO, gene expression omnibus dataset; GO, Gene Ontology; KEGG, Kyoto Encyclopedia of Genes and Genomes; BP, biological process; CC, cellular component; MF, molecular function; PPI, Protein-Protein Interaction; miRNAs, microRNAs; TFs, transcription factors; IL-6, Interleukin-6; IL-18, Interleukin-18; TNF- α , Tumor Necrosis Factor-alpha, DMT1: divalent metal transporter 1; FPN1, ferroportin 1; Dcytb, duodenal cyto-chrome b; TfR: Transferrin receptor.

Data Sharing Statement

The datasets analyzed in this paper can be found in the GEO database. A total of 369 ferroptosis-related genes were performed from the FerrDd Database

Ethics Approval

The study is approved by the Medical Ethics Committees, Zhongnan Hospital of Wuhan University (No.2020099K).

Acknowledgments

We thank Hong Duo (Institute of Hepatobiliary Diseases of Wuhan University, Transplant Center of Wuhan University, Hubei Key Laboratory of Medical Technology on Transplantation, Wuhan, 430071, Hubei, China) for the image technical support

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

Funding was not applicable.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Ferrera MC, Labaki WW, Han MK. Advances in Chronic Obstructive Pulmonary Disease. *Annu Rev Med.* 2021;72(1):119–134. doi:10.1146/annurev-med-080919-112707
2. Xu W, Deng H, Hu S, et al. Role of Ferroptosis in Lung Diseases. *J Inflamm Res.* 2021;14:2079–2090. doi:10.2147/JIR.S307081
3. Labaki WW, Rosenberg SR. Chronic Obstructive Pulmonary Disease. *Ann Intern Med.* 2020;173(3):567.
4. Li Y, Yang Y, Yang Y. Multifaceted Roles of Ferroptosis in Lung Diseases. *Front Mol Biosci.* 2022;9:919187. doi:10.3389/fmolb.2022.919187
5. Cokkinides V, Bandi P, McMahon C, Jemal A, Glynn T, Ward E. Tobacco control in the United States—recent progress and opportunities. *CA Cancer J Clin.* 2009;59(6):352–365. doi:10.3322/caac.20037
6. Hu H, Zhang L, Teng G, Wu Y, Chen Y. A variant in 3'-untranslated region of KRAS compromises its interaction with hsa-let-7g and contributes to the development of lung cancer in patients with COPD. *Int J Chron Obstruct Pulmon Dis.* 2015;10:1641–1649. doi:10.2147/COPD.S83596
7. Ghio AJ, Hilborn ED, Stonehuerner JG, et al. Particulate matter in cigarette smoke alters iron homeostasis to produce a biological effect. *Am J Respir Crit Care Med.* 2008;178(11):1130–1138. doi:10.1164/rccm.200802-334OC
8. Koch A, Bettex M, Tschappeler H, Konig W. The function of the esophagus following cardiomyotomy in childhood achalasia. *Z Kinderchir.* 1983;38(4):206–210. doi:10.1055/s-2008-1059969
9. Shukla SD, Walters EH, Simpson JL, et al. Hypoxia-inducible factor and bacterial infections in chronic obstructive pulmonary disease. *Respirology.* 2020;25(1):53–63. doi:10.1111/resp.13722
10. Boucherat O, Morisette MC, Provencher S, Bonnet S, Maltais F. Bridging Lung Development with Chronic Obstructive Pulmonary Disease. Relevance of Developmental Pathways in Chronic Obstructive Pulmonary Disease Pathogenesis. *Am J Respir Crit Care Med.* 2016;193(4):362–375. doi:10.1164/rccm.201508-1518PP
11. Yang YC, Zhang MY, Liu JY, Jiang YY, Ji XL, Qu YQ. Identification of Ferroptosis-Related Hub Genes and Their Association with Immune Infiltration in Chronic Obstructive Pulmonary Disease by Bioinformatics Analysis. *Int J Chron Obstruct Pulmon Dis.* 2022;17:1219–1236. doi:10.2147/COPD.S348569
12. Jiang X, Stockwell BR, Conrad M. Ferroptosis: mechanisms, biology and role in disease. *Nat Rev Mol Cell Biol.* 2021;22(4):266–282. doi:10.1038/s41580-020-00324-8
13. Dixon SJ, Lemberg KM, Lamprecht MR, et al. Ferroptosis: an iron-dependent form of nonapoptotic cell death. *Cell.* 2012;149(5):1060–1072. doi:10.1016/j.cell.2012.03.042
14. Zhou H, Li F, Niu JY, et al. Ferroptosis was involved in the oleic acid-induced acute lung injury in mice. *Sheng Li Xue Bao.* 2019;71(5):689–697.
15. Wenzel SE, Tyurina YY, Zhao J, et al. PEBP1 Wards Ferroptosis by Enabling Lipoygenase Generation of Lipid Death Signals. *Cell.* 2017;171(3):628–641 e626. doi:10.1016/j.cell.2017.09.044
16. Han C, Liu Y, Dai R, Ismail N, Su W, Li B. Ferroptosis and Its Potential Role in Human Diseases. *Front Pharmacol.* 2020;11:239. doi:10.3389/fphar.2020.00239
17. Ho T, Nichols M, Nair G, et al. Iron in airway macrophages and infective exacerbations of chronic obstructive pulmonary disease. *Respir Res.* 2022;23(1):8. doi:10.1186/s12931-022-01929-7
18. Meng D, Zhu C, Jia R, Li Z, Wang W, Song S. The molecular mechanism of ferroptosis and its role in COPD. *Front Med Lausanne.* 2022;9:1052540. doi:10.3389/fmed.2022.1052540
19. Bao WD, Pang P, Zhou XT, et al. Loss of ferroportin induces memory impairment by promoting ferroptosis in Alzheimer's disease. *Cell Death Differ.* 2021;28(5):1548–1562. doi:10.1038/s41418-020-00685-9
20. Berg S, Luneberg E, Frosch M. Development of an amplification and hybridization assay for the specific and sensitive detection of Mycoplasma fermentans DNA. *Mol Cell Probes.* 1996;10(1):7–14. doi:10.1006/mcpr.1996.0002
21. Gressmann C. Bone development in peromelia (proceedings). *Z Orthop Ihre Grenzgeb.* 1978;116(4):605–606.

22. Ursini F, Maiorino M, Valente M, Ferri L, Gregolin C. Purification from pig liver of a protein which protects liposomes and biomembranes from peroxidative degradation and exhibits glutathione peroxidase activity on phosphatidylcholine hydroperoxides. *Biochim Biophys Acta*. 1982;710(2):197–211. doi:10.1016/0005-2760(82)90150-3
23. Rangasamy T, Cho CY, Thimmulappa RK, et al. Genetic ablation of Nrf2 enhances susceptibility to cigarette smoke-induced emphysema in mice. *J Clin Invest*. 2004;114(9):1248–1259. doi:10.1172/JCI200421146
24. Zhou N, Bao J. FerrDb: a manually curated resource for regulators and markers of ferroptosis and ferroptosis-disease associations. *Database*. 2020;2020. doi:10.1093/database/baaa021
25. Szklarczyk D, Gable AL, Lyon D, et al. STRING v11: protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res*. 2019;47(D1):D607–D613. doi:10.1093/nar/gky1131
26. Shannon P, Markiel A, Ozier O, et al. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res*. 2003;13(11):2498–2504. doi:10.1101/gr.1239303
27. Zhou G, Soufan O, Ewald J, Hancock REW, Basu N, Xia J. NetworkAnalyst 3.0: a visual analytics platform for comprehensive gene expression profiling and meta-analysis. *Nucleic Acids Res*. 2019;47(W1):W234–W241. doi:10.1093/nar/gkz240
28. Kuleshov MV, Jones MR, Rouillard AD, et al. Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Res*. 2016;44(W1):W90–97. doi:10.1093/nar/gkw377
29. Yoo M, Shin J, Kim J, et al. DSigDB: drug signatures database for gene set analysis. *Bioinformatics*. 2015;31(18):3069–3071. doi:10.1093/bioinformatics/btv313
30. Wang L, Tang Y, Chen Y. HIF1A gene rs10873142 polymorphism is associated with risk of chronic obstructive pulmonary disease in a Chinese Han population: a case-control study. *Biosci Rep*. 2018;38(2):79.
31. Guan R, Wang J, Li D, et al. Hydrogen sulfide inhibits cigarette smoke-induced inflammation and injury in alveolar epithelial cells by suppressing PHD2/HIF-1 α /MAPK signaling pathway. *Int Immunopharmacol*. 2020;81:105979. doi:10.1016/j.intimp.2019.105979
32. Semenza GL. Pulmonary vascular responses to chronic hypoxia mediated by hypoxia-inducible factor 1. *Proc Am Thorac Soc*. 2005;2(1):68–70. doi:10.1513/pats.200404-029MS
33. Semenza GL. HIF-1: mediator of physiological and pathophysiological responses to hypoxia. *J Appl Physiol*. 2000;88(4):1474–1480. doi:10.1152/jappl.2000.88.4.1474
34. Zhang HX, Yang JJ, Zhang SA, et al. HIF-1 α promotes inflammatory response of chronic obstructive pulmonary disease by activating EGFR/PI3K/AKT pathway. *Eur Rev Med Pharmacol Sci*. 2018;22(18):6077–6084. doi:10.26355/eurrev_201809_15946
35. Xu MM, Wang J, Xie JX. Regulation of iron metabolism by hypoxia-inducible factors. *Sheng Li Xue Bao*. 2017;69(5):598–610.
36. Tacchini L, Bianchi L, Bernelli-Zazzera A, Cairo G. Transferrin receptor induction by hypoxia. HIF-1-mediated transcriptional activation and cell-specific post-transcriptional regulation. *J Biol Chem*. 1999;274(34):24142–24146. doi:10.1074/jbc.274.34.24142
37. Issemann I, Green S. Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature*. 1990;347(6294):645–650. doi:10.1038/347645a0
38. Lefebvre P, Chinetti G, Fruchart JC, Staels B. Sorting out the roles of PPAR α in energy metabolism and vascular homeostasis. *J Clin Invest*. 2006;116(3):571–580. doi:10.1172/JCI27989
39. Barish GD, Narkar VA, Evans RM. PPAR δ : a dagger in the heart of the metabolic syndrome. *J Clin Invest*. 2006;116(3):590–597. doi:10.1172/JCI27955
40. Caito S, Yang SR, Kode A, et al. Rosiglitazone and 15-deoxy-Delta12,14-prostaglandin J2, PPAR γ agonists, differentially regulate cigarette smoke-mediated pro-inflammatory cytokine release in monocytes/macrophages. *Antioxid Redox Signal*. 2008;10(2):253–260. doi:10.1089/ars.2007.1889
41. Qiu JF, Ma N, He ZY, et al. Erythromycin inhibits cigarette smoke-induced inflammation through regulating the PPAR γ /NF- κ B signaling pathway in macrophages. *Int Immunopharmacol*. 2021;96:107775. doi:10.1016/j.intimp.2021.107775
42. Wang Y, Liao S, Pan Z, et al. Hydrogen sulfide alleviates particulate matter-induced emphysema and airway inflammation by suppressing ferroptosis. *Free Radic Biol Med*. 2022;186:1–16. doi:10.1016/j.freeradbiomed.2022.04.014
43. Moghaddam SJ, Li H, Cho SN, et al. Promotion of lung carcinogenesis by chronic obstructive pulmonary disease-like airway inflammation in a K-ras-induced mouse model. *Am J Respir Cell Mol Biol*. 2009;40(4):443–453. doi:10.1165/rcmb.2008-0198OC
44. Keohavong P, Peter Di Y. Pulmonary Inflammation and KRAS Mutation in Lung Cancer. *Adv Exp Med Biol*. 2021;1303:71–87.
45. Moghaddam SJ, Clement CG, De la Garza MM, et al. Haemophilus influenzae lysate induces aspects of the chronic obstructive pulmonary disease phenotype. *Am J Respir Cell Mol Biol*. 2008;38(6):629–638. doi:10.1165/rcmb.2007-0366OC
46. De la Garza MM, Cumpian AM, Daliri S, et al. COPD-Type lung inflammation promotes K-ras mutant lung cancer through epithelial HIF-1 α mediated tumor angiogenesis and proliferation. *Oncotarget*. 2018;9(68):32972–32983. doi:10.18632/oncotarget.26030
47. Tu X, Zhang J, Yuan W, Wu X, Xu Z, Qing C. Simvastatin Enhanced Anti-tumor Effects of Bevacizumab against Lung Adenocarcinoma A549 Cells via Abating HIF-1 α -Wnt/ β -Catenin Signaling Pathway. *Anticancer Agents Med Chem*. 2023;23(19):2083–2094. doi:10.2174/1871520623666230816090914
48. Khanzada UK, Pardo OE, Meier C, Downward J, Seckl MJ, Arcaro A. Potent inhibition of small-cell lung cancer cell growth by simvastatin reveals selective functions of Ras isoforms in growth factor signalling. *Oncogene*. 2006;25(6):877–887. doi:10.1038/sj.onc.1209117
49. Liu X, Hu Z, Zhou H. N-Acetylcysteine Improves Inflammatory Response in COPD Patients by Regulating Th17/Treg Balance through Hypoxia Inducible Factor-1 α Pathway. *Biomed Res Int*. 2021;2021:6372128. doi:10.1155/2021/6372128
50. Su YC, Jalalvand F, Thegerstrom J, Riesbeck K. The Interplay Between Immune Response and Bacterial Infection in COPD: focus Upon Non-typeable Haemophilus influenzae. *Front Immunol*. 2018;9:2530. doi:10.3389/fimmu.2018.02530
51. Polverino F, Seys LJ, Bracke KR, Owen CA. B cells in chronic obstructive pulmonary disease: moving to center stage. *Am J Physiol Lung Cell Mol Physiol*. 2016;311(4):L687–L695. doi:10.1152/ajplung.00304.2016
52. Chaussy C, Fuchs G, Kahn R, Hunter P, Goodfriend R. Transurethral ultrasonic ureterolithotripsy using a solid-wire probe. *Urology*. 1987;29(5):531–532. doi:10.1016/0090-4295(87)90045-8

International Journal of Chronic Obstructive Pulmonary Disease

Dovepress

Publish your work in this journal

The International Journal of COPD is an international, peer-reviewed journal of therapeutics and pharmacology focusing on concise rapid reporting of clinical studies and reviews in COPD. Special focus is given to the pathophysiological processes underlying the disease, intervention programs, patient focused education, and self management protocols. This journal is indexed on PubMed Central, MedLine and CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-chronic-obstructive-pulmonary-disease-journal>