

# Pathophysiological Mechanisms and Clinical Associations of Non-Alcoholic Fatty Pancreas Disease

Antonia Pagkali<sup>1</sup>, Anastasios Makris<sup>1</sup>, Kalliopi Brofidi<sup>2</sup>, Aris P Agouridis<sup>3,4</sup>, Theodosios D Filippatos<sup>1,2</sup>

<sup>1</sup>School of Medicine, National and Kapodistrian University of Athens, Athens, Greece; <sup>2</sup>Department of Internal Medicine, School of Medicine, University of Crete, Heraklion, Greece; <sup>3</sup>School of Medicine, European University Cyprus, Nicosia, Cyprus; <sup>4</sup>Department of Internal Medicine, German Oncology Center, Limassol, Cyprus

Correspondence: Theodosios D Filippatos, Department of Internal Medicine, School of Medicine, University of Crete, Heraklion, Greece, Tel + (30)2810392960, Fax + (30)2810392359, Email [filtheo@uoc.gr](mailto:filtheo@uoc.gr)

**Abstract:** Non-Alcoholic Fatty Pancreas disease (NAFPD), characterized by fat accumulation in pancreatic tissue, is an emerging clinical entity. However, the clinical associations, the underlying molecular drivers, and the pathophysiological mechanisms of NAFPD have not yet been characterized in detail. The NAFPD spectrum not only includes infiltration and accumulation of fat within and between pancreatic cells but also involves several inflammatory processes, dysregulation of physiological metabolic pathways, and hormonal defects. A deeper understanding of the underlying molecular mechanisms is key to correlate NAFPD with clinical entities including non-alcoholic fatty liver disease, metabolic syndrome, diabetes mellitus, atherosclerosis, as well as pancreatic cancer and pancreatitis. The aim of this review is to examine the pathophysiological mechanisms of NAFPD and to assess the possible causative/predictive risk factors of NAFPD-related clinical syndromes.

**Keywords:** non-alcoholic fatty pancreas disease, NAFPD, steatotic pancreatic disease, metabolic dysfunction-associated steatotic pancreatic disease, MASPD

## Introduction

Pancreatic fat deposition in the absence of excessive alcohol intake, described as non-alcoholic fatty pancreas disease (NAFPD),<sup>1</sup> is an emerging clinical entity.<sup>2</sup> Despite the higher susceptibility to fat deposition of the pancreas compared to the liver,<sup>3,4</sup> the clinical relevance and underlying pathophysiological mechanisms of pancreatic fat infiltration remain unclear in the medical literature. Robertson F. Ogilvie was the first to describe pancreatic fat deposition in obese cadavers, reporting 17% of pancreatic fat storage, as compared to 9% in lean ones.<sup>5</sup> Since then, excessive storage of fat in pancreatic tissue has been described with various terms, including pancreatic lipomatosis, pancreatic steatosis, lipomatous pseudohypertrophy, fatty pancreas, and more.<sup>6,7</sup> Currently, NAFPD is the widely accepted term, and its prevalence varies significantly among populations.<sup>8–11</sup>

There are many sophisticated imaging techniques used as diagnostic tools to evaluate the pancreas, with transcutaneous ultrasound being the most common,<sup>10,12</sup> since it is noninvasive, inexpensive, and easily performed. Nonetheless, endoscopic ultrasound (EUS),<sup>10,13</sup> abdominal computed tomography (CT),<sup>14</sup> or magnetic resonance-based techniques including abdominal magnetic resonance imaging (MRI),<sup>15,16</sup> and magnetic resonance spectroscopy (MRS)<sup>17</sup> can be used to quantify pancreatic fat content with accuracy. A direct comparison of these methods has not been performed yet. MRI enables noninvasive assessment of fat content in solid organs,<sup>18</sup> and with the Iterative Decomposition with Echo Asymmetry and Least squares estimation (IDEAL) method,<sup>19</sup> fat and water can be separated for a more precise organ assessment. EUS allows for a closer access to the pancreas; nonetheless, it is invasive and unsuitable for quantification of the entire pancreatic fat content.<sup>10,13</sup> In the general population and in routine medical checkup settings, invasiveness and cost-effectiveness should always be considered before examining the pancreas. As far as physiological values are

concerned, the highest limit of normal pancreatic fat is 10.4% when assessed by MRI/MRS.<sup>9</sup> However, a meta-analysis showed that the upper normal limit of pancreatic fat in healthy individuals, who participated in MRI studies, was 6.2%,<sup>20</sup> a threshold that is recommended for future research use.


The aim of this review is to explore the pathophysiological mechanisms of NAFPD and its clinical associations, as well as to examine which of these parameters are possible causative or predictive risk factors of NAFPD.

## Pathogenesis

NAFPD is the result of several pathophysiological processes that act simultaneously (Figure 1). These processes are generated mainly from excessive deposition of fat on pancreatic cells and include the following; (1) immediate cellular damage due to direct interaction of fatty tissue with pancreatic cells, with subsequent release of free fatty acids,<sup>1, 21–23</sup> (2) chronic oxidative stress<sup>24,25</sup> and local secretion of detrimental inflammatory markers,<sup>1, 26–28</sup> (3) pathological activation or alterations in lipid-glucose metabolism, beta-oxidation, and other regulators,<sup>29–31</sup> and (4) an imbalance of hormone homeostasis.<sup>32,33</sup>

## Fat Replacement and Fat Accumulation

The main pathogenetic mechanism of NAFPD involves fatty replacement, which can either be intralobular or interlobular, followed by fat accumulation within the pancreatic tissue.<sup>21</sup> In most cases, excessive weight gain gradually leads to the storage of surplus adipose tissue in non-adipose organs, including the pancreas, where fat accumulates intracellularly in both acinar and pancreatic islet cells, causing dysfunction.<sup>21</sup> The fatty replacement phase is characterized by an accumulation of intralobular fat in the form of lipid droplets in both endocrine and acinar cells,<sup>34,35</sup> provoking acinar cell



**Figure 1** Pathogenetic mechanisms behind fatty pancreas. The main pathogenetic mechanism behind non-alcoholic fatty pancreas diseases (NAFPD) is fatty accumulation (intra- or inter-lobular), followed by fatty replacement and, ultimately,  $\beta$  cell dysfunction. The main risk factors associated with NAFPD are obesity and metabolic syndrome (including dyslipidemia), while alcohol, viruses, iron deposition, drugs, gut hormones, psoriasis and pancreatic duct obstruction represent potential secondary hits which participate in cell death and replacement of pancreatic tissue with fat. Mechanisms associated with fat accumulation include reduction in mitochondrial activity, inflammatory cell infiltration with production of inflammatory cytokines (eg TNF $\alpha$ , IL-1 $\beta$ , IL-6, MCP-1), activation of stellate cells, hormonal imbalance (with perturbed leptin and adiponectin levels), as well as STK-25 pathway overexpression, which further exacerbates all the aforementioned mechanisms. NAFPD, in turn, is associated with various conditions including diabetes, pancreatitis, pancreatic cancer, as well as atherosclerosis.

death and their consecutive replacement by adipocytes, while pancreatic  $\beta$ -cells decrease and become dysfunctional.<sup>1,3,36–41</sup> Hence, fatty accumulation follows fatty replacement and acinar cell necrosis.<sup>7,34,41,42</sup> Fatty replacement occurs unevenly throughout the pancreatic tissue, with the anterior aspect being more prone.<sup>7</sup> Of note, the term fatty replacement denotes irreversible damage, not fully correctable with medications or weight reduction.<sup>43</sup> In parallel, interlobular fat, that is fat around large vessels and ducts, involves adipocytes and lipid droplets in inactive stellate cells, which become stimulated.<sup>22,34</sup> In addition, deposition of pancreatic fat ensues secondary to increased levels of free fatty acids in the plasma,<sup>1</sup> a process similar to the hepatic free fatty acid influx noted in non-alcoholic fatty liver disease (NAFLD).<sup>44</sup> Following such pancreatic insults, acinar cell apoptosis leads to rapid pancreatic volume decrease, but eventually, fatty restoration causes volume replacement until the pancreas becomes prominent and enlarged. In an animal model study, accumulation of fat tissue in the pancreas induced apoptosis, stellate cell activation, fibrosis, and, ultimately, a decrease in  $\beta/\alpha$  islet cell ratio with loss of architecture.<sup>45</sup> Likewise, in another animal study, where obese mice were investigated for pancreatic cellular changes, intralobular and interlobular fat storage was responsible for a heavier pancreas, larger pancreatic islet cells, and atypical cellular structure.<sup>28</sup> These modifications may have occurred in an attempt of  $\beta$ -cells for compensation;  $\beta$ -cells compensate against initial cellular and metabolic fluctuations, presenting an enhancement in their secretory activity, an expansion of their  $\beta$ -cell mass, or both.<sup>29</sup> Overall, this might explain why pancreatic cells with chronically elevated fat content will become hypertrophic and hyperplastic.<sup>46</sup> Deposition of fat tissue in the pancreas, in combination with free circulating fatty acids, causes  $\beta$ -cell dysfunction,<sup>23</sup> with a baseline increase in insulin release, an impaired glucose-stimulated insulin secretion of  $\beta$ -cells<sup>29,47,48</sup> and the generation of oxidative stress with consequent organ injuries.<sup>1,23,49</sup> After the increase of basal insulin secretion, there is a subsequent decrease, a state that cannot be repealed with the induction of compensatory proinsulin biosynthesis.<sup>50</sup> Therefore, clinical conditions, such as insulin secretion impairment and diabetes, might arise.<sup>23,46</sup> Many of these mechanisms are supported in a human study, where Tushuizen et al<sup>17</sup> showed a negative correlation between pancreatic fat content with  $\beta$ -cell function in subjects without diabetes.

## Oxidative Stress and Inflammatory Processes

Chronic oxidative stress is implicated in metabolic syndrome, diabetes mellitus and atherosclerosis. Indeed, oxidative stress is referred to as a “second hit” in cases of steatohepatitis,<sup>12</sup> so a similar process in the case of NAFLD may be hypothesized.<sup>51</sup>

Increased body weight provokes chronic cortisol release<sup>24</sup> and activation of the hypothalamic-pituitary-adrenal (HPA) axis,<sup>24,25</sup> thus increasing metabolic syndrome and atherosclerotic disease risk.<sup>52</sup> Fat storage induces local pancreatic inflammatory cell infiltration,<sup>26</sup> further inducing  $\beta$ -cell function deterioration and destruction. Pro-inflammatory cytokines, including tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), interleukin-1 $\beta$  (IL-1 $\beta$ ) and 6 (IL-6), monocyte chemoattractant protein-1 (MCP-1) are released, leading to inflammation, fibrosis, and insulin resistance.<sup>1</sup> Free fatty acids are released via lipolysis of visceral adipose tissue,<sup>22,27,28,53</sup> but also due to inflammatory responses,<sup>27</sup> altogether resulting in histopathological alterations and secondary pancreatic acinar cell atrophy and fibrosis.<sup>38,54</sup> Of importance, the role of copper in pancreatic disease was investigated in an animal study.<sup>55</sup> The authors stated that pancreatic atrophy is more pronounced in males compared with females, while the endocrine male pancreas is also more susceptible to dietary copper deprivation than the female rat.<sup>55</sup> Similarly, copper is also important in NAFLD. In a cross-sectional study that included 100 obese patients with hepatic steatosis detected by ultrasound, it was shown that an altered copper bioavailability predicts early atherosclerosis.<sup>56</sup> Moreover, older studies demonstrated that pancreatic fatty deposition is also accompanied by leukocyte infiltration,<sup>57</sup> although more recent ones did not corroborate these findings.<sup>1</sup>

## Dysregulation of Metabolic Pathways

In terms of dysregulated pathways, fatty accumulation of the pancreas influences multiple regulatory pathways. First, fat deposition impairs pancreatic  $\beta$ -cell cAMP production and, as a result, it inhibits islet cell adaptation to the disturbed metabolic milieu.<sup>29,30</sup> Another probable mechanism is the augmented phospholipid saturation of the mitochondrial membrane, which results in diminished mitochondrial function,<sup>58</sup> peroxisomal fatty acid metabolism, cytochrome P450 CYP-2E1 induction, and the subsequent production of free oxygen radicals, as in the case of NAFLD.<sup>59,60</sup> This

leads to further activation of stellate cells that attract neutrophils and cause an inflammatory reaction.<sup>61</sup> Additionally, in the context of chronic fatty pancreas accumulation, overexpression of the serine/threonine-protein kinase 25 (STK-25), a novel negative regulator of lipid and glucose metabolism, occurs.<sup>62</sup> STK-25 can be overexpressed when feeding mice with high-fat diets, aggravating lipid accumulation in cells and impairing skeletal muscle mitochondrial function, thereby diminishing insulin sensitivity.<sup>62–64</sup> In skeletal myocytes of patients with type 2 diabetes mellitus, STK-25 mRNA levels were significantly elevated, compared with healthy controls.<sup>62</sup> The overexpression of STK25 in NAFLD exacerbates all the aforementioned pathophysiological mechanisms, including inflammatory pancreas infiltration,<sup>45,65</sup> stellate cell activation,  $\beta$ -cell loss, and pancreatic destruction.<sup>45</sup> Indeed, STK25 is considered to play a key role in both the pathogenesis and progression of NAFLD.<sup>66</sup> Overexpression of STK25 induces lipogenesis, triglyceride secretion and suppressed beta-oxidation.<sup>66</sup> A similar mechanism may be adopted by pancreatic cells in cases of STK25 overstimulation highlighting that NAFLD and NAFLD are two closely related conditions. Beta-oxidation is severely impaired during fatty accumulation.<sup>31</sup> When there is an impediment in fatty acid catabolism, fat will also accumulate in zone 3 of hepatocytes, giving rise to NAFLD.<sup>12</sup> In pancreatic tissue, parameters like hyperglycemia and hypertriglyceridemia induce cellular free fatty acid increase, loss of  $\beta$ -cell mass, as well as decreased beta-oxidation.<sup>53</sup> Increased glucose levels augment malonyl-coenzyme A via the tricarboxylic acid cycle, thus inhibiting carnitine palmitoyltransferase-1 and diminishing mitochondrial  $\beta$ -oxidation while stimulating triglyceride production in  $\beta$ -cells.<sup>53</sup> In correlation with that, when obese mice were studied,<sup>1</sup> accumulation of cholesterol and free fatty intoxication were observed. Through this process, triglyceride accumulation in  $\beta$ -cells is promoted, which, in combination with lipid peroxidation, leads acinar and islet cells to destruction.<sup>27,28,53</sup> When triglycerides continue to accumulate inside pancreatic cells, insulin gene expression is downregulated, and glucose-stimulated insulin secretion is impaired.<sup>22,28,53</sup>

## Dysregulation of Hormonal Homeostasis

Adipocytokines, which consist of adipose tissue released cytokines, adipokines and chemokines, are valuable bioactive substances that maintain metabolic homeostasis and protect against metabolic overload.<sup>32,33</sup> Decreased levels of adiponectin in obese patients signify the loss of insulin sensitivity modulation, and the decrease of fatty acid oxidation in body tissues.<sup>33</sup> In contrast, regulation of leptin levels in obese patients is a more complex pathway. It is thought that leptin levels and obesity are reversibly correlated; when leptin increases, it leads to the expression of suppressor-of-cytokine-signaling (SOCS-3) pathway, which thereafter inhibits leptin signaling.<sup>67</sup> In obesity, adipocytokines like leptin and adiponectin may differentially affect pancreatic tissue.<sup>33</sup> It has been shown that adiponectin levels are positively correlated with plasma insulin and negatively associated with plasma triglycerides and insulin resistance.<sup>68</sup>

Notably, it has been demonstrated that the gut-brain axis may also play a role in NAFLD, and more specifically overexpression of ghrelin may contribute to increased intra-pancreatic fat deposition in individuals with acute pancreatitis.<sup>69</sup> Similarly, gut hormones may contribute to NAFLD pathogenesis, but the association between intrahepatic fat deposition and gut hormones remains to be elucidated.<sup>70</sup>

## Human-Based Studies and Correlation with Molecular Abnormalities

Even though the pathophysiological mechanisms between fatty pancreas and  $\beta$ -cell dysfunction/apoptosis have been thoroughly discussed, recent findings from human studies do not come in accordance with previous conclusions. In a recent large cohort study of adult Chinese subjects, who had undergone fatty pancreas evaluation by MRI, an association of fatty pancreas and  $\beta$ -cell function was found.<sup>9</sup> However, this association disappeared after adjusting for BMI and intrahepatic triglycerides. Although this cohort did not show an independent association between fatty pancreas and  $\beta$ -cell function, pancreatic fat accumulation was still associated with insulin resistance, even after adjusting for hepatic fat content and BMI.<sup>9</sup> Likewise, in another human-based comprehensive analysis of interlobular, intralobular and parenchymal pancreatic fat and their effects on  $\beta$ -cell performance, no relationship was found between total and intralobular pancreatic adipose tissue infiltration with  $\beta$ -cell function.<sup>71</sup> In some small case series that compared pancreatic fat among patients with diabetes with that of normal controls, an association of fatty pancreas with weakened  $\beta$ -cell function and insulin secretion was noted.<sup>17,72</sup> Nevertheless, we cannot conclude whether this  $\beta$ -cell dysfunction was established beforehand or if it was a direct consequence of fatty accumulation. Ozturk et al,<sup>73</sup> while studying the

association between NAFLD and atherosclerosis, emphasized that insulin resistance appeared among their study population in the absence of obesity and diabetes mellitus. In a study of 106 subjects with new-onset of type 2 diabetes, no significant correlation was noted between the amount of pancreatic fat deposition and islet cell function.<sup>74</sup> The controversial results of previous studies either drive us towards the existence of more pathophysiological cascades that make fatty accumulation a causative factor of  $\beta$ -cell dysfunction, or to genetic polymorphisms, which are not yet under investigation. Other “hits” that can lead to pancreatic cell death include fibrosis of the pancreas, extreme alcohol intake, viral infections, iron overload, medications, or an obstruction of the pancreatic duct.<sup>42</sup>

## Clinical Associations of NAFPD

### Nonalcoholic Fatty Liver Disease

Nonalcoholic fatty liver disease (NAFLD) occurs when hepatocytes are infiltrated with fat, in the absence of excessive alcohol consumption.<sup>75</sup> As mentioned above, obesity is the main causative risk factor, not only for NAFPD but also for NAFLD,<sup>75</sup> making these two clinical syndromes closely related. In general, fat deposition is linked to multiple diseases like metabolic syndrome, diabetes, and atherosclerosis.<sup>2,76</sup>

In a large cross-sectional study where NAFPD and its risk factors were examined, authors compared pancreatic echogenicity with that of the liver and demonstrated that fatty pancreas and fatty liver composition were found in combination in almost 26% of patients, whereas fatty pancreas alone was found in 16%.<sup>11</sup> In a prospective study almost 68% of the included patients with fatty pancreas presented with fatty liver, but 97% of patients initially diagnosed with fatty liver presented concurrently with a fatty pancreas, making fatty liver a strong predictive factor of fatty pancreas disease.<sup>8</sup> In a Chinese study, 67% of non-diabetic subjects with fatty pancreas had simultaneous fatty liver changes.<sup>8</sup> In this study, fatty liver was an independent risk factor for NAF-PD.<sup>8</sup> Likewise, in another analysis among 60 patients with hepatic steatosis, 57% were found with NAFPD.<sup>12</sup> The amount of fat in liver tissue is significantly correlated with pancreatic fat in numerous other studies,<sup>77-79</sup> signifying that both clinical entities share similar risk factors, pathogenetic mechanisms and disease progression pathways. In a clinical trial including 121 obese/overweight children and adolescents, 48% of those with confirmed fatty liver disease of any severity exhibited NAFPD, while, in subjects with steatohepatitis the percentage of NAFPD increased to 80%.<sup>78</sup>

On the contrary, some studies did not manage to find an association between pancreatic fat and liver fat.<sup>4,17,72,80</sup> In a study of subjects with and without diabetes,<sup>17</sup> those with diabetes had higher liver fat content, but there was no statistically significant association between liver and pancreatic fat content. These contradictory results focusing on the correlation between NAFLD and NAFPD might imply that, despite that both syndromes share similar pathological mechanisms, organ fat accumulation may be distributed differently due to unknown mechanisms.

It is worth mentioning that, based on recent evidence, the association between NAFPD and NAFLD is also mediated via a systemic autoimmune comorbidity, namely psoriasis. Psoriasis is well associated with metabolic disturbances, including obesity, metabolic syndrome and type 2 diabetes, all of which consist risk factors for both NAFPD and NAFLD.<sup>81-83</sup> In psoriatic patients, chronic inflammation, mediated by pro-inflammatory cytokines and adipokines (especially TNF- $\alpha$ ), drives insulin resistance and plays a crucial role not only in hepatic steatosis but potentially also in pancreatic steatosis.<sup>84</sup> Additionally, it has been suggested that psoriasis is associated with pancreatic diseases, such as pancreatitis and pancreatic cancer, as evident in recent published data.<sup>85</sup>

### Obesity and Metabolic Syndrome

Obesity is a multifactorial clinical entity that can lead to the excessive and pathological accumulation of adipose tissue in non-adipose tissues such as the pancreas, liver, heart, skeletal muscle and is associated with diabetes mellitus, cardiovascular disease, hypertension, and hyperlipidemia.<sup>86-88</sup> Since the term NAFPD describes the accumulation of fat into the pancreas, obesity is the main driver of this condition.<sup>75</sup> A large number of studies link central obesity and/or increased BMI to fatty deposition of the pancreas.<sup>9,10,20</sup>

Metabolic syndrome is defined as the presence of at least three of the following risk factors: high blood glucose, increased levels of triglycerides, low levels of HDL cholesterol, large waist circumference and hypertension.<sup>89</sup> Many

studies focusing on patients' metabolic abnormalities try to correlate these with the existence of fatty pancreas. In 2009, Lee et al. studied 293 individuals from an obesity clinic with a confirmed diagnosis of pancreatic fat accumulation of various severity.<sup>90</sup> More than half of the patients (61.4%) were diagnosed with fatty pancreas. Results positively correlated pancreatic fat infiltration with metabolic risk factors, such as increased visceral fat and waist circumference, triglycerides, insulin resistance, correlating fatty pancreas with metabolic syndrome. Similarly, a case-control study showed that fatty pancreas was strongly associated with hypertension, increased blood glucose levels, central obesity, and increased triglycerides, but reported no association with HDL values.<sup>91</sup> A more recent study that systematically reviewed and analyzed data from 2675 individuals found that approximately 33% of them had NAFLD.<sup>20</sup> Among individuals with NAFLD, 44% of them met the criteria for metabolic syndrome.<sup>20</sup> In a prospective study by Sepe et al, fatty pancreas was observed in 27.8% of participants that underwent EUS for various purposes.<sup>10</sup> Notably, the presence of any component of metabolic syndrome was associated with a significant 37% increase in the prevalence of fatty pancreas. Individuals who fulfilled the criteria for metabolic syndrome had a 40% prevalence of fatty pancreas, rendering metabolic syndrome a strong predictive factor of NAFLD. Interestingly, a high-fat diet is associated with triglyceride accumulation mainly in pancreatic cells, rather than hepatocytes, making pancreatic steatosis an early consequence of metabolic syndrome.<sup>3</sup>

On the other hand, Wong et al,<sup>9</sup> showed that, among all metabolic syndrome components, only central obesity and hypertriglyceridemia were independent risk factors of NAFLD. A Turkish clinical trial also reported an association of fatty pancreas with metabolic syndrome, which was influenced, however, by other factors such as BMI and NAFLD.<sup>92</sup>

## Insulin Resistance - Prediabetes - Diabetes

Pancreas has an essential role in regulating glucose metabolism and glandular secretion under various metabolic circumstances. The pathogenesis of type 2 diabetes is characterized by peripheral insulin resistance and impaired glucose secretion, gradually leading to a decline in  $\beta$ -pancreatic cells and, later, to their gradual destruction. A disproportionate accumulation of visceral adipose tissue is a strong risk factor for insulin resistance.<sup>93-95</sup> However, study results remain controversial for the association between fatty pancreas and diabetes mellitus.

In numerous animal studies, islet cell abnormalities are related to pancreatic lipomatosis, and result in increased blood glucose levels.<sup>48,96</sup> A human study reported an almost 12% incidence of diabetes in the NAFLD cohort versus a 5% incidence in the non-NAFLD group. NAFLD patients had an 108% increased risk of diabetes mellitus. Pancreatic fat was also found to be negatively associated with insulin secretion in other analyses,<sup>72</sup> as well as with hyperglycemia and occurrence of diabetes.<sup>17,72,97</sup> Of note, individuals presenting with both fatty pancreas and fatty liver, had the highest levels of insulin resistance compared to those with either condition alone.<sup>9</sup>

Another analysis indicated that fatty pancreas at baseline is associated with an increased incidence of diabetes. However, this finding was not significant in multivariate analyses.<sup>98</sup> This was in line with results from a large cohort study, where, despite the high frequency of fatty pancreas in diabetes, their association did not reach statistical significance.<sup>11</sup> Other investigations did not support the association between fatty pancreas and diabetes mellitus or impairments in glucose levels as well.<sup>10,90</sup>

Conclusively, pancreatic steatosis may be an early event in the pathogenesis of diabetes,<sup>99</sup> but evidence pointing to an independent association is limited.

## Atherosclerosis

NAFLD has been linked with atherosclerosis risk factors including advanced age, BMI, hypertension, and hyperlipidemia.<sup>9,28,72,90,91</sup> The effect of fatty pancreas on glucose and insulin metabolism, chronic low-grade inflammation and chronic stress may increase cardiovascular disease (CVD) risk.<sup>100</sup> This is amplified by the accumulation of lipids in coronary and peripheral blood vessels.

Carotid intima-media thickness (IMT) and carotid-femoral pulse wave velocity (cf-PWV) are widely used as indicators of subclinical atherosclerosis by detecting early functional and structural changes of the vascular wall.<sup>101,102</sup> It has been shown that NAFLD, especially in advanced stages, is correlated with increased risk of atherosclerosis and, potentially, cardiovascular events.<sup>103-107</sup> However, it remains to be answered whether fatty pancreas is a predictive CVD factor. A study that included patients with biopsy-proven NAFLD and healthy controls showed that patients with fatty

pancreas had significantly higher cf-PWV levels, whereas no association was found with carotid IMT. The link between cf-PWV/carotid IMT levels and fatty pancreas, nevertheless, was not observed upon adjusting for relevant variables.<sup>92</sup> In another case–control study, epicardial adipose tissue and aortic IMT were higher among NAFLD subjects, compared to those without NAFLD, while both epicardial adipose tissue and aortic IMT were independent risk factors of subclinical atherosclerosis.<sup>108</sup>

## Pancreatitis

Chronic high-fat dietary intake is associated with inter- as well as intra-pancreatic fat accumulation. As such, NAFLD progression may lead to chronic pancreatitis, and this association is expressed by the term “non-alcoholic steatopancreatitis” (NASP).<sup>1,109</sup> In favor of this hypothesis, a study showed that obese mice are more likely to develop severe forms of pancreatitis than lean ones.<sup>110</sup> Additionally, in a review by Smith et al, preliminary data link acute pancreatitis with NASP.<sup>7</sup> According to the authors, the uneven secretion of proinflammatory cytokines creates an inflammatory environment which promotes bouts of acute pancreatitis.<sup>7</sup> Finally, a recent pilot study of 189 subjects indicated that fatty pancreas is a risk factor of acute pancreatitis.<sup>111</sup>

## Pancreatic Cancer

Obesity is strongly associated with pancreatic steatosis and inconsistently associated with pancreatic cancer; this raises concerns for a potential relationship between fatty pancreas and pancreatic cancer.

A meta-analysis of 4400 patients showed a positive association between increased BMI and risk of pancreatic cancer in both sexes, while central obesity was associated with an increased risk of pancreatic cancer only in women.<sup>112</sup> It has been described that NAFLD can lead to NASP and sequentially to pancreatic cancer.<sup>113</sup> In a study of 102 cases of pancreatic ductal adenocarcinoma, fatty pancreas was significantly higher in cases compared to healthy individuals.<sup>114</sup> Findings of a retrospective cross-sectional study analyzing data from patients with pancreatic carcinoma further supported the role of fatty pancreas in the development of pancreatic cancer.<sup>115</sup> According to Lesmana et al, NAFLD is the only significant risk factor when it comes to pancreatic cancer, while factors like age, gender, diabetes, and pancreatitis failed to show a causative role.<sup>113</sup> Indeed, in a case–control study of patients who underwent total resection due to pancreatic adenocarcinoma, patients with fatty pancreas had higher mortality.<sup>116</sup>

At present, however, there is not enough evidence to decisively support the relationship between NAFLD and pancreatic cancer, and several studies have failed to detect an association. A study by Sepe et al did not find an association between fatty pancreas and pancreatic malignancy.<sup>10</sup> Wu et al also found no association between pancreatic carcinoma markers and NAFLD.<sup>91</sup>

## Pathogenetic Mechanisms and Clinical Entities Associated with NAFLD

The main risk factors associated with NAFLD are obesity and metabolic syndrome (including dyslipidemia), while alcohol, viruses, iron deposition, drugs and pancreatic duct obstruction represent potential secondary hits which participate in cell death and replacement of pancreatic tissue with fat.<sup>41,90,91,100</sup> The main pathogenetic mechanism of NAFLD is fatty replacement, followed by fatty accumulation (intra- or inter-lobular) and, ultimately,  $\beta$ -cell dysfunction.<sup>21</sup> Mechanisms associated with fat accumulation include reduction in mitochondrial activity, inflammatory cell infiltration with production of inflammatory cytokines (eg,  $\text{TNF}\alpha$ , IL-1 $\beta$ , IL-6, MCP-1), activation of stellate cells, hormonal imbalance (with perturbed leptin and adiponectin levels, overexpression of ghrelin), as well as STK-25 pathway overexpression, which further exacerbate all the aforementioned mechanisms.<sup>34,70</sup> B-cell dysfunction, as well as hormonal imbalance, in turn, can lead to insulin resistance, prediabetes and diabetes, while reduction in mitochondrial activity, inflammatory cell infiltration and activation of stellate cells, can lead to pancreatitis and pancreatic cancer.

## Treatment of NAFLD

Despite the growing prevalence of NAFLD, its understanding remains limited and effective management strategies are still being explored. The effectiveness of current treatments for NAFLD is limited, as there is no approved pharmacologic

therapy. To date, the most effective treatments are lifestyle modifications, including dietary changes, regular exercise, and weight-management programs.<sup>117</sup> Calorie restriction has been suggested as the potential means of normalizing  $\beta$ -cell function and reducing pancreatic fat content.

Troglitazone, a member of the thiazolidinedione class with insulin-sensitizing and anti-inflammatory properties<sup>118</sup> which has been withdrawn from the market,<sup>119</sup> has been implicated in NAFLD. Specifically, animal models have shown that troglitazone can reverse changes in tissue structure such as fibrosis and fatty accumulation, while reducing infiltration of inflammatory cells.<sup>120</sup> However, this effect has not been assessed by clinical trials. Glucagon-like peptide-1 (GLP-1) agonists have the potential to lessen the severity of NAFLD, by modulating the ER stress pathway and the subsequent signaling of apoptosis in mice models, and to alter regional adiposity in humans.<sup>121,122</sup> Moreover, the synergistic activity of sitagliptin, a dipeptidyl peptidase (DPP) 4 inhibitor which prolongs the action of GLP-1 and insulin secretion from pancreatic  $\beta$ -cells, with telmisartan, which modulates inflammatory responses and improves endothelial function, exhibits remarkable efficacy in effectively managing the progression of NAFLD in mice.<sup>123</sup> Based on evidence from experimental models, GLP-1 agonists consist the most promising class of drugs for the treatment of NAFLD<sup>122</sup> and prospective trials are needed to verify these data.

It has also demonstrated that bariatric surgery can lead to significant reductions in fat accumulation within the pancreas.<sup>124</sup> Nonetheless, education regarding healthy dietary habits and regular exercise should be the mainstay for reducing the NAFLD risk, as well as for treating those already diagnosed with it.

## Nomenclature of NAFLD

The term “non-alcoholic fatty pancreas disease” is misleading, since many subjects with obesity may simultaneously consume alcohol, while it may also be stigmatizing. Similar to a recent proposal for NAFLD nomenclature,<sup>125–127</sup> we propose the term “Steatotic Pancreatic Disease” (SPD) in order to generally describe the accumulation of fat within pancreatic tissue, and the term “Metabolic dysfunction-associated Steatotic Pancreatic Disease” (MASPD) for describing the accumulation of fat within pancreatic tissue when this is associated with obesity or the metabolic syndrome.

## Conclusion

NAFLD has been associated with a number of clinical entities, including metabolic syndrome, diabetes mellitus and atherosclerosis, but also with pancreatitis and pancreatic cancer. Similar clinical associations are observed in NAFLD. However, due to the scarcity of data on NAFLD, further studies are awaited to better determine related biological processes and metabolic sequelae.

## Disclosure

Theodosios D. Filippatos reports participation in advisory boards for Lilly and lecture honoraria from Boehringer Ingelheim, Mylan, Astra Zeneca, Lilly, Recordati, Bausch Health, Servier, Viartis, Omega-Pharma and Innovis; personal fees from Perrigo, Innovis, and Servier, outside the submitted work. The remaining authors, Antonia Pagkali, Anastasios Makris, Kalliopi Brofidi, and Aris P. Agouridis have no conflict of interest to declare.

## References

1. Mathur A, Marine M, Lu D, et al. Nonalcoholic fatty pancreas disease. *HPB*. 2007;9(4):312–318. doi:10.1080/13651820701504157
2. Filippatos TD. Non-Alcoholic Fatty Pancreas Disease: a Diagnosis of Increasing Importance. *Angiology*. 2022;73(6):495–496. doi:10.1177/00033197211053902
3. Pinnick KE, Collins SC, Lontos C, Gauguier D, Clark A, Fielding BA. Pancreatic ectopic fat is characterized by adipocyte infiltration and altered lipid composition. *Obesity (Silver Spring)*. 2008;16(3):522–530. doi:10.1038/oby.2007.110
4. Kovanlikaya A, Mittelman SD, Ward A, Geffner ME, Dorey F, Gilsanz V. Obesity and fat quantification in lean tissues using three-point Dixon MR imaging. *Pediatr Radiol*. 2005;35(6):601–607. doi:10.1007/s00247-005-1413-y
5. Ogilvie RF. The islands of Langerhans in 19 cases of obesity. *J Pathol Bacteriol*. 1933;37(3):473–481. doi:10.1002/path.1700370314
6. Paul J, Shihaz AVH. Pancreatic Steatosis: a New Diagnosis and Therapeutic Challenge in Gastroenterology. *Arq Gastroenterol*. 2020;57(2):216–220. doi:10.1590/s0004-2803.202000000-27
7. Smits MM, van Geenen EJ. The clinical significance of pancreatic steatosis. *Nat Rev Gastroenterol Hepatol*. 2011;8(3):169–177. doi:10.1038/nrgastro.2011.4



8. Wang CY, Ou HY, Chen MF, Chang TC, Chang CJ. Enigmatic ectopic fat: prevalence of nonalcoholic fatty pancreas disease and its associated factors in a Chinese population. *J Am Heart Assoc.* 2014;3(1):e000297. doi:10.1161/JAHA.113.000297
9. Wong VW, Wong GL, Yeung DK, et al. Fatty pancreas, insulin resistance, and beta-cell function: a population study using fat-water magnetic resonance imaging. *Am J Gastroenterol.* 2014;109(4):589–597. doi:10.1038/ajg.2014.1
10. Sepe PS, Ohri A, Sanaka S, et al. A prospective evaluation of fatty pancreas by using EUS. *Gastrointest Endosc.* 2011;73(5):987–993. doi:10.1016/j.gie.2011.01.015
11. Lesmana CR, Pakasi LS, Inggriani S, Aidawati ML, Lesmana LA. Prevalence of Non-Alcoholic Fatty Pancreas Disease (NAFPD) and its risk factors among adult medical check-up patients in a private hospital: a large cross sectional study. *BMC Gastroenterol.* 2015;15:174. doi:10.1186/s12876-015-0404-1
12. Al-Haddad M, Khashab M, Zyromski N, et al. Risk factors for hyperechogenic pancreas on endoscopic ultrasound: a case-control study. *Pancreas.* 2009;38(6):672–675. doi:10.1097/MPA.0b013e3181a9d5af
13. Muftah AA, Pecha RL, Riojas Barrett M, et al. Pancreatic parenchymal changes seen on endoscopic ultrasound are dynamic in the setting of fatty pancreas: a short-term follow-up study. *Pancreatol.* 2022;22(8):1187–1194. doi:10.1016/j.pan.2022.10.006
14. Ahbab S, Ünsal A, Ataoğlu HE, Can TS, Kayaş D, Savaş Y. Prediabetes and Type 2 Diabetes are Independent Risk Factors for Computed Tomography-Estimated Nonalcoholic Fatty Pancreas Disease. *Clinics.* 2019;74:e1337. doi:10.6061/clinics/2019/e1337
15. Li J, Xie Y, Yuan F, Song B, Tang C. Noninvasive quantification of pancreatic fat in healthy male population using chemical shift magnetic resonance imaging: effect of aging on pancreatic fat content. *Pancreas.* 2011;40(2):295–299. doi:10.1097/MPA.0b013e318201669f
16. Patel NS, Peterson MR, Brenner DA, Heba E, Sirlin C, Loomba R. Association between novel MRI-estimated pancreatic fat and liver histology-determined steatosis and fibrosis in non-alcoholic fatty liver disease. *Aliment Pharmacol Ther.* 2013;37(6):630–639. doi:10.1111/apt.12237
17. Tushuizen ME, Bunck MC, Pouwels PJ, et al. Pancreatic fat content and beta-cell function in men with and without type 2 diabetes. *Diabetes Care.* 2007;30(11):2916–2921. doi:10.2337/dc07-0326
18. Ma J. Dixon techniques for water and fat imaging. *J Magn Reson Imaging.* 2008;28(3):543–558. doi:10.1002/jmri.21492
19. Reeder SB, Pineda AR, Wen Z, et al. Iterative decomposition of water and fat with echo asymmetry and least-squares estimation (IDEAL): application with fast spin-echo imaging. *Magn Reson Med.* 2005;54(3):636–644. doi:10.1002/mrm.20624
20. Singh RG, Yoon HD, Wu LM, Lu J, Plank LD, Petrov MS. Ectopic fat accumulation in the pancreas and its clinical relevance: a systematic review, meta-analysis, and meta-regression. *Metabolism.* 2017;69:1–13. doi:10.1016/j.metabol.2016.12.012
21. Catanzaro R, Cuffari B, Italia A, Marotta F. Exploring the metabolic syndrome: nonalcoholic fatty pancreas disease. *World J Gastroenterol.* 2016;22(34):7660–7675. doi:10.3748/wjg.v22.i34.7660
22. Rugivarodom M, Geeratragoon T, Pausawasdi N, Charatcharoenwitthaya P. Fatty Pancreas: linking Pancreas Pathophysiology to Nonalcoholic Fatty Liver Exploring the metabolic syndrome: nonalcoholic fatty pancreas disease Disease. *J Clin Transl Hepatol.* 2022;10(6):1229–1239. doi:10.14218/JCTH.2022.00085
23. Guglielmi V, Sbraccia P. Type 2 diabetes: does pancreatic fat really matter? *Diabetes Metab Res Rev.* 2018;34(2). doi:10.1002/dmrr.2955
24. Tannenbaum BM, Brindley DN, Tannenbaum GS, Dallman MF, McArthur MD, Meaney MJ. High-fat feeding alters both basal and stress-induced hypothalamic-pituitary-adrenal activity in the rat. *Am J Physiol.* 1997;273(6):E1168–1177. doi:10.1152/ajpendo.1997.273.6.E1168
25. Kyrou I, Tsigos C. Stress mechanisms and metabolic complications. *Horm Metab Res.* 2007;39(6):430–438. doi:10.1055/s-2007-981462
26. Fraulob JC, Ogg-Diamantino R, Fernandes-Santos C, Aguila MB, Mandarim-de-Lacerda CA. A Mouse Model of Metabolic Syndrome: insulin Resistance, Fatty Liver and Non-Alcoholic Fatty Pancreas Disease (NAFPD) in C57BL/6 Mice Fed a High Fat Diet. *J Clin Biochem Nutr.* 2010;46(3):212–223. doi:10.3164/jcbrn.09-83
27. Zhang X, Cui Y, Fang L, Li F. Chronic high-fat diets induce oxidative injuries and fibrogenesis of pancreatic cells in rats. *Pancreas.* 2008;37(3):e31–38. doi:10.1097/MPA.0b013e3181744b50
28. Filippatos TD, Alexakis K, Mavrikaki V, Mikhailidis DP. Nonalcoholic Fatty Pancreas Disease: role in Metabolic Syndrome, “Prediabetes”, Diabetes and Atherosclerosis. *Dig Dis Sci.* 2022;67(1):26–41. doi:10.1007/s10620-021-06824-7
29. Yoshikawa H, Tajiri Y, Sako Y, Hashimoto T, Umeda F, Nawata H. Effects of free fatty acids on beta-cell functions: a possible involvement of peroxisome proliferator-activated receptors alpha or pancreatic/duodenal homeobox. *Metabolism.* 2001;50(5):613–618. doi:10.1053/meta.2001.22565
30. Walz HA, Harndahl L, Wierup N, et al. Early and rapid development of insulin resistance, islet dysfunction and glucose intolerance after high-fat feeding in mice overexpressing phosphodiesterase 3B. *J Endocrinol.* 2006;189(3):629–641. doi:10.1677/joe.1.06522
31. Houten SM, Wanders RJ. A general introduction to the biochemistry of mitochondrial fatty acid beta-oxidation. *J Inherit Metab Dis.* 2010;33(5):469–477. doi:10.1007/s10545-010-9061-2
32. Juge-Aubry CE, Henrichot E, Meier CA. Adipose tissue: a regulator of inflammation. *Best Pract Res Clin Endocrinol Metab.* 2005;19(4):547–566. doi:10.1016/j.beem.2005.07.009
33. Greenberg AS, Obin MS. Obesity and the role of adipose tissue in inflammation and metabolism. *Am J Clin Nutr.* 2006;83(2):461S–465S. doi:10.1093/ajcn/83.2.461S
34. Matsuda A, Makino N, Tozawa T, et al. Pancreatic fat accumulation, fibrosis, and acinar cell injury in the Zucker diabetic fatty rat fed a chronic high-fat diet. *Pancreas.* 2014;43(5):735–743. doi:10.1097/MPA.000000000000129
35. Horii T, Kozawa J, Fujita Y, et al. Lipid droplet accumulation in beta cells in patients with type 2 diabetes is associated with insulin resistance, hyperglycemia and beta cell dysfunction involving decreased insulin granules. *Front Endocrinol.* 2022;13:996716. doi:10.3389/fendo.2022.996716
36. Petrov MS, Taylor R. Intra-pancreatic fat deposition: bringing hidden fat to the fore. *Nat Rev Gastroenterol Hepatol.* 2022;19(3):153–168. doi:10.1038/s41575-021-00551-0
37. Altinel D, Basturk O, Sarmiento JM, et al. Lipomatous pseudohypertrophy of the pancreas: a clinicopathologically distinct entity. *Pancreas.* 2010;39(3):392–397. doi:10.1097/MPA.0b013e3181bd2923
38. Yan MX, Li YQ, Meng M, Ren HB, Kou Y. Long-term high-fat diet induces pancreatic injuries via pancreatic microcirculatory disturbances and oxidative stress in rats with hyperlipidemia. *Biochem Biophys Res Commun.* 2006;347(1):192–199. doi:10.1016/j.bbrc.2006.06.063
39. Lee Y, Lingvay I, Szczepaniak LS, Ravazzola M, Orci L, Unger RH. Pancreatic steatosis: harbinger of type 2 diabetes in obese rodents. *Int J Obes Lond.* 2010;34(2):396–400. doi:10.1038/ijo.2009.245
40. Lopez JM, Bombi JA, Valderrama R, et al. Effects of prolonged ethanol intake and malnutrition on rat pancreas. *Gut.* 1996;38(2):285–292. doi:10.1136/gut.38.2.285

41. Chang ML. Fatty Pancreas-Centered Metabolic Basis of Pancreatic Adenocarcinoma: from Obesity, Diabetes and Pancreatitis to Oncogenesis. *Biomedicines*. 2022;10(3). doi:10.3390/biomedicines10030692
42. Silva L, Fernandes MSS, Lima EA, Stefano JT, Oliveira CP, Jukemura J. Fatty Pancreas: disease or Finding? *Clinics (Sao Paulo)*. 2021;76:e2439. doi:10.6061/clinics/2021/e2439
43. Patel S, Bellon EM, Haaga J, Park CH. Fat replacement of the exocrine pancreas. *AJR Am J Roentgenol*. 1980;135(4):843–845. doi:10.2214/ajr.135.4.843
44. den Boer M, Voshol PJ, Kuipers F, Havekes LM, Romijn JA. Hepatic steatosis: a mediator of the metabolic syndrome. Lessons from animal models. *Arterioscler Thromb Vasc Biol*. 2004;24(4):644–649. doi:10.1161/01.ATV.0000116217.57583.6e
45. Nunez-Duran E, Chanclon B, Sutt S, et al. Protein kinase STK25 aggravates the severity of non-alcoholic fatty pancreas disease in mice. *J Endocrinol*. 2017;234(1):15–27. doi:10.1530/JOE-17-0018
46. Pinte L, Balaban DV, Baicus C, Jinga M. Non-alcoholic fatty pancreas disease - practices for clinicians. *Rom J Intern Med*. 2019;57(3):209–219. doi:10.2478/rjim-2019-0005
47. Nascimento FA, Barbosa-da-silva S, Fernandes-Santos C, Mandarim-de-Lacerda CA, Aguila MB. Adipose tissue, liver and pancreas structural alterations in C57BL/6 mice fed high-fat-high-sucrose diet supplemented with fish oil (n-3 fatty acid rich oil). *Exp Toxicol Pathol*. 2010;62(1):17–25. doi:10.1016/j.etp.2008.12.008
48. Lee Y, Hirose H, Ohneda M, Johnson JH, McGarry JD, Unger RH. Beta-cell lipotoxicity in the pathogenesis of non-insulin-dependent diabetes mellitus of obese rats: impairment in adipocyte-beta-cell relationships. *Proc Natl Acad Sci U S A*. 1994;91(23):10878–10882. doi:10.1073/pnas.91.23.10878
49. Chitturi S, Abeygunasekera S, Farrell GC, et al. NASH and insulin resistance: insulin hypersecretion and specific association with the insulin resistance syndrome. *Hepatology*. 2002;35(2):373–379. doi:10.1053/jhep.2002.30692
50. Bollheimer LC, Skelly RH, Chester MW, McGarry JD, Rhodes CJ. Chronic exposure to free fatty acid reduces pancreatic beta cell insulin content by increasing basal insulin secretion that is not compensated for by a corresponding increase in proinsulin biosynthesis translation. *J Clin Invest*. 1998;101(5):1094–1101. doi:10.1172/JCI420
51. Lin M, Weng SY, Chai KF, Mao ZJ. Lipidomics as a tool of predicting progression from non-alcoholic fatty pancreas disease to type 2 diabetes mellitus. *RSC Adv*. 2019;9(71):41419–41430. doi:10.1039/c9ra07071k
52. Bonora E. The metabolic syndrome and cardiovascular disease. *Ann Med*. 2006;38(1):64–80. doi:10.1080/07853890500401234
53. Yu TY, Wang CY. Impact of non-alcoholic fatty pancreas disease on glucose metabolism. *J Diabetes Investig*. 2017;8(6):735–747. doi:10.1111/jdi.12665
54. Fernandes-Santos C, Evangelista Carneiro R, de Souza Mendonca L, Barbosa Aguila M, Mandarim-de-Lacerda CA. Rosiglitazone aggravates nonalcoholic Fatty pancreatic disease in C57BL/6 mice fed high-fat and high-sucrose diet. *Pancreas*. 2009;38(3):e80–86. doi:10.1097/MPA.0b013e3181987d9d
55. Fields M, Lewis CG. Impaired endocrine and exocrine pancreatic functions in copper-deficient rats: the effect of gender. *J Am Coll Nutr*. 1997;16(4):346–351. doi:10.1080/07315724.1997.10718696
56. Tarantino G, Porcu C, Arciello M, Andreozzi P, Balsano C. Prediction of carotid intima-media thickness in obese patients with low prevalence of comorbidities by serum copper bioavailability. *J Gastroenterol Hepatol*. 2018;33(8):1511–1517. doi:10.1111/jgh.14104
57. Walters MN. Adipose atrophy of the exocrine pancreas. *J Pathol Bacteriol*. 1966;92(2):547–557. doi:10.1002/path.1700920232
58. de Wilde J, Mohren R, van den Berg S, et al. Short-term high fat-feeding results in morphological and metabolic adaptations in the skeletal muscle of C57BL/6J mice. *Physiol Genomics*. 2008;32(3):360–369. doi:10.1152/physiolgenomics.00219.2007
59. Emery MG, Fisher JM, Chien JY, et al. CYP2E1 activity before and after weight loss in morbidly obese subjects with nonalcoholic fatty liver disease. *Hepatology*. 2003;38(2):428–435. doi:10.1053/jhep.2003.50342
60. Chtioui H, Semela D, Ledermann M, Zimmermann A, Dufour JF. Expression and activity of the cytochrome P450 2E1 in patients with nonalcoholic steatosis and steatohepatitis. *Liver Int*. 2007;27(6):764–771. doi:10.1111/j.1478-3231.2007.01524.x
61. Ikejima K, Okumura K, Kon K, Takei Y, Sato N. Role of adipocytokines in hepatic fibrogenesis. *J Gastroenterol Hepatol*. 2007;22(Suppl 1):S87–92. doi:10.1111/j.1440-1746.2007.04961.x
62. Nerstedt A, Cansby E, Andersson CX, et al. Serine/threonine protein kinase 25 (STK25): a novel negative regulator of lipid and glucose metabolism in rodent and human skeletal muscle. *Diabetologia*. 2012;55(6):1797–1807. doi:10.1007/s00125-012-2511-7
63. Chursa U, Nunez-Duran E, Cansby E, et al. Overexpression of protein kinase STK25 in mice exacerbates ectopic lipid accumulation, mitochondrial dysfunction and insulin resistance in skeletal muscle. *Diabetologia*. 2017;60(3):553–567. doi:10.1007/s00125-016-4171-5
64. Amrutkar M, Cansby E, Chursa U, et al. Genetic Disruption of Protein Kinase STK25 Ameliorates Metabolic Defects in a Diet-Induced Type 2 Diabetes Model. *Diabetes*. 2015;64(8):2791–2804. doi:10.2337/db15-0060
65. Amrutkar M, Cansby E, Nunez-Duran E, et al. Protein kinase STK25 regulates hepatic lipid partitioning and progression of liver steatosis and NASH. *FASEB J*. 2015;29(4):1564–1576. doi:10.1096/fj.14-264937
66. Amrutkar M, Chursa U, Kern M, et al. STK25 is a critical determinant in nonalcoholic steatohepatitis. *FASEB J*. 2016;30(10):3628–3643. doi:10.1096/fj.201600562R
67. Munzberg H, Flier JS, Bjorbaek C. Region-specific leptin resistance within the hypothalamus of diet-induced obese mice. *Endocrinology*. 2004;145(11):4880–4889. doi:10.1210/en.2004-0726
68. Akagiri S, Naito Y, Ichikawa H, et al. A Mouse Model of Metabolic Syndrome; Increase in Visceral Adipose Tissue Precedes the Development of Fatty Liver and Insulin Resistance in High-Fat Diet-Fed Male KK/Ta Mice. *J Clin Biochem Nutr*. 2008;42(2):150–157. doi:10.3164/jcbs.2008022
69. Al-Ani Z, Ko J, Petrov MS. Intra-pancreatic fat deposition across the pancreatitis spectrum and the influence of gut hormones. *Dig Liver Dis*. 2023;55(8):1081–1090. doi:10.1016/j.dld.2023.02.013
70. Finelli C, Padula MC, Martelli G, Tarantino G. Could the improvement of obesity-related co-morbidities depend on modified gut hormones secretion? *World J Gastroenterol*. 2014;20(44):16649–16664. doi:10.3748/wjg.v20.i44.16649
71. Begovatz P, Koliaki C, Weber K, et al. Pancreatic adipose tissue infiltration, parenchymal steatosis and beta cell function in humans. *Diabetologia*. 2015;58(7):1646–1655. doi:10.1007/s00125-015-3544-5

72. Heni M, Machann J, Staiger H, et al. Pancreatic fat is negatively associated with insulin secretion in individuals with impaired fasting glucose and/or impaired glucose tolerance: a nuclear magnetic resonance study. *Diabetes Metab Res Rev*. 2010;26(3):200–205. doi:10.1002/dmrr.1073
73. Ozturk K, Uygun A, Guler AK, et al. Nonalcoholic fatty liver disease is an independent risk factor for atherosclerosis in young adult men. *Atherosclerosis*. 2015;240(2):380–386. doi:10.1016/j.atherosclerosis.2015.04.009
74. Li YX, Sang YQ, Sun Y, et al. Pancreatic Fat is not significantly correlated with beta-cell Dysfunction in Patients with new-onset Type 2 Diabetes Mellitus using quantitative Computed Tomography. *Int J Med Sci*. 2020;17(12):1673–1682. doi:10.7150/ijms.46395
75. van Geenen EJ, Smits MM, Schreuder TC, van der Peet DL, Bloemena E, Mulder CJ. Nonalcoholic fatty liver disease is related to nonalcoholic fatty pancreas disease. *Pancreas*. 2010;39(8):1185–1190. doi:10.1097/MPA.0b013e3181f6fce2
76. Wu S, Wu F, Ding Y, Hou J, Bi J, Zhang Z. Association of non-alcoholic fatty liver disease with major adverse cardiovascular events: a systematic review and meta-analysis. *Sci Rep*. 2016;6:33386. doi:10.1038/srep33386
77. Sijens PE, Edens MA, Bakker SJ, Stolk RP. MRI-determined fat content of human liver, pancreas and kidney. *World J Gastroenterol*. 2010;16(16):1993–1998. doi:10.3748/wjg.v16.i16.1993
78. Della Corte C, Mosca A, Majo F, et al. Nonalcoholic fatty pancreas disease and Nonalcoholic fatty liver disease: more than ectopic fat. *Clin Endocrinol (Oxf)*. 2015;83(5):656–662. doi:10.1111/cen.12862
79. Patel NS, Peterson MR, Lin GY, et al. Insulin Resistance Increases MRI-Estimated Pancreatic Fat in Nonalcoholic Fatty Liver Disease and Normal Controls. *Gastroenterol Res Pract*. 2013;2013:498296. doi:10.1155/2013/498296
80. Schwenzer NF, Machann J, Martirosian P, et al. Quantification of pancreatic lipomatosis and liver steatosis by MRI: comparison of in/opposed-phase and spectral-spatial excitation techniques. *Invest Radiol*. 2008;43(5):330–337. doi:10.1097/RLI.0b013e31816a88c6
81. Rathod A, Neema S, Radhakrishnan S, Vendhan S, Tripathy DM, Vasudevan B. Palmoplantar Plaque Psoriasis is Associated with Diabetes, Hypertension, Obesity, and Metabolic Syndrome-A Case-Control Study. *Indian Dermatol Online J*. 2022;13(5):606–610. doi:10.4103/idoj.idoj\_59\_22
82. Jensen P, Skov L. Psoriasis and Obesity. *Dermatology*. 2016;232(6):633–639. doi:10.1159/000455840
83. Hao Y, Zhu YJ, Zou S, et al. Metabolic Syndrome and Psoriasis: mechanisms and Future Directions. *Front Immunol*. 2021;12:711060. doi:10.3389/fimmu.2021.711060
84. Balato N, Napolitano M, Ayala F, Patruno C, Megna M, Tarantino G. Nonalcoholic fatty liver disease, spleen and psoriasis: new aspects of low-grade chronic inflammation. *World J Gastroenterol*. 2015;21(22):6892–6897. doi:10.3748/wjg.v21.i22.6892
85. Bialecka A. Evaluation of pancreatic function in patients with plaque psoriasis. *Postepy Dermatol Alergol*. 2023;40(3):372–376. doi:10.5114/ada.2023.129400
86. Haslam DW, James WP. Obesity. *Lancet*. 2005;366(9492):1197–1209. doi:10.1016/S0140-6736(05)67483-1
87. Van Gaal LF, Mertens IL, De Block CE. Mechanisms linking obesity with cardiovascular disease. *Nature*. 2006;444(7121):875–880. doi:10.1038/nature05487
88. Panuganti KK, Nguyen M, Kshirsagar RK. Obesity. *Statpearls, Treasure Island*. 2022.
89. Grundy SM, Cleeman JI, Daniels SR, et al. Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation*. 2005;112(17):2735–2752. doi:10.1161/CIRCULATIONAHA.105.169404
90. Lee JS, Kim SH, Jun DW, et al. Clinical implications of fatty pancreas: correlations between fatty pancreas and metabolic syndrome. *World J Gastroenterol*. 2009;15(15):1869–1875. doi:10.3748/wjg.15.1869
91. Wu WC, Wang CY. Association between non-alcoholic fatty pancreatic disease (NAFPD) and the metabolic syndrome: case-control retrospective study. *Cardiovasc Diabetol*. 2013;12:77. doi:10.1186/1475-2840-12-77
92. Ozturk K, Dogan T, Celikkanat S, et al. The association of fatty pancreas with subclinical atherosclerosis in nonalcoholic fatty liver disease. *Eur J Gastroenterol Hepatol*. 2018;30(4):411–417. doi:10.1097/MEG.0000000000001059
93. Seppala-Lindroos A, Vehkavaara S, Hakkinen AM, et al. Fat accumulation in the liver is associated with defects in insulin suppression of glucose production and serum free fatty acids independent of obesity in normal men. *J Clin Endocrinol Metab*. 2002;87(7):3023–3028. doi:10.1210/jcem.87.7.8638
94. Gabriely I, Ma XH, Yang XM, et al. Removal of visceral fat prevents insulin resistance and glucose intolerance of aging: an adipokine-mediated process? *Diabetes*. 2002;51(10):2951–2958. doi:10.2337/diabetes.51.10.2951
95. Kotronen A, Seppala-Lindroos A, Bergholm R, Yki-Jarvinen H. Tissue specificity of insulin resistance in humans: fat in the liver rather than muscle is associated with features of the metabolic syndrome. *Diabetologia*. 2008;51(1):130–138. doi:10.1007/s00125-007-0867-x
96. Milburn JL Jr, Hirose H, Lee YH, et al. Pancreatic beta-cells in obesity. Evidence for induction of functional, morphologic, and metabolic abnormalities by increased long chain fatty acids. *J Biol Chem*. 1995;270(3):1295–1299. doi:10.1074/jbc.270.3.1295
97. Lingvay I, Esser V, Legendre JL, et al. Noninvasive quantification of pancreatic fat in humans. *J Clin Endocrinol Metab*. 2009;94(10):4070–4076. doi:10.1210/jc.2009-0584
98. Yamazaki H, Tsuboya T, Katanuma A, et al. Lack of Independent Association Between Fatty Pancreas and Incidence of Type 2 Diabetes: 5-Year Japanese Cohort Study. *Diabetes Care*. 2016;39(10):1677–1683. doi:10.2337/dc16-0074
99. Raeder H, Haldorsen IS, Erslund L, et al. Pancreatic lipomatosis is a structural marker in nondiabetic children with mutations in carboxyl-ester lipase. *Diabetes*. 2007;56(2):444–449. doi:10.2337/db06-0859
100. Lim S, Meigs JB. Links between ectopic fat and vascular disease in humans. *Arterioscler Thromb Vasc Biol*. 2014;34(9):1820–1826. doi:10.1161/ATVBAHA.114.303035
101. Chung GE, Choi SY, Kim D, et al. Nonalcoholic fatty liver disease as a risk factor of arterial stiffness measured by the cardioankle vascular index. *Medicine (Baltimore)*. 2015;94(12):e654. doi:10.1097/MD.0000000000000654
102. Ozturk K, Guler AK, Cakir M, et al. Pulse Wave Velocity, Intima Media Thickness, and Flow-mediated Dilatation in Patients with Normotensive Normoglycemic Inflammatory Bowel Disease. *Inflamm Bowel Dis*. 2015;21(6):1314–1320. doi:10.1097/MIB.0000000000000355
103. Rasool A, Dar W, Latief M, Dar I, Sofi N, Khan MA. Nonalcoholic fatty liver disease as an independent risk factor for carotid atherosclerosis. *Brain Circ*. 2017;3(1):35–40. doi:10.4103/bc.bc\_28\_16
104. Fracanzani AL, Tiraboschi S, Pisano G, et al. Progression of carotid vascular damage and cardiovascular events in non-alcoholic fatty liver disease patients compared to the general population during 10 years of follow-up. *Atherosclerosis*. 2016;246:208–213. doi:10.1016/j.atherosclerosis.2016.01.016

105. Li N, Zhang GW, Zhang JR, et al. Non-alcoholic fatty liver disease is associated with progression of arterial stiffness. *Nutr, Metab Cardiovasc Dis.* 2015;25(2):218–223. doi:10.1016/j.numecd.2014.10.002
106. Zheng J, Zhou Y, Zhang K, et al. Association between nonalcoholic fatty liver disease and subclinical atherosclerosis: a cross-sectional study on population over 40 years old. *BMC Cardiovasc Disord.* 2018;18(1):147. doi:10.1186/s12872-018-0877-2
107. Sahin S, Yerlikaya MG, Ozderya A, et al. Non-Alcoholic Fatty Pancreas Disease is Associated with SYNTAX Score in Acute Coronary Syndrome. *Acta Cardiol Sin.* 2022;38(6):683–690. doi:10.6515/ACS.202211\_38(6).20220424A
108. Kul S, Karadeniz A, Dursun I, et al. Non-Alcoholic Fatty Pancreas Disease is Associated with Increased Epicardial Adipose Tissue and Aortic Intima-Media Thickness. *Acta Cardiol Sin.* 2019;35(2):118–125. doi:10.6515/ACS.201903\_35(2).20181009A
109. Pitt HA. Hepato-pancreato-biliary fat: the good, the bad and the ugly. *HPB.* 2007;9(2):92–97. doi:10.1080/13651820701286177
110. Zyromski NJ, Mathur A, Pitt HA, et al. A murine model of obesity implicates the adipokine milieu in the pathogenesis of severe acute pancreatitis. *Am J Physiol Gastrointest Liver Physiol.* 2008;295(3):G552–558. doi:10.1152/ajpgi.90278.2008
111. Sbeit W, Khoury T. Fatty Pancreas Represents a Risk Factor for Acute Pancreatitis: a Pilot Study. *Pancreas.* 2021;50(7):990–993. doi:10.1097/MPA.0000000000001867
112. Arslan AA, Helzlsouer KJ, Kooperberg C, et al. Anthropometric measures, body mass index, and pancreatic cancer: a pooled analysis from the Pancreatic Cancer Cohort Consortium (PanScan). *Arch Intern Med.* 2010;170(9):791–802. doi:10.1001/archinternmed.2010.63
113. Lesmana CRA, Gani RA, Lesmana LA. Non-alcoholic fatty pancreas disease as a risk factor for pancreatic cancer based on endoscopic ultrasound examination among pancreatic cancer patients: a single-center experience. *JGH Open.* 2018;2(1):4–7. doi:10.1002/jgh3.12032
114. Hori M, Takahashi M, Hiraoka N, et al. Association of pancreatic Fatty infiltration with pancreatic ductal adenocarcinoma. *Clin Transl Gastroenterol.* 2014;5(3):e53. doi:10.1038/ctg.2014.5
115. Khoury T, Sbeit W. Fatty Pancreas and Pancreatic Cancer: an Overlooked Association? *J Clin Med.* 2022;11(3). doi:10.3390/jcm11030763
116. Mathur A, Zyromski NJ, Pitt HA, et al. Pancreatic steatosis promotes dissemination and lethality of pancreatic cancer. *J Am Coll Surg.* 2009;208(5):989–994. doi:10.1016/j.jamcollsurg.2008.12.026
117. Zhang CL, Wang JJ, Li JN, Yang Y. Nonalcoholic fatty pancreas disease: an emerging clinical challenge. *World J Clin Cases.* 2021;9(23):6624–6638. doi:10.12998/wjcc.v9.i23.6624
118. Rizos CV, Liberopoulos EN, Mikhailidis DP, Elisaf MS. Pleiotropic effects of thiazolidinediones. *Expert Opin Pharmacother.* 2008;9(7):1087–1108. doi:10.1517/14656566.9.7.1087
119. Gottlieb S. Company played down drug's risks, report says. *BMJ.* 2001;322(7288).
120. Jia DM, Fukumitsu KI, Tabaru A, Akiyama T, Otsuki M. Troglitazone stimulates pancreatic growth in congenitally CCK-A receptor-deficient OLETF rats. *Am J Physiol Regul Integr Comp Physiol.* 2001;280(5):R1332–1340. doi:10.1152/ajpregu.2001.280.5.R1332
121. Fang T, Huang S, Chen Y, Chen Z, Chen J, Hu W. Glucagon Like Peptide-1 Receptor Agonists Alter Pancreatic and Hepatic Histology and Regulation of Endoplasmic Reticulum Stress in High-fat Diet Mouse Model. *Exp Clin Endocrinol Diabetes.* 2021;129(9):625–633. doi:10.1055/a-1240-4936
122. Akoumianakis I, Zagaliotis A, Konstantaraki M, Filippatos TD. GLP-1 analogs and regional adiposity: a systematic review and meta-analysis. *Obes Rev.* 2023;e13574. doi:10.1111/obr.13574
123. Souza-Mello V, Gregório BM, Relvas-Lucas B, da Silva Faria T, Aguila MB, Mandarim-de-Lacerda CA. Pancreatic ultrastructural enhancement due to telmisartan plus sitagliptin treatment in diet-induced obese C57BL/6 mice. *Pancreas.* 2011;40(5):715–722. doi:10.1097/MPA.0b013e3182153922
124. Salman AA, Salman MA, Said M, et al. Improvement of Pancreatic Steatosis and Indices of Insulin Resistance After Metabolic Surgery. *Front Med Lausanne.* 2022;9:894465. doi:10.3389/fmed.2022.894465
125. Rinella ME, Lazarus JV, Ratziu V, et al. A multi-society Delphi consensus statement on new fatty liver disease nomenclature. *Ann Hepatol.* 2023;101133. doi:10.1016/j.aohep.2023.101133
126. Rinella ME, Lazarus JV, Ratziu V, et al. A multi-society Delphi consensus statement on new fatty liver disease nomenclature. *J Hepatol.* 2023. doi:10.1016/j.jhep.2023.06.003
127. Rinella ME, Lazarus JV, Ratziu V, et al. A multi-society Delphi consensus statement on new fatty liver disease nomenclature. *Hepatology.* 2023. doi:10.1097/hep.0000000000000520

## Diabetes, Metabolic Syndrome and Obesity

Dovepress

### Publish your work in this journal

Diabetes, Metabolic Syndrome and Obesity is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert opinion and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-journal>