

# Stress management interventions for HIV-infected individuals: review of recent intervention approaches and directions for future research

Jennifer L Brown<sup>1,2</sup>  
Peter A Vanable<sup>3</sup>

<sup>1</sup>Department of Behavioral Sciences and Health Education, <sup>2</sup>Center for AIDS Research, Emory University, Atlanta, GA, USA; <sup>3</sup>Department of Psychology, Syracuse University, Syracuse, NY, USA

**Abstract:** To promote psychological wellbeing and adaptive coping among people living with human immunodeficiency virus/acquired immune deficiency syndrome (PLWHA), a number of stress management interventions have been designed and evaluated. This paper reviews recent stress management intervention approaches designed to improve the coping skills of PLWHA and reduce psychological distress. First, a summary of findings from previous narrative reviews and meta-analyses of the stress management intervention literature for PLWHA is provided. Next, recent stress management interventions for PLWHA that fall into one of four categories are reviewed: (a) interventions to improve coping and modify other health behaviors (ie, highly active antiretroviral medication adherence, sexual behaviors), (b) meditation, mindfulness, and relaxation-based stress management approaches, (c) computer-delivered interventions, and (d) interventions that target specific populations including older individuals, individuals with childhood sexual abuse histories, and women. A critique of recent stress management interventions for PLWHA is provided as well as directions for future research.

**Keywords:** stress management intervention, stress, coping, cognitive behavioral stress management, review, HIV, HIV-positive, HIV-infected, AIDS

## Introduction

The Centers for Disease Control and Prevention estimates that there are now more than 1 million people living with human immunodeficiency virus (HIV)/acquired immune deficiency syndrome (PLWHA) in the United States with approximately 53,600 new HIV infections diagnosed annually.<sup>1,2</sup> Highly active antiretroviral therapy (HAART) has improved the long-term health outlook of PLWHA and has markedly decreased HIV-related mortality.<sup>3,4</sup> However, HIV poses a number of unique psychosocial challenges associated with living with a chronic, stigmatized disease.<sup>5-7</sup> These challenges are compounded by the fact that HIV-infected individuals experience high rates of mental health difficulties and substance abuse.<sup>8-10</sup> To reduce psychological distress and improve coping among PLWHA there is a clear need to develop efficacious stress management interventions.

Despite the improved medical prognosis following an HIV diagnosis, PLWHA face a number of ongoing coping challenges. PLWHA must cope with a variety of challenging stressors, including concerns surrounding serostatus disclosure, challenges posed by HAART adherence and medication side effects, and navigating decisions about sexual activity and pregnancy desires.<sup>7</sup> While some PLWHA are able to effectively manage their care and lead fulfilling lives, a significant proportion report difficulties

Correspondence: Jennifer L Brown  
Department of Behavioral Sciences and Health Education, Rollins School of Public Health, Emory University, 1518 Clifton Road NE, Room 426, Atlanta, GA 30322, USA  
Tel +1 404 727 8673  
Fax +1 404 712 9738  
Email [jennifer.brown@emory.edu](mailto:jennifer.brown@emory.edu)

copied with stress.<sup>11</sup> Elevated psychosocial stressors, coupled with poor stress management skills, can exacerbate existing psychiatric illnesses or heighten an individual's risk for a new disorder including major depression, alcohol or drug dependence, and anxiety disorders.<sup>12,13</sup> Indeed, research suggests elevated rates of psychiatric illness among PLWHA relative to noninfected samples.<sup>9</sup> For instance, a recent investigation found a 19% prevalence rate for major depressive disorder among PLWHA, compared to only 5% in a noninfected comparison sample.<sup>14</sup>

Elevated psychological distress may contribute to poor disease management and negative health outcomes among PLWHA.<sup>15,16</sup> Prior investigations demonstrate that heightened psychological distress is associated with accelerated disease progression,<sup>17</sup> as indicated by CD4 decline, increased viral load,<sup>15</sup> and fewer natural killer cells.<sup>18</sup> Lower engagement in protective health behaviors may mediate the association between psychological functioning and disease outcomes.<sup>19</sup> Mental health difficulties may also be associated with suboptimal medication adherence<sup>20,21</sup> and less engagement in preventative health practices, including missed medical appointments<sup>22</sup> and sexual risk taking.<sup>23,24</sup> Thus, there is increasing evidence that psychological distress is associated with poor health outcomes and less engagement in protective health behaviors among PLWHA.

Stress management interventions for PLWHA address an important public health need. Stress management training is increasingly viewed as integral to the broader goal of assuring that PLWHA maintain adequate self-care for their illness.<sup>25</sup> Insofar as stress management interventions can reduce distress and, potentially, improve disease management and health outcomes, research that evaluates the efficacy of stress management interventions among PLWHA is of considerable importance. To date, a number of promising stress management interventions have been developed and evaluated. This review provides a summary of key findings from the previous narrative review and meta-analyses that have examined the efficacy of stress management interventions for PLWHA.<sup>26–28</sup> Next, the existing reviews are updated by highlighting recent stress management interventions designed to reduce psychological distress and improve coping skills among PLWHA. Specifically, this review describes core components of recent intervention approaches, reviews intervention efficacy findings, and offers guidance for future research. Thus, this review seeks to summarize the state of the science and outline key research priorities for stress management interventions for PLWHA.

## Summary of previous stress management reviews

To date, one narrative review<sup>26</sup> and two meta-analyses<sup>27,28</sup> have reviewed the stress management intervention literature for PLWHA. In what follows, this review provides a succinct summary of findings from each of the reviews. For each of the previous reviews, this review summarizes the format and content of the interventions, reports on the efficacy of reviewed interventions, and notes intervention characteristics associated with differential intervention efficacy.

Brown and Vanable<sup>26</sup> conducted a narrative review of 21 stress management interventions for PLWHA that included both cognitive (eg, modification of automatic thoughts) and behavioral skills training (eg, progressive muscle relaxation).<sup>29–55</sup> Across the reviewed cognitive behavioral stress management (CBSM) studies, 95% of programs were delivered in a group format with group sizes ranging from four to ten members (mode = seven group members). Most interventions consisted of multiple sessions (range = 6–20 sessions; mode = ten sessions) and were between 1 hour and 2.5 hours in length (mode = 2 hour session). Stress management interventions focused primarily on HIV-infected men who have sex with men and more than half (57%) of the reviewed studies excluded PLWHA because of current or past psychiatric, substance abuse, or personality disorder histories.

Overall, there was evidence that stress management programs were effective in reducing perceived stress levels and improving overall psychological adjustment (ie, depression and anxiety symptoms, global psychological functioning) and psychosocial functioning (ie, quality of life, social support). Although a primary aim of all of the reviewed studies was to improve adaptive coping, there was little consistency in approaches used to assess coping and findings were, at best, mixed. Among a small subset of studies that included measures of coping self-efficacy, there is some evidence to suggest that stress management interventions improve PLWHA's self-efficacy to effectively cope with stressors. When examining health status markers, there was little evidence to suggest that the interventions improved biological markers of immune system functioning including stress hormones, CD4 counts, natural killer cells, naïve T cells, and cytotoxic T cells. In summary, CBSM interventions for PLWHA show promise as an approach to reducing psychological distress. However, the authors identified a number of methodological limitations in the reviewed studies and also noted that the use of group-delivered, multisession interventions may

not be feasible to implement in resource-limited HIV medical centers.

In addition to the narrative review conducted by Brown and Venable,<sup>26</sup> two recent meta-analyses have been published examining the efficacy of cognitive-behavioral<sup>27</sup> and stress management interventions.<sup>28</sup> Table 1 provides the effect size estimates for outcomes from each of the meta-analyses. The meta-analysis conducted by Crepaz and colleagues<sup>27</sup> included 15 cognitive-behavioral randomized controlled trials (RCTs) designed to modify maladaptive cognitions, reduce negative affect, and improve immune functioning among PLWHA.<sup>29,32,35,39,41,43,44,47,50,51,53,56-61</sup> Fourteen of these interventions included coping skills and stress management training; the remaining study focused exclusively on cognitive-behavioral therapeutic approaches to alleviate depressed mood and reduce anxiety, excluding modules on coping. This meta-analysis found that cognitive-behavioral interventions resulted in significant decreases in depression, anxiety, anger, and stress across studies (Table 1).<sup>27</sup> Effect sizes across outcomes were typically small; however, there was a large effect for the interventions' efficacy in reducing anger. In contrast, findings provided no evidence that the reviewed interventions led to improvements in immune functioning as measured by CD4 count.<sup>27</sup> An evaluation of participant characteristics and research methodology features associated with differential intervention efficacy indicated that interventions were more efficacious in improving depression and anxiety outcomes under the following conditions:

**Table 1** Summary of cognitive behavioral and stress management intervention effect size estimates from Crepaz et al<sup>27</sup> and Scott-Sheldon et al<sup>28</sup>

	Effect size estimate	
	Crepez et al <sup>27</sup> (N = 15 RCTs)	Scott-Sheldon et al <sup>28</sup> (N = 35 RCTs)
Stress and coping outcomes		
Coping	–	0.10
Stress	0.43	–
Psychological and psychosocial outcomes		
Social support	–	0.01
Depression	0.33	0.28
Anxiety	0.30	0.28
Anger	1.00	–
Distress	–	0.19
Quality of life	–	0.16
Fatigue	–	0.38
Health status markers		
CD4 counts	0.08	0.04
Viral load	–	–0.09
Hormonal level	–	–0.14

**Abbreviation:** RCTs, randomized controlled trials.

(a) less than 50% of the sample consisted of men who have sex with men, (b) participants were not excluded due to depressive symptoms, (c) stress management training was provided, (d) interventions were led by trained research staff, (e) ten or more sessions were conducted for the intervention, and (f) the study's retention rate was greater than 80%.

A second meta-analysis conducted by Scott-Sheldon and colleagues<sup>28</sup> included 35 stress management RCTs.<sup>29,32,35,36,39,41,44,46,53,59-83</sup> In contrast to the Crepez et al study,<sup>27</sup> this meta-analysis utilized broad inclusion criteria for studies including psychosocial, behavioral, or educational stress management approaches to promote adjustment among PLWHA. The meta-analysis included studies testing cognitive-behavioral interventions, as well as less traditional stress management approaches, including massage therapy, relaxation education, and tai chi, among others. Among the reviewed studies, stress management interventions resulted in significant improvement on the outcomes of anxiety, depression, psychological distress, fatigue, and quality of life (Table 1).<sup>28</sup> All effect sizes for these outcomes were small. However, stress management programs did not result in significant changes for coping, social support, CD4 counts, viral load, or hormonal outcomes.<sup>28</sup> Stress management programs resulted in greater reduction in anxiety symptoms when samples included more women, individuals with higher anxiety at baseline, and younger participants.<sup>28</sup> In addition, intervention approaches that included medication adherence training were less efficacious than other interventions.<sup>28</sup> Furthermore, when entering all of the moderators into a multiple regression model, inclusion of medication adherence training as part of an intervention remained as the only significant predictor of poorer intervention outcomes.<sup>28</sup>

Consistent with findings from the narrative review,<sup>26</sup> findings from both meta-analyses<sup>27,28</sup> suggest that stress management programs for PLWHA result in the greatest change for psychological outcomes, such as depression and anxiety. In contrast, stress management interventions showed less consistent effects for improvement in coping and health status markers such as CD4 counts. Efficacious interventions typically included multiple, group-based sessions that provided a combination of psychoeducation and cognitive and behavioral strategies to improve stress management skills. In what follows, this review highlights recently published stress management interventions for PLWHA.


## Literature search method and criteria for inclusion

Database searches of PsycINFO, PsycARTICLES, Medline, and PubMed were conducted to identify published articles

in peer-reviewed journals that evaluated stress management interventions for PLWHA published subsequent to the three previous reviews.<sup>26–28</sup> Combinations of the following search terms were used to identify relevant articles: stress management, stress, coping, relaxation, intervention, HIV, HIV-positive, HIV-infected, and acquired immune deficiency syndrome. A search of references cited in relevant studies was conducted to identify additional articles. English language articles published in peer-reviewed journals were screened for inclusion. Studies were included if they met all of the following criteria: (a) the intervention's primary goal was to improve stress management skills, (b) the study's sample was composed of only PLWHA, and (c) the intervention did not focus exclusively on modifying other health behaviors (eg, sexual risk reduction, improvement in HAART adherence). For this review, studies that were not conducted with United States samples were excluded. Figure 1 depicts the selection process for articles included in the review.

## Recent stress management intervention approaches

The goal of this segment of the review is to summarize stress management interventions for PLWHA that have been published since the narrative review<sup>26</sup> and meta-analyses.<sup>27,28</sup> First, key intervention content included in the reviewed interventions is outlined. Next, the efficacy of stress management programs that fall into one of four categories is reviewed: (a) interventions to improve coping and modify other health behaviors (ie, HAART medication adherence, sexual behaviors), (b) meditation, mindfulness, and relaxation-based stress management approaches, (c) computer-delivered interventions, and (d) interventions that target specific populations including older individuals, individuals with childhood sexual abuse histories, and women. Table 2 summarizes key study characteristics for each of the reviewed intervention



**Figure 1** Selection process for inclusion in the review.

trials including methodological features and core intervention content areas.

## Intervention components

All of the reviewed interventions sought to improve PLWHA's coping skills by encouraging the use of cognitive and behavioral stress management strategies (Table 2). For some of the reviewed studies, a core intervention component was provision of psychoeducation about the nature and health consequences of stress (Table 2). Across studies, the majority of stress management programs sought to modify general coping strategies that could be implemented across a range of stressful situations. Additionally, a minority of studies also targeted changes in specific, HIV-related health behavior domains, such as sexual risk behaviors and antiretroviral medication adherence.

Across the interventions, cognitive and behavioral approaches were designed to facilitate adaptive coping and reduce the negative effects of stress. Emotional regulation strategies and reducing overall psychological distress were often specified as goals of the interventions. As a behavioral strategy, the majority of studies (77%) included a relaxation training component (Table 2). Furthermore, across interventions, the use of other active coping strategies (eg, problem solving) was stressed as more adaptive than avoidant coping strategies (eg, substance use).

In most of the reviewed recent coping interventions, stress management skills training also included modules on the use of cognitive strategies to modify PLWHA's appraisal of stressors and encourage the use of active problem solving strategies. For instance, cognitive distortions and automatic thoughts about HIV-related stressors were often identified and targeted via cognitive restructuring. In studies evaluating variations of coping effectiveness training, the focus was on the stress appraisal process and matching the stressor's level of perceived changeability with the use of either problem- or emotion-focused coping strategies. Similarly, problem solving approaches taught PLWHA to identify characteristics of specific stressors, brainstorm potential solutions, select a coping strategy, and evaluate the effectiveness of the chosen solution for the problem situation.

## Stress management combined with health behavior change Adherence

High levels of adherence are required to achieve optimal viral suppression and to prevent the development of drug-resistant HIV strains.<sup>64,84,85</sup> Three recent studies have examined the

**Table 2** Key characteristics of recent stress management interventions for persons living with human immunodeficiency virus

Study citations	Sample gender and sexual orientation; ethnicity	Study exclusion criteria	Study conditions	RCT	Number (length) of intervention sessions	Core intervention components	Assessment time points
64	N = 52 men, 11 women; ethnicity NR	HAART, C/P, MH	E: individual SM, MAT C: standard care	✓	E: 4 (75 min)	HBC, GB	Pre, post
86	N = 226 men, 23 women; 55% Caucasian, 19% African-American, 15% Hispanic, 11% other	HAART, C/P	E: individual SM, MAT C: WLC	✓	E: 5 (60 min)	HBC, GB, GC, SAM, PS	Pre, 3 mos, 6 mos, 9 mos, 12 mos, 15 mos
87	N = 130 MSM; ethnicity NR	HAART, C/P, MH	E: group SM, MAT C: MAT, condensed group SM	✓	E: 10 (135 min), 1 (60 min), 2 (30 min) C: 1 (60 min), 2 (30 min), 1 day session E: 15 (90 min)	HBC, GB, GC, R, SAM	Pre, post, 9 mos, 15 mos
94	N = 674 men, 262 women; 45% African-American, 32% Caucasian, 15% Hispanic, 8% other	SB	E: individual SM C: WLC	✓	E: 8 (120 min), 1 (360 min) C: 1 day session	HBC, GB, GC, SAM, R	Pre, 5 mos, 10 mos, 15 mos, 20 mos, 25 mos
97	N = 43 men, 5 women; 50% African-American, 21% Hispanic, 23% Caucasian, 6% other	HIVSX	E: group SM C: condensed group SM	✓	E: 8 (120 min), 1 (360 min) C: 1 day session	R, GB	Pre, post
98	N = 75 men, 8 women; 52% Caucasian, 31% African-American, 15% Hispanic	MH, C/P	E: group SM C: psychoeducational videos	✓	E: 5 (90 min) C: 5 (90 min)	R, GB	Pre, post, 10 wks, 22 wks
99	N = 152 men, 100 women; 75% African-American, 23% Caucasian, 2% Hispanic	MH, C/P	E: group SM C1: group Tai-Chi C2: group spiritual growth C3: WLC	✓	E: 10 (90 min) C1: 10 (90 min) C2: 10 (90 min)	R, GB	Pre, post, 6 mos

(Continued)

Table 2 (Continued)

Study citations	Sample gender and sexual orientation; ethnicity	Study exclusion criteria	Study conditions	RCT	Number (length) of intervention sessions	Core intervention components	Assessment time points
101	N = 60 women; 70% African-American, 21% Caucasian, 9% other, 8% Hispanic	C/P	E: computerized SM C: time-delayed computerized SM	✓	E: 1 (90 min) C: 1 (90 min)	GB, GC, SAM, R	Pre, 1 mos
102	N = 197 men, 98 women; 49% African-American	C/P	E: group SM C1: support group C2: individual therapy on request	✓	E: 12 (90 min) C1: 12 (90 min)	GB, GC, SAM, R	Pre, post, 4 mos, 8 mos
104–106	N = 129 women, 117 men; 68% African-American, 17% Hispanic, 10% Caucasian, 5% other	C/P, MH	E: group SM C: support group	✓	E: 15 (90 min) C: 15 (90 min)	HBC, SAM, GB, GC	Pre, post, 4 mos, 8 mos, 12 mos
108	N = 39 women; 90% black, 5% Hispanic	MH, C/P	E: group SM C: condensed group SM	✓	E: 10 (135 min) C: 1 (5 hours)	R, GB, GC, SS, HBC	Pre, 9 mos
109: Study 1	N = 451 women; ethnicity NR		E: group SM C1: individual psychoeducation	✓	E: 10 (135 min) C: 10 (135 min)	GB, GC, R, SS	Pre, post, 6 mos, 12 mos
109: Study 2	N = 482 women; ethnicity NR		E: group SM, HBC C1: individual psychoeducation	✓	E: 10 (135 min), 6 (150 min) C: 10 (135 min)	GB, GC, R, SS, HBC	Pre, post, 6 mos, 12 mos, 24 mos

**Note:** When multiple papers were written on a single intervention trial, study citation numbers for these papers are listed in succession.

**Abbreviations:** C, comparison/control condition; C/P, excluded because of cognitive impairment or psychosis; E, experimental stress management intervention; GB, general behavioral coping skills (eg, variety of behavioral approaches); GC, general cognitive therapeutic approaches (eg, cognitive restructuring); HAART, excluded based on current medication status; HBC, health behavior change modules (adherence, substance use, sexual risk behaviors); HIV-Info, human immunodeficiency virus informational condition; HIVSX, excluded based on human immunodeficiency virus symptoms or acquired immune deficiency syndrome diagnosis; MAT, medication adherence training; MH, excluded due to mental health difficulties; min, minutes; mos, months; MSM, men who have sex with men; NR, not reported; P, psychoeducation about stress management; Post, postintervention assessment; Pre, preintervention assessment; PS, problem solving; R, relaxation; RCT, randomized controlled trial; SAM, stressor appraisal and matching of coping approach; SB, stressor appraisal and matching of coping approach; SS, stress management; SM, social support enhancement and utilization; wks, weeks; WLC, waitlist control.

efficacy of supplementing stress management interventions with content to address HAART medication adherence.<sup>64,86,87</sup> Balfour and colleagues<sup>64</sup> examined the efficacy of a four-session intervention to reduce depressive symptoms and enhance readiness to start HAART medication. Results indicated that the intervention reduced both depressive symptoms and enhanced PLWHA's readiness to start HAART medication.<sup>64</sup> A second study sought to improve stress management skills and enhance PLWHA's ability to cope with HAART medication side effects.<sup>86</sup> Individuals who completed the five individual stress management sessions were less likely to report nonadherence and were more likely to seek out additional information about medication side effects and utilize social support to cope with HAART side effects.<sup>86</sup> A third study compared a 10-week CBSM intervention supplemented with four brief medication adherence training sessions delivered by a pharmacist to a medication adherence training only group.<sup>87</sup> There were no differences in viral load between the two groups; however, PLWHA in the intervention condition who had a detectable viral load at baseline had decreased viral load relative to the medication adherence training only group.<sup>87</sup> Additionally, decreases in depressed mood among the intervention group participants explained the decreased viral load across the 15-month follow-up period.<sup>87</sup>

### Sexual risk reduction

Despite increased secondary HIV prevention efforts to reduce sexual risk behaviors among PLWHA,<sup>88</sup> reports suggest that some PLWHA have difficulties engaging in consistent safer sex practices.<sup>89,90</sup> Unprotected sexual encounters pose the risk of HIV transmission to uninfected partners and increased possibility of negative health consequences for PLWHA through acquisition of other sexually transmitted infections or drug-resistant HIV strains.<sup>91–93</sup> To improve stress management skills and reduce sexual risk behaviors among PLWHA, one intervention examined the efficacy of a 15-session individual intervention relative to a waitlist control condition.<sup>94</sup> There were no improvements in psychological functioning across a variety of measures including anxiety and depressive symptoms, burnout, perceived stress, positive affect, positive states of mind, coping self-efficacy, and perceived social support.<sup>94</sup> Furthermore, there were no intervention effects for decreasing psychological distress among PLWHA who had mild to moderate depressive symptoms when starting the study.<sup>94</sup> However, the intervention resulted in decreased frequency of unprotected sexual acts with partners who were HIV-negative or of unknown serostatus.<sup>94</sup>

## Meditation- and relaxation-based stress management interventions

Mindfulness-based stress reduction (MBSR) is an empirically validated treatment consisting of 8–10 weeks of mindfulness meditation training.<sup>95</sup> This intervention approach focuses on developing greater in the moment self-awareness through guided exercises.<sup>96</sup> A meta-analysis examining the efficacy of MBSR for a variety of medical (eg, heart, pain) and psychiatric (eg, anxiety disorders) conditions found medium effect sizes, indicating that MBSR interventions result in improvements in psychological distress, health functioning, and coping skills.<sup>96</sup> Although MBSR has been used with a variety of medical populations, one recent study suggests that the mindfulness-based approach may be helpful for PLWHA. Creswell and colleagues<sup>97</sup> examined the impact of an eight-session MBSR intervention relative to a one day control seminar condition for increasing CD4 lymphocyte counts. PLWHA in the control condition demonstrated greater declines in CD4 lymphocyte counts relative to those in the MBSR condition.<sup>97</sup> Additional analyses indicated that PLWHA who attended more MBSR sessions had greater CD4 lymphocyte counts.<sup>97</sup>

A second study examined the efficacy of a group-based mantram intervention for PLWHA.<sup>98</sup> A mantram is a spiritual word or phrase that is repeated throughout the day as a form of meditation to focus one's attention.<sup>98</sup> Primary outcomes from this RCT were reviewed in Scott-Sheldon and colleagues' meta-analysis;<sup>28</sup> this secondary analysis examined positive reappraisal and distance coping as mediators of the intervention effects for reducing anger.<sup>98</sup> Findings indicated that the mantram intervention reduced trait-anger by enhancing positive reappraisal coping.<sup>98</sup>

A third study compared the efficacy of 10-week interventions of CBSM, tai chi, and spiritual growth relative to a waitlist control group.<sup>99</sup> Compared to the control group, individuals in the CBSM and tai chi interventions utilized less emotion-focused coping strategies and had improved lymphocyte proliferative functioning.<sup>99</sup> However, there were no differences between groups for quality of life, impact of HIV life events, neuroendocrine functioning, and HIV-specific health status.<sup>99</sup>

## Computer-delivered interventions

Stress management interventions have typically been administered in a group format led by a clinician (eg, psychologist, social worker) or an experienced health educator. Such an approach may not be feasible in resource-limited HIV clinic settings. To address this concern,

computer delivery is a promising modality for implementing stress management interventions that can be cost effective and flexibly administered. In a recent pilot study, a brief, interactive cognitive-behavioral computerized stress management intervention for HIV-infected women was evaluated relative to a delayed intervention control group.<sup>100,101</sup> All participants successfully navigated the computerized program, demonstrating intervention feasibility and most found the intervention to be acceptable.<sup>100</sup> While women demonstrated improved stress management knowledge, there were few differences between the intervention and control groups on psychological distress, perceived stress, and coping self-efficacy outcomes.<sup>101</sup>

## Stress management for specific populations

### Older adults

Given the improved long-term prognosis of PLWHA with HAART medication, there is a clear need for stress management programs to address the psychosocial stressors experienced by older individuals. One study tested a twelve-session coping intervention relative to an interpersonal support group intervention and a mental health service upon request condition for older PLWHA.<sup>102</sup> Results from this RCT indicated that individuals in both the coping intervention and interpersonal support group reported fewer depressive symptoms than the treatment upon request control group.<sup>102</sup>

### Individuals with childhood sexual abuse histories

Rates of childhood sexual abuse are high among PLWHA, and it is often associated with greater psychological distress in adulthood.<sup>103</sup> To improve coping skills and address psychological sequelae associated with childhood sexual abuse, the efficacy of a 15-session CBSM intervention for both male and female PLWHA with childhood sexual abuse histories relative to a time-matched support group or waitlist control group was evaluated.<sup>104–106</sup> In addition to core cognitive and behavioral stress management approaches, the intervention sessions also incorporated content to address individuals' trauma histories.<sup>104–106</sup> This intervention resulted in decreased symptoms of traumatic stress and reductions in the use of avoidance coping for individuals in the treatment group compared to patients assigned to a waitlist condition.<sup>105</sup> In addition, individuals in both the intervention and support group conditions endorsed fewer intrusive thoughts linked to the trauma.<sup>105</sup> Although not a direct focus for the intervention, participants in the stress management intervention reported

fewer instances of unprotected sex with all sexual partners and with partners who were HIV-negative or of unknown serostatus.<sup>106</sup> While the intervention did not directly focus on reducing substance use, longitudinal analyses also indicated that individuals in the stress management condition reduced the quantity of alcohol consumed and decreased their use of cocaine; however there were no changes in marijuana use.<sup>104</sup> Thus, this stress management intervention that incorporated content to improve coping and also address trauma history was efficacious in decreasing psychological distress and other health risk behaviors.

### Women

Rates of HIV infection are on the rise among several subgroups in the United States, including low-income, minority women.<sup>107</sup> However, relatively few stress management programs have been developed for HIV-infected women. In addition to the computer-delivered intervention developed by Brown and colleagues for HIV-infected women,<sup>100,101</sup> two intervention trials evaluated the efficacy of a CBSM intervention tailored to the unique psychosocial stressors faced by HIV-infected women.<sup>108,109</sup> The first study examined the impact of a 10-week CBSM intervention on cervical intraepithelial neoplasia, the neoplastic changes that occur following sexual transmission of human papillomaviruses and are precursors to cervical cancer.<sup>108</sup> Women in the intervention condition endorsed decreased perceived life stress and had lower odds of cervical intraepithelial neoplasia over 9 months relative to the one psychoeducational control session.<sup>108</sup> Weiss and colleagues<sup>109</sup> summarized the findings from two linked intervention trials of a CBSM intervention for HIV-infected women. The first trial compared a 10-week intervention to an individual psychoeducational condition.<sup>109</sup> Women in the intervention condition endorsed decreased depression and anxiety symptoms, less use of denial coping, and decreased health distress. In addition, participants reported greater social support, improved coping skills, improved quality of life, and increased medication adherence. The second trial was a continuation of the initial trial that enhanced the first intervention by adding six additional sessions focused on modifying other health behaviors (eg, medication adherence, reducing sexual risk behavior and substance use, increasing physical activity). Women in the enhanced CBSM condition endorsed decreased depressive symptoms, improved mental health quality of life, decreased risky sexual behaviors and alcohol use, and enhanced medication adherence, nutrition, and physical activity.<sup>109</sup> Thus, the core CBSM intervention decreased psychological distress, and when supplemented



with additional health promotion sessions, resulted in improvements across a variety of health behaviors.

## Conclusion

There is considerable promise in the use of stress management interventions to improve the lives of PLWHA. Combined with findings from previous reviews of the literature, there is a growing body of evidence demonstrating that stress management interventions for PLWHA reduce psychological distress and improve psychosocial functioning. Recent intervention studies have targeted other important health behaviors, utilized meditation and mindfulness approaches, used technology-delivered intervention formats, and designed interventions for specific subpopulations of PLWHA. The latest stress management interventions highlight the need to tailor intervention approaches to specific populations and address related challenges faced by PLWHA, including consistent safer sexual behaviors and adherence to medication regimens.

Across the reviewed recent interventions, the majority were delivered via a group format ( $n = 8$ ), with three individual-delivered interventions, and one computer-delivered intervention. Most recent interventions consisted of multiple sessions (mode = ten sessions) that were typically over an hour in length. Although group-delivered training provides the added benefit of fostering social support, some PLWHA may be reticent to attend groups due to confidentiality concerns or dislike of group meetings. Thus, the usefulness of group stress management interventions is limited to the subset of patients who are willing to participate in a group-based program. Group interventions also require significant financial resources to implement and may not be feasible in some resource-limited clinic settings. Moving forward, research should examine strategies to enhance the dissemination and uptake of evidence-based stress management interventions, particularly in resource-limited care settings. Such research should also examine the cost effectiveness of interventions and ways to improve retention in stress management interventions.

Few stress management interventions have addressed the unique psychosocial needs of HIV-infected patients using briefer, more cost-effective formats. An important gap in the literature is to examine the efficacy of briefer psycho-educational stress management approaches that can be of use to a broader range of patients. To date, only one brief stress management intervention delivered via computer has been pilot tested.<sup>101</sup> While this approach was feasible and acceptable to PLWHA, the intervention demonstrated only modest improvements in stress management knowledge.<sup>101</sup>

Thus, additional research to understand characteristics of PLWHA for whom briefer stress management approaches are most effective is needed.

Given the multitude of challenges faced by PLWHA, integrated, multidisciplinary HIV treatment centers offer the potential to conduct ongoing assessment of individuals' psychological distress, psychosocial functioning, and health behavior engagement. The use of ongoing assessment combined with adaptive intervention approaches<sup>110</sup> could guide the selection of an appropriate stress management intervention approach for a particular individual. Adaptive clinical trials allow for planned, reactive treatment changes throughout the trial based on initial treatment response.<sup>110,111</sup> Thus, for PLWHA who do not demonstrate improved psychological functioning after receiving a stress management intervention, such an approach could facilitate appropriate treatment referrals for more intensive psychiatric or substance use services.

In sum, the stress management literature indicates that stress management interventions facilitate positive adjustment and improve coping skills to effectively manage stress. Future research should examine mechanisms responsible for these positive changes and adapt the interventions to best meet the needs of the broader HIV patient population. Particularly important is the need to design and test more cost-effective, brief treatment approaches that can be implemented in under-resourced treatment settings.

## Acknowledgment

Jennifer L Brown is supported by K12 GM000680 from the National Institute of General Medical Sciences.

## Disclosure

The authors report no conflicts of interest in this work.

## References

1. Centers for Disease Control and Prevention (CDC). HIV prevalence estimates – United States, 2006. *MMWR Morb Mortal Wkly Rep*. 2008; 57(39):1073–1076.
2. Centers for Disease Control and Prevention (CDC). HIV Surveillance Report, 2008. 2010. Available from: <http://www.cdc.gov/hiv/topics/surveillance/resources/reports/>. Accessed August 16, 2010.
3. Schackman BR, Gebo KA, Walensky RP, et al. The lifetime cost of current human immunodeficiency virus care in the United States. *Med Care*. 2006;44(11):990–997.
4. Palella FJ Jr, Delaney KM, Moorman AC, et al. Declining morbidity and mortality among patients with advanced human immunodeficiency virus infection. HIV Outpatient Study Investigators. *N Eng J Med*. 1998; 338(13):853–860.
5. Bower JE, Kemeny ME, Fawzy FI. Group interventions for individuals with serious medical illness. In: Chesney M, Antoni MH, editors. *Innovative Approaches to Health Psychology: Prevention and Treatment Lessons from AIDS*. Washington, DC: American Psychological Association; 2002:197–218.

6. Kalichman SC, Catz SL. Stressors in HIV infection. In: Nott KH, Vedhara K, editors. *Psychosocial and Biomedical Interactions in HIV Infection*. Amsterdam, Netherlands: Harwood Academic Publishers; 2000:31–60.
7. Bravo P, Edwards A, Rollnick S, Elwyn G. Tough decisions faced by people living with HIV: a literature review of psychosocial problems. *AIDS Rev*. 2010;12(2):76–88.
8. Klinkenberg WD, Sacks S. Mental disorders and drug abuse in persons living with HIV/AIDS. *AIDS Care*. 2004;16(Suppl 1):S22–S42.
9. Bing EG, Burnam MA, Longshore D, et al. Psychiatric disorders and drug use among human immunodeficiency virus-infected adults in the United States. *Arch Gen Psychiatry*. 2001;58(8):721–728.
10. Galvan FH, Bing EG, Fleishman JA, et al. The prevalence of alcohol consumption and heavy drinking among people with HIV in the United States: results from the HIV Cost and Services Utilization Study. *J Stud Alcohol*. 2002;63(2):179–186.
11. Heckman TG, Anderson ES, Sikkema KJ, Kochman A, Kalichman SC, Anderson T. Emotional distress in nonmetropolitan persons living with HIV disease enrolled in a telephone-delivered, coping improvement group intervention. *Health Psychol*. 2004;23(1):94–100.
12. Griffin KW, Rabkin JG. Psychological distress in people with HIV/AIDS: Prevalence rates and methodological issues. *AIDS Behav*. 1997;1(1):29–42.
13. Catz S, Gore-Felton C, McClure JB. Psychological distress among minority and low-income women living with HIV. *Behav Med*. 2002;28(2):53–60.
14. Morrison MF, Petitto JM, Ten Have T, et al. Depressive and anxiety disorders in women with HIV infection. *Am J Psychiatry*. 2002;159(5):789–796.
15. Ironson G, O’Cleirigh C, Fletcher M, et al. Psychosocial factors predict CD4 and viral load change in men and women with human immunodeficiency virus in the era of highly active antiretroviral treatment. *Psychosom Med*. 2005;67(6):1013–1021.
16. Jenkins SR, Coons HL. Psychosocial stress and adaptation processes for women coping with HIV/AIDS. In: O’Leary A, Jemmott LS, editors. *Women and AIDS: Coping and Care*. New York, NY: Plenum Press; 1996:33–86.
17. Chida Y, Vedhara K. Adverse psychosocial factors predict poorer prognosis in HIV disease: a meta-analytic review of prospective investigations. *Brain Behav Immun*. 2009;23(4):434–445.
18. Leserman J, Jackson ED, Petitto JM, et al. Progression to AIDS: the effects of stress, depressive symptoms, and social support. *Psychosom Med*. 1999;61(3):397–406.
19. Gore-Felton C, Koopman C. Behavioral mediation of the relationship between psychosocial factors and HIV disease progression. *Psychosom Med*. 2008;70(5):569–574.
20. Mellins CA, Havens JF, McDonnell C, et al. Adherence to antiretroviral medications and medical care in HIV-infected adults diagnosed with mental and substance abuse disorders. *AIDS Care*. 2009;21(2):168–177.
21. Weaver KE, Llabre MM, Durán RE, et al. A stress and coping model of medication adherence and viral load in hiv-positive men and women on highly active antiretroviral therapy (HAART). *Health Psychol*. 2005;24(4):385–392.
22. Andersen M, Hockman E, Smereck G, et al. Retaining women in HIV medical care. *J Assoc Nurses AIDS Care*. 2007;18(3):33–41.
23. Gore-Felton C, Koopman C, Turner-Cobb JM, Duran R, Israelski D, Spiegel D. The influence of social support, coping and mood on sexual risk behavior among HIV+ men and women. *J Health Psychol*. 2002;7(6):713–722.
24. Morin SF, Myers JJ, Shade SB, Koester K, Maiorana A, Rose CD. Predicting HIV transmission risk among HIV-infected patients seen in clinical settings. *AIDS Behav*. 2007;11(5 Suppl):S6–S16.
25. Ironson G, Antoni MH, Schneiderman N, et al. Coping: Interventions for optimal disease management. In: Chesney M, Antoni MH, editors. *Innovative Approaches to Health Psychology: Prevention and Treatment Lessons from AIDS*. Washington, DC: APA; 2002:167–195.
26. Brown JL, Vanable PA. Cognitive-behavioral stress management interventions for persons living with HIV: a review and critique of the literature. *Ann Behav Med*. 2008;35(1):26–40.
27. Crepaz N, Passin WF, Herbst JH, et al. Meta-analysis of cognitive-behavioral interventions on HIV-positive persons’ mental health and immune functioning. *Health Psychol*. 2008;27(1):4–14.
28. Scott-Sheldon LA, Kalichman SC, Carey MP, Fielder RL. Stress management interventions for HIV+ adults: a meta-analysis of randomized controlled trials, 1989 to 2006. *Health Psychol*. 2008;27(2):129–139.
29. Côté JK, Pepler C. A randomized trial of a cognitive coping intervention for acutely ill HIV-positive men. *Nurs Res*. 2002;51(4):237–244.
30. Sikkema KJ, Hansen NB, Ghebremichael M, et al. A randomized controlled trial of a coping group intervention for adults with HIV who are AIDS bereaved: longitudinal effects on grief. *Health Psychol*. 2006;25(5):563–570.
31. Hansen NB, Tarakeshwar N, Ghebremichael M, Zhang H, Kochman A, Sikkema KJ. Longitudinal effects of coping on outcome in a randomized controlled trial of a group intervention for HIV-positive adults with AIDS-related bereavement. *Death Stud*. 2006;30(7):609–636.
32. Sikkema KJ, Hansen NB, Kochman A, Tate DC, Difrancesco W. Outcomes from a randomized controlled trial of a group intervention for HIV positive men and women coping with AIDS-related loss and bereavement. *Death Stud*. 2004;28(3):187–209.
33. Sikkema KJ, Hansen NB, Meade CS, Kochman A, Lee RS. Improvements in health-related quality of life following a group intervention for coping with AIDS-bereavement among HIV-infected men and women. *Qual Life Res*. 2005;14(4):991–1005.
34. Coates TJ, McKusick L, Kuno R, Stites DP. Stress reduction training changed number of sexual partners but not immune function in men with HIV. *Am J Public Health*. 1989;79(7):885–887.
35. Kelly JA, Murphy DA, Bahr GR, et al. Outcome of cognitive-behavioral and support group brief therapies for depressed, HIV-infected persons. *Am J Psychiatry*. 1993;150(11):1679–1686.
36. Carrico AW, Antoni MH, Duran RE, et al. Reductions in depressed mood and denial coping during cognitive behavioral stress management with HIV-positive gay men treated with HAART. *Ann Behav Med*. 2006;31(2):155–1164.
37. Heckman TG, Kochman A, Sikkema KJ, et al. A pilot coping improvement intervention for late middle-aged and older adults living with HIV/AIDS in the USA. *AIDS Care*. 2001;13(1):129–139.
38. Chesney MA, Folkman S, Chambers DB. Coping effectiveness training for men living with HIV: preliminary findings. *Int J STD AIDS*. 1996;7 Suppl 2:75–82.
39. Chesney MA, Chambers DB, Taylor JM, Johnson LM, Folkman S. Coping effectiveness training for men living with HIV: results from a randomized clinical trial testing a group-based intervention. *Psychosom Med*. 2003;65(6):1038–1046.
40. Heckman TG, Barcikowski R, Ogles BM, et al. A telephone-delivered coping improvement group intervention for middle-aged and older adults living with HIV/AIDS. *Ann Behav Med*. 2006;32(1):27–38.
41. Mulder CL, Emmelkamp PM, Antoni MH, Mulder JW. Cognitive-behavioral and experiential group psychotherapy for HIV-infected homosexual men: a comparative study. *Psychosom Med*. 1994;56(5):423–431.
42. Mulder CL, Antoni MH, Emmelkamp PM, et al. Psychosocial group intervention and the rate of decline of immunological parameters in asymptomatic HIV-infected homosexual men. *Psychother Psychosom*. 1995;63(3–4):185–192.
43. McCain NL, Zeller JM, Cella DF, Urbanski PA. The influence of stress management training in HIV disease. *Nurs Res*. 1996;45(4):246–253.
44. Chan I, Kong P, Leung P, et al. Cognitive-behavioral group program for Chinese heterosexual HIV-infected men in Hong Kong. *Patient Educ Couns*. 2005;56(1):78–84.
45. Ironson G, Weiss S, Lydston D, et al. The impact of improved self-efficacy on HIV viral load and distress in culturally diverse women living with AIDS: the SMART/EST Women’s Project. *AIDS Care*. 2005;17(2):222–236.

46. Lechner SC, Antoni MH, Lydston D, et al. Cognitive-behavioral interventions improve quality of life in women with AIDS. *J Psychosom Res.* 2003;54(3):253–261.
47. Molassiotis A, Callaghan P, Twinn SF, Lam SW, Chung WY, Li CK. A pilot study of the effects of cognitive-behavioral group therapy and peer support/counseling in decreasing psychological distress and improving quality of life in Chinese patients with symptomatic HIV disease. *AIDS Patient Care STDS.* 2002;16(2):83–96.
48. Antoni MH, Cruess S, Cruess DG, et al. Cognitive-behavioral stress management reduces distress and 24-hour urinary free cortisol output among symptomatic HIV-infected gay men. *Ann Behav Med.* 2000;22(1):29–37.
49. Cruess S, Antoni M, Cruess D, et al. Reductions in herpes simplex virus type 2 antibody titers after cognitive behavioral stress management and relationships with neuroendocrine function, relaxation skills, and social support in HIV-positive men. *Psychosom Med.* 2000;62(6):828–837.
50. Cruess S, Antoni MH, Hayes A, et al. Changes in mood and depressive symptoms and related change processes during cognitive-behavioral stress management in HIV-infected men. *Cognit Ther Res.* 2002;26(3):373–392.
51. Lutgendorf SK, Antoni MH, Ironson G, et al. Cognitive-behavioral stress management decreases dysphoric mood and herpes simplex virus-Type 2 antibody titers in symptomatic HIV-seropositive gay men. *J Consult Clin Psychol.* 1997;65(1):31–43.
52. Lutgendorf SK, Antoni MH, Ironson G, et al. Changes in cognitive coping skills and social support during cognitive behavioral stress management intervention and distress outcomes in symptomatic human immunodeficiency virus (HIV)-seropositive gay men. *Psychosom Med.* 1998;60(2):204–214.
53. Inouye J, Flannelly L, Flannelly KJ. The effectiveness of self-management training for individuals with HIV/AIDS. *J Assoc Nurses AIDS Care.* 2001;12(5):71–82.
54. Lee MR, Cohen L, Hadley SW, Goodwin FK. Cognitive-behavioral group therapy with medication for depressed gay men with AIDS or symptomatic HIV infection. *Psychiatr Serv.* 1999;50(7):948–952.
55. Fawzy FI, Namir S, Wolcott DL. Structured group intervention model for AIDS patients. *Psychiatr Med.* 1989;7(2):35–45.
56. Cruess DG, Antoni MH, Kumar M, et al. Cognitive-behavioral stress management buffers decreases in dehydroepiandrosterone sulfate (DHEA-S) and increases in the cortisol/DHEA-S ratio and reduces mood disturbance and perceived stress among HIV-seropositive men. *Psychoneuroendocrinology.* 1999;24(5):537–549.
57. Evans S, Fishman B, Spielman L, Haley A. Randomized trial of cognitive behavior therapy versus supportive psychotherapy for HIV-related peripheral neuropathic pain. *Psychosomatics.* 2003;44(1):44–50.
58. Laperriere A, Ironson GH, Antoni MH, et al. Decreased depression up to one year following CBSM+ intervention in depressed women with AIDS: the smart/EST women's project. *J Health Psychol.* 2005;10(2):223–231.
59. Markowitz JC, Kocsis JH, Fishman B, et al. Treatment of depressive symptoms in human immunodeficiency virus-positive patients. *Arch Gen Psychiatry.* 1998;55(5):452–457.
60. McCain NL, Munjas BA, Munro CL, et al. Effects of stress management on PNI-based outcomes in persons with HIV disease. *Res Nurs Health.* 2003;26(2):102–117.
61. Perry S, Fishman B, Jacobsberg L, Young J, Frances A. Effectiveness of psychoeducational interventions in reducing emotional distress after human immunodeficiency virus antibody testing. *Arch Gen Psychiatry.* 1991;48(2):143–147.
62. Auerbach JE, Oleson TD, Solomon GF. A behavioral medicine intervention as an adjunctive treatment for HIV-related illness. *Psychol Health.* 1992;6(4):325–334.
63. Baigis J, Korniewicz DM, Chase G, Butz A, Jacobson D, Wu AW. Effectiveness of a home-based exercise intervention for HIV-infected adults: a randomized trial. *J Assoc Nurses AIDS Care.* 2002;13(2):33–45.
64. Balfour L, Kowal J, Silverman A, et al. A randomized controlled psycho-education intervention trial: Improving psychological readiness for successful HIV medication adherence and reducing depression before initiating HAART. *AIDS Care.* 2006;18(7):830–838.
65. Birk TJ, McGrady A, MacArthur RD, Khuder S. The effects of massage therapy alone and in combination with other complementary therapies on immune system measures and quality of life in human immunodeficiency virus. *J Altern Complement Med.* 2000;6(5):405–414.
66. Bormann JE, Gifford AL, Shively M, et al. Effects of spiritual mantram repetition on HIV outcomes: a randomized controlled trial. *J Behav Med.* 2006;29(4):359–376.
67. Brazier A, Mulkins A, Verhoef M. Evaluating a yogic breathing and meditation intervention for individuals living with HIV/AIDS. *Am J Health Promot.* 2006;20(3):192–195.
68. Caddick ND. *Reducing Emotional Distress Using the Stress Inoculation Model for HIV-Infected People in the Pre-AIDS Stage.* Ann Arbor, MI: ProQuest Information and Learning; 1996.
69. Danielson AL. *Emotional Status and Social Support Relations to Immune Status in Women Co-Infected with HIV and HPV: Preliminary Evidence for Beneficial Psychosocial Effects of a CBSM Intervention Tailored for This Population.* Ann Arbor, MI: ProQuest Information and Learning; 2003.
70. Eller LS. Effects of two cognitive-behavioral interventions on immunity and symptoms in persons with HIV. *Ann Behav Med.* 1995;17(4):339–348.
71. Galantino ML, Shepard K, Krafft L, et al. The effect of group aerobic exercise and t'ai chi on functional outcomes and quality of life for persons living with acquired immunodeficiency syndrome. *J Altern Complement Med.* 2005;11(6):1085–1092.
72. Gifford AL, Laurent DD, Gonzales VM, Chesney MA, Lorig KR. Pilot randomized trial of education to improve self management skills of men with symptomatic HIV/AIDS. *J Acquir Immune Defic Syndr Hum Retrovirol.* 1998;18(2):136–144.
73. Kalichman SC. The other side of the healthy relationships intervention: mental health outcomes and correlates of sexual risk behavior change. *AIDS Educ Prevention.* 2005;17(1 Suppl A):66–75.
74. LaPerriere AR, Antoni MH, Schneiderman N, et al. Exercise intervention attenuates emotional distress and natural killer cell decrements following notification of positive serologic status for HIV-1. *Biofeedback Self Regul.* 1990;15(3):229–242.
75. Lutgendorf SK. *Cognitive Behavioral Stress Management in a Symptomatic HIV-1 Seropositive Population: Effects on Mood, Coping, Immune and Neuroendocrine Factors.* Ann Arbor, MI: ProQuest Information and Learning; 1995.
76. Miles MS, Holditch-Davis D, Eron J, Black BP, Pedersen C, Harris DA. An HIV self-care symptom management intervention for African American mothers. *Nurs Res.* 2003;52(6):350–360.
77. Neidig JL, Smith BA, Brashers DE. Aerobic exercise training for depressive symptom management in adults living with HIV infection. *J Assoc Nurses AIDS Care.* 2003;14(2):30–40.
78. Schlenzig C, Jager H, Rieder H, Hammel G, Popescu M. Supervised physical exercise leads to psychological and immunological improvement in pre-AIDS patients. Proceedings of the 5th International AIDS Conference; June 1989; Montreal, Canada.
79. Stout-Shaffer SR. *Effects of Relaxation Training on Physiological and Psychological Measures of Distress and Quality of Life in HIV-Seropositive Subjects.* Ann Arbor, MI: ProQuest Information and Learning; 2000.
80. Stringer WW, Berezovskaya M, O'Brien WA, Beck CK, Casaburi R. The effect of exercise training on aerobic fitness, immune indices, and quality of life in HIV+ patients. *Med Sci Sports Exerc.* 1998;30(1):11–16.
81. Taylor DN. Effects of a behavioral stress-management program on anxiety, mood, self-esteem, and T-cell count in HIV positive men. *Psychol Rep.* 1995;76(2):451–457.
82. Terry L, Sprinz E, Ribeiro JP. Moderate and high intensity exercise training in HIV-1 seropositive individuals: a randomized trial. *Int J Sports Med.* 1999;20(2):142–146.

83. Weiss JJ, Mulder CL, Antoni MH, de Vroome EMM, Garssen B, Goodkin K. Effects of a supportive-expressive group intervention on long-term psychosocial adjustment in HIV-infected gay men. *Psychoth Psychosom.* 2003;72(3):132–140.
84. Paterson DL, Swindells S, Mohr J, et al. Adherence to protease inhibitor therapy and outcomes in patients with HIV infection. *Ann Intern Med.* 2000;133(1):21–30.
85. Clavel F, Hance AJ. HIV drug resistance. *N Eng J Med.* 2004;350(10):1023–1035.
86. Johnson MO, Dilworth SE, Taylor JM, Neilands TB. Improving coping skills for self-management of treatment side effects can reduce antiretroviral medication nonadherence among people living with HIV. *Ann Behav Med.* 2011;41(1):83–91.
87. Antoni MH, Carrico AW, Durán RE, et al. Randomized clinical trial of cognitive behavioral stress management on human immunodeficiency virus viral load in gay men treated with highly active antiretroviral therapy. *Psychosom Med.* 2006;68(1):143–151.
88. Brown JL, DiClemente RJ. Secondary HIV prevention: novel intervention approaches to impact populations most at risk. *Curr HIV/AIDS Rep.* August 12, 2011. [Epub ahead of print]. doi: 10.1007/s11904-011-0092-6.
89. Eaton LA, Kalichman SC. Changes in transmission risk behaviors across stages of HIV disease among people living with HIV. *J Assoc Nurses AIDS Care.* 2009;20(1):39–49.
90. Weinhardt LS. HIV diagnosis and risk behavior. In: Kalichman SC, editor. *Positive Prevention: Reducing HIV Transmission Among People Living with HIV/AIDS.* New York, NY: Springer Science; 2005.
91. Kalichman SC, Pellowski J, Turner C. Prevalence of sexually transmitted co-infections in people living with HIV/AIDS: systematic review with implications for using HIV treatments for prevention. *Sex Transm Infect.* 2011;87(3):183–190.
92. Brenner B, Routy JP, Quan Y, et al. Persistence of multidrug-resistant HIV-1 in primary infection leading to superinfection. *AIDS.* 2004;18(12):1653–1660.
93. Campbell MS, Gottlieb GS, Hawes SE, et al. HIV-1 superinfection in the antiretroviral therapy era: are seroconcordant sexual partners at risk? *PLoS One.* 2009;4(5):e5690.
94. Carrico AW, Chesney M, Johnson MO, et al. Randomized controlled trial of a cognitive-behavioral intervention for HIV-positive persons: an investigation of treatment effects on psychosocial adjustment. *AIDS Behav.* 2009;13(3):555–563.
95. Kabat-Zinn J. An outpatient program in behavioral medicine for chronic pain patients based on the practice of mindfulness meditation: theoretical considerations and preliminary results. *Gen Hosp Psychiatry.* 1982;4(1):33–47.
96. Grossman P, Niemann L, Schmidt S, Walach H. Mindfulness-based stress reduction and health benefits. A meta-analysis. *J Psychosom Res.* 2004;57(1):35–43.
97. Creswell JD, Myers HF, Cole SW, Irwin MR. Mindfulness meditation training effects on CD4+ T lymphocytes in HIV-1 infected adults: a small randomized controlled trial. *Brain Behav Immun.* 2009;23(2):184–188.
98. Bormann JE, Carrico AW. Increases in positive reappraisal coping during a group-based mantram intervention mediate sustained reductions in anger in HIV-positive persons. *Int J Behav Med.* 2009;16(1):74–80.
99. McCain NL, Gray DP, Elswick RK, et al. A randomized clinical trial of alternative stress management interventions in persons with HIV infection. *J Consult Clin Psychol.* 2008;76(3):431–441.
100. Brown JL, Venable PA, Carey MP, Elin L. The development of a computer-administered cognitive-behavioral intervention to promote stress management among HIV+ women. *J Cogn Psychother.* 2010;24(4):265–280.
101. Brown JL, Venable PA, Carey MP, Elin L. Computerized stress management training for HIV+ women: a pilot intervention study. *AIDS Care.* 2011. In press. doi: 10.1080/09540121.2011.569699.
102. Heckman TG, Sikkema KJ, Hansen N, Kochman A, Heh V, Neufeld S. A randomized clinical trial of a coping improvement group intervention for HIV-infected older adults. *J Behav Med.* 2011;34(2):102–111.
103. Liebschutz JM, Feinman G, Sullivan L, Stein M, Samet J. Physical and sexual abuse in women infected with the human immunodeficiency virus: increased illness and health care utilization. *Arch Intern Med.* 2000;160(11):1659–1664.
104. Meade CS, Drabkin AS, Hansen NB, Wilson PA, Kochman A, Sikkema KJ. Reductions in alcohol and cocaine use following a group coping intervention for HIV-positive adults with childhood sexual abuse histories. *Addiction.* 2010;105(11):1942–1951.
105. Sikkema KJ, Hansen NB, Kochman A, et al. Outcomes from a group intervention for coping with HIV/AIDS and childhood sexual abuse: reductions in traumatic stress. *AIDS Behav.* 2007;11(1):49–60.
106. Sikkema KJ, Wilson PA, Hansen NB, et al. Effects of a coping intervention on transmission risk behavior among people living with HIV/AIDS and a history of childhood sexual abuse. *J Acquir Immune Defic Syndr.* 2008;47(4):506–513.
107. Centers for Disease Control and Prevention (CDC). HIV/AIDS among women: fact sheet. 2006. Available from: <http://www.cdc.gov/hiv/topics/women/resources/factsheets/women.htm>. Accessed October 6, 2011.
108. Antoni MH, Pereira DB, Marion I, et al. Stress management effects on perceived stress and cervical intraepithelial neoplasia in low-income HIV infected women. *J Psychosom Res.* 2008;65(4):389–401.
109. Weiss SM, Tobin JN, Antoni M, et al. Enhancing the health of women living with HIV: the SMART/EST Women's Project. *Int J Womens Health.* 2011;3:63–77.
110. Luce BR, Kramer JM, Goodman SN, et al. Rethinking randomized clinical trials for comparative effectiveness research: the need for transformational change. *Ann Intern Med.* 2009;151(3):206–209.
111. Sullivan P, Goldmann D. The promise of comparative effectiveness research. *JAMA.* 2011;305(4):400–401.

## Neurobehavioral HIV Medicine

### Publish your work in this journal

Neurobehavioral HIV Medicine is an international, peer-reviewed, open access journal focusing on advances in research in HIV/AIDS, with specific reference to the neurological, psychiatric and behavioral consequences of the disease, concomitant infections and specific antiretroviral therapy. The manuscript

Submit your manuscript here: <http://www.dovepress.com/journal-of-neurobehavioral-hiv-medicine-journal>

management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress