


CD44+ Circulating Tumor Endothelial Cells Indicate Poor Prognosis in Pancreatic Ductal Adenocarcinoma After Radical Surgery: A Pilot Study

Cheng Xing
Yatong Li
Cheng Ding
Shunda Wang 
Hanyu Zhang
Lixin Chen
Pengyu Li
Menghua Dai

Department of General Surgery, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing, 100730, People's Republic of China

Background: Circulating tumor endothelial cells (CTECs) are cells that originate from tumor endothelial cells (TECs) of blood vessels and are shed into peripheral blood. Some studies have shown that CTECs are associated with tumor angiogenesis, growth and indicate prognosis in patients with malignant solid tumor. However, the role of CTECs especially the phenotype of CTECs in pancreatic adenocarcinoma (PDAC) is still not clear. We investigated the relationship between CTECs and patients' prognosis.

Methods: A total of 73 patients with resectable PDAC were enrolled in our research and underwent radical surgery. Peripheral venous blood samples were collected before surgery, on postoperative day (POD) 7 and on postoperative month (POM) 1, respectively. We used integrated subtraction enrichment and immunostaining-fluorescence in situ hybridization (SE-iFISH) platform to identify and enumerate CTECs. Immunofluorescence was used to identify CTECs expressing CD44 and vimentin.

Results: In patients with early tumor recurrence (DFS < 6 months), the preoperative CD44+ CTEC levels showed significantly higher ($P = 0.023$). Univariate and multivariate analysis showed that history of diabetes [HR 2.656 (1.194–5.908), $P = 0.017$], numbers of positive lymph nodes [HR 1.871 (1.388–2.522), $P < 0.001$], preoperative numbers of CD44+ CTECs [HR 1.216 (1.064–1.390), $P = 0.004$], and POM1 CA19-9 level [HR 1.002 (1.001–1.002), $P < 0.001$] were independent prognostic factors for DFS.

Conclusion: The detection of CD44+CTECs in patients with resectable PDAC preoperatively could be an independent predictor of shorter DFS after radical surgery.

Keywords: circulating tumor endothelial cells, pancreatic adenocarcinoma, prognostic factor, CD44

Introduction

Pancreatic ductal adenocarcinoma (PDAC) is one of the primary causes of cancer-related mortality worldwide.^{1,2} The only potentially curative treatment for PDAC is surgical resection. While patients with localized disease have been treated with integrated therapy based on radical surgery, the 5-year survival rate still remains 7%-8%.³ Many of these patients developed early postoperative metastatic recurrences due to micrometastatic foci occurred at the time of surgery. Many patients are understaged at diagnosis for the reason that these micrometastatic foci are not identified preoperatively.

Correspondence: Menghua Dai
Department of General Surgery, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences & Peking Union Medical College, No. 1, Shuai Fu Yuan, Dongcheng District, Beijing, 100730, People's Republic of China
Tel +86-10-69152600
Email daimh@pumch.cn

Circulating tumor endothelial cells (CTECs)^{4,5} are cells that originate from tumor endothelial cells (TECs)^{6,7} of blood vessels and are shed into peripheral blood. It has been extensively investigated that TECs shows clinical significance in tumor growth and metastasis.^{8–10} Previous studies demonstrated that CTCs may play a part in tumor angiogenesis.^{11,12} CTCs have been recently reported to express multiple biomarkers such as tumor or stemness markers in patients with breast cancer¹³ and non-small-cell lung cancer (NSCLC).^{14,15}

Several recent studies indicated that some circulating tumor cells (CTCs) have characteristics similar to circulating tumor stem cells (CTSCs).¹⁶ Since cancer growth depends on cancer stem cells (CSCs), which are commonly chemoresistant, CTSCs might be a more sensitive prognostic factor comparing to CTCs.¹⁷ Cluster of differentiation 44 (CD44) was a useful stemness marker as reported previously.¹⁸ A study demonstrated that gastric cancer patients with CTCs expressing CD44 were more inclined to develop disease recurrence and metastasis.¹⁹ Furthermore, another study showed that CTCs labeled with CD44 were independent prognostic factor of decreased disease-free survival (DFS) and overall survival (OS).²⁰ A hypothesis has been proposed that CD44+ CTCs represent a more aggressive subset of CTCs with a more stem cell-like phenotype. Vimentin was considered as an important epithelial–mesenchymal transition (EMT) markers, and was correlated with cancer recurrence as well as decreased overall survival.^{21,22} Some studies found that CTCs expressing vimentin were more invasive and could promote metastases.²³ However, the role of the stemness phenotype or mesenchymal phenotype of CTCs in patients with PDAC is still unclear.

The aim of the present study was to detect different phenotypes of CTCs in the peripheral blood of patients with PDAC, and to determine the prognostic value of CTCs.

Methods

Patients and Samples

From November 2017 to October 2020, patients with resectable PDAC who underwent radical surgery including pancreaticoduodenectomy, distal pancreatectomy with splenectomy, or total pancreatectomy were considered eligible for this study. 6 mL of peripheral venous blood samples were collected 1 day before surgery, postoperative day (POD) 7 and in postoperative month

(POM) 1, respectively. All enrolled patients signed consent forms before blood sample collection. The study was approved by the Ethics Review Committees of Peking Union Medical College Hospital (PUMCH), and performed in compliance to the Declaration of Helsinki Principles. Inclusion criteria: (1) patients who underwent radical pancreatectomy (pancreaticoduodenectomy, distal pancreatectomy with splenectomy or total pancreatectomy); (2) patients' postoperative pathological diagnosis was PDAC; (3) patients gave informed consent, complied with sample collection and follow-up. Exclusion criteria: (1) tumor was found unresectable or distal metastasis preoperatively or intraoperatively; (2) patients' postoperative pathological diagnosis was not PDAC; (3) patients withdrawn informed consent, or patients were unable to comply with sample collection or follow-up. We collected the data regarding patients' demographics, perioperative factors, pathologic details, surgical outcomes, survival, neoadjuvant or adjuvant therapy. The diagnosis of PDAC was confirmed by 2 independent pathologists. Pathological information including tumor size, differentiate degree, nodal status, margin status, perineural and perivascular invasion was recorded. Patients were followed up every 3–6 months postoperatively by the outpatient department. Contrast computed tomography of chest, abdomen and pelvis were routinely performed every 3–6 months to monitor the recurrence of tumor. The physicians were blinded to CTCs or CTCs information to ensure that the treatment plan was independent.

Subtraction Enrichment (SE)

We use Subtraction Enrichment Kit from Cytelligen (San Diego, CA, USA) to collect CTCs and CTCs. The procedure was performed according to the manufacturer's protocol. The details of procedure was described in our previous study.²⁴

Tumor Marker

Immunostaining-Chromosome Fluorescence in situ Hybridization (i-FISH)

6-channel tri-marker (CD44/vimentin/CD31)-iFISH was used to identify CTCs or CTCs according to the manufacturer's protocol (Cytelligen).⁵ The dried monolayer cells on coated slides were rinsed and incubated with PBS for 3 minutes. Then, the cells were hybridized with centromere probe 8 (CEP8) (Abbott Laboratories, Chicago, IL,

USA) for 4 hours. S500 StatSpinThermoBrite Slide Hybridization/Denaturation System (Abbott Molecular, Abbott Park, IL, USA) was used to identify aneuploid tumor cells. Samples were subsequently incubated with the indicated post-fluorescence labeled monoclonal antibodies (1:200 dilution), including Alexa Fluor (AF) 594-CD45 (ATCC, Manassas, VA, USA, Clone 9.4), CD44 (MiltenyiBiotec, San Diego, CA, USA), Cy5-CD31 (Abcam, Burlingame, CA, USA, Catalog No. EP3095), and Cy7-vimentin (Abcam, Catalog No. EPR3776) for 20 min in the dark.²⁵ After washing, we use mounting media with DAPI (Vector Laboratories, Burlingame, CA, USA) to mount the samples, then scanned the images of CTCs/CTECs for analysis.

Image Scanning and Cell Counting

Metafer-iFISH system (Carl Zeiss, MetaSystems, and Cytelligen⁵) was used to scan images and analyze CTCs and CTECs. CTCs were defined as DAPI+/CD45-/CD31-/CD44 (+ or -)/vimentin (+ or -) with aneuploid CEP8, CTECs were defined as DAPI+/CD45-/CD31+/CD44 (+ or -)/vimentin (+ or -) with aneuploid CEP8. Circulating tumor microemboli (CTMs) were defined as multiple CTCs (≥ 3) aggregated into clusters. Circulating tumor endothelial microemboli (CTEMs) were defined as multiple CTECs (≥ 3) aggregated into clusters. Automated CTC classification and statistical analysis were applied upon cell size, cell cluster, and chromosome ploidy.

Statistical Analyses

Categorical variables were expressed as a percentage and compared using χ^2 test or Fisher's exact test. Continuous variables were expressed as the mean \pm standard deviation or the median (interquartile range) and compared using Student's *t*-test or Mann-Whitney *U*-test. Kaplan-Meier method and comparison using a Log rank test were applied to estimate survival data. Prognostic factors for disease-free survival (DFS) or overall survival (OS) were identified using univariate and multivariate Cox proportional hazards models. In univariate Cox regression analysis, all variables with *P* value < 0.10 were included in multivariate Cox model with backward selection. Unless otherwise specified, 2-sided statistical tests were used and a *P* value less than 0.05 was considered statistically significant. The hazard ratios (HR) and their 95% confidence intervals (95% CI) were calculated to show the variation. Spearman correlation coefficients were used to evaluate the correlation between pathological characteristics and

CTEC levels. The optimal cutoff values of independent prognostic factors were assessed by the X-Tile software (version 3.6.1).²⁶ SPSS version 22.0 (IBM, Armonk, NY, USA) was used for statistical analysis.

Results

Detection of CTECs and CTCs by SE-iFISH


CTECs and CTCs were enriched and identified in the peripheral blood of 73 patients by applying the sorting method. CTECs and CTCs can be classified by the platform with the detection of aneuploid CEP8. The CTCs identification criteria were as follows (Figure 1A): nuclear DAPI+, CD45-, CD31-, aneuploid CEP8 positive, and CTCs tumor marker (CD44 and vimentin) positive or negative. CTECs with the same aneuploidy of CEP8 were also found under the fluorescence microscope. The definition of CTECs was similar to CTCs except the endothelial cell marker CD31 was strongly positive (Figure 1B).

Participant Characteristics and Categorical Analysis of CTECs and CTCs

During the study period, we evaluated 125 patients whose preoperative diagnosis were considered PDAC. Fifty-two cases did not meet the inclusion criteria: 29 patients had unresectable tumors, 19 patients' postoperative pathological diagnosis were not PDAC, 3 patients died from postoperative complication within 1 month, 1 patient was lost to follow-up. A total of 73 cases were enrolled into the study. The flow diagram is presented in Figure 2. The median (range) follow-up duration was 10.8 (1.2–31.8) months. Patients' demographic characteristics, surgical details and pathological data are summarized in Table 1.

Table 2 and Figure 3 describe the dynamics of CA 19-9, CTCs and CTECs in different phases. The mean total CA19-9 level, numbers of vimentin+ CTCs and numbers of vimentin+ CTECs showed a decreasing trend at POD7 and then increased at POM1, respectively. The mean total numbers of CTCs, CD44+ CTCs, CTECs and CD44+ CTECs increased at POD7 and then decreased at POM1, respectively. We divided the patients into early recurrence (ER) group (DFS < 6 months) and late recurrence (LR) group (DFS ≥ 6 months) according to the DFS. We compared the level of CTCs and CTECs between the ER group and the LR group. The mean preoperative CD44 + CTECs level was significantly higher in the ER group

A



B




Figure 1 Detection of different subtypes of CTCs and CTECs expressing CD44 and vimentin in PDAC patients by SE-iFISH. **(A)** A CTC (DAPI+/CD45-/CD31-/vimentin-/CEP8+) has a positive expression of CD44; A CTC cluster (DAPI+/CD45-/CD31-/vimentin-/CEP8+) has a positive expression of CD44; A CTC (DAPI+/CD45-/CD31-/vimentin+/CEP8+) has a negative expression of CD44; A CTC cluster (DAPI+/CD45-/CD31+/vimentin+/CEP8+) has a negative expression of CD44. **(B)** A CTEC (DAPI+/CD45-/CD31+/vimentin-/CEP8+) has a positive expression of CD44; A CTEC cluster (DAPI+/CD45-/CD31+/vimentin+/CEP8+) has a positive expression of CD44; A CTEC (DAPI+/CD45-/CD31+/vimentin-/CEP8+) has a negative expression of CD44; A CTEC cluster (DAPI+/CD45-/CD31+/vimentin+/CEP8+) has a negative expression of CD44.




Figure 2 Flow diagram of patient enrollment.

than it in the LR group (3.00 vs 0.56, $P = 0.023$). The mean preoperative vimentin+ CTC levels (0.06 vs 0.63, $P = 0.012$) and POD7 vimentin+ CTC levels (0 vs 0.21, $P = 0.022$) showed significantly lower in the ER group than it in the LR group. The POM1 CA19-9 level was significantly higher in the ER group than it in the LR group (415.77U/mL vs 28.52U/mL, $P = 0.002$).

Detection of CD44+ CTECs Associated with Poor DFS in Enrolled Patients

Univariate Cox regression analysis was applied to investigate the association between DFS or OS, as well as demographic characteristics, pathological data, perioperative details, CTCs and CTECs levels. History of smoke ($P = 0.030$), history of diabetes ($P = 0.011$), neoadjuvant chemotherapy ($P = 0.020$), type of operation (total pancreatectomy, $P = 0.001$), differentiation of tumor (poor, $P = 0.042$), positive lymph nodes numbers ($P < 0.001$), preoperative CD44+ CTECs level ($P = 0.001$), POD7 CD44+ CTCs level ($P = 0.011$), POM1 CA19-9 level ($P < 0.001$) were identified as statistically significant influential factors for DFS (Tables 3 and 4). History of diabetes ($P = 0.002$), neoadjuvant chemotherapy ($P = 0.009$), type of operation (total pancreatectomy, $P = 0.001$), preoperative CD44+ CTECs level ($P = 0.010$), POD7 CD44+ CTCs level ($P = 0.011$), POM1 CA19-9 level ($P = 0.007$) was identified as statistically significant influential factors for OS (Tables 5 and 6).

According to the pre-specified criteria, potential prognostic factors with P values < 0.10 in the univariate analysis were included in the multivariate Cox regression model. History of diabetes [HR 2.656 (1.194–5.908), $P = 0.017$], positive lymph nodes number [HR 1.871 (1.388–2.522), $P < 0.001$], preoperative CD44+ CTECs level [HR 1.216 (1.064–1.390), $P = 0.004$] and POM1 CA19-9 level [HR 1.002 (1.001–1.002), $P < 0.001$] were identified as independent prognostic factors for DFS (Table 3). We determined the optimal cutoff value of the 3 independent prognostic factors for DFS. The cutoff value for positive lymph nodes number was 2 (Figure S1A). The cutoff value for preoperative CD44+ CTECs number was 3 (Figure S1B), and it for POM1 CA19-9 level was 89.6 U/mL (Figure S1C). History of diabetes [HR 7.227 (1.916–27.265), $P = 0.004$] and POM1 CA19-9 level [HR 1.001 (1.000–1.002), $P = 0.026$] were identified as independent prognostic factors for OS (Tables 5 and 6). The cutoff value for POM1 CA19-9 level was 131.9U/mL (Figure S1D). Then, we estimated incidence of disease recurrence or death in different risk-stratified subgroups using Kaplan–Meier method. History of diabetes (11.2 months vs 5.8 months, $P = 0.009$, Figure 4A), positive lymph nodes > 2 (11.3 months vs 5.3 months, $P < 0.001$, Figure 4B), preoperative CD44+ CTECs > 3 (11.1 months vs 5.1 months, $P = 0.002$, Figure 4C), POM1 CA19-9 > 89.6 U/mL (11.3 months vs 4.2 months, $P < 0.001$, Figure 4D) were significantly associated with increased risk of tumor recurrence. History of diabetes (16.6 months vs 11.7

Table 1 Demographic Characteristics, Surgical Details and Pathological Details

Baseline Characteristics	
Age (yr, \pm SD)	59.41 \pm 9.62
Gender (Female)	33 (45.2%)
BMI (kg/m ²)	23.27 \pm 3.24
Smoke	27 (37.0%)
Alcohol	20 (27.4%)
Pancreatitis	4 (5.5%)
Diabetes	18 (24.7%)
Cardiovascular disease	4 (5.5%)
Obstructive jaundice	21 (28.8%)
Neoadjuvant chemotherapy	11 (15.1%)
Neoadjuvant radiotherapy	0
Surgical Details	
Type of operation	
Pancreaticoduodenectomy	48 (65.8%)
Distal pancreatectomy	23 (31.5%)
Total pancreatectomy	2 (2.7%)
Laparoscopic	26 (35.6%)
Operation time (min)	330.60 \pm 94.00
Blood loss (mL)	678.08 \pm 548.03
Transfusion (RBCs)	27 (37%)
Postoperative complications	29 (39.7%)
Adjuvant chemotherapy	67 (91.8%)
Adjuvant radiotherapy	7 (9.6%)
Pathological Details	
Tumor size	3.58 \pm 1.76
Differentiation (poor)	31 (42.5%)
Margin status (not R0)	26 (35.6%)
Vascular infiltration	25 (34.2%)
Perineural infiltration	61 (83.6%)
Carcinoma embolus	34 (46.6%)
Total lymph nodes number	22.70 \pm 11.62
Positive lymph nodes number	1.11 \pm 1.45

Abbreviation: RBCs, red blood cells.

months, $P = 0.001$, Figure 4E) and POM1 CA19-9 > 131.9U/mL (15.4 months vs 7.6 months, $P < 0.001$, Figure 4F) were significantly associated with increased risk of death. Spearman correlation analysis showed no significant correlation between preoperative CD44 +CTECs and pathological characteristics (Supplemental Material Table S1)

Discussion

In the present pilot study, we identified both CTECs and CTCs in patients with PDAC, and investigated their potential clinical impact. We found that CD44+ CTECs might

be related to early recurrence and poor prognosis in patients with PDAC after radical surgery. Many potential factors may affect the prognosis of pancreatic cancer, including patient factors (CA19-9 level, lymph nodes, history of diabetes, etc) and treatment-related factors (surgical margin status, postoperative adjuvant therapy, etc).²⁷ Our study identified 3 independent risk factors (history of diabetes, positive lymph nodes and POM1 CA19-9 level) associated with DFS and 1 independent risk factor (history of diabetes) associated with OS, which is agreement with the previous findings. Meanwhile, various studies reported that CTCs and CTECs were associated with the prognosis of pancreatic cancer.^{12–15} The present study demonstrated preoperative CD44+ CTECs number was significantly higher in patients whose DFS < 6 months and could be an independent factor for shorter DFS. CTECs are cells that originate from tumor endothelial cells (TECs) of blood vessels and are shed into peripheral blood. As indicated in some studies, elevated CTECs count may be a prognostic factor in adults with malignant diseases, such as breast cancer¹³ and lung cancer.^{14,15} Some other studies explored the role of specific phenotype CTECs in clinical utilities, such as drug therapy effect evaluation. PD-L1+CTECs were found to be associated with a shorter progression-free survival in NSCLC patients receiving PD-L1 immunotherapy, and PD-L1 therapy would facilitate the karyotype shifting of CTECs.²⁸ The levels of aneuploid CTEC may be influenced by neoadjuvant chemotherapy in patients with locally advanced breast cancer.²⁹ CTECs expressing stemness marker of CD44 in PDAC patients remain to be investigated.¹⁰ Poruk KE et al proved that CTCs labeled with stemness marker such as CD44 are independent predictors of decreased disease-free and overall survival.²⁰ The CD44+ CTECs might also present a characteristics of stem cells or tumor initiating cells (TIC), suggest a possible mechanism for metastatic spread. Our finding based on clinical data provide fundamental evidence for the role of CD44+ CTECs in PDAC.

In the process from TECs to CTECs, the inducible endothelial-to-mesenchymal transition (Endo-MT) may occur, similar to the epithelial-to-mesenchymal transition (EMT) for CTCs.³⁰ Cao et al proved that CTECs could bind to metastatic cancer cells and inhibit apoptosis of tumor cells.³¹ CTECs has also shown its potential in the treatment of malignant tumors. Previous reports indicated that CTEC levels were associated with the clinical outcome in patients of breast tumor under chemotherapy combined with anti-VEGF treatment.³² The conclusion

Table 2 CTCs and CTECs Details

		Total	DFS<6 Months (n=16)	DFS≥6 Months (n=57)	P value
CA 19–9	Preoperative	238.45±327.35	270.87±273.99	229.35±342.47	0.657
	POD 7	59.45±81.38	75.48±64.03	56.10±84.82	0.521
	POM 1	117.40±427.72	415.77±846.19	28.52±84.82	0.002
CTCs	Preoperative	11.49±10.72	15.44±12.12	10.39±10.14	0.096
	POD 7	49.33±93.43	68.31±161.06	44.00±64.36	0.361
	POM 1	30.42±53.69	49.23±84.48	25.11±41.00	0.336
CD44+ CTCs	Preoperative	0.33±0.80	0.50±1.09	0.28±0.70	0.336
	POD 7	9.55±73.23	39.94±156.29	1.02±3.74	0.335
	POM 1	0.43±1.21	0.54±1.66	0.40±1.07	0.721
Vimentin+ CTCs	Preoperative	0.51±1.43	0.06±0.25	0.63±1.59	0.012
	POD 7	0.16±0.60	0.00±0.00	0.21±0.67	0.022
	POM 1	0.64±1.48	0.46±1.66	0.69±1.44	0.631
CTMs	Preoperative	0.52±1.40	0.56±1.54	0.51±1.37	0.894
	POD 7	0.74±1.23	0.94±1.28	0.68±1.22	0.473
	POM 1	0.46±0.91	0.62±1.04	0.41±0.88	0.487
CTECs	Preoperative	6.90±11.13	10.13±10.57	6.00±11.21	0.193
	POD 7	12.40±22.78	13.13±26.22	12.19±21.98	0.886
	POM 1	6.19±12.84	10.08±19.80	5.07±10.04	0.218
CD44+ CTECs	Preoperative	1.10±2.27	3.00±3.81	0.56±1.19	0.023
	POD 7	2.22±7.87	5.50±15.85	1.30±2.90	0.308
	POM 1	1.09±3.59	1.85±6.06	0.87±2.53	0.580
Vimentin+ CTECs	Preoperative	1.33±6.21	0.88±3.50	1.46±6.80	0.744
	POD 7	0.30±1.77	0.00±0.00	0.39±2.00	0.447
	POM 1	0.90±3.76	0.31±0.75	1.07±4.25	0.527
CTEMs	Preoperative	0.74±2.93	1.94±5.93	0.40±1.03	0.320
	POD 7	0.23±0.75	0.13±0.34	0.26±0.83	0.522
	POM 1	0.31±1.23	0.08±0.27	0.38±1.38	0.443

Note: Bold value: $P < 0.05$, considered statistically significant.

Abbreviations: CTCs, circulating tumor cells; CTMs, circulating tumor microemboli; CTECs, circulating tumor endothelial cells; CTEMs, circulating tumor endothelial microemboli.

also supports that CTEC levels may be associated with prognosis of PDAC patients receiving gemcitabine chemotherapy.³³ However, Kindler et al proved that gemcitabine combined bevacizumab³⁴ or axitinib (a VEGF inhibitor)³⁵ does not improve advanced pancreatic cancer patients' survival through Phase III clinical trial. Based

on the present study, we speculate that the conclusion of these clinical trials may be related to the fact that CTECs and their various subtypes were not identified in the past. Subgroup analysis based on CTEC may lead to a different conclusion. Some studies^{36,37} also proposed that tumor cells may become less sensitive to anti-




Figure 3 The dynamics of (A) CA 19-9 levels, numbers of (B) CTCs, (C) CD44+ CTC, (D) vimentin+ CTC, (E) CTM, and numbers of (F) CTECs (G) CD44+ CTEC, (H) vimentin+ CTEC, (I) CTEM in different phases (preoperative, POD7 and POM1). The patients were divided into early recurrence (ER) group (DFS<6 months) and late recurrence (LR) group (DFS≥6 months) according to the DFS.

angiogenic agents in hypoxia or nutrient deprivation. Signals from the stromal compartment may play a major role in refractoriness of anti-angiogenic therapy and could potentially acquired resistance to VEGF inhibitors.³⁸ We hypothesized that CTECs especially CD44+CTECs may lead to distant metastases, cross-talk with the stromal cells in tumor microenvironment (TME), then facilitate tumor growth, angiogenesis and drug resistance. Recent studies showed that the inhibition of CD44 signaling could lead to effective therapeutic

responses in PDAC models.³⁹ Detection and characterization of CD44+ CTECs show potential to assist in evaluating the efficacy of the classic chemotherapy regimen combining angiogenesis inhibitors and anti-CD44 therapy.

It is controversial whether CTCs and CTECs can be driven into the blood to disseminate tumor cells by surgical manipulation.⁴⁰ In present study, most phenotype of CTCs and CTECs showed an increasing trend at POD7 and then decreased at POM1. But CTCs and CTECs at

Table 3 Univariate and Multivariate Cox Regression Analyses of Prognosis Factors (Clinicopathological Factors) for Disease Free Survival

Prognostic Factor	Univariate Analysis		Multivariate Analysis		Cutoff Value
	HR (95% CI)	P value	HR (95% CI)	P value	
Age (years)	1.000 (0.968–1.033)	0.984			
Gender (Female)	1.670 (0.921–3.027)	0.091			
BMI (kg/m ²)	0.998 (0.915–1.089)	0.967			
Obstructive jaundice	1.158 (0.613–2.186)	0.652			
Smoke	2.070 (1.072–3.997)	0.030			
Alcohol	1.054 (0.542–2.051)	0.876			
Pancreatitis	0.825 (0.198–3.432)	0.791			
Diabetes	2.422 (1.223–4.795)	0.011	2.656 (1.194–5.908)	0.017	
Cardiovascular disease	1.693 (0.513–5.583)	0.387			
Neoadjuvant chemotherapy	2.714 (1.173–6.279)	0.020			
Type of operation		0.003			
Pancreaticoduodenectomy	Ref.				
Distal pancreatectomy	1.314 (0.681–2.538)	0.416			
Total pancreatectomy	20.262 (3.614–113.587)	0.001			
Laparoscopic	0.761 (0.411–1.410)	0.385			
Operation time (min)	0.999 (0.995–1.002)	0.470			
Blood loss (mL)	1.000 (1.000–1.001)	0.827			
Transfusion (RBCs)	1.116 (0.602–2.068)	0.727			
Postoperative complication	1.057 (0.562–1.989)	0.863			
Adjuvant chemotherapy	1.800 (0.431–7.511)	0.420			
Adjuvant radiotherapy	0.562 (0.201–1.578)	0.274			
Tumor size	1.029 (0.910–1.165)	0.647			
Differentiation (poor)	1.862 (1.023–3.389)	0.042			
Margin status (not R0)	1.174 (0.632–2.182)	0.611			
Vascular infiltration	1.018 (0.538–1.926)	0.957			
Perineural infiltration	1.852 (0.661–5.189)	0.241			
Carcinoma embolus	1.048 (0.575–1.911)	0.879			
Total lymph nodes (numbers)	0.992 (0.961–1.024)	0.607			
Positive lymph nodes (numbers)	1.749 (1.373–2.228)	<0.001	1.871 (1.388–2.522)	<0.001	2

Note: Bold value: $P < 0.05$, considered statistically significant.

Abbreviation: RBCs: red blood cells.

POD7 or POM1 did not correlate with prognosis. We speculate that the postoperative elevation may be due to an inflammatory response or decreased immunity of

patients. Moreover, in the ER subgroup, the CD44+ CTECs level was significantly higher preoperatively while the CA 19–9 levels significantly increased to

Table 4 Univariate and Multivariate Cox Regression Analyses of Prognosis Factors (CA19-9, CTCs and CTECs) for Disease Free Survival

Prognostic Factor	Univariate Analysis		Multivariate Analysis		Cutoff Value
	HR (95% CI)	P value	HR (95% CI)	P value	
Preoperative CA19-9	1.000 (0.999–1.001)	0.686			
Preoperative CTC	0.999 (0.974–1.025)	0.956			
Preoperative CD44+ CTC	0.974 (0.665–1.426)	0.892			
Preoperative Vimentin+ CTC	0.889 (0.630–1.254)	0.502			
Preoperative CTM	1.027 (0.847–1.245)	0.786			
Preoperative CTEC	0.999 (0.975–1.024)	0.940			
Preoperative CD44+ CTEC	1.207 (1.077–1.353)	0.001	1.216 (1.064–1.390)	0.004	3
Preoperative Vimentin+ CTEC	1.005 (0.967–1.004)	0.810			
Preoperative CTECM	1.062 (0.968–1.166)	0.203			
POD7 CA19-9	1.001 (0.998–1.004)	0.633			
POD7 CTC	1.000 (0.996–1.004)	0.939			
POD7 CD44+ CTC	1.005 (1.001–1.008)	0.011			
POD7 Vimentin+ CTC	0.738 (0.344–1.583)	0.435			
POD7 CTM	0.927 (0.735–1.170)	0.523			
POD7 CTEC	0.980 (0.958–1.002)	0.072			
POD7 CD44+ CTEC	1.028 (0.986–1.072)	0.197			
POD7 Vimentin+ CTEC	1.074 (0.922–1.236)	0.322			
POD7 CTECM	0.571 (0.316–1.030)	0.063			
POM1 CA19-9	1.001 (1.001–1.002)	<0.001	1.002 (1.001–1.002)	<0.001	89.6
POM1 CTC	1.003 (0.996–1.010)	0.399			
POM1 CD44+ CTC	0.889 (0.643–1.228)	0.474			
POM1 Vimentin+ CTC	0.940 (0.698–1.266)	0.683			
POM1 CTM	1.285 (0.940–1.755)	0.116			
POM1 CTEC	0.987 (0.957–1.018)	0.420			
POM1 CD44+ CTEC	0.954 (0.840–1.083)	0.463			
POM1 Vimentin+ CTEC	0.868 (0.680–1.108)	0.257			
POM1 CTECM	0.730 (0.445–1.199)	0.214			

Note: Bold value: $P < 0.05$, considered statistically significant.

Abbreviations: CTCs, circulating tumor cells; CTMs, circulating tumor microemboli; CTECs, circulating tumor endothelial cells; CTEMs, circulating tumor endothelial microemboli.

a high level in POM1. CD44+ CTEC may have potential in detecting pre-existing micrometastatic foci.

Several limitations should be taken into account when interpreting these results in present study. Patients' follow-up

may be further extended to observe significant association between CTEC levels and OS. Considering the convenience and invasiveness for the patients in this study, the blood samples were collected from peripheral vein instead of portal

Table 5 Univariate and Multivariate Cox Regression Analyses of Prognosis Factors (Clinicopathological Factors) for Overall Survival

Prognostic Factor	Univariate Analysis		Multivariate Analysis		Cutoff Value
	HR (95% CI)	P value	HR (95% CI)	P value	
Age (years)	0.967 (0.915–1.021)	0.226			
Gender (Female)	2.325 (0.778–6.952)	0.131			
BMI (kg/m ²)	0.944 (0.800–1.115)	0.497			
Obstructive jaundice	0.424 (0.094–1.916)	0.265			
Smoke	46.586 (0.546–3976.136)	0.090			
Alcohol	0.537 (0.120–2.404)	0.417			
Pancreatitis	1.610 (0.206–12.596)	0.650			
Diabetes	6.307 (1.917–20.758)	0.002	7.227 (1.916–27.265)	0.004	
Cardiovascular disease	0.045 (0.000–857.323)	0.537			
Neoadjuvant chemotherapy	5.083 (1.514–17.063)	0.009			
Type of operation		0.005			
Pancreaticoduodenectomy	Ref.				
Distal pancreatectomy	1.368 (0.411–4.547)	0.609			
Total pancreatectomy	18.353 (3.162–106.523)	0.001			
Laparoscopic	0.212 (0.046–0.978)	0.047			
Operation time (min)	0.999 (0.994–1.005)	0.823			
Blood loss (mL)	1.000 (0.999–1.001)	0.605			
Transfusion (RBCs)	1.601 (0.545–4.700)	0.392			
Postoperative complication	0.774 (0.242–2.481)	0.667			
Adjuvant chemotherapy	0.406 (0.089–1.847)	0.243			
Adjuvant radiotherapy	0.451 (0.059–3.473)	0.445			
Tumor size	1.131 (0.917–1.396)	0.249			
Differentiation (poor)	2.460 (0.878–7.938)	0.084			
Margin status (not R0)	0.997 (0.333–2.984)	0.996			
Vascular infiltration	1.306 (0.435–3.914)	0.634			
Perineural infiltration	2.090 (0.272–16.091)	0.479			
Carcinoma embolus	0.834 (0.287–2.424)	0.739			
Total lymph nodes (numbers)	1.021 (0.979–1.065)	0.331			
Positive lymph nodes (numbers)	1.435 (0.993–2.073)	0.054			

Note: Bold value: $P < 0.05$, considered statistically significant.

Abbreviation: RBCs, red blood cells.

vein. As a result, the count of CTCs and CTECs might be diminished by the percolatory function of the lung capillary bed. The present study was a retrospective study, a selection bias may exist. High volume multicenter study should be

designed to further validate the present conclusion. Future efforts need to be focused on the origin of CTECs and its role in tumor progression using the technique of single-cell sequencing.

Table 6 Univariate and Multivariate Cox Regression Analyses of Prognosis Factors (CA19-9, CTCs and CTECs) for Overall Survival

Prognostic Factor	Univariate Analysis		Multivariate Analysis		Cutoff Value
	HR (95% CI)	P value	HR (95% CI)	P value	
Preoperative CA19-9	1.000 (0.999–1.002)	0.454			
Preoperative CTC	1.011 (0.970–1.054)	0.603			
Preoperative CD44+ CTC	1.122 (0.551–2.283)	0.751			
Preoperative Vimentin+ CTC	0.506 (0.127–2.018)	0.335			
Preoperative CTM	1.004 (0.725–1.391)	0.979			
Preoperative CTEC	1.009 (0.971–1.049)	0.648			
Preoperative CD44+ CTEC	1.204 (1.044–1.387)	0.010			
Preoperative Vimentin+ CTEC	0.987 (0.904–1.079)	0.780			
Preoperative CTECM	1.076 (0.985–1.175)	0.102			
POD7 CA19-9	1.001 (0.996–1.007)	0.659			
POD7 CTC	1.003 (0.999–1.007)	0.195			
POD7 CD44+ CTC	1.005 (1.001–1.008)	0.011			
POD7 Vimentin+ CTC	0.949 (0.303–2.973)	0.929			
POD7 CTM	1.105 (0.768–1.590)	0.589			
POD7 CTEC	1.001 (0.978–1.026)	0.915			
POD7 CD44+ CTEC	1.040 (0.999–1.083)	0.058			
POD7 Vimentin+ CTEC	0.977 (0.571–1.673)	0.933			
POD7 CTECM	0.664 (0.230–1.918)	0.450			
POMI CA19-9	1.001 (1.000–1.002)	0.007	1.001 (1.000–1.002)	0.026	131.9
POMI CTC	1.004 (0.991–1.017)	0.567			
POMI CD44+ CTC	1.030 (0.625–1.696)	0.907			
POMI Vimentin+ CTC	0.759 (0.382–1.510)	0.432			
POMI CTM	0.993 (0.511–1.930)	0.984			
POMI CTEC	1.014 (0.979–1.051)	0.435			
POMI CD44+ CTEC	1.060 (0.942–1.193)	0.335			
POMI Vimentin+ CTEC	0.825 (0.402–1.694)	0.600			
POMI CTECM	0.568 (0.074–4.334)	0.585			

Note: Bold value: $P < 0.05$, considered statistically significant.

Abbreviations: CTCs, circulating tumor cells; CTMs, circulating tumor microemboli; CTECs, circulating tumor endothelial cells; CTEMs, circulating tumor endothelial microemboli.

Conclusion

In summary, our preliminary study demonstrated that preoperative CD44+ CTECs could be an independent factor for shorter DFS in patients with PDAC. We

speculate that CD44+ CTECs may have association with greater angiogenic ability, resulting in greater invasive and metastasis ability of the tumor, leading to a worse prognosis.




Figure 4 The DFS were significantly different between two comparative groups divided by the independent prognostic factors: (A) history of diabetes, (B) positive lymph nodes > 2, (C) preoperative CD44+ CTECs > 3, (D) POM1 CA19-9 > 89.6 U/mL. The OS showed significantly different between two comparative groups divided by (E) history of diabetes, (F) POM1 CA19-9 > 131.9U/mL.

Data Sharing Statement

The datasets analyzed during the current study would be available from the corresponding author on reasonable request.

Ethics Approval

All the patients signed informed consent for the publication of this study, following the regulations outlined in the Declaration of Helsinki. Ethical approval of this study was obtained by the ethics committee of Peking Union Medical College Hospital.

Acknowledgment

We thank all of the staff from Peking Union Medical College Hospital.

Funding

This study was supported by the project of National Natural Science Foundation of China (82073238&81903150), the project of CAMS Innovation Fund for Medical Sciences (2016-I2M-3-005), the project of National Major Research and Development Programs of the Ministry of Science and Technology of China (2017YFC1308602) and the project of Beijing Natural Science Foundation Program (7194305).

Disclosure

The authors declare no conflicts of interest.

References

- Siegel RL, Miller KD, Jemal A. Cancer statistics, 2020. *CA Cancer J Clin.* 2020;70:7–30. doi:10.3322/caac.21590
- Wong MCS, Jiang JY, Liang M, Fang Y, Yeung MS, Sung JY. Global temporal patterns of pancreatic cancer and association with socioeconomic development. *Sci Rep.* 2017;7(1):3165. doi:10.1038/s41598-017-02997-2
- Noone AM, Krapcho M, Miller D, et al. SEER cancer statistics review, 1975–2015, based on November 2017 SEER data submission, posted to the SEER web site. Bethesda, MD: National Cancer Institute; 2018 cited Apr.
- Lin PP. Aneuploid CTC and CEC. *Diagnostics (Basel).* 2018;8(2):26. doi:10.3390/diagnostics8020026
- Lin PP, Gires O, Wang DD, Li L, Wang H. Comprehensive in situ co-detection of aneuploid circulating endothelial and tumor cells. *Sci Rep.* 2017;7(1):9789. doi:10.1038/s41598-017-10763-7
- Akino T, Hida K, Hida Y, et al. Cytogenetic abnormalities of tumor-associated endothelial cells in human malignant tumors. *Am J Pathol.* 2009;175(6):2657–2667. doi:10.2353/ajpath.2009.090202
- Hida K, Klagsbrun M. A new perspective on tumor endothelial cells: unexpected chromosome and centrosome abnormalities. *Cancer Res.* 2005;65(7):2507–2510. doi:10.1158/0008-5472.CAN-05-0002
- Dudley AC. Tumor endothelial cells. *Cold Spring Harb Perspect Med.* 2012;2(3):a006536. doi:10.1101/cshperspect.a006536
- Hida K, Maishi N, Annan DA, Hida Y. Contribution of tumor endothelial cells in cancer progression. *Int J Mol Sci.* 2018;19(5):1272. doi:10.3390/ijms19051272
- Zhao Y, Li J, Li D, et al. Tumor biology and multidisciplinary strategies of oligometastasis in gastrointestinal cancers. *Semin Cancer Biol.* 2020;60:334–343. doi:10.1016/j.semcancer.2019.08.026
- Bertolini F, Shaked Y, Mancuso P, Kerbel RS. The multifaceted circulating endothelial cell in cancer: towards marker and target identification. *Nat Rev Cancer.* 2006;6(11):835–845. doi:10.1038/nrc1971
- Cima I, Kong SL, Sengupta D, et al. Tumor-derived circulating endothelial cell clusters in colorectal cancer. *Sci Transl Med.* 2016;8(345):345ra89. doi:10.1126/scitranslmed.aad7369
- Bidard FC, Mathiot C, Degeorges A, et al. Clinical value of circulating endothelial cells and circulating tumor cells in metastatic breast cancer patients treated first line with bevacizumab and chemotherapy. *Ann Oncol.* 2010;21(9):1765–1771. doi:10.1093/annonc/mdq052
- Wang J, Xiao J, Wei X, et al. Circulating endothelial cells and tumor blood volume as predictors in lung cancer. *Cancer Sci.* 2013;104(4):445–452. doi:10.1111/cas.12097
- Lei Y, Sun N, Zhang G, et al. Combined detection of aneuploid circulating tumor-derived endothelial cells and circulating tumor cells may improve diagnosis of early stage non-small-cell lung cancer. *Clin Transl Med.* 2020;10(3):e128. doi:10.1002/ctm2.128
- Grover PK, Cummins AG, Price TJ, et al. Circulating tumor cells: the evolving concept and the inadequacy of their enrichment by EpCAM-based methodology for basic and clinical cancer research. *Ann Oncol.* 2014;25:1506–1516. doi:10.1093/annonc/mdu018
- Aktas B, Tewes M, Fehm T, et al. Stem cell and epithelial-mesenchymal transition markers are frequently overexpressed in circulating tumor cells of metastatic breast cancer patients. *Breast Cancer Res.* 2009;11(4):R46. doi:10.1186/bcr2333
- Fan CW, Chen T, Shang YN, et al. Cancer-initiating cells derived from human rectal adenocarcinoma tissues carry mesenchymal phenotypes and resist drug therapies. *Cell Death Dis.* 2013;4(10):e828. doi:10.1038/cddis.2013.337
- Li M, Zhang B, Zhang Z, et al. Stem cell-like circulating tumor cells indicate poor prognosis in gastric cancer. *Biomed Res Int.* 2014;2014:981261.
- Poruk KE, Blackford AL, Weiss MJ, et al. Circulating tumor cells expressing markers of tumor-initiating cells predict poor survival and cancer recurrence in patients with pancreatic ductal adenocarcinoma. *Clin Cancer Res.* 2017;23(11):2681–2690. doi:10.1158/1078-0432.CCR-16-1467
- Yu M, Bardia A, Wittner BS, et al. Circulating breast tumor cells exhibit dynamic changes in epithelial and mesenchymal composition. *Science.* 2013;339(6119):580–584. doi:10.1126/science.1228522
- Huber MA, Kraut N, Beug H. Molecular requirements for epithelial-mesenchymal transition during tumor progression. *Curr Opin Cell Biol.* 2005;17(5):548–558. doi:10.1016/j.ceb.2005.08.001
- Poruk KE, Valero V 3rd, Saunders T, et al. Circulating tumor cell phenotype predicts recurrence and survival in pancreatic adenocarcinoma. *Ann Surg.* 2016;264(6):1073–1081. doi:10.1097/SLA.0000000000001600
- Wu G, Zhu R, Li Y, Zhao Y, Dai M. Prognostic significance of circulating tumor microemboli in patients with pancreatic ductal adenocarcinoma. *Oncol Lett.* 2018;15(5):7376–7382. doi:10.3892/ol.2018.8264
- Lin P, Fischer T, Weiss T, Farquhar MG. Calnuc, an EF-hand Ca(2+) binding protein, specifically interacts with the C-terminal alpha5-helix of G(alpha)i3. *Proc Natl Acad Sci U S A.* 2000;97(2):674–679. doi:10.1073/pnas.97.2.674
- Camp RL, Dolled-Filhart M, Rimm DL. X-tile: a new bio-informatics tool for biomarker assessment and outcome-based cut-point optimization. *Clin Cancer Res.* 2004;10(21):7252–7259. doi:10.1158/1078-0432.CCR-04-0713

27. Barhli A, Cros J, Bartholin L, Neuzillet C. Prognostic stratification of resected pancreatic ductal adenocarcinoma: past, present, and future. *Dig Liver Dis.* 2018;50(10):979–990. doi:10.1016/j.dld.2018.08.009
28. Zhang L, Zhang X, Liu Y, et al. PD-L1+ aneuploid circulating tumor endothelial cells (CTECs) exhibit resistance to the checkpoint blockade immunotherapy in advanced NSCLC patients. *Cancer Lett.* 2020;469:355–366. doi:10.1016/j.canlet.2019.10.041
29. Ma G, Jiang Y, Liang M, et al. Dynamic monitoring of CD45-/CD31 +/DAPI+ circulating endothelial cells aneuploid for chromosome 8 during neoadjuvant chemotherapy in locally advanced breast cancer. *Ther Adv Med Oncol.* 2020;12:1758835920918470. doi:10.1177/1758835920918470
30. Piera-Velazquez S, Mendoza FA, Jimenez SA. Endothelial to mesenchymal transition (EndoMT) in the pathogenesis of human fibrotic diseases. *J Clin Med.* 2016;5(4):45. doi:10.3390/jcm5040045
31. Cao Z, Livas T, Kyprianou N. Anoikis and EMT: lethal “Liaisons” during cancer progression. *Crit Rev Oncog.* 2016;21(3–4):155–168. doi:10.1615/CritRevOncog.2016016955
32. Calleri A, Bono A, Bagnardi V, et al. Predictive potential of angiogenic growth factors and circulating endothelial cells in breast cancer patients receiving metronomic chemotherapy plus bevacizumab. *Clin Cancer Res.* 2009;15(24):7652–7657. doi:10.1158/1078-0432.CCR-09-1493
33. Kondo S, Ueno H, Hashimoto J, et al. Circulating endothelial cells and other angiogenesis factors in pancreatic carcinoma patients receiving gemcitabine chemotherapy. *BMC Cancer.* 2012;12:268. doi:10.1186/1471-2407-12-268
34. Kindler HL, Niedzwiecki D, Hollis D, et al. Gemcitabine plus bevacizumab compared with gemcitabine plus placebo in patients with advanced pancreatic cancer: phase III trial of the Cancer and Leukemia Group B (CALGB 80303). *J Clin Oncol.* 2010;28(22):3617–3622. doi:10.1200/JCO.2010.28.1386
35. Kindler HL, Ioka T, Richel DJ, et al. Axitinib plus gemcitabine versus placebo plus gemcitabine in patients with advanced pancreatic adenocarcinoma: a double-blind randomised Phase 3 study. *Lancet Oncol.* 2011;12(3):256–262. doi:10.1016/S1470-2045(11)70004-3
36. Crawford Y, Ferrara N. Tumor and stromal pathways mediating refractoriness/resistance to anti-angiogenic therapies. *Trends Pharmacol Sci.* 2009;30(12):624–630. doi:10.1016/j.tips.2009.09.004
37. Allen E, Miéville P, Warren CM, et al. Metabolic symbiosis enables adaptive resistance to anti-angiogenic therapy that is dependent on mTOR signaling. *Cell Rep.* 2016;15(6):1144–1160. doi:10.1016/j.celrep.2016.04.029
38. Huijbers EJ, van Beijnum JR, Thijssen VL, Sabrkhanly S, Nowak-Sliwinska P, Griffioen AW. Role of the tumor stroma in resistance to anti-angiogenic therapy. *Drug Resist Updat.* 2016;25:26–37. doi:10.1016/j.drug.2016.02.002
39. Molejon MI, Tellechea JI, Moutardier V, et al. Targeting CD44 as a novel therapeutic approach for treating pancreatic cancer recurrence. *Oncoscience.* 2015;2(6):572–575. doi:10.18632/oncoscience.172
40. Sergeant G, Roskams T, van Pelt J, Houtmeyers F, Aerts R, Topal B. Perioperative cancer cell dissemination detected with a real-time RT-PCR assay for EpCAM is not associated with worse prognosis in pancreatic ductal adenocarcinoma. *BMC Cancer.* 2011;11:47. doi:10.1186/1471-2407-11-47

Cancer Management and Research

Dovepress

Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient.

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/cancer-management-and-research-journal>