

Genetic Variations of *CARMN* Modulate Glioma Susceptibility and Prognosis in a Chinese Han Population

Min Xi¹, Gang Zhang¹, Liang Wang², Hu Chen², Li Gao², Luyi Zhang¹, Zhangkai Yang¹, Hangyu Shi¹

¹Department of Neurosurgery, Xi'an Children's Hospital, Xi'an, 710043, People's Republic of China; ²Department of Neurosurgery, Tangdu Hospital, Air Force Medical University, Xi'an, 710038, People's Republic of China

Correspondence: Hangyu Shi, Department of Neurosurgery, Xi'an Children's Hospital, #69, Xijuyuan Lane, Lianhu District, Xi'an, 710043, Shaanxi, People's Republic of China, Tel/Fax +86-15202910508, Email shihangyu723@aliyun.com

Background: This study aimed to evaluate the relationship between *CARMN* polymorphisms and glioma risk and prognosis in a Chinese Han population.

Methods: Seven single nucleotide polymorphisms (SNPs) in *CARMN* were genotyped among 592 glioma patients and 502 healthy controls. Log-additive models were used for risk assessment by the odds ratios (ORs) and 95% confidence intervals (CIs). Univariate and multivariate Cox regression analysis was applied to calculate Hazard ratios (HRs) and 95% CIs for prognosis assessment.

Results: *CARMN* rs13177623 was a protective factor for glioma susceptibility (OR = 0.78, $p = 0.043$). In addition, rs13177623, rs11168100, rs12654195 and rs17796757 were associated with the risk of glioma among the subgroup stratified by age or gender. We also found that G_{rs13177623}G_{rs12654195} haplotype was related to the decreased risk of glioma (OR = 0.61, $p = 0.005$). Importantly, rs13177623 [overall survival (OS): HR = 0.83, $p = 0.047$, and progression free survival (PFS): HR = 0.82, $p = 0.031$], rs12654195 (OS: HR = 0.64, $p = 0.005$ and PFS: HR = 0.65, $p = 0.007$) and rs11168100 (OS: HR = 0.71, $p = 0.035$) were associated with a better prognosis for glioma, especially in grade I-II glioma. In patients with grade III-IV glioma, rs17796757 polymorphism presented an improved OS.

Conclusion: Our results firstly reported the contribution of *CARMN* variants (rs11168100, rs12654195, rs13177623, and rs17796757) to the susceptibility and prognosis of glioma in a Chinese Han population, which provided a novel insight on the relationship between *CARMN* gene and glioma tumorigenesis.

Keywords: glioma, *CARMN* variants, susceptibility, prognosis, genetic variations

Introduction

Glioma is the most common intracranial malignant tumor derived from glial cells, accounting for the majority of all primary brain and central nervous system tumors.¹ It is characterized by the significant mortality and morbidity of approximately 101,600 new cases and 61,000 deaths in China each year.² Malignant glioma is a devastating type of brain and other nervous system tumors because of its high malignancy, extremely high mortality rate, and recurrence risk.³ Despite improvements in therapeutics including surgery in combination with chemo- and/or radiotherapy, the five-year relative survival rate following diagnosis of a malignant brain still grim.⁴ The etiology of glioma remains poorly understood to date, but environmental exposure and genetic factors are identified to increase glioma risk. In recent years, the role of inherited genetic variants in glioma has been highly addressed, which revealed single nucleotide polymorphisms (SNPs) in genes contribute to the susceptibility and prognosis of glioma.⁵⁻⁷

Long non-coding RNAs (lncRNAs) are a class more than 200 nucleotides non-protein coding RNA, that regulate gene or miRNA expression at the transcriptional, post-transcriptional and epigenetic levels.⁸ LncRNAs participate in different stages of glioma formation, invasion, and progression.⁷ Recent evidence indicates that genetic variations in functional lncRNAs may play important roles in the occurrence and development of glioma, such as genetic polymorphisms in lncRNA-PTENP1 and lncRNA H19.^{9,10}

Cardiac mesoderm enhancer-associated non-coding RNA (*CARMN*) is a newly identified lncRNA, also named MIR143HG, and has been reported to be the precursor of miR-143 and miR-145, which linked to gliomagenesis.^{11,12} MicroRNA-145-5p downregulation has been shown to play important roles in the oncogenesis and progression of many cancer types including glioblastoma.¹³ Furthermore, miR-143 inhibited glioma cells migration and invasion through cytoskeletal rearrangement.¹⁴ Ropivacaine suppressed glioma progression by regulating circSCAF11 and miR-145-5p.¹⁵ These suggested *CARMN*, the host gene of miR-143 and miR-145, might have an important role in the occurrence and development of glioma. Nevertheless, no association studies between *CARMN* polymorphisms and glioma have been published to date.

Considering the effect of genetic variants on glioma, we hypothesized that *CARMN* polymorphisms might contribute to glioma development and prognosis. Here, we conducted a case–control study to evaluate the role of *CARMN* polymorphisms in glioma and found that four SNPs were significantly related to glioma risk and patients survival in a Chinese Han population.

Materials and Methods

Study Subjects

In this study, 592 glioma patients and 502 healthy controls enrolled from the department of Neurosurgery at Xi'an Children's Hospital and Tangdu Hospital. All included patients had recently diagnosed and histopathologically confirmed glioma according to the World Health Organization (WHO) classification. All subjects had a Han Chinese ethnic background. All glioma patients were newly diagnosed and confirmed by histopathology. The blood samples were collected before radiotherapy and chemotherapy or surgery. Patients with a self-reported cancer history, serious systemic diseases or other complex diseases were excluded. Age and gender matched healthy controls were enrolled from annual checkup at the same hospitals. The controls had no any cancers or chronic diseases and no brain and central nervous system diseases. Demographic and clinical information was collected from structured questionnaires and/or medical records. All the patients were followed up every 3 months by return visit, telephone and letter. During the follow-up period, the survival time was recorded until death or the last follow-up. This study was approved by the institute ethics committee of the Xi'an Children's Hospital (No. 20200014) and in accordance with the Helsinki Declaration. Written informed consent was obtained from each participant.

Genotyping of *CARMN* Polymorphisms

Peripheral blood samples (5 mL) were collected from all of the study participants. Genomic DNA was extracted using the commercially available GoldMag-Mini Whole Blood Genomic DNA Purification Kit (GoldMag Co. Ltd., Xi'an, China), and stored at -80°C until analysis. The candidate variants in *CARMN* were selected based on a minor allele frequency (MAF) of $> 5\%$ in Chinese populations of the 1000 Genomes Project data (<http://www.internationalgenome.org/>), a pairwise linkage disequilibrium (LD) $r^2 \geq 0.80$, in conformance with Hardy–Weinberg equilibrium (HWE, $p > 0.05$) and the genotyping call rate $> 95\%$. Seven *CARMN* SNPs (rs11168100, rs12654195, rs13177623, rs17796757, rs353299, rs353300 and rs353303) were included for genotyping in the current study.¹⁶ Agena MassARRAY platform (Agena, San Diego, CA, USA) was applied to determine the genotypes of *CARMN* polymorphisms as described previously.^{17,18} The MassARRAY platform is based on MALDI-TOF (matrix-assisted laser desorption/ionization-time of flight) mass spectrometry in a high-throughput and cost-effective manner. Primers for amplification and extension were designed by Agena on-line design software (<https://agenacx.com/online-tools/>), as shown in [Supplementary Table 1](#). The steps for SNPs genotyping were based on manufacturer's protocol, as following: 1) targeted regions for the multiplex assay were amplified by PCR; 2) PCR products were treated through shrimp alkaline phosphatase (SAP) to neutralize unincorporated nucleotides; 3) single base extension reaction were then performed to extend the PCR fragments by one base into the SNP site; 4) the mass of the resultant extended fragments were measured by MALDI-TOF, resulting in a spectrum of distinct mass peaks for the multiplex reaction. The process of genotyping was in double-blinded by two laboratory personnel. For quality control, 10% random sample was repeated genotyping, and the reproducibility was 100%.

Statistical Analyses

SPSS 18.0 (SPSS, Chicago, IL, USA) and PLINK 1.07 package was used for statistical analyses. Differences between cases and controls in demographic characteristics were evaluated by χ^2 test or independent samples *t*-test where appropriate. The frequencies of allele and genotype of *CARMN* polymorphisms in cases and controls were calculated by χ^2 test. HWE was tested for controls with the χ^2 test. The association between *CARMN* genetic variants and glioma risk was estimated by the odds ratios (ORs) and 95% confidence intervals (CIs) after adjusting for age and sex using logistic regression under allele genotype, dominant, recessive and log-additive models, respectively. The pairwise linkage disequilibrium (LD) were measured by the Lewontin's coefficient *D'* using the Haploview v4.2 software, and haplotype association tests for glioma susceptibility were carried out using logistic regression analysis. Univariate and multivariate Cox regression analysis was applied to calculate Hazard ratios (HRs) and 95% CIs for evaluating the association of *CARMN* polymorphisms with glioma prognosis. Survival analysis of glioma patients was assessed by Kaplan–Meier survival curves and the log rank test. A two-sided *p* values of <0.05 were regarded as statistically significant.

Results

Participant Characteristics

The subjects included 592 glioma samples (40.53 ± 13.90, 326 males and 266 females) and 502 cancer-free controls (40.46 ± 18.08, 275 males and 227 females). The frequency distribution of age and sex was matched between cases and controls (*p* = 0.934 and *p* = 0.924, respectively). Other clinical details of patients with glioma such as WHO grade, surgical method, radiotherapy, chemotherapy and survival condition were presented in Table 1.

Table 1 Features of Glioma Patients and Health Controls

Features	Cases (n = 592)	Controls (n = 502)	<i>p</i>
Age (Mean ± SD, years)	40.53 ± 13.90	40.46 ± 18.08	0.934 ^a
≥ 40	329	249	
< 40	263	253	
Gender			0.924 ^b
Male	326	275	
Female	266	227	
WHO grade			
I–II	378		
III–IV	214		
Surgical method			
STR & NTR	185		
GTR	407		
Radiotherapy			
No	58		
Conformal radiotherapy	159		
Gamma knife	375		
Chemotherapy			
No	349		
Yes	243		
Survival condition			
Survival	41		
Lost	24		
Death	527		

Notes: ^a*p* values was calculated by independent samples *t*-test. ^b*p* values was calculated by Chi-square tests.

Abbreviations: WHO, World Health Organization; NTR, near-total resection; STR, sub-total resection; GTR, gross-total resection.

Details of *CARMN* Genetic Polymorphisms

Seven genetic polymorphism in *CARMN* was genotyping and the call rate was > 99.7%. Details of *CARMN* genetic polymorphisms were displayed in [Supplementary Table 2](#). The genotype frequencies of all variants in the controls were in HWE ($p > 0.05$), which suggesting selected samples could represent the whole population. We used HaploRegv4.1 to annotate the potential function of these selected SNPs ([Supplementary Table 2](#)). The results found that six intronic SNPs were associated with the regulation of promoter and/or enhancer histones, DNase, proteins bound, or changed motifs, suggesting they might exert biological functions in this way in patients.

Genetic Effects *CARMN* Variants of on Glioma Susceptibility

The allele and genotype distribution for *CARMN* variants was summarized in [Table 2](#) and [Supplementary Table 3](#). Logistic regression analysis adjusted for age and sex was performed to examine the role of *CARMN* variants in glioma risk. We found that *CARMN* rs13177623 was a protective factor for glioma susceptibility, and GA-AA genotype of rs13177623 had a reduced glioma risk compared with GG genotype (OR = 0.78, 95% CI: 0.61–1.01, $p = 0.043$; [Table 2](#)). There was no statistically significant association between other *CARMN* variants (rs353299, rs353303, rs12654195, rs11168100, rs17796757 and rs353300) and risk for glioma (all p values > 0.05, [Supplementary Table 3](#)) in the overall participants.

We further investigated the correlation of *CARMN* variants with glioma risk by stratifying for age, sex and pathological grade. Stratified analyses by age ([Table 3](#)) displayed that rs13177623 had a lower risk of glioma (OR = 0.67, 95% CI: 0.48–0.94, $p = 0.022$) among the subgroup at age ≥ 40 years. *CARMN* rs11168100 and rs12654195 were associated with decreased the risk of glioma (OR = 0.47, 95% CI: 0.26–0.85, $p = 0.012$ and OR = 0.55, 95% CI: 0.31–0.96, $p = 0.034$, respectively), while rs17796757 increased the risk (OR = 1.50, 95% CI: 1.02–2.19, $p = 0.038$) among the subjects at age < 40 years. In stratified analyses by sex, rs13177623 was significantly associated with decreased risk in males under the allele (OR = 0.77, 95% CI: 0.60–0.99, $p = 0.045$) and dominant (OR = 0.72, 95% CI: 0.52–0.99, $p = 0.043$) models. However, no significant association was observed in females (all $p > 0.05$). These results suggested that *CARMN* rs13177623 polymorphism might be male specific for glioma risk. When stratified by the WHO grade, patients with III-IV glioma had a significantly lower frequency of rs13177623 GA genotype compared with patients with I-II glioma (OR = 0.66, 95% CI: 0.46–0.95, $p = 0.027$, [Supplementary Table 4](#)).

We also examined the impacts of the haplotypes on glioma susceptibility. Linkage disequilibrium (LD) is a nonrandom allele association, and generated by mutation and recombination. LD is measured by the LD coefficient D' : $D' = 1$ is defined as complete linkage disequilibrium; $D' = 0$ is called linkage equilibrium; and $D' < 1$ indicated that gene recombination had occurred. If there is a linkage disequilibrium between SNPs, these SNPs can form a linkage disequilibrium block. As shown in [Figure 1](#), three LD blocks (rs13177623–rs12654195, rs11168100–rs353303 and rs353300–rs353299) were constructed from the seven variants in *CARMN* by coefficient $D' = 0.97$. In addition, we

Table 2 Correlation Between *CARMN* Variants and the Susceptibility to Glioma

SNP ID	Model	Genotype	Control	Case	OR (95% CI)	p
rs13177623	Allele	G	718	889	1	0.059
		A	286	295	0.83 (0.69–1.01)	
	Genotype	GG	256	338	1	0.055
		GA	206	213	0.78 (0.61–1.01)	
		AA	40	41	0.78 (0.49–1.24)	
		GA-AA	246	254	0.78 (0.61–0.99)	
	Dominant	GG	256	338	1	0.043
		GA-AA	246	254	0.78 (0.61–0.99)	
	Recessive	GG-GA	462	551	1	0.512
		AA	40	41	0.86 (0.55–1.35)	
Additive	GG+GA+AA	—	—	—	0.84 (0.69–1.01)	0.062

Notes: p values were calculated by logistic regression analysis with adjustments for age and gender. Bold $p < 0.05$ means the data is statistically significant.

Abbreviations: SNP, single nucleotide polymorphism; OR, odds ratio; CI, confidence interval.

Table 3 Correlation of *CARMN* Variants with Glioma Risk Stratified by Age and Gender

SNP ID	Model	OR (95% CI)	p	OR (95% CI)	p
		≥ 40		< 40	
rs11168100	Allele	1.01 (0.79–1.30)	0.946	0.86 (0.66–1.12)	0.254
	Homozygote	1.32 (0.72–2.44)	0.372	0.54 (0.29–1.02)	0.057
	Heterozygote	0.85 (0.60–1.20)	0.344	1.38 (0.94–2.02)	0.101
	Dominant	0.91 (0.65–1.27)	0.583	1.15 (0.80–1.64)	0.462
	Recessive	1.44 (0.80–2.58)	0.226	0.47 (0.26–0.85)	0.012
	Additive	1.02 (0.79–1.31)	0.901	0.92 (0.70–1.20)	0.517
rs12654195	Allele	1.10 (0.86–1.41)	0.467	0.89 (0.69–1.15)	0.377
	Homozygote	1.52 (0.83–2.80)	0.173	0.66 (0.36–1.19)	0.164
	Heterozygote	0.94 (0.66–1.33)	0.710	1.46 (0.99–2.15)	0.057
	Dominant	1.02 (0.73–1.42)	0.923	1.22 (0.85–1.75)	0.284
	Recessive	1.58 (0.88–2.81)	0.123	0.55 (0.31–0.96)	0.034
	Additive	1.11 (0.86–1.43)	0.432	0.97 (0.74–1.26)	0.799
rs13177623	Allele	0.89 (0.69–1.16)	0.406	0.77 (0.58–1.02)	0.064
	Homozygote	1.59 (0.73–3.45)	0.241	0.56 (0.29–1.08)	0.084
	Heterozygote	0.67 (0.48–0.94)	0.022	1.05 (0.71–1.55)	0.810
	Dominant	0.74 (0.53–1.03)	0.076	0.92 (0.64–1.33)	0.663
	Recessive	1.90 (0.89–4.06)	0.098	0.55 (0.29–1.04)	0.067
	Additive	0.89 (0.68–1.17)	0.419	0.85 (0.65–1.12)	0.252
rs17796757	Allele	0.97 (0.75–1.24)	0.784	1.18 (0.91–1.54)	0.202
	Homozygote	1.24 (0.68–2.27)	0.486	1.09 (0.59–2.02)	0.778
	Heterozygote	0.79 (0.56–1.12)	0.186	1.50 (1.02–2.19)	0.038
	Dominant	0.86 (0.61–1.19)	0.358	1.41 (0.98–2.02)	0.063
	Recessive	1.39 (0.78–2.48)	0.266	0.89 (0.50–1.61)	0.709
	Additive	0.97 (0.76–1.26)	0.842	1.19 (0.90–1.56)	0.218
Gender		Male		Female	
rs13177623	Allele	0.77 (0.60–0.99)	0.045	0.92 (0.69–1.22)	0.554
	Homozygote	0.66 (0.36–1.22)	0.188	0.96 (0.47–1.99)	0.920
	Heterozygote	0.73 (0.52–1.02)	0.067	0.85 (0.59–1.24)	0.405
	Dominant	0.72 (0.52–0.99)	0.043	0.87 (0.61–1.24)	0.440
	Recessive	0.76 (0.42–1.37)	0.360	1.03 (0.51–2.09)	0.940
	Additive	0.78 (0.60–1.00)	0.049	0.92 (0.69–1.22)	0.555

Notes: p values were calculated by logistic regression analysis with adjustments for age and gender. Bold p < 0.05 means the data is statistically significant.

Abbreviations: SNP, single nucleotide polymorphism; OR, odds ratio; CI, confidence interval.

found that G_{rs13177623}G_{rs12654195} haplotype was related to the decreased risk of glioma (OR = 0.61, 95% CI: 0.43–0.86, p = 0.005, Table 4).

Genetic Effects *CARMN* Variants of on Glioma Prognosis

During follow-up, there were 527 patients died of glioma, 41 patients survived and 24 patients lost. We next explored the contribution of *CARMN* variants to the overall survival (OS) and progression free survival (PFS) of glioma patients. The Kaplan–Meier survival curves indicated that the genotype of rs12654195 variant might be associated with OS (Log-rank p = 0.026) and PFS (Log-rank p = 0.027) of glioma patients, as shown in Figure 2. In addition, rs17796757 polymorphism had the effect on OS (Log-rank p = 0.039) of patients with grade III–IV grade III–IV glioma, while rs12654195 variant on OS (Log-rank p = 0.008) and PFS (Log-rank p = 0.011) of patients with grade I–II glioma (Supplementary Figure 1).




Figure 1 The linkage disequilibrium structure of seven SNPs in the *CARMN* gene. Three LD blocks (rs13177623–rs12654195, rs11168100–rs353303 and rs353300–rs353299) were constructed from the seven variants in *CARMN* by coefficient D' 0.97. The numbers in squares are D' values.

The results of univariate Cox proportional hazard model revealed that GG genotype of rs12654195 had a better OS (HR = 0.71, 95% CI: 0.52–0.96, $p = 0.025$) and PFS (HR = 0.69, 95% CI: 0.51–0.95, $p = 0.021$) of glioma patients compared with TT genotype (Table 5). In patients with grade III–IV glioma, rs17796757 was significantly related to the

Table 4 Correlation of *CARMN* Haplotypes with Glioma Susceptibility

Blocks	SNPs	Haplotype	Frequency		Crude Analysis		Adjusted by Age and Gender	
			Case	Control	OR (95% CI)	p	OR (95% CI)	p
Block 1	rs13177623 rs12654195	AG	0.248	0.281	0.84 (0.70–1.02)	0.080	0.84 (0.70–1.02)	0.079
	rs13177623 rs12654195	GG	0.916	0.947	0.61 (0.43–0.86)	0.005	0.61 (0.43–0.86)	0.005
	rs13177623 rs12654195	GT	0.666	0.663	1.02 (0.85–1.22)	0.867	1.02 (0.85–1.22)	0.863
Block 2	rs11168100 rs353303	AG	0.408	0.413	0.98 (0.82–1.16)	0.797	0.98 (0.82–1.16)	0.796
	rs11168100 rs353303	TA	0.313	0.330	0.92 (0.77–1.11)	0.387	0.92 (0.77–1.11)	0.386
	rs11168100 rs353303	AA	0.723	0.744	0.90 (0.74–1.09)	0.272	0.90 (0.74–1.09)	0.270
Block 3	rs353300 rs353299	TT	0.854	0.861	0.95 (0.75–1.21)	0.677	0.95 (0.75–1.21)	0.678
	rs353300 rs353299	TC	0.338	0.351	0.95 (0.79–1.13)	0.538	0.94 (0.79–1.13)	0.534
	rs353300 rs353299	CC	0.515	0.508	1.03 (0.87–1.22)	0.732	1.03 (0.87–1.22)	0.727

Notes: p values were calculated using logistic regression analysis with and without adjustment by gender and age. Bold $p < 0.05$ indicates statistical significance.

Abbreviations: OR, odds ratio; CI, confidence interval.




Figure 2 Kaplan–Meier survival curve for significant association of rs12654195 with OS (A) and PFS (B) of glioma patients.
Abbreviations: OS, overall survival; PFS, progression free survival.

improved OS (AT vs AA, HR = 0.71, 95% CI: 0.52–0.95, $p = 0.024$). In patients with grade I–II glioma, GT genotype and TT genotype of rs12654195 presented an increased OS (HR = 0.75, 95% CI: 0.59–0.94, $p = 0.011$, and HR = 0.66, 95% CI: 0.44–0.99, $p = 0.043$, respectively) and PFS (HR = 0.76, 95% CI: 0.61–0.96, $p = 0.020$, and HR = 0.66, 95% CI: 0.44–0.99, $p = 0.046$, respectively).

Further, the correlation of *CARMN* variants and PFS or OS was evaluated using a multivariate Cox proportional hazard model, adjusted for age, gender, WHO grade, surgical method, radiotherapy and chemotherapy (Table 6). We found rs13177623 GA genotype carriers had an improved OS (HR = 0.83, 95% CI: 0.69–1.00, $p = 0.047$) and PFS (HR = 0.82, 95% CI: 0.68–0.98, $p = 0.031$) for glioma. Rs12654195 (GG vs TT, OS: HR = 0.64, 95% CI: 0.47–0.87, $p = 0.005$ and PFS: HR = 0.65, 95% CI: 0.48–0.89, $p = 0.007$) and rs11168100 (TT vs AA, OS: HR = 0.71, 95% CI: 0.51–0.98, $p = 0.035$) homozygous carriers were also associated with a better prognosis for glioma. For the subgroup of patients with grade III–IV glioma, rs17796757 polymorphism presented an increased OS (AT vs AA, HR = 0.70, 95% CI: 0.51–0.95, $p = 0.025$). For the subgroup of patients with grade I–II glioma, the heterozygous of rs13177623 and rs11168100 were significantly associated with improved OS (HR = 0.78, 95% CI: 0.62–0.98, $p = 0.030$ and HR = 0.73, 95% CI: 0.58–0.92, $p = 0.008$, respectively) and PFS (HR = 0.79, 95% CI: 0.63–0.99, $p = 0.044$ and HR = 0.76, 95% CI: 0.60–0.96, $p = 0.020$, respectively). In addition, improved OS and PFS for grade I–II glioma was also seen for the homozygote (OS: HR = 0.62, 95% CI: 0.42–0.94, $p = 0.024$, and PFS: HR = 0.62, 95% CI: 0.41–0.94, $p = 0.024$) and heterozygous (OS: HR = 0.70, 95% CI: 0.56–0.88, $p = 0.002$, and PFS: HR = 0.72, 95% CI: 0.57–0.91, $p = 0.006$) of rs12654195 variant.

Discussion

The present study explored the possible correlation of seven polymorphisms in *CARMN* with the risk and prognosis of glioma among a Han Chinese population. Our results revealed that rs11168100, rs12654195, rs13177623, and rs17796757 variants were associated with the susceptibility to glioma and the OS and PFS of patients. In addition, we also found that G_{rs13177623}G_{rs12654195} haplotype was a protective factor for glioma susceptibility. To the best of our knowledge, this is the first to assess the role of *CARMN* polymorphisms in glioma risk and prognosis.

CARMN gene, located on chromosome 5q32, is affiliated with the non-coding RNA class.¹⁹ The expression of *CARMN* was significantly dysregulated in various cancers and involved in carcinogenesis. For example, Lin et al reported that *CARMN* inhibited tumor proliferation and metastasis by suppressing MAPK and Wnt signaling pathways in hepatocellular carcinoma.²⁰ *CARMN* suppressed miR-21 through methylation to inhibit cell invasion and migration.²¹ *CARMN* have reported expressing stably homologous miRNAs: miR-143 and miR-145.²² Previous studies have demonstrated miR-143/145 regulate the proliferation, migration and invasion of glioma cells and could be potential

Table 5 Univariate Analysis of the Association Between *CARMN* Variants and OS and PFS of Glioma Patients

SNP ID	Genotype	OS					PFS				
		Total	Events	SR (1-/3-Year)	HR (95% CI)	<i>p</i>	Total	Events	SR (1-/3-Year)	HR (95% CI)	<i>p</i>
rs12654195	TT	260	237	0.267/0.073	1	0.097	259	236	0.118/0.077	1	0.184
	GT	271	240	0.369/0.097	0.86 (0.72–1.03)		269	239	0.212/0.091	0.89 (0.74–1.06)	
	GG	61	50	0.344/0.132	0.71 (0.52–0.96)		59	48	0.305/0.153	0.69 (0.51–0.95)	
III–IV grade											
rs17796757	AA	98	93	0.245/0.041	1	0.024	97	92	0.124/0.049	1	0.161
	AT	90	80	0.356/0.097	0.71 (0.52–0.95)		88	79	0.170/0.097	0.81 (0.60–1.09)	
	TT	26	24	0.385/0.051	0.75 (0.48–1.17)		26	24	0.247/0.041	0.82 (0.52–1.28)	
I–II grade											
rs12654195	TT	158	144	0.224/0.072	1	0.011	157	143	0.104/0.076	1	0.020
	GT	184	158	0.418/0.123	0.75 (0.59–0.94)		184	158	0.244/0.114	0.76 (0.61–0.96)	
	GG	36	28	0.333/0.187	0.66 (0.44–0.99)		35	27	0.314/-	0.66 (0.44–0.99)	

Notes: Log-rank *p* values were calculated using the Chi-Square test. Bold *p* < 0.05 indicates statistical significance.

Abbreviations: OS, overall survival; PFS, progression free survival; SR, survival rate; HR, hazard ratio; CI, confidence interval.

Table 6 Multivariate Analysis of the Association Between *CARMN* Variants and OS and PFS of Glioma Patients

SNP ID	Genotype	OS		PFS	
		HR (95% CI)	p	HR (95% CI)	p
rs13177623	GG				
	GA	0.83 (0.69–1.00)	0.047	0.82 (0.68–0.98)	0.031
	AA	0.72 (0.51–1.03)	0.070	0.75 (0.53–1.05)	0.096
rs12654195	TT				
	GT	0.87 (0.72–1.04)	0.129	0.84 (0.70–1.01)	0.059
	GG	0.64 (0.47–0.87)	0.005	0.65 (0.48–0.89)	0.007
rs11168100	AA				
	AT	0.88 (0.73–1.06)	0.167	0.84 (0.70–1.01)	0.067
	TT	0.71 (0.51–0.98)	0.035	0.74 (0.54–1.01)	0.060
III–IV grade					
rs17796757	AA				
	AT	0.70 (0.51–0.95)	0.025	0.75 (0.55–1.03)	0.079
	TT	0.70 (0.44–1.10)	0.123	0.75 (0.47–1.18)	0.213
I–II grade					
rs13177623	GG				
	GA	0.78 (0.62–0.98)	0.030	0.79 (0.63–0.99)	0.044
	AA	0.88 (0.56–1.38)	0.579	0.85 (0.54–1.33)	0.469
rs12654195	TT				
	GT	0.70 (0.56–0.88)	0.002	0.72 (0.57–0.91)	0.006
	GG	0.62 (0.42–0.94)	0.024	0.62 (0.41–0.94)	0.024
rs11168100	AA				
	AT	0.73 (0.58–0.92)	0.008	0.76 (0.60–0.96)	0.020
	TT	0.82 (0.53–1.27)	0.375	0.77 (0.50–1.21)	0.264

Notes: p values were calculated by Cox multivariate analysis with adjustments for gender, age, WHO grade, surgical method, use of radiotherapy and chemotherapy. Bold p < 0.05 indicates statistical significance.

Abbreviations: OS, overall survival; PFS, progression free survival; HR, hazard ratio; CI, confidence interval.

therapeutic target for anti-invasion therapies of glioma patients.^{11,23} Recently, LncRNA *CARMN* inhibited the proliferation of glioblastoma cells by sponging miR-504.²⁴ These suggested that *CARMN* could be of pathogenic importance in glioma.

Our study was the first to evaluate the correlation of *CARMN* variants with susceptibility and prognosis of glioma. We found *CARMN* rs13177623 was related to the decreased risk of glioma. Previous studies have indicated that the incidence rates of glioma tended to be associated with age and gender.²⁵ Age stratified analysis showed rs13177623 had a lower risk of glioma at age ≥ 40 years, while rs11168100, rs12654195 and rs17796757 were associated with the susceptibility to glioma at age < 40 years. These indicate that the contribution of *CARMN* polymorphisms to glioma risk was associated with age exposures. In stratified analyses by gender, rs13177623 was significantly associated with decreased risk in males, but not in females, which suggesting the effect of rs13177623 polymorphism on glioma risk presented sex difference. Moreover, our study also evaluated the effect of *CARMN* polymorphisms on the prognosis of glioma patients. We found that rs13177623, rs12654195 and rs11168100 were associated with a better prognosis for glioma, especially in grade I–II glioma. In patients with grade III–IV glioma, rs17796757 polymorphism presented an improved OS. Previous studies supported that SNPs differentially might influence the expression and function of lncRNAs.^{26–28} Therefore, *CARMN* variants might contribute to the risk and prognosis of glioma by affecting the function of *CARMN*. However, further functional study is necessary to explore the role of these polymorphisms in the etiology of glioma.

Inevitably, our study had several limitations. Firstly, the inherent selection bias cannot be exclude because this study based on a hospital-based case–control study. Therefore, we recruited subjects matched by age, gender, and residential

area to reduce the bias. Secondly, we did not assess the potential function of these polymorphisms in *CARMN*. Further functional experiments should be required to investigate the role of *CARMN* variants in glioma occurrence and development. Thirdly, some environmental factors such as occupational exposure and dietary were not available; the interaction of these factors with *CARMN* genotypes should be performed in a larger survey.

Conclusion

In summary, we firstly reported the contribution of *CARMN* variants (rs11168100, rs12654195, rs13177623, and rs17796757) to the susceptibility and prognosis of glioma in a Chinese Han population. Our study provides a novel insight on the relationship between *CARMN* gene and glioma tumorigenesis. These findings add to the growing body of evidence linking lncRNAs polymorphisms to glioma etiology. In addition, further studies are required to validate our results.

Data Sharing Statement

All the data regarding the findings are available within the manuscript. Anyone who is interested in the information should contact the corresponding author.

Acknowledgments

The authors thank all patients for providing blood samples and all the research staff for their contributions to this project.

Disclosure

The authors declared no conflicts of interest in this work.

References

1. Goodenberger ML, Jenkins RB. Genetics of adult glioma. *Cancer Genet.* 2012;205(12):613–621. doi:10.1016/j.cancergen.2012.10.009
2. Chen W, Zheng R, Baade PD, et al. Cancer statistics in China, 2015. *CA Cancer J Clin.* 2016;66(2):115–132. doi:10.3322/caac.21338
3. Lin D, Wang M, Chen Y, et al. Trends in intracranial glioma incidence and mortality in the United States, 1975–2018. *Front Oncol.* 2021;11:748061. doi:10.3389/fonc.2021.748061
4. Ostrom QT, Cioffi G, Gittleman H, et al. CBTRUS statistical report: primary brain and other central nervous system tumors diagnosed in the United States in 2012–2016. *Neuro Oncology.* 2019;21(Suppl 5):v1–v100. doi:10.1093/neuonc/noz150
5. Guo H, Liao Y, Lin A, et al. Association between LIN28A gene polymorphisms and glioma susceptibility in Chinese children. *J Moffitt Cancer Center.* 2021;28:10732748211040009.
6. Guan Q, Yuan L, Lin A, et al. KRAS gene polymorphisms are associated with the risk of glioma: a two-center case-control study. *Transl Pediatr.* 2021;10(3):579–586. doi:10.21037/tp-20-359
7. He J, Yuan L, Lin H, et al. Genetic variants in m (6) A modification core genes are associated with glioma risk in Chinese children. *Mol Ther Oncolytics.* 2021;20:199–208. doi:10.1016/j.omto.2020.12.013
8. Wang KC, Chang HY. Molecular mechanisms of long noncoding RNAs. *Mol Cell.* 2011;43(6):904–914. doi:10.1016/j.molcel.2011.08.018
9. Yang S, Fu ZZ, Zhang YQ, Fu BH, Dong LX. Rs7853346 polymorphism in lncRNA-PTENP1 and rs1799864 polymorphism in CCR2 are associated with radiotherapy-induced cognitive impairment in subjects with glioma via regulating PTENP1/miR-19b/CCR2 signaling pathway. *Biochem Genet.* 2021. doi:10.1007/s10528-021-10145-9
10. Deng Y, Zhou L, Yao J, et al. Associations of lncRNA H19 polymorphisms at MicroRNA binding sites with glioma susceptibility and prognosis. *Mol Ther Nucleic Acids.* 2020;20:86–96. doi:10.1016/j.omtn.2020.02.003
11. Zhao H, Du P, Peng R, et al. Long noncoding RNA OR7E156P/miR-143/HIF1A axis modulates the malignant behaviors of glioma cell and tumor growth in mice. *Front Oncol.* 2021;11:690213. doi:10.3389/fonc.2021.690213
12. Wang Y, Zhang C, Lu W, Chen R, Yu M. Establishment of a prognostic-related microRNAs risk model for glioma by bioinformatics analysis. *Ann Trans Med.* 2021;9(12):1022. doi:10.21037/atm-21-2402
13. Zhang Y, Ta WW, Sun PF, Meng YF, Zhao CZ. Diagnostic and prognostic significance of serum miR-145-5p expression in glioblastoma. *Int J Clin Exp Pathol.* 2019;12(7):2536–2543.
14. Chen WY, Lang ZQ, Ren C, Yang P, Zhang B. miR-143 acts as a novel Big mitogen-activated protein kinase 1 suppressor and may inhibit invasion of glioma. *Oncol Rep.* 2019;42(3):1194–1204. doi:10.3892/or.2019.7218
15. Yin D, Liu L, Shi Z, Zhang L, Yang Y. Ropivacaine inhibits cell proliferation, migration and invasion, whereas induces oxidative stress and cell apoptosis by circSCAF11/miR-145-5p axis in glioma. *Cancer Manag Res.* 2020;12:11145–11155. doi:10.2147/CMAR.S274975
16. Guo Y, Cao Y, Gong S, et al. Correlation analysis between CARMEN variants and alcohol-induced osteonecrosis of the femoral head in the Chinese population. *BMC Musculoskelet Disord.* 2020;21(1):547. doi:10.1186/s12891-020-03553-2
17. Gabriel S, Ziaugra L, Tabbaa D. SNP genotyping using the Sequenom MassARRAY iPLEX platform. *Curr Protoc Hum Genet.* 2009;60(1):2–12.
18. Dai ZJ, Liu XH, Ma YF, et al. Association between single nucleotide polymorphisms 7in DNA polymerase kappa gene and breast cancer risk in Chinese Han population: a STROBE-compliant observational study. *Medicine.* 2016;95(2):e2466. doi:10.1097/MD.0000000000002466
19. Ounzain S, Micheletti R, Arnan C, et al. CARMEN, a human super enhancer-associated long noncoding RNA controlling cardiac specification, differentiation and homeostasis. *J Mol Cell Cardiol.* 2015;89:98–112. doi:10.1016/j.yjmcc.2015.09.016

20. de Vries PS, Brown MR, Bentley AR, et al. Multiancestry genome-wide association study of lipid levels incorporating gene-alcohol interactions. *Am J Epidemiol.* 2019;188(6):1033–1054. doi:10.1093/aje/kwz005
21. Xun W, Cen W, Dahai Y. LncRNA miR143HG suppresses miR-21 through methylation to inhibit cell invasion and migration. *Laryngoscope.* 2020;130(11):E640–E645. doi:10.1002/lary.28474
22. Zhao Q, Sun X, Liu C, Li T, Cui J, Qin C. Expression of the microRNA-143/145 cluster is decreased in hepatitis B virus-associated hepatocellular carcinoma and may serve as a biomarker for tumorigenesis in patients with chronic hepatitis B. *Oncol Lett.* 2018;15(5):6115–6122. doi:10.3892/ol.2018.8117
23. Guo Y, He X, Zhang M, et al. Reciprocal control of ADAM17/EGFR/Akt signaling and miR-145 drives GBM invasiveness. *J Neurooncol.* 2020;147(2):327–337. doi:10.1007/s11060-020-03453-4
24. Wang P, Bao W, Liu X, Xi W. LncRNA miR143HG inhibits the proliferation of glioblastoma cells by sponging miR-504. *Int J Neurosci.* 2021;1–9. doi:10.1080/00207454.2020.1865950
25. Le Rhun E, Weller M. Sex-specific aspects of epidemiology, molecular genetics and outcome: primary brain tumours. *ESMO open.* 2020;5(Suppl 4):e001034. doi:10.1136/esmoopen-2020-001034
26. Fu Y, Zhang Y, Cui J, et al. SNP rs12982687 affects binding capacity of lncRNA UCA1 with miR-873-5p: involvement in smoking-triggered colorectal cancer progression. *CCS.* 2020;18(1):37. doi:10.1186/s12964-020-0518-0
27. Feng T, Feng N, Zhu T, et al. A SNP-mediated lncRNA (LOC146880) and microRNA (miR-539-5p) interaction and its potential impact on the NSCLC risk. *J Exp Clin Cancer Res.* 2020;39(1):157. doi:10.1186/s13046-020-01652-5
28. Wang L, Xing Q, Feng T, He M, Yu W, Chen H. SNP rs710886 A>G in long noncoding RNA PCAT1 is associated with the risk of endometriosis by modulating expression of multiple stemness-related genes via microRNA-145 signaling pathway. *J Cell Biochem.* 2020;121(2):1703–1715. doi:10.1002/jcb.29406

Pharmacogenomics and Personalized Medicine

Dovepress

Publish your work in this journal

Pharmacogenomics and Personalized Medicine is an international, peer-reviewed, open access journal characterizing the influence of genotype on pharmacology leading to the development of personalized treatment programs and individualized drug selection for improved safety, efficacy and sustainability. This journal is indexed on the American Chemical Society's Chemical Abstracts Service (CAS). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/pharmacogenomics-and-personalized-medicine-journal>