

Plant-Derived Vesicles: A New Era for Anti-Cancer Drug Delivery and Cancer Treatment

Yuying Zhao¹, Hanxu Tan², Juping Zhang^{1,3}, Bo Pan¹, Neng Wang², Tongkai Chen⁴, Yafei Shi^{1,2}, Zhiyu Wang^{1,3,5}

¹State Key Laboratory of Dampness Syndrome of Chinese Medicine, The Second Affiliated Hospital of Guangzhou University of Chinese Medicine, Guangzhou, People's Republic of China; ²The Research Center for Integrative Medicine, School of Basic Medical Sciences, Guangzhou University of Chinese Medicine, Guangzhou, Guangdong, People's Republic of China; ³Guangdong-Hong Kong-Macau Joint Laboratory on Chinese Medicine and Immune Disease Research, Guangzhou University of Chinese Medicine, Guangzhou, Guangdong, People's Republic of China; ⁴Science and Technology Innovation Center, Guangzhou University of Chinese Medicine, Guangzhou, Guangdong, People's Republic of China; ⁵Guangdong Provincial Key Laboratory of Clinical Research on Traditional Chinese Medicine Syndrome, Guangdong Provincial Hospital of Chinese Medicine, Guangdong Provincial Academy of Chinese Medical Sciences, Guangzhou, Guangdong, People's Republic of China

Correspondence: Zhiyu Wang; Yafei Shi, Email wangzhiyu@gzucm.edu.cn; shiyafei@gzucm.edu.cn

Abstract: Lipid-structured vesicles have been applied for drug delivery system for over 50 years. Based on their origin, lipid-structured vesicles are divided into two main categories, namely synthetic lipid vesicles (SLNVEs) and vesicles of mammalian origin (MDVEs). Although SLNVEs can stably transport anti-cancer drugs, their biocompatibility is poor and degradation of exogenous substances is a potential risk. Unlike SLNVEs, MDVEs have excellent biocompatibility but are limited by a lack of stability and a risk of contamination by dangerous pathogens from donor cells. Since the first discovery of plant-derived vesicles (PDVEs) in carrot cell supernatants in 1967, emerging evidence has shown that PDVEs integrate the advantages of both SLNVEs and MDVEs. Notably, 55 years of dedicated research has indicated that PDVEs are an ideal candidate vesicle for drug preparation, transport, and disease treatment. The current review systematically focuses on the role of PDVEs in cancer therapy and in particular compares the properties of PDVEs with those of conventional lipid vesicles, summarizes the preparation methods and quality control of PDVEs, and discusses the application of PDVEs in delivering anti-cancer drugs and their underlying molecular mechanisms for cancer therapy. Finally, the challenges and future perspectives of PDVEs for the development of novel therapeutic strategies against cancer are discussed.

Keywords: plant-derived vesicles, cancer therapy, drug delivery, liposome nanoparticles, mammalian-derived exosomes


Introduction

Mammalian-derived vesicles (MDVEs) are small vesicles present in mammalian body fluids that are used to transmit biological signals between cells. However, when these vesicles were first discovered, it was thought that they were cellular debris or cellular litter.¹ With the development of new technologies, it was discovered that the vesicles not only existed in mammalian body fluids but were also found in multiple plants, which were termed as plant-derived vesicles (PDVEs).² A number of studies showed that plants were able to produce vesicles in response to a variety of biotic and abiotic environmental stresses, including pathogen infection and attack.^{3,4}

PDVEs are structurally and functionally similar to MDVEs and carry large amounts of proteins, lipids, nucleic acids, and metabolites⁵ that are necessary for intercellular signal communication.⁶ Notably, the signal transfer is cross-species.⁷ The nucleic acids and proteins in PDVEs can alter the physiological activities of animal cells.⁸ Unlike MDVEs, PDVEs are easier and faster to extract, are cost effective, and do not involve ethical issues.⁹ Compared to synthetic lipid vesicles (SLNVEs), PDVEs are much safer. Emerging evidence suggests that PDVEs have good prospects in the development of nano-delivery systems (Figure 1).

The tumor microenvironment (TME) is completely different from the normal physiological environment and is a prerequisite to support cancer cell proliferation and metastasis.¹⁰ However, this unique TME also poses a challenge for cancer treatment. For example, the low pH of the TME can degrade chemotherapeutic drugs before their uptake.¹¹ The

Graphical Abstract



	Type of vesicles	Source	Stability in Stomach acid	Cross different barriers	Active substance	Toxicity
Mammalian-derived vesicles		Blood Cancer	Unstable	Intestinal Barrier Blood Brain Barrier Placental Barrier	DNA RNA Enzymes Proteins Lipids	Potential Toxicity
Plant-derived vesicles		Plant	Stable	Intestinal Barrier Blood Brain Barrier Placental Barrier	Enzymes Proteins Lipids	Safe
liposome nanoparticles		Industrial Synthesis	Stable	Intestinal Barrier Blood Brain Barrier Placental Barrier	Phospholipids	Potential Toxicity

Figure 1 Priorities of plant-derived vesicles compared to mammalian-derived exosomes and liposome nanoparticles.

inflammatory microenvironment significantly promotes cancer proliferation, metastasis, and drug resistance.¹² The relatively hypoxic environment facilitates neoangiogenesis and provides a constant nutrient supply for cancer cell growth.¹³ In light of these characteristics, nanodrug delivery systems were developed to design intelligent nanocarriers to increase drug targeting and improve therapeutic effects.¹⁴ Currently, MDVEs and SLNVEs have been widely used in cancer therapy. However, a number of challenges still remain, such as trans-biological barrier capabilities, acid tolerance, cancer cell-specific targeting, safety, toxicity, potential environmental hazards, and large-scale manufacturing costs. The development of PDVEs has potential to overcome these problems. PDVEs can remain stable in gastric acid mimics¹⁵ and penetrate the intestinal barrier within 6 h.¹⁶ Natural PDVEs have active cancer targeting properties and can be improved with acquired modifications.^{17,18} Furthermore, there are currently no reported instances of PDVEs causing explicit harm to organisms. Thus, it can be tentatively concluded that they are viable carriers for drug delivery, environmentally friendly, and cost-effective.^{19,20}

Emerging studies have validated the unique therapeutic effects of PDVEs in multiple malignancies. For example, *Petasites japonicus*-derived vesicles were found to act as adjuvants to activate the immune response, and Han et al demonstrated that *Petasites japonicus*-derived vesicles treated dendritic cells strongly induced the proliferation and differentiation of naive T cells to Th1-type T cells and cytotoxic CD8+ T cells, leading to increased secretion of

interferon- γ and interleukin-2.²¹ Bitter melon-derived vesicles were also reported to inhibit oral squamous cell carcinoma proliferation, as well as overcome drug resistance.²² Interestingly, PDVEs do not seem to exert significant anti-cancer effects on normal cells.²³ PDVEs warrant continued investigation in anti-cancer drug research and development in the future. Here, in a systematic review, we present the properties, preparations, and applications of PDVEs for cancer treatment.

Properties of PDVEs Compared to Traditional Vesicles Comparison with Liposome Nanoparticles

SNLVEs consist of a wide range of particulate systems, including dendrimers, micelles, liposomes, and polymeric nanoparticles. Liposomes are one representative type of SNLVEs, and they have been applied in cancer therapy for a long time due to their good biocompatibility²⁴ owing to their similar composition to cell membranes.²⁵ Previous research has confirmed the applications of liposomes in various fields, such as encapsulating DNA, RNA, water-soluble drugs, and hydrophobic drugs.^{26,27} However, liposomes are still limited due to their synthetic nature. It was shown that liposomes induced cellular stress, apoptosis, activation of inflammatory vesicles,²⁸ cytokine storms,²⁹ and immune responses.³⁰ Notably, liposomes were reported to cause severe DNA and RNA damage by upregulating the expression of Heat shock protein 70-2 α and Heat shock protein 90.³¹ In clinical practice, a drug for acute myelogenous leukemia named CPX-351, which is primarily composed of a dual-drug liposomal encapsulation of cytarabine and daunorubicin, caused nonhematologic adverse events to varying degrees.³²

Compared to liposomes, PDVEs have demonstrated significant advantages in several key areas. Firstly, the inherent nature of PDVEs ensures that they are safer for the body than SNLVEs,³³ and more environmentally friendly.³⁴ Additionally, the RNA and proteins carried within PDVEs facilitate communication with the body. Unlike SNLVEs, which merely serve as carriers, PDVEs inherently possess therapeutic effects.³⁵ Furthermore, PDVEs showed an advantage in cellular uptake. A recent study found that the uptake efficiency of PDVEs reached over 80%, while only 40% of liposomes were internalized.³⁶ In summary, in contrast to traditional liposomes, PDVEs have showcased remarkable advantages in various pivotal domains, particularly in the realm of cancer treatment, offering new possibilities for future disease therapies.

Comparison with Mammalian-Derived Vesicles

MDVEs are shed from the cell membrane and carry with them the biological information and metabolic waste of the donor cells.^{29,30} According to vesicle size, MDVEs are classified into apoptotic vesicles (100–5000 nm in diameter), microvesicles (100–1000 nm in diameter), and the well-known exosomes (30–150 nm in diameter). It was found that MDVEs are widely distributed in mammalian tissues and body fluids, therefore providing a transport mechanism supporting biological signal communication between tissues and organs. MDVEs present remarkable biocompatibility and are widely applied in cancer treatment. For example, plasma-derived exosomes protected miRNAs from nuclease degradation and successfully promoted apoptosis of HepG2 hepatocellular carcinoma cells.³⁷ Additionally, exosome-encapsulated doxorubicin was reported to inhibit cancer growth with reduced cardiotoxicity.³⁸ Notably, cancer-derived exosomes are considered as adjuvants in immunotherapy due to the existence of cancer antigen information.³⁹ In addition to exosomes, apoptotic bodies derived from live cells represent a novel drug delivery system. The apoptotic body vehicle can deliver the ammunition to tumor and achieve deep penetration by macro-phage-hitchhiking.⁴⁰ Adipocyte-derived lipid droplets are also a type of mammalian-derived vesicle, and are increasingly paid great attention for anti-cancer therapies.⁴¹ Furthermore, beyond the direct use of vesicles derived from cells, emerging studies have leveraged cell membranes to construct delivery systems *ex vivo*. For instance, since platelets target postsurgical wounds and circulating tumor cells, platelet-derived membrane vesicles are candidates for drug delivery in post-surgery cancer therapy. Nano-vesicles using platelet membranes have been employed to encapsulate oxaliplatin as a synergistic treatment for post-surgery tumor recurrence and metastasis.⁴² However, MDVEs secreted by pathological cells have the potential to induce the spread of diseases.⁴³ Furthermore, there are moral and ethical arguments concerning the commercial utilization of

MDVEs, and the manufacturing cost of MDVEs is high due to the limited donor availability and low yield. Therefore, MDVEs have not yet met the requirements for the development of clinical drugs.

PDVEs share many similarities with MDVEs, such as a size between 30 and 500 nm, a predominantly circular or cup shape, a negatively charged surface, and an ability for uptake and crossing biological barriers facilitated by phospholipid bilayers. However, unlike MDVEs, PDVEs cannot penetrate the placental barrier, suggesting a better safety profile.^{36,44} In addition, PDVEs have abundant donor resources, allowing for large-scale production at low manufacturing costs. Ginger root has been reported to contain $0.5\text{--}2 \times 10^{14}$ vesicles/kg, which is 10 times higher than that of MDVEs.⁹ In addition, PDVEs are non-toxic and are a part of most daily diets as additives. More importantly, as a carrier, PDVEs can protect a drug from gastric acid degradation, while MDVEs are not tolerant of low pH, except for milk-derived MDVEs.³⁶

Preparation of PDVEs

The properties of PDVEs make them promising for cancer treatment. However, it is difficult to produce high purity PDVEs on a large scale,⁴⁵ given that vesicles tend to adhere to each other through macromolecules, such as starch, cellulose, and tannins.⁴⁶ Therefore, the extracted PDVEs should be highly dispersed and reproducible for subsequent studies and clinical applications. Generally, the extraction methods of PDVEs are divided into two broad categories, namely physically based techniques and chemically based techniques (Table 1).

Physically Based Techniques

The physically based methods include differential centrifugation, density gradient centrifugation, and ultrafiltration membrane separation. Differential centrifugation is considered as the standard for PDVE extraction and is the most widely used. It is based on the principle that large particles will settle at the bottom layer, while small particles will float at a higher layer under centrifugal force, and the target vesicles can be isolated by adjusting the centrifugal force.^{47,48} The method is easy to implement and cost-effective and is suitable for the mass production of PDVEs. However, vesicles may be disrupted by the ultra-intense centrifugal forces. A high-density hypotonic solution added to the bottom of the

Table 1 The Advantages and Disadvantages of Various Preparation Methods

Physically Based Techniques		
Methods	Advantage	Disadvantage
Differential Centrifugation	<ol style="list-style-type: none"> 1. The standard for PDVE extraction. 2. Easy to implement. 3. Cost-effective. 4. Suitable for the mass production. 	<ol style="list-style-type: none"> 1. Affecting vesicle integrity. 2. The separation conditions for each type of vesicle require special customization, making standardization difficult.
Density Gradient Centrifugation	<ol style="list-style-type: none"> 1. Simple. 2. High yielding. 3. Less time-consuming. 	<ol style="list-style-type: none"> 1. Poorly homogeneous. 2. Easily clumped.
Ultrafiltration Membrane Separation	<ol style="list-style-type: none"> 1. Simple. 2. High purity. 	<ol style="list-style-type: none"> 1. Affecting the separation performance of the membrane. 2. Highly experience-dependent. 3. Membrane fouling.
Chemically based techniques		
Co-precipitation	<ol style="list-style-type: none"> 1. Use of a kit, eliminating the need for large-scale equipment, ensuring high feasibility. 2. High adaptability. 3. High repeatability. 	<ol style="list-style-type: none"> 1. High cost. 2. Low purity.
Immunoaffinity Separation	<ol style="list-style-type: none"> 1. Time-saving. 2. High purity. 	<ol style="list-style-type: none"> 1. High cost. 2. Not suitable for mass production.

centrifuge tube can protect vesicles from being shattered. Additionally, experimental conditions for differential centrifugation are not standardized and are not interchangeable for each PDVE.^{44,49} Thus, a personalized centrifugation protocol for each type of PDVE is time-consuming.

To save on extraction time, density gradient centrifugation was developed. Specifically, a series of a continuous or discontinuous density medium are placed in a centrifuge tube according to the gradient, and vesicles with different densities are stratified by centrifugal force.^{50,51} The density gradient centrifugation method is simple, high yielding, and less time-consuming than many other methods. However, the vesicles obtained by this method are poorly homogeneous and easily clumped, which is not satisfactory for subsequent studies. Thus, differential centrifugation and density gradient centrifugation methods are usually applied together to improve the quality of the vesicles. Generally, a certain amount of vesicles is first extracted by differential centrifugation, and purer vesicles are further obtained by density gradient centrifugation according to the size and density of vesicles.⁵²

In recent years, ultrafiltration membrane separation, a method separating vesicles by size, has also been used to separate PDVEs. A solution is filtered sequentially through porous membranes of different sizes under a certain pressure, and the large particles are physically intercepted by the porous membrane, while small particles are unimpeded.⁵³ The purity of the vesicles extracted by this method is satisfactory. However, the concentration polarization phenomenon appearing during filtration changes the separation performance of the membrane. Also, membrane fouling usually occurs. The phenomenon of concentration polarization and membrane fouling can be effectively improved by adjusting the flow rate, pressure, and concentration of filtration, a process that is highly experience-dependent. In addition, clearance of soluble protein impurities was shown to avoid membrane fouling prior to ultrafiltration separation.⁵⁴

Chemically Based Techniques

Chemical-based techniques include co-precipitation and immunoaffinity separation methods. Co-precipitation separation is performed by adding a precipitant to a mixture to precipitate the target vesicle and then separating the precipitate to obtain the target vesicle. During the separation of vesicles, surface adsorption of the precipitate utilizing positive and negative charge attraction is the most common separation method.⁵⁵ Most kits available on the market for vesicle isolation are based on the principle of co-precipitation isolation. However, the method is costly and the vesicles that are obtained are generally impure and need further purification. Immunoaffinity separation is a method based on the specific binding of antibodies to antigens on the vesicle membrane. This method is fast, specific, and pure, but only small amounts can be extracted, and it is a high cost method.⁵⁶

Taken together, physically based techniques are suitable for mass processing and for obtaining pure vesicles, but the separation parameters are complex and time-consuming. Chemically based techniques are simple and reproducible, but the cost is high. Currently, much attention has been paid to combining both physical and chemical isolation techniques to obtain high purity and reproducible PDVEs.⁵⁷

Applications of PDVEs in Cancer Therapy

A number of anti-cancer drugs have exhibited strong cytotoxic killing effects *in vitro*, but the *in vivo* inhibitory effects were unsatisfactory. Furthermore, the significant systemic toxicity due to the poor bioavailability also limited their clinical application. The emergence of nano-drug delivery systems tends to overcome this problem. The nanocarriers can chemically or physically conjugate the drug and deliver it to the designated lesion site to improve therapeutic efficacy.⁵⁸ The efficiency of drug targeted delivery to the lesion is determined by the properties of the carriers, including physical properties, surface modifications, cellular uptake, active targeting ability, acid tolerance, and the trans-biological barrier characteristics. PDVEs were first utilized for drug delivery in 2014. The GfDVEs encapsulated methotrexate and were used for ameliorating dextran sulfate sodium induced mouse colitis.⁴⁴ In recent years, PDVEs with anti-inflammatory,⁵⁹ anti-cancer,²³ anti-bacterial,⁶⁰ and anti-oxidant aging effects have been found.^{61–63} They are widely used to prevent or treat cancer,⁶⁴ colitis,⁶⁵ alcoholic liver disease,⁶⁶ and even COVID-19.⁶⁷ PDVEs are promising as drug delivery systems for cancer treatment because of their intrinsic targeting,⁶⁸ acid tolerance,⁶⁹ easy penetration of physiological barriers,¹⁸ storage stability,⁷⁰ non-toxicity to the organism,³⁶ and low fabrication cost (Figure 2).




Figure 2 Applications of plant-derived vesicles in cancer therapy as a carrier.

As a Carrier

Properties

PDVEs are spherical, elliptical, or cup-shaped in shape, depending on the plant species and the extraction method (Table 2). PDVEs have a negative surface charge, which can effectively prolong their residence time in the blood.⁷¹ The size of PDVEs from different species is inconsistent and mainly distributed between 30 and 500 nm. For example, the size of Beta vulgaris-derived vesicles is 50 nm,⁷² that of asparagus-derived vesicles is 119 nm,⁷³ that of ginger-derived vesicles (GiDVEs) is 220–290 nm,⁶⁶ and that of ginseng-derived vesicles is 344.8 nm.⁷⁴ In addition, the isolation method also influences the size. For example, the size of GiDVEs extracted by differential centrifugation was 403 nm, while that of vesicles extracted by co-precipitation separation with a 15% concentration of polyethylene glycol-6000 (PEG6000)

Table 2 Physicochemical Characterization of Plant-Derived EVs

Source	Scientific Name	Isolation Method	Size (nm)	Zeta (mV)	Shape	Ref.
Apple	<i>Malus domestica</i>	Differential centrifugation/filtration/ ultracentrifugation	152± 32.3	/	Round	[76]
Arabidopsis	<i>Arabidopsis thaliana</i>	Vacuum infiltration/centrifugation/differential centrifugation	52± 170	/	Spherical	[54]
		Differential centrifugation	100–150	Negative	Cupped	[77]
Blood orange	<i>Citrus sinensis</i>	Differential centrifugation/filtration/ ultracentrifugation	178.8± 4.3	/	/	[78]
Blueberry	<i>Vaccinium myrtillus</i> , <i>Vaccinium corymbosum</i>	Differential centrifugation/filtration/ ultracentrifugation	198± 112	/	Spherical	[79]
		Differential centrifugation/filtration/ ultracentrifugation	/	/	Spherical or oval	[80]
Broccoli	<i>Brassica oleracea</i>	Density gradient centrifugation	18–120	–17	Spherical	[65]
Cabbage	<i>Brassica oleracea</i> var. <i>capitata</i>	Polyethylene glycol-based precipitation/ ultracentrifugation	100	–14.8	Spherical	[53]
Carrot	<i>Daucus carota</i>	Differential centrifugation/filtration	150	–10.2	Spherical	[81]
		Differential centrifugation/density gradient centrifugation	200	–25	Spherical or cupped	[7]
Clementine	<i>Citrus clementina</i>	Density gradient centrifugation	75–345	/	/	[46]
Coconut	<i>Cocos nucifera</i>	Differential centrifugation/filtration/ ultracentrifugation	/	/	Spherical or oval	[80]
		Differential centrifugation/filtration/ ultracentrifugation	Coconut water: 13– 60 Coconut milk: 30– 100	/	Spherical	[82]
Cucumber	<i>Cucumis sativus</i>	Differential centrifugation/filtration/ ultracentrifugation	167± 3	–32± 0.13	Spherical	[83]
Garlic	<i>Allium sativum</i>	Differential centrifugation/filtration/ ultracentrifugation	113–153	/	Spherical	[84]

(Continued)

Table 2 (Continued).

Source	Scientific Name	Isolation Method	Size (nm)	Zeta (mV)	Shape	Ref.
Ginger	<i>Zingiber officinale</i>	Differential centrifugation/filtration/ ultracentrifugation	124.5	/	Spherical	[85]
		Differential centrifugation	230	Negative	/	[86]
		Density gradient centrifugation	292	-12.9– -2.1	Spherical or cupped	[87]
		Differential centrifugation/filtration/ ultracentrifugation or density gradient centrifugation	120–150	/	Spherical	[52]
		Density gradient centrifugation	294	-29.7	Spherical	[66]
		Differential centrifugation/filtration/ ultracentrifugation	/	/	Spherical or oval	[80]
		Differential centrifugation/density gradient centrifugation	100–1000	-25	Spherical or cupped	[7]
		Differential centrifugation/density gradient centrifugation	232.7	Negative	Spherical	[88]
		Density gradient centrifugation	50–150	/	Spherical	[68]
		Differential centrifugation/density gradient centrifugation	100–600	/	/	[60]
Ginseng	<i>Panax ginseng</i>	Differential centrifugation/ultracentrifugation	344.8	-25.4	Spherical	[74]
		Differential centrifugation/filtration/ ultracentrifugation	92.04± 4.8	/	Spherical	[89]
Grape	<i>Vitis vinifera</i>	Differential centrifugation/filtration/ ultracentrifugation	30–200	/	Spherical	[90]
		Density gradient centrifugation	380	-27	Spherical	[35]
		Density gradient centrifugation	500–1000	-40	Spherical or cupped	[7]
Grapefruit	<i>Citrus paradisi</i>	Differential centrifugation/filtration	200.5	/	/	[36]
		UV irradiation/bath sonication/ultracentrifugation	110± 10	/	/	[91]
		Differential centrifugation/filtration	102.4	-38.15	Spherical	[92]
		Density gradient centrifugation	200	Negative	Spherical	[93]
		Differential centrifugation/filtration/ ultracentrifugation	/	/	Spherical or oval	[80]
		Density gradient centrifugation	50–100	-40	Spherical or cupped	[7]
		Differential centrifugation/filtration/ ultracentrifugation	147.7± 2.4	/	/	[78]
		Density gradient centrifugation	210.8 ± 48.62	-49.2– -1.52	Spherical or cupped	[44]

(Continued)

Table 2 (Continued).

Source	Scientific Name	Isolation Method	Size (nm)	Zeta (mV)	Shape	Ref.
		Differential centrifugation/filtration/ ultracentrifugation	105–400	–25	Spherical or cupped	[90]
Kiwi	<i>Actinidia chinensis</i>	Differential centrifugation/filtration/ ultracentrifugation	/	/	Spherical or oval	[80]
		Differential centrifugation/filtration/ ultracentrifugation	161.6± 0.6	/	/	[78]
Lemon	<i>Citrus limon</i>	Density gradient centrifugation	50–70	/	/	[64]
		Differential centrifugation/density gradient centrifugation	180.5± 3.2	/	Spherical	[94]
		Differential centrifugation/filtration/ ultracentrifugation	193.5± 2.2	/	/	[78]
Mistletoe	<i>Viscum album</i>	Vacuum infiltration/centrifugation/differential centrifugation	280± 115	/	Spherical	[54]
Nut	<i>Juglans regia</i>	Differential centrifugation/filtration	90–330	–21– –23	Spherical	[95]
	<i>Juglans californica</i>	Differential centrifugation/filtration	230	–21– –23	Spherical	[95]
	<i>Corylus avellana</i>	Differential centrifugation/filtration	90–560	–21– –23	Spherical	[95]
Orange	<i>Citrus sinensis</i>	Density gradient centrifugation/filtration/ ultracentrifugation	62± 12; 247± 61	/	Spherical	[15]
		Differential centrifugation/filtration/ ultracentrifugation	/	/	Spherical or oval	[80]
Pear	<i>Pyrus communis</i>	Differential centrifugation/filtration/ ultracentrifugation	/	/	Spherical or oval	[80]
Peas	<i>Pisum sativum</i>	Differential centrifugation/filtration/ ultracentrifugation	/	/	Spherical or oval	[80]
Periwinkle	<i>Vinca minor</i>	Vacuum infiltration/centrifugation/differential centrifugation	38± 200	/	Spherical	[54]
Soybean	<i>Glycine soja</i>	Differential centrifugation/filtration/ ultracentrifugation	/	/	Spherical or oval	[80]
Strawberry	<i>Fragaria ananassa</i>	Differential centrifugation/filtration/ ultracentrifugation	30–191	/	Round or cupped	[61]
Sunflower	<i>Helianthus annuus</i>	Vacuum infiltration/centrifugation procedure/ differential centrifugation/ultracentrifugation	50–200	/	Spherical	[96]
Tobacco	<i>Nicotiana tabacum</i>	Vacuum infiltration/centrifugation/differential centrifugation	70± 20	/	Spherical	[54]
Tomato	<i>Lycopersicon esculentum</i>	Differential centrifugation/filtration/ ultracentrifugation	110± 10	/	Spherical	[9]
		Differential centrifugation/filtration/ ultracentrifugation	100–1000	/	Spherical or oval	[80]

was 252 nm, which decreased with increase in PEG concentration.⁵⁵ The vesicle size determines the vesicle distribution in vivo; thus, different extraction methods should be carefully selected according to specific needs. Drug loading into PDVEs can be accomplished by simply co-incubating PDVEs with the drug, which is known as the passive loading technique. Active loading techniques such as ultrasonic treatment and freeze-thaw cycles can also be used to temporarily disrupt the PDVE membrane and increase the drug loading rate.⁷⁵

pH Tolerance

In clinical practice, oral administration has the highest patient compliance compared to other administration routes. However, a drug can be degraded by gastric acid, which results in a decline of bioavailability. Therefore, it is crucial to improve the acid tolerance of anti-cancer drugs. Several studies performed in vitro acid tolerance tests on *Kaempferia parviflora*-derived vesicles, and the results showed that *Kaempferia parviflora*-derived vesicles were stable in simulated gastric acid, maintaining their structural integrity with the size slightly enlarged. The only significant change was that the surface charge changed from negative to positive, due to the neutralization effect by hydrogen ions in the gastric acid.⁶⁹ Notably, not all PDVEs change surface charges in gastric acid. The size and the electric potential of tea flower-derived vesicles remained unchanged in a gastric simulant, small intestinal simulant, and colonic stimulant.⁹⁸ An illustrative example of PDVEs applied for oral drug administration is sorafenib. The bioavailability of sorafenib is significantly limited when orally administered due to degradation in gastric acid. However, when encapsulated within *Kiwifruit*-derived PDVEs, sorafenib exhibited enhanced stability in simulated gastric conditions, subsequently showing improved absorption and therapeutic efficacy.⁹⁹ The TME is also acidic, and PDVEs with acid tolerance can protect anti-cancer drugs in the microenvironment until they are taken up by cancer cells. In addition, Zhang et al reported that PDVEs were not only acid-tolerant but also resistant to pepsin catabolism.⁷ In conclusion, PDVEs resist gastric acid degradation in the stomach and avoid degradation by pepsin, which makes them excellent carriers for oral drug administration.

Crossing Barriers

Biological barriers block foreign substances to maintain homeostasis but also prevent drugs from reaching a lesion. When administered orally, most drugs are unable to enter the bloodstream because of the intestinal barrier. When treating brain-related diseases, drugs have difficulties entering the brain due to the blood-brain barrier and instead reach other organs, leading to potentially severe side effects.¹⁰⁰ Based on the phospholipid surface and nanostructure, PDVEs can transport a drug through a biological barrier and improve the bioavailability of the drug.

One prominent example showcasing the potential of PDVEs in surmounting biological barriers is their application in delivering dexamethasone (Dex) for neuroinflammation-related disease treatment. Dex is known for its anti-inflammatory properties, but its therapeutic potential is limited due to its poor bioavailability and inability to cross the blood-brain barrier efficiently. Researchers utilized *Allium tuberosum*-derived PDVEs to encapsulate Dex. Upon administration, these PDVEs were shown high efficiency in crossing the blood-brain barrier, and delivered Dex directly to the affected brain regions. This led to a significant alleviation of inflammation in microglial, demonstrating the potency of PDVEs as drug delivery vehicles for brain diseases.¹⁰¹

To verify the ability of PDVEs to penetrate the intestinal barrier, grape-derived vesicles (GDVEs) were stained with fluorescent dyes. The results showed that GDVEs accumulated in the intestine during the first 6 h after oral administration and were decreased within 48 h.³⁵ Most brain cancers require surgical intervention since the restrictive blood-brain barrier usually prevents drugs from reaching a therapeutic concentration. However, it was found that GDVEs carrying miR17 could significantly inhibit brain cancer growth, without any toxic effects.⁹² Notably, although PDVEs could cross biological barriers, they could not penetrate the placental barrier when injected intravenously into pregnant mice, suggesting the high safety of PDVEs in pregnant women.¹⁰² Taken together, PDVEs act as intelligent Trojan horses, encasing a drug and crossing a biological barrier to transport it to the lesion. Meanwhile, they are safe for the fetus in the uterus.

Surface Modifications and Active Targeting

Regardless of the delivery method, the drug will be transported everywhere in the body through the circulatory system after administration, which inevitably decreases the drug concentration at the cancer lesion and increases the risk of toxic

side effects in other organs. Therefore, it is important to enhance the cancer-targeting ability of PDVEs to improve the bioavailability of drugs at the target and reduce their toxic side effects. This is crucial for increasing the efficacy of anti-cancer drugs.

Currently, there are two main approaches for improving PDVEs through surface modifications. One is loading the vesicle surface with ligands capable of binding with cancer-specific receptors. For example, a therapeutically siRNA is fused into the 3WJ core, modified with folic acid, coupled to cholesterol, and incubated with GiDVEs membranes. By cholesterol anchoring to the membrane, the complex is formed with 80% drug loading.⁸⁵ It was shown that the vesicles efficiently delivered survivin siRNA to a human oral epidermal-like (KB) cancer model and that folic acid ligands significantly promoted the uptake of GiDVEs by the KB cells. Another approach is to modify vesicles based on the TME. For example, the leukocyte membrane in a cancer-inflammatory environment was utilized to wrap around a GfDVE,⁹³ which avoided the clearance by the immune system and allowed the vesicles to directly deliver to the site of inflammatory cancer. Therefore, the surface of PDVEs is amenable for modification, such as loading small molecules or covering with bionic membranes. This property makes it possible to use PDVEs for enhanced targeting of cancer cells and warrants more in-depth investigation.

Cellular Uptake

The ability to be taken up by mammalian cells is a prerequisite for PDVEs to exchange information between plants and animals. As far as is known, PDVEs are taken up by mammalian cells through multiple pathways, depending on the type of plant donors and the type of animal recipient cells. For example, GfDVEs enter the intestinal cells through micropinocytosis,³⁵ whereas GiDVEs are specifically internalized by enterocytes via caveolin-mediated endocytosis and micropinocytosis.¹⁰³ The entry of GiDVEs into hepatocytes is ATP-dependent, and its uptake rate is slow at 4°C and increases with increasing temperature.⁶⁶ Furthermore, it was found that the main pathway of macrophage internalization of GiDVEs was not macrocytic drinking but clathrin-mediated endocytosis. Furthermore, the intracellular GiDVEs directly induced macrophage polarization and improved the cancer immune suppression microenvironment, resulting in an enhanced therapeutic efficacy. This phenomenon suggests that PDVEs can transport their contents into the cytoplasm, thus altering the physiological state of the recipient cells.⁷⁴

However, current research on the uptake mechanism of PDVEs by mammalian cells is still incomplete, and alterations of gene or protein expression of the recipient cells remain unknown. Elucidation of the molecular mechanisms involved in the internalization process will provide more efficient delivery and better therapeutic effects.

As a Therapeutic Agent

In nature, plants and animals are interdependent, and research on the information exchange between the two biological communities is ongoing. Recently, it was found that the nucleic acids, proteins and enzymes within PDVEs positively regulated the physiological homeostasis of animals when absorbed by their cells.^{103,104} Thus, PDVEs can act not only as carriers but also as therapeutic agents (Table 3). Emerging studies show that PDVEs have anti-cancer effects but have little influence on normal tissues,¹⁰⁵ which closely correlated with the contents of the PDVEs.¹⁰⁶ Here, we systematically describe the contents of PDVEs and their involvement in anti-cancer strategies, such as chemotherapy, starvation therapy, and immunotherapy (Figure 3).

Contents

Like MDVEs, PDVEs contain various biological information of donors, including lipids, nucleic acids, proteins, and metabolites. To clarify the therapeutic mechanism of PDVEs, a number of studies have identified several types of inclusions in PDVEs.

In contrast to MDVEs, the lipids of PDVEs are diverse. For example, the lipids of GiDVEs consisted primarily of 42% phosphatidic acid (PA), 27% digalactosyl diacylglycerol, and 19% monogalactosyl diacylglycerol.⁸⁷ In contrast, the lipids of GfDVEs consisted primarily of 24% phosphatidylethanolamine, 23% phosphatidylcholine (PC), and 13% phosphatidylinositol.¹¹¹ The lipids of GDVEs were composed of 53% PA and 26% phosphatidylethanolamine.³⁵ Moreover, lipids from different PDVEs have been shown to have different functions. PAs from GiDVEs prolonged

Table 3 Applications of PDVEs in Cancer Therapy

Source	Cancer type	Mechanism	Cargo	Administration Way	Ref
Bitter melon	Oral squamous cell carcinoma	Generate ROS Up-regulate JUN protein Downregulate NLRP3 expression	/	Intravenous injection	[22]
Cannabis	In vitro: Hepatocellular carcinomas (HepG2; Huh-7)	Arrest the G0/G1 phase of the cell growth cycle Induce cell death by activating mitochondrial-dependent apoptosis signaling pathways	/	/	[107]
Garlic	In vitro: Kidney cancer cells (A498); Lung adenocarcinoma (A549)	Arrest the S phase of the cell growth cycle Upregulate gene expression levels of pro-apoptotic genes such as p53, Bax, Cas3, and Cas9 Downregulate angiogenic VEGF protein expression levels	/	/	[23]
Ginger	Oral epidermoid carcinoma	Target tumor cells	siRNA	Intravenous injection	[85]
	Colon cancer	Reduce side effects Dual targeting of epithelial cells and macrophages	/	Oral administration	[86]
	Colon cancer	High drug loading rate pH-dependent drug release Target tumor cells	DOX	Oral administration	[17]
	Colon cancer	Dual targeting of intestinal epithelial cells and macrophages Increase survival and proliferation of intestinal epithelial cells Reduce proinflammatory cytokines (TNF- α , IL-6 and IL-1 β) Increase anti-inflammatory factors (IL-10 and IL-22)	/	Oral administration	[87]
Ginseng	BI6F10 Melanoma	Promote the transformation of M2 type to M1 type Generate ROS	/	Intraperitoneally injection	[74]
	Colon adenocarcinoma (COAD) Triple negative breast cancer (TNBCs)	Promote secretion of CCL5 and CXCL9 from macrophages Recruit CD8 ⁺ T cells into the tumor microenvironment Target macrophage cells	PD-1 mAb	Intraperitoneally injection	[108]
Grape	Colon cancer	Gastrointestinal fluid tolerance Able to penetrate into epithelial mucosa and be taken up by intestinal stem cells Promote proliferation of Lgr5 ⁺ intestinal stem cells	/	Oral administration	[109]
Grapefruit	Colon cancer (CT26; SW620)	Target tumor cells Extend the circulation time in the body	JSI-124	Intravenous injection	[36]

(Continued)

Table 3 (Continued).

Source	Cancer type	Mechanism	Cargo	Administration Way	Ref
	In vitro: Breast adenocarcinoma (MCF7); Human melanoma (A375); Lung adenocarcinoma (A549)	Downregulate Cyclin B1 and Cyclin B2 expression levels Up-regulate cell cycle inhibitor p21 Arrest cell cycle in the G2/M phase	/	/	[110]
	Brain Tumor (GL-26)	Target tumor cells High RNA loading rate	miR17	Intranasal administration	[92]
	Colon cancer (CT26); Breast cancer (4T1)	Targeting the tumor inflammatory environment using leukocyte membrane wrapping	DOX	Intravenous injection	[93]
Lemon	In vitro: Lung adenocarcinoma (A549); Colon cancer (SW480); Chronic myeloid leukemia cells (LAMA84) in vivo: Chronic myeloid leukemia cells (LAMA84)	Target tumor cells Activate TRAIL-mediated apoptosis Down-regulate the expression of apoptosis genes Survivin and Bcl-xl Inhibit angiogenesis	/	Intraperitoneally injection	[64]
Petasites japonicus	/	Promote the expression of surface molecules and production of Th1-polarizing pro-inflammatory cytokines in BMDCs Induce the Maturation of DCs via the Activation of MAPK and NF- κ B Signaling Pathways Induce Naïve T Cells toward Th1 Polarization and Activated CD8+ T Cells	/	/	[21]
Tea	Breast cancer	Generate ROS Trigger mitochondrial damage Arrest cell growth cycle	/	Intravenous injection	[98]

Abbreviations: SLNVEs, Synthetic lipid vesicles; MDVEs, Mammalian-derived vesicles; PDVEs, Plant-derived vesicles; TME, Tumor microenvironment; mRNA, Messenger ribonucleic acid; DNA, Deoxyribonucleic acid; siRNA, Small interfering ribonucleic acid; miRNA, Micro ribonucleic acid; GfDVEs, Grapefruit-derived vesicles; TRAIL, Tumor-necrosis factor-related apoptosis inducing ligand; GiDVEs, Ginger-derived vesicles; PEG6000, Polyethylene glycol-6000; GDVEs, Grape-derived vesicles; KB, Human oral epidermal-like; PA, Phosphatidic acid; PC, Phosphatidylcholine; GaDVEs, Garlic-derived vesicles; TAMs, Tumor-associated macrophages; TNF- α , Tumor-necrosis factor- α ; ROS, Reactive oxygen species; TCM, Traditional Chinese medicine.

the resident time of PDVEs in the small intestine and were absorbed by the intestinal microbiota,⁶⁰ whereas PAs from GDVEs were able to mediate cell proliferation in the colon.³⁵

miRNAs represent another important component in PDVEs. A total of 198 miRNAs and 39 novel miRNAs were identified in ginger, among which 136 miRNAs were also found in GiDVEs. A molecular mechanism study showed that most co-expressed miRNAs between ginger and its vesicles were responsible for regulating stress responses, root development, and other physiological processes in plants.⁶⁸ In addition, high-throughput sequencing techniques and bioinformatics analyses have shown that plant-derived miRNAs have the potential to regulate human metabolic activities and even treat human diseases.⁶⁸ For example, miRNAs in GiDVEs regulated intestinal flora homeostasis and alleviated lipopolysaccharide induced colitis.¹⁰³ Furthermore, miRNAs extracted from coconut water-derived vesicles can target the metabolism-related mRNAs in human genome.⁸²

Finally, proteins are also considered as significant signaling molecules in PDVEs. Proteomic analysis identified 745 proteins in tea flower-derived vesicles. The Gene Ontology database and gene and genome (KEGG) annotation pathway analysis revealed that the proteins in the vesicles were mainly related to cellular composition, metabolism, and growth.




Figure 3 Applications of plant-derived vesicles in cancer therapy as a therapeutic agent.

Among them, 16 proteins had oxidation-related functions, suggesting that these vesicles may trigger intracellular oxidative stress.⁹⁸

In summary, the diverse biological components within PDVEs play pivotal roles in both plant and human physiology. The distinct lipid profiles of various PDVEs not only signify their uniqueness but also underline specific therapeutic potentials, especially in gastrointestinal applications. The presence of numerous miRNAs in PDVEs, particularly those with origins in plants like ginger, emphasizes their potential in regulating a myriad of cellular processes, from stress responses in plants to metabolic activities in humans. Moreover, the abundance of proteins, especially those with antioxidant properties, accentuates the potential of PDVEs in mediating cellular events, possibly even leading to responses like oxidative stress.

Cytotoxic Killing Therapy

Apoptosis and cell cycle arrest are the main evaluating indicators for the development of anti-cancer drugs. It was demonstrated that garlic-derived vesicles (GaDVEs) induced apoptosis in cancer cells by activating caspase-mediated intrinsic pathways. The expression levels of pro-apoptotic genes such as TP53, BAX, CASP3, and CASP9 were significantly increased. In contrast, the expression levels of anti-apoptotic genes were significantly decreased in kidney and lung cancer cells following treatment with GaDVEs.²³ In addition, GaDVEs blocked the growth cycle of cancer cells, leading to S phase arrest. GADD45A, a gene regulating cell cycle arrest and DNA damage,¹¹² is an essential signal in the cellular network response to external damage. Yang et al found that lemon-derived vesicles induced upregulation of GADD45A in 3D cultures of SGC-7901 spheroids, resulting in cell cycle arrest.⁵⁷

However, it is essential to note that the cytotoxic killing effect of PDVEs can be varied depending on the donor species. In one study,¹¹⁰ Cho et al analyzed vesicles derived from four Citrus species: *C. sinensis*, *C. limon*, *C. paradisi*, and *C. aurantium*, and showed that all four of these PDVEs inhibited the proliferation of A375 (human melanoma), A549 (human lung cancer), and MCF-7 (human breast cancer) cells. However, only vesicles from *C. paradisi* were able to block the cell cycle of melanoma cells in the G2/M phase by enhancing the gene expression of CDKN1A (encoding p21) and reducing the levels of cyclin B1 and cyclin B2. In addition to plant species differences, the anti-cancer effects of PDVEs produced from different locations of the same plant may also vary. For example, vesicles from the roots of

Dendropanax morbifera inhibited melanoma by downregulating the expression of TYR, TRP-1, and TRP-2 in B16BL6 cells, but vesicles from leaves were more effective.⁷⁴

The contents in PDVEs also determine the cytotoxic killing effects against cancer cells. It was shown that the PDVEs from Cannabis sp. could be classified into high cannabidiol vesicles and low cannabidiol vesicles based on the concentration of cannabidiol. Compared to the low cannabidiol vesicles, the high cannabidiol vesicles significantly inhibited the cell viability of hepatocellular carcinoma cell lines HepG2 and Huh-7 in a time- and dose-dependent manner but had no inhibitory effect on normal growth of HUVECs. Molecular mechanism investigations further indicated that the high cannabidiol vesicles blocked the cell cycle at the G0/G1 phase and activated the mitochondria-dependent apoptotic signaling pathway in hepatocellular carcinoma cells.¹⁰⁷

Several experiments have demonstrated the high selectivity of PDVEs for cancer cells. First, the difference in the composition of cancer cell membranes and normal cell membranes affects the uptake pathway of cells, resulting in different amounts of PDVEs entering the cancer cells and normal cells. One study showed that HepG2 cells took up a two-fold increase of asparagus-derived vesicles compared to LO2 cells. Thus, asparagus-derived vesicles upregulated apoptotic factors and induced apoptosis in hepatocellular carcinoma cells but caused less damage to normal hepatocytes.⁷³ Second, PDVEs were more likely to activate apoptotic pathways to which cancer cells are susceptible, such as TRAIL-mediated apoptosis.¹¹³ Specifically, vesicles in citrus (Rutaceae) sap exerted their anti-cancer effects by stimulating the TRAIL-mediated apoptotic pathway⁹⁷ but had no toxic effects on normal cells.

In conclusion, the potential of PDVEs in anti-cancer therapies is vast, showcasing not only their ability to selectively target cancer cells but also the importance of their donor species and specific plant parts. Whether through specific molecular pathways like caspase-mediated intrinsic or TRAIL-mediated apoptosis, or through the differential composition influencing uptake by cancer versus normal cells, the versatility of PDVEs is evident. The findings also underscore the significance of studying individual constituents, such as cannabidiol concentration, in influencing the cytotoxic effects of these vesicles.

Immune Modulatory Therapy

With the recognition of the body's innate ability to combat cancer, Immune modulatory therapy took center stage. This approach seeks to harness and amplify the body's immune system's natural capability to detect and destroy cancer cells. By modulating immune checkpoints or introducing tailored immune cells, this therapy offers a more targeted approach with fewer side effects than cytotoxic agents.

The TME is infiltrated by a large number of inflammatory cells, including TAMs, lymphocytes, natural killer cells, dendritic cells, and mast cells. TAMs are the most predominant subset.¹¹⁴ TAMs are plastic and can be M1- or M2-polarized upon stimulation by different cytokines. Generally, the M1 phenotype has anti-cancer effects, while the M2 phenotype has pro-cancer properties.¹¹⁵ PDVEs can modulate the immune microenvironment by altering the polarization of TAMs.

Cao et al demonstrated that GiDVEs promoted the polarization from the M2 phenotype to the M1 phenotype.⁷⁴ The results showed that GiDVE treatment significantly increased the secretion of M1 phenotype-associated cytokines, including tumor-necrosis factor- α (TNF- α), interleukin-12 and interleukin-6, while the expression of the M2 phenotype marker CD206 was significantly reduced.⁷⁴ To clarify the active components regulating macrophage polarization, TAMs were treated with DNase I, RNase I, proteinase K, or sonicated GiDVEs. The results showed that proteinase K and sonicated GiDVE-treated TAMs were unable to induce macrophage polarization. This indicates that the proteins in GiDVEs are the main components responsible for macrophage polarization, and their structural integrity is essential for regulating this process.

In the last decade, immune checkpoint inhibitors have received much attention in cancer therapy for their ability to promote cytotoxic T cell activity to kill cancer cells. However, immune checkpoint inhibitors are ineffective in solid cancers because of the insufficient infiltration of T cells; thus, these are referred to as cold tumors. PDVEs have been reported to enhance the therapeutic effects of immune checkpoint antibodies by reprogramming the microenvironment of cold tumors. The results showed that GiDVEs reprogrammed TAMs to increase CCL5 and CXCL9 secretion and recruited CD8⁺ T cells into the TME, converting the cold TME to a hot one. When combined with programmed

death-1 inhibitors, the cancer progression in both colon and breast cancer models was inhibited, with no detectable systemic toxicity.¹⁰⁸

In summary, the profound effects of GiDVEs on TAM polarization and the subsequent modulation of the tumor microenvironment underscore their potential as vital tools in oncology. The specific protein components in GiDVEs, pivotal for macrophage polarization, emphasize the importance of understanding the molecular components driving these effects. Further, the ability of PDVEs to turn “cold” tumor environments “hot” by reprogramming TAMs, combined with immune checkpoint inhibitors, establishes a promising avenue in tackling even solid tumors.

Oxidative Therapy

However, certain tumors evade the immune system or develop resistance. This led to the exploration of Oxidative therapy, a more recent innovation. Cancer cells, due to their altered metabolism, are often more susceptible to oxidative stress than normal cells. Oxidative therapy capitalizes on this vulnerability by inducing a surge of reactive oxygen species (ROS) in the tumor environment, leading to cancer cell death while sparing most normal cells.

Reactive oxygen species (ROS), including peroxides, superoxides, hydroxyl radicals, singlet oxygen, and α -oxygen, are natural by-products of oxygen metabolism and play an essential role in cell signaling and maintenance of organism homeostasis. However, excessive levels of ROS can cause severe damage to cells.¹⁰⁵ Therefore, ROS bursts in the TME can trigger oxidative stress in cancer cells and achieve anti-cancer effects. It was reported that edible tea flower-derived vesicles inhibited metastatic breast cancer by generating ROS. The increased intracellular ROS levels triggered mitochondrial damage and blocked the cell cycle, thus suppressing breast cancer cell proliferation and migration. In contrast, the tea flower-derived vesicles led to little ROS production and minimal mitochondrial damage in normal cells.⁹⁸ A similar phenomenon was also observed with lemon-derived vesicles.⁵⁷

Elevated ROS may also improve the susceptibility of drug-resistant cancer cells. It was found that bitter melon-derived vesicles improved the chemosensitivity of 5-fluorouracil-resistant oral squamous cell carcinoma cells and thus enhanced the therapeutic effect of 5-fluorouracil *in vitro* and *in vivo*.²² However, when treated with N-acetylcysteine, the chemosensitizing effects disappeared, suggesting that the synergistic effects were dependent on ROS production. Notably, bitter melon-derived PDVEs scavenged excessive ROS production caused by X-ray irradiation in normal H9C2 cells and thus rebalanced the mitochondrial membrane potential and attenuated DNA damage.¹¹⁶ Therefore, the intracellular ROS level may be the underlying reason for the highly selective action of PDVEs against cancer cells.

In essence, the dual role of ROS, both as a potential enhancer of cancer therapies and as a possible protective agent against cellular damage, is evident. The selective action of PDVEs, underpinned by intracellular ROS levels, provides a fascinating insight into their anti-cancer potential and their ability to safeguard non-malignant cells from unwanted oxidative stress. Such a discerning approach offers a promising direction in optimizing cancer treatments while preserving the integrity of surrounding tissues.

Other Therapeutic Effects

In the TME, the relative hypoxic environment stimulates angiogenesis, which ensures the delivery of nutrients required by cancer cell proliferation. Therefore, angiogenesis inhibition can block the supply of nutrients and thus inhibit cancer growth. It was reported that GaDVEs inhibited cancer growth by reducing VEGF expression *in vitro*, without affecting normal human dermal fibroblasts.²³ However, such evidence is still limited, and further investigation is necessary to provide more evidence.

In addition to anti-cancer effects, PDVEs also exhibit remarkable preventive effects against carcinogenesis. It was found that oral administration of tea leaf-derived vesicles reduced the levels of pro-inflammatory cytokines, including TNF- α , interleukin-6, and interleukin-12, while increasing the expression of the anti-inflammatory cytokine interleukin-10, and thus effectively inhibited inflammatory bowel responses. At the same time, the expression of tight junction-related proteins zona occludens protein 1 and mucin 2 were elevated to restore the damaged colonic barrier. Furthermore, the vesicles enhanced the diversity and abundance of the intestinal microbiota, thereby preventing or alleviating inflammatory bowel disease and colitis-associated colon cancer.¹¹⁷

In summary, the intricate interplay within the TME underscores the significance of angiogenesis as a potential therapeutic target. Preliminary findings highlight the inhibitory effects of GaDVEs on cancer growth through VEGF modulation. Yet, the journey to fully understand their potential is still underway, necessitating more rigorous studies. On another front, PDVEs are emerging as potent preventive agents against carcinogenesis. Their ability to modulate the inflammatory milieu, bolster barrier integrity, and foster a healthful gut microbiome place them at the forefront of innovative strategies against inflammatory diseases and associated cancers. The multipronged effects of PDVEs accentuate their vast potential in cancer research and treatment.

Discussion and Perspectives

Here, we systematically summarized the preparation, contents, and application of PDVEs in cancer prevention and treatment. Although SLNVEs meet the basic requirements for drug delivery,¹¹⁸ they are still challenged by a number of problems such as cytotoxicity, non-specific tissue distribution, and therapeutic efficacy. SLNVEs tend to be phagocytosed by the reticuloendothelial system, resulting in low bioavailability and hypersensitivity reactions.¹¹⁹ In contrast, the natural properties of PDNVEs allow them to avoid these side effects. Compared to MDVEs, PDVEs have no zoonotic pathogens, can be manufactured from abundant resources, are easily extracted, and are environmentally friendly.¹⁹

Although PDVEs have great potential in anticancer therapy, a series of issues remain to be solved. First, there is no standardization of PDVEs in the extraction process and no representative markers to qualify PDVEs. The differences in plant donor species make it difficult to establish quality control criteria for PDVEs. Macroscopic control can be performed based on commonalities of PDVEs, such as describing their shape, size, and potential, or identifying their compositions including lipids, proteins, and nucleic acids. For example, transmission electron microscopy is typically used to detect the ultrastructure of PDVEs, and dynamic light scattering is used to determine the size and potential of PDVEs. In addition, it is vital to identify the composition of PDVEs. Unlike other vesicles, PDVEs are composed of multiple lipids. By determining the lipid composition of the PDVEs extracted to date, it was found that each PDVE contains phospholipids and glycerol lipids, but none contain cholesterol.¹²⁰ It is known that vesicles of mammalian origin have well-defined protein markers, including the transmembrane proteins CD63, CD81, and CD9. PDVEs do not yet have clear protein markers, but all PDVEs inevitably require some hydrolase to cleave the cell wall during release,¹²¹ and therefore, this hydrolase may be a candidate for protein markers.

Second, PDVEs may carry allergens from the donor. Stanly et al evaluated the allergen information carried in strawberry-derived vesicles.¹²² Immunological testing revealed that several proteins in strawberry-derived vesicles, such as Fra a 1, Fra a 3, and Fra a 4, could competitively bind with immunoglobulin E, suggesting that PDVEs may induce allergic reactions in human beings. In addition to allergens, plant exposure to toxic substances during growth may also remain in the PDVEs,⁷⁸ such as organophosphorus, organonitrogen, organoarsenic, and dinitroaniline. Therefore, a strict selection of plant donors can effectively avoid potential risks, such as organic plants, in-season plants, and non-GMO plants.

In addition to the aforementioned lack of standardized extraction methods and potential allergenic issues associated with PDVEs, there are several other challenges confronting PDVE research: (1) The mechanisms of PDVE formation and release remain unclear; (2) There is no unified naming convention for PDVEs; (3) Standardized biomarkers for PDVEs are lacked; (4) Techniques for PDVE preservation and transportation are still in their infancy; (5) The distribution and metabolism of PDVEs within the body are not well-understood. Despite the burgeoning research in PDVEs, further efforts are required globally from the scientific community to fully harness its potential as a biological therapy or drug carrier. Specifically, it is necessary to elucidate the roles of PDVEs in plant growth, stress responses, and death, as well as the mechanisms underlying PDVE formation and release. Meanwhile, it is urgent to develop more cost-effective and time-saving extraction methods suitable for large-scale production, along with robust preservation and transportation techniques for clinical applications. In addition, we need to establish a comprehensive, intuitive, and easily adopted naming system that differentiates PDVEs based on type, origin, location within the plant, and state. More importantly, identification of generic biomarkers for PDVEs to standardize subsequent extractions is significant for mechanism exploration. Last but not least, the impacts of PDVEs, when introduced into the body via various administration routes, on critical systems such as the immune, hematological, and nervous systems, as well as their distribution patterns and

metabolic pathways in the body, is worth to be intensively investigated on multiple diseases, such as malignancies and auto-immune diseases.

Finally, Traditional Chinese medicine (TCM) is a valuable resource to screen and prepare efficient PDVEs. TCM has been appreciated for cancer prevention and treatment in Asian countries for thousands of years. Although a number of formulas have been validated in clinical settings, the underlying molecular mechanisms remain unknown. In recent years, with the discovery of PDVEs, it was hypothesized that PDVEs are largely produced and precipitated from decoctions, and the proteins, nucleic acids, lipids, and metabolites in them may contribute to the therapeutic effects. The hypothesis is also supported by recent findings, which suggested that most of an herb's active ingredients are covered in PDVEs.¹²³ It was found that ginger contained 237 miRNAs, 136 of which were found in GiDVEs and were mainly involved in modulating physiological processes such as a stress response. A total of 745 proteins were found in the vesicles from tea flower and were demonstrated to be crucial components responsible for biological processes including gene expression, signaling pathway transmission, and cellular metabolism.⁹⁸ Currently, characterization and molecular elucidation of PDVEs in herbs or formulas have become a topic of interest in TCM study.

Taken together, the available evidence suggests that PDVEs offer tremendous advantages in anti-cancer therapy, both as carriers and as therapeutic agents. PDVEs offer cancer patients a new strategy for targeted cytotoxic therapy, immunomodulatory therapy, oxidative therapy, as well as neoangiogenesis inhibition therapy. However, further research is still warranted to optimize PDVEs concerning their quality control, safety, and high-throughput production.

Acknowledgments

This work was supported by the National Natural Science Foundation of China [82074165, 82374446, 81973526]; the State Key Laboratory of Dampness Syndrome of Chinese Medicine[SZ2021ZZ19]; Science and Technology Planning Project of Guangdong Province [2022B1515230002, 2021A0505030059, 2017B030314166]; The 2020 Guangdong Provincial Science and Technology Innovation Strategy Special Fund (Guangdong-Hong Kong-Macau Joint Lab) [2020B1212030006]; and the Foundation for Young Scholars of Guangzhou University of Chinese Medicine [QNYC20190101].

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Liu Y, Wang Y, Lv Q, Li X. Exosomes: from garbage bins to translational medicine. *Int J Pharm.* 2020;583:119333. doi:10.1016/j.ijpharm.2020.119333
2. Halperin W, Jensen WA. Ultrastructural changes during growth and embryogenesis in carrot cell cultures. *J Ultrastruct Res.* 1967;18(3):428–443. doi:10.1016/s0022-5320(67)80128-x
3. An Q, Ehlers K, Kogel KH, van Bel AJ, Hüchelhoven R. Multivesicular compartments proliferate in susceptible and resistant MLA12-barley leaves in response to infection by the biotrophic powdery mildew fungus. *New Phytol.* 2006;172(3):563–576. doi:10.1111/j.1469-8137.2006.01844.x
4. An Q, Hüchelhoven R, Kogel KH, van Bel AJ. Multivesicular bodies participate in a cell wall-associated defence response in barley leaves attacked by the pathogenic powdery mildew fungus. *Cell Microbiol.* 2006;8(6):1009–1019. doi:10.1111/j.1462-5822.2006.00683.x
5. Alfieri M, Leone A, Ambrosone A. Plant-derived nano and microvesicles for human health and therapeutic potential in nanomedicine. *Pharmaceutics.* 2021;13(4):498. doi:10.3390/pharmaceutics13040498
6. Garaeva L, Kamyshinsky R, Kil Y, et al. Delivery of functional exogenous proteins by plant-derived vesicles to human cells in vitro. *Sci Rep.* 2021;11(1). doi:10.1038/s41598-021-85833-y
7. Mu J, Zhuang X, Wang Q, et al. Interspecies communication between plant and mouse gut host cells through edible plant derived exosome-like nanoparticles. *Mol Nutr Food Res.* 2014;58(7):1561–1573. doi:10.1002/mnfr.201300729
8. Fujita D, Arai T, Komori H, et al. Apple-derived nanoparticles modulate expression of organic-anion-transporting polypeptide (OATP) 2B1 in caco-2 cells. *Mol Pharm.* 2018;15(12):5772–5780. doi:10.1021/acs.molpharmaceut.8b00921
9. Bokka R, Ramos AP, Fiume I, et al. Biomanufacturing of tomato-derived nanovesicles. *Foods.* 2020;9(12):1852. doi:10.3390/foods9121852

10. Hinshaw DC, Shevde LA. The tumor microenvironment innately modulates cancer progression. *Cancer Res.* 2019;79(18):4557–4566. doi:10.1158/0008-5472.Can-18-3962
11. Boedtkjer E, Pedersen SF. The acidic tumor microenvironment as a driver of cancer. *Annu Rev Physiol.* 2020;82:103–126. doi:10.1146/annurev-physiol-021119-034627
12. Greten FR, Grivennikov SI. Inflammation and cancer: triggers, mechanisms, and consequences. *Immunity.* 2019;51(1):27–41. doi:10.1016/j.immuni.2019.06.025
13. Jiang X, Wang J, Deng X, et al. The role of microenvironment in tumor angiogenesis. *J Exp Clin Cancer Res.* 2020;39(1):204. doi:10.1186/s13046-020-01709-5
14. Lee Y, Thompson DH. Stimuli-responsive liposomes for drug delivery. *Wiley Interdiscip Rev Nanomed Nanobiotechnol.* 2017;9(5). doi:10.1002/wnan.1450
15. Bruno SP, Paolini A, D’Oria V, et al. Extracellular vesicles derived from citrus sinensis modulate inflammatory genes and tight junctions in a human model of intestinal epithelium. *Front Nutr.* 2021;8. doi:10.3389/fnut.2021.778998
16. Kumar A, Sundaram K, Teng Y, et al. Ginger nanoparticles mediated induction of Foxa2 prevents high-fat diet-induced insulin resistance. *Theranostics.* 2022;12(3):1388–1403. doi:10.7150/thno.62514
17. Zhang M, Xiao B, Wang H, et al. Edible ginger-derived nano-lipids loaded with doxorubicin as a novel drug-delivery approach for colon cancer therapy. *Mol Ther.* 2016;24(10):1783–1796. doi:10.1038/mt.2016.159
18. Mao Y, Han M, Chen C, et al. A biomimetic nanocomposite made of a ginger-derived exosome and an inorganic framework for high-performance delivery of oral antibodies. *Nanoscale.* 2021;13(47):20157–20169. doi:10.1039/d1nr06015e
19. Cai Y, Zhang L, Zhang Y, Lu R. Plant-derived exosomes as a drug-delivery approach for the treatment of inflammatory bowel disease and colitis-associated cancer. *Pharmaceutics.* 2022;14(4):822. doi:10.3390/pharmaceutics14040822
20. Karamanidou T, Tsouknidas A. Plant-derived extracellular vesicles as therapeutic nanocarriers. *Int J Mol Sci.* 2021;23(1):191. doi:10.3390/ijms23010191
21. Han JM, Song H-Y, Lim S-T, Kim K-I, Seo HS, Byun E-B. Immunostimulatory potential of extracellular vesicles isolated from an edible plant, *Petasites japonicus*, via the induction of murine dendritic cell maturation. *Int J Mol Sci.* 2021;22(19):10634. doi:10.3390/ijms221910634
22. Yang M, Luo Q, Chen X, Chen F. Bitter melon derived extracellular vesicles enhance the therapeutic effects and reduce the drug resistance of 5-fluorouracil on oral squamous cell carcinoma. *J Nanobiotechnology.* 2021;19(1):259. doi:10.1186/s12951-021-00995-1
23. Özkan İ, Koçak P, Yıldırım M, et al. Garlic (*Allium sativum*)-derived SEVs inhibit cancer cell proliferation and induce caspase mediated apoptosis. *Sci Rep.* 2021;11(1). doi:10.1038/s41598-021-93876-4
24. Crommelin DJA, van Hoogevest P, Storm G. The role of liposomes in clinical nanomedicine development. What now? Now what? *J Control Release.* 2020;318:256–263. doi:10.1016/j.jconrel.2019.12.023
25. Grimaldi N, Andrade F, Segovia N, et al. Lipid-based nanovesicles for nanomedicine. *Chem Soc Rev.* 2016;45(23):6520–6545. doi:10.1039/c6cs00409a
26. Sonju JJ, Dahal A, Singh SS, Jois SD. Peptide-functionalized liposomes as therapeutic and diagnostic tools for cancer treatment. *J Control Release.* 2021;329:624–644. doi:10.1016/j.jconrel.2020.09.055
27. Li X, Diao W, Xue H, et al. Improved efficacy of doxorubicin delivery by a novel dual-ligand-modified liposome in hepatocellular carcinoma. *Cancer Lett.* 2020;489:163–173. doi:10.1016/j.canlet.2020.06.017
28. Hu M, Palić D. Micro- and nano-plastics activation of oxidative and inflammatory adverse outcome pathways. *Redox Biol.* 2020;37:101620. doi:10.1016/j.redox.2020.101620
29. Yue X, Dai Z. Recent advances in liposomal nanohybrid cerasomes as promising drug nanocarriers. *Adv Colloid Interface Sci.* 2014;207:32–42. doi:10.1016/j.cis.2013.11.014
30. Raemdonck K, Braeckmans K, Demeester J, De Smedt SC. Merging the best of both worlds: hybrid lipid-enveloped matrix nanocomposites in drug delivery. *Chem Soc Rev.* 2014;43(1):444–472. doi:10.1039/c3cs60299k
31. Radmanesh F, Razi M, Shalzar-Jalali A. Curcumin nano-micelle induced testicular toxicity in healthy rats; evidence for oxidative stress and failed homeostatic response by heat shock proteins 70-2a and 90. *Biomed Pharmacother.* 2021;142:111945. doi:10.1016/j.biopha.2021.111945
32. Lancet JE, Uy GL, Cortes JE, et al. CPX-351 (cytarabine and daunorubicin) liposome for injection versus conventional cytarabine plus daunorubicin in older patients with newly diagnosed secondary acute myeloid leukemia. *J Clin Oncol.* 2018;36(26):2684–2692. doi:10.1200/jco.2017.77.6112
33. Anand U, Carpena M, Kowalska-Góralaska M, et al. Safer plant-based nanoparticles for combating antibiotic resistance in bacteria: a comprehensive review on its potential applications, recent advances, and future perspective. *Sci Total Environ.* 2022:821. doi:10.1016/j.scitotenv.2022.153472
34. Feng J, Xiu Q, Huang Y, Troyer Z, Li B, Zheng L. Plant-derived vesicle-like nanoparticles as promising biotherapeutic tools: present and future. *Adv Mater.* 2023;35(24):e2207826. doi:10.1002/adma.202207826
35. Ju S, Mu J, Dokland T, et al. Grape exosome-like nanoparticles induce intestinal stem cells and protect mice from DSS-induced colitis. *Molecular Therapy.* 2013;21(7):1345–1357. doi:10.1038/mt.2013.64
36. Wang Q, Zhuang X, Mu J, et al. Delivery of therapeutic agents by nanoparticles made of grapefruit-derived lipids. *Nat Commun.* 2013;4:1867. doi:10.1038/ncomms2886
37. Pomatto MAC, Bussolati B, D’Antico S, et al. Improved loading of plasma-derived extracellular vesicles to encapsulate antitumor miRNAs. *Mol Ther Methods Clin Dev.* 2019;13:133–144. doi:10.1016/j.omtm.2019.01.001
38. Li Y, Gao Y, Gong C, et al. A33 antibody-functionalized exosomes for targeted delivery of doxorubicin against colorectal cancer. *Nanomedicine.* 2018;14(7):1973–1985. doi:10.1016/j.nano.2018.05.020
39. Xiao L, Erb U, Zhao K, Hackert T, Zoller M. Efficacy of vaccination with tumor-exosome loaded dendritic cells combined with cytotoxic drug treatment in pancreatic cancer. *Oncoimmunology.* 2017;6(6):e1319044–1319088. doi:10.1080/2162402X.2017.1319044
40. Sheng S, Yu X, Xing G, et al. An apoptotic body-based vehicle with navigation for photothermal-immunotherapy by precise delivery and tumor microenvironment regulation. *Adv Funct Mater.* 2023;33(5):2212118. doi:10.1002/adfm.202212118
41. Liang T, Wen D, Chen G, et al. Adipocyte-derived anticancer lipid droplets. *Adv Mater.* 2021;33(26):2100629. doi:10.1002/adma.202100629

42. Yu Y, Cheng Q, Ji X, et al. Engineered drug-loaded cellular membrane nanovesicles for efficient treatment of postsurgical cancer recurrence and metastasis. *Sci Adv.* 2022;8(49):eadd3599. doi:10.1126/sciadv.add3599
43. Wang Y, Xie Y, Zhang A, Wang M, Fang Z, Zhang J. Exosomes: an emerging factor in atherosclerosis. *Biomed Pharmacother.* 2019;115:108951. doi:10.1016/j.biopha.2019.108951
44. Wang B, Zhuang X, Deng ZB, et al. Targeted drug delivery to intestinal macrophages by bioactive nanovesicles released from grapefruit. *Mol Ther.* 2014;22(3):522–534. doi:10.1038/mt.2013.190
45. Cong M, Tan S, Li S, et al. Technology insight: plant-derived vesicles-How far from the clinical biotherapeutics and therapeutic drug carriers? *Adv Drug Deliv Rev.* 2022;182:114108. doi:10.1016/j.addr.2021.114108
46. Stanly C, Moubarak M, Fiume I, Turiak L, Pocsfalvi G. Membrane transporters in citrus clementina fruit juice-derived nanovesicles. *Int J Mol Sci.* 2019;20(24):6205. doi:10.3390/ijms20246205
47. Baldini N, Torreggiani E, Roncuzzi L, Perut F, Zini N, Avnet S. Exosome-like nanovesicles isolated from citrus limon l. Exert antioxidative effect. *Curr Pharm Biotechnol.* 2018;19(11):877–885. doi:10.2174/1389201019666181017115755
48. Pocsfalvi G, Turiak L, Ambrosone A, et al. Protein biocargo of citrus fruit-derived vesicles reveals heterogeneous transport and extracellular vesicle populations. *J Plant Physiol.* 2018;229:111–121. doi:10.1016/j.jplph.2018.07.006
49. Chen X, Liu B, Li X, et al. Identification of anti-inflammatory vesicle-like nanoparticles in honey. *J Extracell Vesicles.* 2021;10(4):e12069. doi:10.1002/jev2.12069
50. Stanly C, Fiume I, Capasso G, Pocsfalvi G. Isolation of exosome-like vesicles from plants by ultracentrifugation on sucrose/deuterium oxide (D2O) density cushions. *Methods Mol Biol.* 2016;1459:259–269. doi:10.1007/978-1-4939-3804-9_18
51. Li P, Kaslan M, Lee SH, Yao J, Gao Z. Progress in Exosome Isolation Techniques. *Theranostics.* 2017;7(3):789–804. doi:10.7150/thno.18133
52. Chen X, Zhou Y, Yu J. Exosome-like nanoparticles from ginger rhizomes inhibited NLRP3 inflammasome activation. *Mol Pharm.* 2019;16(6):2690–2699. doi:10.1021/acs.molpharmaceut.9b00246
53. You JY, Kang SJ, Rhee WJ. Isolation of cabbage exosome-like nanovesicles and investigation of their biological activities in human cells. *Bio Mater.* 2021;6(12):4321–4332. doi:10.1016/j.bioactmat.2021.04.023
54. Woith E, Melzig MF. Extracellular vesicles from fresh and dried plants-simultaneous purification and visualization using gel electrophoresis. *Int J Mol Sci.* 2019;20(2):357. doi:10.3390/ijms20020357
55. Kalarikkal SP, Prasad D, Kasiappan R, Chaudhari SR, Sundaram GM. A cost-effective polyethylene glycol-based method for the isolation of functional edible nanoparticles from ginger rhizomes. *Sci Rep.* 2020;10(1). doi:10.1038/s41598-020-61358-8
56. Kang YT, Hadlock T, Lo TW, et al. Dual-isolation and profiling of circulating tumor cells and cancer exosomes from blood samples with melanoma using immunoaffinity-based microfluidic interfaces. *Advan Sci.* 2020;7(19):2001581. doi:10.1002/advs.202001581
57. Yang M, Liu X, Luo Q, Xu L, Chen F. An efficient method to isolate lemon derived extracellular vesicles for gastric cancer therapy. *J Nanobiotechnology.* 2020;18(1):100. doi:10.1186/s12951-020-00656-9
58. Moura RP, Pacheco C, Pêgo AP, Des Rieux A, Sarmento B. Lipid nanocapsules to enhance drug bioavailability to the central nervous system. *J Control Release.* 2020;322:390–400. doi:10.1016/j.jconrel.2020.03.042
59. Raimondo S, Urzi O, Meraviglia S, et al. Anti-inflammatory properties of lemon-derived extracellular vesicles are achieved through the inhibition of ERK /NF- κ B signalling pathways. *J Cell Mol Med.* 2022;26:4195–4209. doi:10.1111/jcmm.17404
60. Sundaram K, Miller DP, Kumar A, et al. Plant-derived exosomal nanoparticles inhibit pathogenicity of porphyromonas gingivalis. *iScience.* 2019;21:308–327. doi:10.1016/j.isci.2019.10.032
61. Perut F, Roncuzzi L, Avnet S, et al. Strawberry-derived exosome-like nanoparticles prevent oxidative stress in human mesenchymal stromal cells. *Biomolecules.* 2021;11(1):87. doi:10.3390/biom11010087
62. Peršurić Ž, Pavelić SK. Bioactives from bee products and accompanying extracellular vesicles as novel bioactive components for wound healing. *Molecules.* 2021;26(12):3770. doi:10.3390/molecules26123770
63. Lee R, Ko HJ, Kim K, et al. Anti-melanogenic effects of extracellular vesicles derived from plant leaves and stems in mouse melanoma cells and human healthy skin. *J Extracell Vesicles.* 2019;9(1). doi:10.1080/20013078.2019.1703480
64. Raimondo S, Naselli F, Fontana S, et al. Citrus limon-derived nanovesicles inhibit cancer cell proliferation and suppress CML xenograft growth by inducing TRAIL-mediated cell death. *Oncotarget.* 2015;6(23):19514–19527. doi:10.18632/oncotarget.4004
65. Deng Z, Rong Y, Teng Y, et al. Broccoli-derived nanoparticle inhibits mouse colitis by activating dendritic cell AMP-activated protein kinase. *Mol Ther.* 2017;25(7):1641–1654. doi:10.1016/j.ymthe.2017.01.025
66. Zhuang X, Deng ZB, Mu J, et al. Ginger-derived nanoparticles protect against alcohol-induced liver damage. *J Extracell Vesicles.* 2015;4:28713. doi:10.3402/jev.v4.28713
67. Kalarikkal SP, Sundaram GM. Edible plant-derived exosomal microRNAs: exploiting a cross-kingdom regulatory mechanism for targeting SARS-CoV-2. *Toxicol Appl Pharmacol.* 2021;414:115425. doi:10.1016/j.taap.2021.115425
68. Teng Y, Ren Y, Sayed M, et al. Plant-derived exosomal MicroRNAs shape the gut microbiota. *Cell Host Microbe.* 2018;24(5):637–652.e8. doi:10.1016/j.chom.2018.10.001
69. Raza K, Nemidkanam V, Chaichanawongsaraj N, Raza K, Khan YA. Characterizing Kaempferia parviflora extracellular vesicles, a nanomedicine candidate. *PLoS One.* 2022;17(1). doi:10.1371/journal.pone.0262884
70. Kim K, Park J, Sohn Y, et al. Stability of plant leaf-derived extracellular vesicles according to preservative and storage temperature. *Pharmaceutics.* 2022;14(2). doi:10.3390/pharmaceutics14020457
71. Vader P, Mol EA, Pasterkamp G, Schiffelers RM. Extracellular vesicles for drug delivery. *Adv Drug Deliv Rev.* 2016;106(Pt A):148–156. doi:10.1016/j.addr.2016.02.006
72. Mahdipour E. Beta vulgaris juice contains biologically active exosome-like nanoparticles. *Tissue Cell.* 2022;76:101800. doi:10.1016/j.tice.2022.101800
73. Zhang L, He F, Gao L, et al. Engineering exosome-like nanovesicles derived from asparagus cochinchinensis can inhibit the proliferation of hepatocellular carcinoma cells with better safety profile. *Int J Nanomedicine.* 2021;16:1575–1586. doi:10.2147/IJN.S293067
74. Cao M, Yan H, Han X, et al. Ginseng-derived nanoparticles alter macrophage polarization to inhibit melanoma growth. *J Immunother Cancer.* 2019;7(1):326. doi:10.1186/s40425-019-0817-4

75. Kim J, Li S, Zhang S, Wang J. Plant-derived exosome-like nanoparticles and their therapeutic activities. *Asian J Pharm Sci.* 2022;17(1):53–69. doi:10.1016/j.ajps.2021.05.006
76. Trentini M, Zanotti F, Tiengo E, et al. An apple a day keeps the doctor away: potential role of miRNA 146 on macrophages treated with exosomes derived from apples. *Biomedicines.* 2022;10(2):415. doi:10.3390/biomedicines10020415
77. Rutter BD, Innes RW. Extracellular vesicles isolated from the leaf apoplast carry stress-response proteins. *Plant Physiol.* 2017;173(1):728–741. doi:10.1104/pp.16.01253
78. Logozzi M, Di Raimo R, Mizzone D, Fais S. Nanovesicles from organic agriculture-derived fruits and vegetables: characterization and functional antioxidant content. *Int J Mol Sci.* 2021;22(15):8170. doi:10.3390/ijms22158170
79. De Robertis M, Sarra A, V D, et al. Blueberry-derived exosome-like nanoparticles counter the response to TNF- α -induced change on gene expression in EA.hy926 cells. *Biomolecules.* 2020;10(5):742. doi:10.3390/biom10050742
80. Xiao J, Feng S, Wang X, et al. Identification of exosome-like nanoparticle-derived microRNAs from 11 edible fruits and vegetables. *PeerJ.* 2018;6:e5186. doi:10.7717/peerj.5186
81. Kim DK, Rhee WJ. Antioxidative effects of carrot-derived nanovesicles in cardiomyoblast and neuroblastoma cells. *Pharmaceutics.* 2021;13(8):1203. doi:10.3390/pharmaceutics13081203
82. Zhao Z, Yu S, Li M, Gui X, Li P. Isolation of exosome-like nanoparticles and analysis of MicroRNAs derived from coconut water based on small RNA high-throughput sequencing. *J Agric Food Chem.* 2018;66(11):2749–2757. doi:10.1021/acs.jafc.7b05614
83. Abraham AM, Wiemann S, Ambreen G, et al. Cucumber-derived exosome-like vesicles and plant crystals for improved dermal drug delivery. *Pharmaceutics.* 2022;14(3):476. doi:10.3390/pharmaceutics14030476
84. Liu B, Li X, Yu H, et al. Therapeutic potential of garlic chive-derived vesicle-like nanoparticles in NLRP3 inflammasome-mediated inflammatory diseases. *Theranostics.* 2021;11(19):9311–9330. doi:10.7150/thno.60265
85. Li Z, Wang H, Yin H, Bennett C, Zhang HG, Guo P. Arrowtail RNA for ligand display on ginger exosome-like nanovesicles to systemic deliver siRNA for cancer suppression. *Sci Rep.* 2018;8(1):14644. doi:10.1038/s41598-018-32953-7
86. Zhang M, Collins JF, Merlin D. Do ginger-derived nanoparticles represent an attractive treatment strategy for inflammatory bowel diseases? *Nanomedicine.* 2016;11(23):3035–3037. doi:10.2217/nmm-2016-0353
87. Zhang M, Viennois E, Prasad M, et al. Edible ginger-derived nanoparticles: a novel therapeutic approach for the prevention and treatment of inflammatory bowel disease and colitis-associated cancer. *Biomaterials.* 2016;101:321–340. doi:10.1016/j.biomaterials.2016.06.018
88. Zhang M, Wang X, Han MK, Collins JF, Merlin D. Oral administration of ginger-derived nanolipids loaded with siRNA as a novel approach for efficient siRNA drug delivery to treat ulcerative colitis. *Nanomedicine.* 2017;12(16):1927–1943. doi:10.2217/nmm-2017-0196
89. Cho EG, Choi SY, Kim H, et al. Panax ginseng-derived extracellular vesicles facilitate anti-senescence effects in human skin cells: an eco-friendly and sustainable way to use ginseng substances. *Cells.* 2021;10(3):486. doi:10.3390/cells10030486
90. Pérez-Bermúdez P, Blesa J, Soriano JM, Marcilla A. Extracellular vesicles in food: experimental evidence of their secretion in grape fruits. *Eur J Pharm Sci.* 2017;98:40–50. doi:10.1016/j.ejps.2016.09.022
91. Teng Y, Mu J, Hu X, et al. Grapefruit-derived nanovectors deliver miR-18a for treatment of liver metastasis of colon cancer by induction of M1 macrophages. *Oncotarget.* 2016;7(18):25683–25697. doi:10.18632/oncotarget.8361
92. Zhuang X, Teng Y, Samykutty A, et al. Grapefruit-derived nanovectors delivering therapeutic miR17 through an intranasal route inhibit brain tumor progression. *Mol Ther.* 2016;24(1):96–105. doi:10.1038/mt.2015.188
93. Wang Q, Ren Y, Mu J, et al. Grapefruit-derived nanovectors use an activated leukocyte trafficking pathway to deliver therapeutic agents to inflammatory tumor sites. *Cancer Res.* 2015;75(12):2520–2529. doi:10.1158/0008-5472.CAN-14-3095
94. Lei C, Teng Y, He L, et al. Lemon exosome-like nanoparticles enhance stress survival of gut bacteria by RNase P-mediated specific tRNA decay. *iScience.* 2021;24(6):102511. doi:10.1016/j.isci.2021.102511
95. Aquilano K, Ceci V, Gismondi A, et al. Adipocyte metabolism is improved by TNF receptor-targeting small RNAs identified from dried nuts. *Communic Bio.* 2019;2:317. doi:10.1038/s42003-019-0563-7
96. Regente M, Corti-Monzón G, Maldonado AM, Pinedo M, Jorrín J, de la Canal L. Vesicular fractions of sunflower apoplast fluids are associated with potential exosome marker proteins. *FEBS Lett.* 2009;583(20):3363–3366. doi:10.1016/j.febslet.2009.09.041
97. Johnstone RW, Frew AJ, Smyth MJ. The TRAIL apoptotic pathway in cancer onset, progression and therapy. *Nat Rev Cancer.* 2008;8(10):782–798. doi:10.1038/nrc2465
98. Chen Q, Li Q, Liang Y, et al. Natural exosome-like nanovesicles from edible tea flowers suppress metastatic breast cancer via ROS generation and microbiota modulation. *Acta Pharmaceutica Sinica B.* 2022;12(2):907–923. doi:10.1016/j.actph.2021.08.016
99. Fang Z, Song M, Lai K, Cui M, Yin M, Liu K. Kiwi-derived extracellular vesicles for oral delivery of sorafenib. *Eur J Pharm Sci.* 2023;191:106604. doi:10.1016/j.ejps.2023.106604
100. Langen UH, Ayloo S, Gu C. Development and cell biology of the blood-brain barrier. *Annu Rev Cell Dev Biol.* 2019;35:591–613. doi:10.1146/annurev-cellbio-100617-062608
101. Ishida T, Kawada K, Jobu K, et al. Exosome-like nanoparticles derived from *Allium tuberosum* prevent neuroinflammation in microglia-like cells. *J Pharm Pharmacol.* 2023;75(10):1322–1331. doi:10.1093/jpp/rgad062
102. Zhang M, Viennois E, Xu C, Merlin D. Plant derived edible nanoparticles as a new therapeutic approach against diseases. *Tissue Barriers.* 2016;4(2):e1134415. doi:10.1080/21688370.2015.1134415
103. Yin L, Yan L, Yu Q, et al. Characterization of the MicroRNA profile of ginger exosome-like nanoparticles and their anti-inflammatory effects in intestinal caco-2 cells. *J Agric Food Chem.* 2022;70(15):4725–4734. doi:10.1021/acs.jafc.1c07306
104. Sriwastva MK, Deng ZB, Wang B, et al. Exosome-like nanoparticles from Mulberry bark prevent DSS-induced colitis via the AhR/COP8 pathway. *EMBO Rep.* 2022;23(3):e53365. doi:10.15252/embr.202153365
105. Kim K, Yoo HJ, Jung JH, et al. Cytotoxic effects of plant sap-derived extracellular vesicles on various tumor cell types. *J Funct Biomater.* 2020;11(2). doi:10.3390/jfb11020022
106. Potestà M, Minutolo A, Gismondi A, et al. Cytotoxic and apoptotic effects of different extracts of *Moringa oleifera* Lam on lymphoid and monocytoid cells. *Exp Ther Med.* 2019;18(1):5–17. doi:10.3892/etm.2019.7544
107. Tajik T, Baghaei K, Moghadam VE, Farrokhi N, Salami SA. Extracellular vesicles of cannabis with high CBD content induce anticancer signaling in human hepatocellular carcinoma. *Biomed Pharmacother.* 2022;152. doi:10.1016/j.biopha.2022.113209

108. Han X, Wei Q, Lv Y, et al. Ginseng-derived nanoparticles potentiate immune checkpoint antibody efficacy by reprogramming the cold tumor microenvironment. *Mol Ther.* 2022;30(1):327–340. doi:10.1016/j.ymthe.2021.08.028
109. Rahimi Ghiasi M, Rahimi E, Amirkhani Z, Salehi R. Leucine-rich repeat-containing G-protein coupled receptor 5 gene overexpression of the rat small intestinal progenitor cells in response to orally administered grape exosome-like nanovesicles. *Adv Biomed Res.* 2018;7:125. doi:10.4103/abr.abr_114_18
110. Stanly C, Alfieri M, Ambrosone A, Leone A, Fiume I, Pocsfalvi G. Grapefruit-derived micro and nanovesicles show distinct metabolome profiles and anticancer activities in the A375 human melanoma cell line. *Cells.* 2020;9(12):2722. doi:10.3390/cells9122722
111. Tang Z, Jun Y, Lv Y, et al. Aptamer-conjugated and doxorubicin-loaded grapefruit-derived nanovectors for targeted therapy against HER2(+) breast cancer. *J Drug Target.* 2020;28(2):186–194. doi:10.1080/1061186x.2019.1624970
112. Tamura RE, de Vasconcellos JF, Sarkar D, Libermann TA, Fisher PB, Zerbini LF. GADD45 proteins: central players in tumorigenesis. *Curr Mol Med.* 2012;12(5):634–651. doi:10.2174/156652412800619978
113. Nesterov A, Nikrad M, Johnson T, Kraft AS. Oncogenic Ras sensitizes normal human cells to tumor necrosis factor-alpha-related apoptosis-inducing ligand-induced apoptosis. *Cancer Res.* 2004;64(11):3922–3927. doi:10.1158/0008-5472.Can-03-2219
114. Pan Y, Yu Y, Wang X, Zhang T. Tumor-associated macrophages in tumor immunity. *Front Immunol.* 2020;11:583084. doi:10.3389/fimmu.2020.583084
115. Wu K, Lin K, Li X, et al. Redefining tumor-associated macrophage subpopulations and functions in the tumor microenvironment. *Front Immunol.* 2020;11:1731. doi:10.3389/fimmu.2020.01731
116. Cui -W-W, Ye C, Wang K-X, et al. Momordica charantia-derived extracellular vesicles-like nanovesicles protect cardiomyocytes against radiation injury via attenuating DNA damage and mitochondria dysfunction. *Front Cardiovasc Med.* 2022;9. doi:10.3389/fcvm.2022.864188
117. Zu M, Xie D, Canup BSB, et al. ‘Green’ nanotherapeutics from tea leaves for orally targeted prevention and alleviation of colon diseases. *Biomaterials.* 2021;279:121178. doi:10.1016/j.biomaterials.2021.121178
118. Srivastava A, Rathore S, Munshi A, Ramesh R. Organically derived exosomes as carriers of anticancer drugs and imaging agents for cancer treatment. *Semin Cancer Biol.* 2022;86:80–100. doi:10.1016/j.semcancer.2022.02.020
119. Asghar S, Khan IU, Salman S, Khalid SH, Ashfaq R, Vandamme TF. Plant-derived nanotherapeutic systems to counter the overgrowing threat of resistant microbes and biofilms. *Adv Drug Deliv Rev.* 2021;179. doi:10.1016/j.addr.2021.114019
120. Dad HA, Gu TW, Zhu AQ, Huang LQ, Peng LH. Plant exosome-like nanovesicles: emerging therapeutics and drug delivery nanoplatforms. *Mol Ther.* 2021;29(1):13–31. doi:10.1016/j.ymthe.2020.11.030
121. Woith E, Guerriero G, Hausman JF, et al. Plant extracellular vesicles and nanovesicles: focus on secondary metabolites, proteins and lipids with perspectives on their potential and sources. *Int J Mol Sci.* 2021;22(7):3719. doi:10.3390/ijms22073719
122. Stanly C, Kim H, Antonucci G, et al. Crosstalk between the immune system and plant-derived nanovesicles: a study of allergen transporting. *Front Bioeng Biotechnol.* 2021;9. doi:10.3389/fbioe.2021.760730
123. Man F, Meng C, Liu Y, et al. The study of ginger-derived extracellular vesicles as a natural nanoscale drug carrier and their intestinal absorption in rats. *AAPS Pharm Sci Tech.* 2021;22(6). doi:10.1208/s12249-021-02087-7