Open Access Full Text Article

ORIGINAL RESEARCH

Protein kinase $C\alpha$ downregulation via siRNA-PKC α released from foldable capsular vitreous body in cultured human retinal pigment epithelium cells

Xiaoqing Chen Yaqin Liu Zhaoxin Jiang Lian Zhou Jian Ge Qianying Gao

State Key Laboratory of Ophthalmology, Zhongshan Ophthalmic Center, Sun Yat-Sen University, Guangzhou, People's Republic of China

Correspondence: Qianying Gao Zhongshan Ophthalmic Center, State Key Lab of Ophthalmology, 54 Xianlie Road, Guangzhou, Guangdong 510060, People's Republic of China Tel +86 20 8733 0490 Fax +86 20 8733 1350 Email gaoqy@mail.sysu.edu.cn **Abstract:** We previously found that downregulation of protein kinase C α (PKC α) can inhibit retinal pigment epithelium (RPE) cell proliferation involved in the development of proliferative vitreoretinopathy (PVR). In this study, we tested whether PKC α could be downregulated via small interfering RNA (siRNA)-PKC α released from foldable capsular vitreous body (FCVB) in cultured human RPE cells. SiRNA-PKC α content, determined by ultraviolet (UV) spectrophotometer, was released from FCVB containing 200, 300, 400, 500, and 600 nm siRNA-PKC α in a time-dependent manner from 1 to 96 hours and a dose-dependent manner at five concentrations. The content (y) had a good linear relationship with time (x), especially in the 600 nm siRNA-PKC α group (y = 16.214x, R^2 = 0.9809). After treatment with siRNA-PKC α released from FCVBs, the PKC α was significantly decreased by RT-PCR, Western blot, and immunofluorescence analysis in RPE cells. These results indicate that PKC α was significantly downregulated by siRNA-PKC α released from FCVB in human RPE cells and provide us with a new avenue to prevent PVR.

Keywords: small interfering RNA–protein kinase C α , foldable capsular vitreous body, drug delivery system, retinal pigment epithelium, protein kinase C α

Introduction

Proliferative vitreoretinopathy (PVR) is a severe complication of retinal detachment and the most common reason for the failure of retinal detachment surgery.^{1,2} The risk of PVR varies from 10% to 40% depending on the original diseases.^{3,4} The pathology of PVR involves a complex network of relationships of cellular and humoral factors, in which proliferating retinal pigment epithelium (RPE) cells are thought to be an initiating factor.⁵⁻⁸ Protein kinase C (PKC) is an intracellular enzyme family, which transduces extracellular signals into intracellular responses. It has been well documented that the PKC family is involved in a series of life phenomena such as cell communication, secretion, ion channel regulation, cell proliferation and differentiation, and cancer, implying it is an ideal target for blocking signal transduction.^{9–12} Our previous study found that, of the ten kinds of PKC isoforms expressed in RPE cells, only PKCα is downstream of the confluence of the complex PVR network, which can promote RPE cell proliferation by downregulation of p27^{Kip1}.^{13,14} All of these studies strongly suggest that PKCα is an ideal and special blocking target for PVR.

Drug delivery to the posterior segments of the eye presents considerable challenges, because drugs administered by the conventional routes of topical or systemic dosing are usually ineffective due to the existence of critical barriers (such as corneal, blood–retinal, and blood–aqueous barriers) and short drug half-lives.^{15,16} Considering these challenges, development of a drug delivery system (DDS) is becoming increasingly important in the treatment of vitreoretinal diseases not only to facilitate drug efficacy but also to attenuate adverse effects.¹⁷

In our previous studies, we discovered that foldable capsular vitreous body (FCVB) is a public release vehicle DDS and can sustainably and mechanically release dexamethasone sodium phosphate (DexP) via the capsule apertures.¹⁸ In the clinical trial (ClinicalTrials.gov number, NCT00910702) evaluating the flexibility, efficacy, and safety of an FCVB in the treatment of severe retinal detachment, the FCVB was shown to be flexible, effective, and safe as a vitreous substitute over a 3-month implantation time. The FCVB used in the clinical trial consists of a thin vitreous-shaped capsule with a tube-valve system made using computer and industrial technology.¹⁹⁻²¹

RNA interference (RNAi) is an ancient and common biological phenomenon discovered in vivo in recent years.²² It is commonly used as a powerful tool in biological and biomedical research and has been used experimentally to prevent ocular neovascularization and inflammation.^{23–25}

In our previous research, small interfering RNA (siRNA)-PKC α), a specific inhibitor of PKC α , was shown to be able to inhibit RPE cell proliferation.¹⁴ Therefore, in this study, we investigate whether PKC α could be downregulated via siRNA-PKC α released from FCVB in cultured human RPE cells, providing us with a new avenue to prevent PVR.

Materials and methods Characteristics of siRNA-PKCα

SiRNA-PKC α was purchased from Ribuo Biotech (Guangzhou, China). The sequence of siRNA-PKC α was: positive-sense strand 5'GGCGUCCUGUUGUAUGAAAdAdT3', antisense strand 3'dTdACCGCAGGACAACAUACUUU5'. Chemically synthesized siRNA-PKC α was mixed the suspension of two deoxyribose nucleotides to increase the stability in its 3' end. Lyophilized powder of siRNA-PKC α was dissolved in TE buffer (110 mM Tris·HCl, 1 mM EDTA, pH = 8.00) into the stock concentration of 20 μ M, and stored at -20°C.

Basic material and fabrication of the FCVB

FCVB is made of tailor-made modified liquid silicone rubber, which consists of a vitreous-like capsule and a tube-valve system (shown in Figure 1A). The basic material, Dow



Figure I Small interfering RNA–protein kinase C α (SiRNA-PKC α) sustained-release studies in vitro. (A) Foldable capsular vitreous body (FCVB) consists of a thin (60 μ m) vitreous-like capsule mimicked by computer with a tube-valve system. (B) SiRNA-PKC α at concentrations of 200, 300, 400, 500, and 600 nM in TE buffer was injected into the capsules of FCVB, which were immersed in cups of modified Frans diffusion cells. Two hundred (200) μ L liquid was aspirated at time intervals of 1, 2, 3, 6, 9, 12, 24, 48, 72, and 96 hours.

Corning Class VI elastomer, was purchased from Dow Corning Company (Midland, MI). The details of FCVB were described in our previous study.¹⁹

In vitro drug sustained-release studies

In a typical experiment, siRNA-PKC α at concentrations of 200, 300, 400, 500, and 600 nm in TE buffer was injected into the capsules of the FCVBs, and the FCVBs were separately immersed in five cups of modified Franz diffusion cells (shown in Figure 1B) and 4 mL of TE buffer as release medium was transferred into each cup. These cups were kept in an orbital shaker maintained at 37°C with constant agitation. Two hundred microliters (200 µL) of the release medium was aspirated for measurement at 1, 2, 3, 6, 9, 12, 24, 48, 72, and 96 hours, and then the same amount of fresh medium was added into the cups. The collected samples were stored in a refrigerator. The siRNA-PKC α content in the liquid was then detected by UV spectrophotometer (DU800; Beckman Coulter, Brea, CA).

SiRNA-PKC α sample assay in vitro

First, a standard curve was made to demonstrate the effectiveness of the assay method. In brief, the 20 μ M siRNA-PKC α stock solution was serially diluted with TE buffer to the desired concentrations: 20, 40, 60, 80, 100, 200, 300, 400, 500, and 600 nm. Samples of 100 μ L were transferred into the cuvette chamber. Samples were assayed at the absorbance of 260 nm, and 100 μ L of TE buffer was added to blank the background. Calibrations were made using Excel software (2003; Microsoft, Redmond, WA).

Micromorphology of the FCVB

The capsule of the FCVB was cut into a 1-cm diameter disc, and then the sample was cleaned, coated with gold, and fixed on a specimen stub. The image of the specimen surface was captured with a scanning electron microscope (XL 30 ESEM FEG; FEI-Philips, Eindhoven, the Netherlands). Examinations were carried out at 10 kV and in a humid atmosphere of approximately 1.8–2.0 Torr. Individual parameters are visible in the scale bar of each image.

Human RPE cell culture

Human RPE cells were isolated within 24 hours of death from donor eyes, which were obtained from the Eye Bank of the Zhongshan Ophthalmic Center as previously described.²⁶ The project was approved by the Ethics Committee of the Zhongshan Ophthalmic Center and followed the tenets of the Declaration of Helsinki. RPE cells were cultured in Dulbecco's modified Eagle's medium (DMEM; Invitrogen-Gibco, Karlsruhe, Germany) containing 10% fetal bovine serum, penicillin G (100 μ g/mL), streptomycin sulfate (100 mg/mL), and L-glutamate in Corning culture plates. The cells were maintained at 37°C in 5% CO₂ in a humidified atmosphere, and those between the third and sixth passages were used for all experiments.

Treatment of human RPE cells with siRNA-PKC α released from FCVB

Human RPE cells grown in transwells (Corning Inc, Corning, NY) were incubated for 12 hours. The culture media was then replaced with DMEM without serum or antibiotics, and filter inserts (8 µm pore size) carrying FCVBs containing 400, 500, and 600 nm siRNA-PKCa were transferred to transwells. Lipofectamine 2000 (5 μ L) (Invitrogen, Karlsruhe, Germany) was added to each well at time points of 48, 24, and 12 hours for each group, and the medium in each well was replaced with DMEM with 10% fetal bovine serum 6 hours later. Eighteen hours later, cells were harvested for Western blot and reverse transcription-polymerase chain reaction (RT-PCR). For immunofluorescence analysis, cells which were grown on cover lips in transwells were subjected to the same treatments as Western blot and RT-PCR. Cells treated by DMEM with 10% fetal bovine serum and transfected with 100 nm siRNA for 24 hours were used for blank control and positive control, respectively.

RT-PCR

Total RNA was extracted using Trizol reagent (Life Technologies, Gaithersburg, MD). The integrity of the RNA was checked by 2% agarose gel electrophoresis. Approximately 2 µg RNA was reverse-transcribed following the protocol of the Super Script[™] first-strand synthesis system. CDNAs encoding PKCa and GAPDH genes were amplified by PCR as follows: denaturation at 94°C for 30 seconds, annealing at 63°C for 30 seconds and elongation at 72°C for 45 seconds. Primer sequence was designed using Primer 3. For PKCa, the forward primer was 5'-ATCCGCAGTGGAATGAGTC CTTTACAT-3', and the reverse primer was 5'-TTGGAAG GTTGTTTCCTGTCTTCAGAG-3'. A housekeeping gene, glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was used as the internal control. The forward primer was 5'-ACCCAGAAGACTGTGGATGG-3', and the reverse primer was 5'-TGCTGTAGCCAAATTCGTTG-3'. PCR products were analyzed by agarose (2%) gel electrophoresis. Quantifications of signal intensity were confirmed using a specific computer program (Image J 1.43U software, Wayne Rasband, National Institutes of Health, USA).

Preparation of cell extracts

The medium was removed and the plates were washed twice with ice-cold phosphate buffered saline (PBS). The human RPE cells were lysed with sample buffer that contained 60 mM Tris, pH 6.8, 2% (w/v) sodium dodecyl sulfate (SDS), 100 mM 2-mercaptoethanol, and 0.01% (w/v) bromophenol blue.²⁷ The lysate was then incubated on ice for 30 minutes. The lysate was scraped using a cell scraper and harvested using a pipettor, then centrifuged at 4°C for 30 minutes. The supernatant was collected and boiled for 5 minutes and stored at -20° C.

Western blot analysis

Cellular extracts from human RPE cells were processed for Western blot analysis.28 Briefly, 60 µg of protein per well was loaded on a 10% SDS-polyacrylamide gel electrophoresis (PAGE) gel. The protein was electro-transferred to polyvinylidene difluoride membranes (Millipore, Billerica, MA) for 1 hour at 350 mA, then blocked with Tris-buffered saline (TBS) containing 5% nonfat milk and 0.1% Tween-20 (TBST) for 1 hour and incubated with mouse anti-PKCa (BD Systems, Torrance, CA) overnight. After three washes with TBST, the membranes were incubated with horseradish peroxidase conjugated secondary antibody for 1 hour at room temperature and washed with TBST. Localization of the antibody was detected by chemiluminescence using the ECL kit (Cell Signaling, Danvers, MA) following the manufacturer's instructions. β-actin (Boster Biologic Technology Ltd, Wuhan, China) was used as an internal control.

Immunofluorescence analysis

After treatment, cells were fixed for 15 minutes in 4% paraformaldehyde, and then rinsed three times in PBS. All reagent incubations were performed in a humidified chamber. The mouse anti-PKC α was incubated for 16 hours in PBS at room temperature; horse serum was used as the negative control instead of the primary antibody. After washing four times for 10 minutes in PBS, fluorescein isothiocyanate-labeled secondary antibodies were incubated for 10 minutes at 37°C. Then Hoechst 33342 was incubated for 10 minutes at room temperature. After three rinses in PBS, cover slips were mounted onto glass slides and analyzed on a Zeiss laser scanning confocal microscope (LSCM 510 META; Oberkochen, Germany).

Statistical analysis

All data were expressed as mean \pm standard deviation (SD). All experiments were repeated at least 4 times, and similar results were obtained. One-way analysis of variance (ANOVA), the test of homogeneity of variances, and a post hoc test (Bonferroni test) were used to determine significant differences between the treated groups and the control groups. P < 0.05 was considered statistically significant.

Results Selectivity and acceptability of the standard curve

A representative result of the standard curve is shown in Figure 2A. The standard curve showed a very good linear relationship (y = 2976.3x, $R^2 = 1$), demonstrating that the method selected had good selectivity and acceptability.

In vitro release of siRNA-PKC α

The cumulative release of siRNA-PKC α from FCVB is shown in Figure 2B–2G. SiRNA-PKC α released from the capsules of FCVB increased with time in a 96-hour experimental time period. The content (y) had a good linear relationship with time (x), especially in the 600 nm group, indicating that the FCVB released siRNA-PKC α stably with time (y = 16.214x, $R^2 = 0.9809$) (P < 0.05).

The contents released from FCVB increased accordingly, with siRNA-PKC α concentration rising in the FCVB. The released content was much greater in the 600 nm group at the same release time than in the other groups. All data indicated that siRNA-PKC α was released from the capsules of FCVB in a dose-dependent manner in vitro in 5 concentrations during a 96-hour experiment time period.

Scanning electron microscopy image of the FCVB

Before the experiment, scattered mini apertures of 300 nm could be seen in the capsule of FCVB by scanning electron microscopy. Figure 3A shows 2 of the typical apertures. After a 96-hour experimental time period, 300-nm mini apertures still existed in the capsules, as shown in Figure 3B. No great changes were observed in the capsule apertures by scanning electron microscopy.

PKC α downregulation after treatment by siRNA-PKC α released from FCVB

As shown in Figure 4A1 and 4A2, RT-PCR results showed that PKC α messenger RNA (mRNA) was significantly



Figure 2 Standard curve and in vitro release profile in the cumulative amount of five different concentrations. (**A**) The standard curve had a very good linear relationship. With the absorbance (x) increase, the small interfering RNA-protein kinase $C\alpha$ (SiRNA-PKC α) concentration (y) rose accordingly (y = 2976.3x, $R^2 = 1$). **B–G**) Time and dose-dependent release. The siRNA-PKC α released from the foldable capsular vitreous body (FCVB) increased in a linear manner with time. Of the five different concentrations, the 600 nM group had a best linear relationship (y = 16.214x, $R^2 = 0.9809$). When siRNA-PKC α concentration rose inside the FCVBs, the content released from the capsules rose accordingly. The tendency began from 1 hour, and lasted until 96 hours.

Notes: Results shown were representative of four repeated experiments. The concentrations of different time points are statistically different (P < 0.05). The data are expressed as the mean \pm standard deviation (SD).

downregulated following 24-hour release siRNA-PKC α treatments in the 400 nm, 500 nm, and 600 nm release groups. Consistent with a change at the mRNA level, the PKC α protein after treatment decreased 45.1% ± 6.73%, 50.8% ± 4.96%, and 54.3% ± 3.62% in the 400 nm, 500 nm, and 600 nm release groups, respectively, and 64.0% ± 4.5%



Figure 3 Scanning electron microscope images of the capsule of the foldable capsular vitreous body. Mini apertures of 300 nm in the capsules were observed before the release (\mathbf{A}) and at the end of the experiment time (\mathbf{B}) (arrows).

in the positive control of 100 nm siRNA-PKC α treated group compared with that in the untreated control group (Figure 4B1 and 4B2). In the untreated RPE cells, both the mRNA and protein level remained constant. These data indicate that siRNA-PKC α can actually be released from the FCVBs and further decrease expression of PKC α in cultured human RPE cells.

We further examined the expression changes of PKC α using immunocytochemistry. The cultured human RPE cells formed a monolayer with typical polygonal cellular arrays, as shown in Figure 5. Compatible with findings from RT-PCR and Western blot analysis, staining of PKC α decreased significantly after exposure to both 100 nm siRNA-PKC α and siRNA-PKC α released from FCVBs. Compared with the positive control of 100 nm siRNA-PKC α treated group, the stain did not change greatly in the experimental groups.



Figure 4 Expression of protein kinase C α (PKC α) messenger RNA (mRNA) and protein downregulation by released small interfering RNA (siRNA)-PKC α in retinal pigment epithelium (RPE) cells. **A1**, **A2**) RT-PCR showed that released siRNA-PKC α reduced the mRNA levels of PKC α significantly compared with control groups. *P < 0.05 vs the corresponding value for control groups. Data are the mean ± standard deviation of quadruplicates from an experiment that was repeated with similar results (siRNA100 means RPE cells treated with 100 nM siRNA-PKC α ; R400, R500, and R600 mean RPE cells treated with siRNA-PKC α released from foldable capsular vitreous bodies containing 400, 500, and 600 nM siRNA-PKC α). **B1**, **B2**) Western blot showed that released siRNA-PKC α reduced the protein levels of PKC α significantly compared with control groups.

Discussion

In this study, we determined the siRNA-PKC α release of a nonbiodegradable device (FCVB) and its effect on PKC α in cultured human RPE cells. Our present study indicates that FCVB can mechanically and sustainably release siRNA-PKC α in a time-dependent and dose-dependent

manner during a 96-hour experimental time period. Furthermore, PKC α was decreased in RPE cells treated with siRNA-PKC α released from FCVBs. However, compared with the positive control group, expression of the target protein did not decrease remarkably with our experimental systems even in the presence of transfection reagents.



Figure 5 Confocal images of protein kinase C α (PKC α) expression in retinal pigment epithelium (RPE) cells by different treatments. PKC α staining was much lighter in the small interfering RNA (siRNA)-PKC α treatment than in the control, with no significant difference in the released siRNA-PKC α treatment groups. (**A**) blank control. (**B**) RPE cells treated with 100 nM siRNA-PKC α . (**C**–**E**) RPE cells treated with siRNA-PKC α released from foldable capsular vitreous bodies containing 400, 500, and 600 nM siRNA-PKC α , respectively. (1) PKC α staining in the cytoplasm and nuclei of RPE cells. (2) Staining of the nuclei with Hoechst 33342. (3) An overlay of images I and 2. Scale bars, 20 µm. Magnification × 1000.

This is likely attributable to the fact that our FCVB drug delivery system released siRNA-PKC α progressively and to the fact that siRNA-PKC α of the positive control group reached peak concentration quickly. The present results were consistent with a previous study that FCVB can sustainably, mechanically release DexP via the apertures of the capsule in a time-dependent and dose-dependent manner.¹⁸

In this study, siRNA-PKC α was shown to be fairly permeable through the capsule apertures of FCVB and to downregulate PKC α in RPE cells. We speculated that there were two main factors involved in the mechanism. The first is the concentration gradient of siRNA-PKC α . With the concentration gradient increase, siRNA-PKC α released from the FCVBs subsequently increased, as shown in

Dovepress

Figure 2B-2G. Second, the diffusion coefficient was related to the concentration gradient. The two factors mentioned previously have been reported by Acharya and colleagues in a homogeneous poly(lactic-co-glycolic acid) microstructures drug release system.²⁹

The current intraocular DDS confronts some major disadvantages. For example, they are opaque, may interfere with vision, and result in a nonuniform drug distribution.³⁰ In contrast to the current intraocular DDS, FCVB has good transparency and induces very little refractive shift,³¹ permitting the siRNA-PKC α in solution to make direct contact with the retina and achieve targeted drug delivery. Therefore, the FCVB can sustainably and mechanically release drugs without changing their chemical properties, and may provide a common vehicle for releasing different drugs.

The ideal intravitreal DDS is a kind of material that is not only an intravitreal drug delivery system but also a vitreous tamponade agent.³⁰ Previous studies demonstrated that FCVB was a fine vitreous substitute that closely mimicked vitreous morphology and restored its physiological functions, such as support and cellular barriers, during a 3-month implantation period without the obvious complications commonly induced by silicone oil.³² In our study, siRNA-PKCα was released from FCVB in a time-dependent and a dose-dependent manner in vitro. SiRNA-PKCα was detected until 96 hours, indicating that FCVB can be used as a DDS in addition to serving as a vitreous substitute.

Instability is one of the essential obstacles of the RNAi therapeutic applications.³³ SiRNA is easily degraded by nucleases in vitro and in vivo. Andrieu-Soler and his colleagues have demonstrated that there was high nuclease activity in the vitreous.³⁴ To solve the problem, siRNA-PKC α combined with FCVB shows great advantages. On the one hand, chemically synthesized siRNA-PKCa was mixed with the suspension of 2 deoxyribose nucleotides to increase the stability in its 3' end; on the other hand, siRNA-PKC α was injected into the capsule of FCVB, protecting it from contacting nucleases in vitro and in vivo. In addition, Ambati J and colleagues reported that 21-nt or longer siRNAs suppress hemanogiogenesis and lymphangiogenesis in various mouse models of neovascularization in a sequence- and targetindependent manner through activating cell surface Toll-like receptor 3.35,36 Therefore, it is critical to address the safety issue using in vivo systems.

The underlying factor in many vision-threatening vitreoretinal diseases is unwanted intraocular cell proliferation.³⁷ In the pathology of PVR, proliferating RPE cells are thought to be an initiating factor.^{5–8} In a previous study, we found that PKC α affects the cell cycle progression and proliferation in RPE cells through the downregulation of p27kip1.¹⁴ Since PKC α was downregulated after exposure to siRNA-PKC α released from FCVB, it may be a rational approach for targeted therapies against RPE cell proliferation and PVR disease.

Intravitreal sustained-release systems are exciting new modalities of drug delivery that offer effective treatment of some visually devastating diseases.³⁸ SiRNA-PKC α combined with FCVB shows great prospects for preventing PVR, and it will greatly reduce the suffering and economic burden of patients. As a public release vehicle, FCVB has the potential to release several drugs at the same time, which will meet the requirements of drug combinations for many diseases.

In conclusion, FCVB can sustainably and mechanically release siRNA-PKC α via capsule apertures in a timedependent and dose-dependent manner in addition to serving as a vitreous substitute. The in vitro study demonstrated that PKC α was inhibited after exposure to siRNA-PKC α released from FCVBs. The present research offers a new avenue for the treatment of diseases of the posterior segment of eyes, especially proliferative eye diseases.

Conclusion

In this study, we developed a FCVB drug delivery system to release siRNA-PKC α to inhibit PKC α expression in RPE cells in vitro. We found that PKC α was significantly downregulated by siRNA-PKC α released from FCVB, providing a new avenue to prevent PVR.

Acknowledgments

This study was supported by Natural National Science Foundation of China (NSFC-30973258) and the National Basic Research Program of China ("973" program, Number 2007CB512200). The Eye Bank of Zhongshan Ophthalmic Center is thanked for providing eye cups for primary cell culture.

Disclosure

The authors report no conflicts of interest in this work.

References

- Sunalp MA, Wiedemann P, Sorgente N, Ryan SJ. Effect of adriamycin on experimental proliferative vitreoretinopathy in the rabbit. *Exp Eye Res.* 1985;41(1):105–115.
- Cowley M, Conway BP, Campochiaro PA, Kaiser D, Gaskin H. Clinical risk factors for proliferative vitreoretinopathy. *Arch Ophthalmol.* 1989; 107(8):1147–1151.
- Kirchhof B. Strategies to influence PVR development. Graefes Arch Clin Exp Ophthalmol. 2004;242(8):699–703.

- 4. Cardillo JA, Stout JT, LaBree L, et al. Post-traumatic proliferative vitreoretinopathy. The epidemiologic profile, onset, risk factors, and visual outcome. *Ophthalmology*. 1997;104(7):1166–1173.
- Murphy TL, Sakamoto T, Hinton DR, et al. Migration of retinal pigment epithelium cells in vitro is regulated by protein kinase C. *Exp Eye Res.* 1995;60(6):683–695.
- Harris MS, Sakamoto T, Kimura H, et al. Hypericin inhibits cell growth and induces apoptosis in retinal pigment epithelial cells: possible involvement of protein kinase C. Curr Eye Res. 1996;15(3):255–262.
- Sheu SJ, Sakamoto T, Osusky R, et al. Transforming growth factorbeta regulates human retinal pigment epithelial cell phagocytosis by influencing a protein kinase C-dependent pathway. *Graefes Arch Clin Exp Ophthalmol.* 1994;232(11):695–701.
- Sakamoto T, Hinton DR, Sakamoto H, et al. Collagen gel contraction induced by retinal pigment epithelial cells and choroidal fibroblasts involves the protein kinase C pathway. *Curr Eye Res.* 1994; 13(6):451–459.
- 9. Nishizuka Y. The molecular heterogeneity of protein kinase C and its implications for cellular regulation. *Nature*. 1988;334(6184):661–665.
- Moriarty P, Dickson AJ, Erichsen JT, Boulton M. Protein kinase C isoenzyme expression in retinal cells. *Ophthalmic Res*. 2000;32(2–3): 57–60.
- Evans VG. Multiple pathways to apoptosis. *Cell Biol Int.* 1993;17(5): 461–476.
- Kizaki H, Suzuki K, Tadakuma T, Ishimura Y. Adenosine receptormediated accumulation of cyclic AMP-induced T-lymphocyte death through internucleosomal DNA cleavage. *J Biol Chem.* 1990; 265(9):5280–5284.
- Yu K, Ma P, Ge J, et al. Expression of protein kinase C isoforms in cultured human retinal pigment epithelial cells. *Graefes Arch Clin Exp Ophthalmol.* 2007;245(7):993–999.
- Gao Q, Tan J, Ma P, et al. PKC alpha affects cell cycle progression and proliferation in human RPE cells through the downregulation of p27kip1. *Mol Vis.* 2009;15:2683–2695.
- Hughes PM, Olejnik O, Chang-Lin JE, Wilson CG. Topical and systemic drug delivery to the posterior segments. *Adv Drug Deliv Rev.* 2005;57(14):2010–2032.
- 16. Cunha-Vaz JG. The blood-ocular barriers: past, present, and future. *Doc Ophthalmol*. 1997;93(1–2):149–157.
- Yasukawa T, Ogura Y, Tabata Y, Kimura H, Wiedemann P, Honda Y. Drug delivery systems for vitreoretinal diseases. *Prog Retin Eye Res.* 2004; 23(3):253–281.
- Liu Y, Ke Q, Chen J, et al. Sustained mechanical release of dexamethasone sodium phosphate from a foldable capsular vitreous body. *Invest Ophthalmol Vis Sci.* 2010;51(3):1636–1642.
- Liu Y, Jiang Z, Gao Q, et al. Technical standards of a foldable capsular vitreous body in terms of mechanical, optical, and biocompatible properties. *Artif Organs*. 2010;34(10):836–845.
- Lin X, Ge J, Gao Q, et al. Evaluation of the flexibility, efficacy, and safety of a foldable capsular vitreous body in the treatment of severe retinal detachment. *Invest Ophthalmol Vis Sci.* 2011;52(1):374–381.
- Gao Q, Mou S, Ge J, et al. A new strategy to replace the natural vitreous by a novel capsular artificial vitreous body with pressure-control valve. *Eye (Lond)*. 2008;22(3):461–468.

International Journal of Nanomedicine

Publish your work in this journal

The International Journal of Nanomedicine is an international, peerreviewed journal focusing on the application of nanotechnology in diagnostics, therapeutics, and drug delivery systems throughout the biomedical field. This journal is indexed on PubMed Central, MedLine, CAS, SciSearch®, Current Contents®/Clinical Medicine,

Submit your manuscript here: http://www.dovepress.com/international-journal-of-nanomedicine-journal

- Bernstein E, Caudy AA, Hammond SM, Hannon GJ. Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature*. 2001;409(6818):363–366.
- Novina CD, Sharp PA. The RNAi revolution. *Nature*. 2004;430(6996): 161–164.
- Reich SJ, Fosnot J, Kuroki A, et al. Small interfering RNA (siRNA) targeting VEGF effectively inhibits ocular neovascularization in a mouse model. *Mol Vis*. 2003;9:210–216.
- Nakamura H, Siddiqui SS, Shen X, et al. RNA interference targeting transforming growth factor-beta type II receptor suppresses ocular inflammation and fibrosis. *Mol Vis.* 2004;10:703–711.
- Cinatl J Jr, Michaelis M, Fleckenstein C, et al. West Nile virus infection induces interferon signaling in human retinal pigment epithelial cells. *Invest Ophthalmol Vis Sci.* 2006;47(2):645–651.
- Campa MJ, Wang MZ, Howard B, Fitzgerald MC, Patz EF Jr. Protein expression profiling identifies macrophage migration inhibitory factor and cyclophilin a as potential molecular targets in non-small cell lung cancer. *Cancer Res.* 2003;63(7):1652–1656.
- Weaver SA, Schaefer AL, Dixon WT. Western blotting for detection of glucocorticoid receptors in the brain and pituitary gland from adrenal intact pigs. *Brain Res.* 2000;869(1–2):130–136.
- Acharya G, Shin CS, Vedantham K, et al. A study of drug release from homogeneous PLGA microstructures. *J Control Release*. 2010; 146(2):201–206.
- Colthurst MJ, Williams RL, Hiscott PS, Grierson I. Biomaterials used in the posterior segment of the eye. *Biomaterials*. 2000;21(7):649–665.
- Gao Q, Chen X, Ge J, et al. Refractive shifts in four selected artificial vitreous substitutes based on Gullstrand-Emsley and Liou-Brennan schematic eyes. *Invest Ophthalmol Vis Sci.* 2009;50(7):3529–3534.
- Chen J, Gao Q, Liu Y, et al. Clinical device-related article evaluation of morphology and functions of a foldable capsular vitreous body in the rabbit eye. J Biomed Mater Res B Appl Biomater. 2011;97(2):396–404.
- Fattal E, Barratt G. Nanotechnologies and controlled release systems for the delivery of antisense oligonucleotides and small interfering RNA. *Br J Pharmacol.* 2009;157(2):179–194.
- Andrieu-Soler C, Bejjani RA, de Bizemont T, Normand N, BenEzra D, Behar-Cohen F. Ocular gene therapy: a review of nonviral strategies. *Mol Vis.* 2006;12:1334–1347.
- Kleinman ME, Yamada K, Takeda A, et al. Sequence- and targetindependent angiogenesis suppression by siRNA via TLR3. *Nature*. 2008;452(7187):591–597.
- Cho WG, Albuquerque RJ, Kleinman ME, et al. Small interfering RNA-induced TLR3 activation inhibits blood and lymphatic vessel growth. *Proc Natl Acad Sci U S A*. 2009;106(17):7137–7142.
- Cheng L, Hostetler K, Valiaeva N, et al. Intravitreal crystalline drug delivery for intraocular proliferation diseases. *Invest Ophthalmol Vis Sci.* 2010;51(1):474–481.
- Guidetti B, Azéma J, Malet-Martino M, Martino R. Delivery systems for the treatment of proliferative vitreoretinopathy: materials, devices and colloidal carriers. *Curr Drug Deliv*. 2008;5(1):7–19.

Dovepress

Journal Citation Reports/Science Edition, EMBase, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/ testimonials.php to read real quotes from published authors.