

Ex vivo laser confocal microscopy findings of cultured *Acanthamoeba* trophozoites

Natsuko Yamazaki¹
Akira Kobayashi¹
Hideaki Yokogawa¹
Yasuhisa Ishibashi²
Yosaburo Oikawa³
Masaharu Tokoro⁴
Kazuhisa Sugiyama¹

¹Department of Ophthalmology, Kanazawa University Graduate School of Medical Science, Kanazawa, Japan;

²Department of Ophthalmology, East Washinomiya Hospital, Kuki, Japan;

³Department of Medical Zoology, Kanazawa Medical University, Kahoku, Japan; ⁴Department of Parasitology, Kanazawa University Graduate School of Medical Science, Kanazawa, Japan

Purpose: The purpose of the current study was to investigate ex vivo laser confocal microscopic findings of cultured *Acanthamoeba* trophozoites obtained from *Acanthamoeba* keratitis patients.

Methods: Eight cultured samples of *Acanthamoeba* trophozoites from eight eyes of seven patients (mean age, 26.9 years; age range, 18–52 years) were used. Seven samples were from corneal scrapings of *Acanthamoeba* keratitis patients and one sample was from the solution in a soft contact lens case. Ex vivo laser confocal microscopy was performed to qualitatively evaluate the shape and degree of light reflection of the living *Acanthamoeba* trophozoites.

Results: Ex vivo laser confocal microscopy demonstrated highly reflective, high-contrast *Acanthamoeba* trophozoites with no walls (mean size, 25.4 μm ; range, 17.1–58.5 μm). The shapes of the trophozoites were highly pleomorphic, and some showed characteristic acanthopodia by laser confocal microscopy.

Conclusion: Ex vivo laser confocal microscopy was effective in demonstrating cultured *Acanthamoeba* trophozoites of various shapes and sizes. The observations of the current study may be helpful when similar structures are identified under in vivo conditions.

Keywords: *Acanthamoeba*, trophozoite, laser confocal microscopy

Introduction

Acanthamoeba is a ubiquitous, free-living amoeba found in water (eg, swimming pools, hot tubs, tap water, and contact lens solution), air, and soil, but *Acanthamoeba* keratitis is a relatively new entity.¹ The first case of *Acanthamoeba* keratitis was reported in 1974,¹ and the first case in Japan was reported in 1988.² As the use of soft contact lenses increased in the early 1980s, the incidence of reported *Acanthamoeba* keratitis increased dramatically. *Acanthamoeba* keratitis is relatively uncommon, but is a potentially blinding corneal infection. Clinical diagnosis is very difficult, especially in the early phase of the disease, and it is often misdiagnosed and treated as a herpes simplex infection.³ A definitive diagnosis is made by confirmation of *Acanthamoeba* in corneal lesions by direct examination, by corneal biopsy, or by culture. However, these methods are invasive and time-consuming, and are not always routinely available. The invasive methods are often postponed until there is a high index of suspicion for the disease and when there has been no response to treatments for bacterial, viral, and/or fungal keratitis.⁴ Unfortunately, delayed diagnosis of *Acanthamoeba* keratitis often results in poorer patient outcomes.

In vivo confocal microscopy has been used as a noninvasive technique for the observation of normal and pathological corneal microstructures. The usefulness of

Correspondence: Akira Kobayashi
13-1 Takara-machi, Kanazawa-shi,
Ishikawa-ken 920-8641 Japan
Tel +81 76 265 2403
Email kobaya@kenroku.kanazawa-u.ac.jp

this device in diagnosis and monitoring the improvement of *Acanthamoeba* keratitis has been reported.^{4–11} Previously, we¹² and others¹³ reported ex vivo confocal microscopic images of cultured *Acanthamoeba* cysts. As a result, *Acanthamoeba* cysts were observed as highly reflective round- or stellate-shaped high-contrast particles (10–20 microns in diameter).^{12,13} In this study, we investigated ex vivo laser confocal microscopic findings of cultured *Acanthamoeba* trophozoites.

Patients and methods

The present study was approved by the Ethical Committee of the Kanazawa University Graduate School of Medical Science and followed the tenets of the Declaration of Helsinki.

Sample collection and culturing of *Acanthamoeba* trophozoites

Eight cultured samples of trophozoites from eight eyes of seven patients (mean age, 26.9 years; age range, 18–52 years) were used. Seven samples were from corneal scrapings of *Acanthamoeba* keratitis patients and one sample was from the solution from a soft contact lens case (Case 3). All patients were seen and treated at the Department of Ophthalmology, Kanazawa University Graduate School of Medical Science between August 2006 and November 2008. The demographic data and treatments are shown in Table 1. The culture medium was an amoeba saline containing 0.012% NaCl, 0.00035% KCl, 0.0003% CaCl₂, and 0.0004% MgCl₂ · 7 H₂O in 0.05 mM Tris-HCl with a pH of 6.8, supplemented with *Escherichia coli*.¹⁴ The detailed culture method was described previously.¹⁴

Ex vivo laser confocal microscopy

After applying a large drop of contact gel (Comfort Gel ophthalmic ointment®; Bosch and Lomb, GmbH, Berlin, Germany) on the front surface of the microscope lens and ensuring no air bubbles had formed, a Tomo-Cap® (Heidelberg Engineering GmbH, Dossenheim, Germany) was mounted on the holder to cover the microscope lens. A suspension of cultured *Acanthamoeba* trophozoites was then dropped into a small space created in front of the Tomo-Cap using Scotch® tape. Ex vivo laser confocal microscopy (Heidelberg Retina Tomograph 2-Rostock Cornea Module, HRT 2-RCM; Heidelberg Engineering GmbH) was performed to qualitatively evaluate the shape and degree of light reflection of the living *Acanthamoeba* trophozoites. The HRT 2-RCM uses a 60× water-immersion objective lens (Olympus Europa GmbH, Hamburg, Germany) and utilizes a 670-nm diode laser as the light source, with a 400-µm² area of observation.

Results

Light microscopic and ex vivo laser confocal microscopic observation

Light microscopic observation of cultured *Acanthamoeba* trophozoites showed amorphous *Acanthamoeba* trophozoites (Figure 1A). In all culture samples, ex vivo laser confocal microscopy demonstrated highly reflective, high-contrast *Acanthamoeba* trophozoites with no walls (mean size, 25.4 µm; range, 17.1–58.5 µm) (Figure 1B–D). The shapes of the trophozoites were highly pleomorphic and some showed characteristic acanthopodia by HRT2-RCM (Figure 1E). In some culture samples, clusters of *Acanthamoeba* cysts were sporadically observed as highly reflective round- or

Table 1 Demographic and clinical data for patients with *Acanthamoeba* keratitis

Case	Age (years)	Sex	Eye	Stage	CL	Treatment
1	20	F	B	Early	2WSCL	Topical 0.1% micafungin, topical 0.02% chlorhexidine, oral itraconazole
2	23	M	R	Early	2WSCL	Topical 0.1% micafungin, topical 0.1% chlorhexidine, oral itraconazole
3	18	M	L	Early	2WSCL	Topical 0.1% micafungin, topical 0.1% chlorhexidine, oral itraconazole
4	52	M	R	Advanced	HCL	Topical 0.05% micafungin, topical 0.05% chlorhexidine, oral itraconazole
5	32	M	R	Early	2WSCL	Topical 1% voriconazole, topical 0.02% chlorhexidine, oral itraconazole
6	24	F	L	Early	1MSCL	Topical 0.05% micafungin, topical 0.05% chlorhexidine, oral itraconazole
7	19	M	L	Early	2WSCL	Topical 0.05% micafungin, topical 0.05% chlorhexidine, oral itraconazole

Abbreviations: M, male; F, female; R, right eye; L, left eye; B, bilateral eyes; 2WSCL, two-week disposable soft contact lens; 1MSCL, one-month disposable soft contact lens; HCL, hard contact lens.




Figure 1 Light and confocal microscopic observation of cultured *Acanthamoeba* trophozoites. (A) Light microscopic observation of cultured *Acanthamoeba* trophozoites (arrows) (one scale = 10 μm). Note that the shape of the *Acanthamoeba* trophozoites was amorphous. (B) Ex vivo laser confocal microscopic observation of cultured *Acanthamoeba* trophozoites from Case 1. The *Acanthamoeba* trophozoites were observed as highly reflective, high-contrast amorphous structures. No walls were observed around the trophozoites (bar = 100 μm). (C) Ex vivo laser confocal microscopic observation of cultured *Acanthamoeba* trophozoites from Case 2 (bar = 100 μm). (D) Ex vivo laser confocal microscopic observation of cultured *Acanthamoeba* trophozoites from Case 6 (bar = 100 μm). (E) Acanthopodia (arrowhead), which are characteristic of *Acanthamoeba* trophozoites, were observed in some of the trophozoite images (Case 6) (bar = 100 μm). (F) Clusters of *Acanthamoeba* cysts were also observed as highly reflective, high-contrast stellate-shaped particles 10–20 μm in diameter (Case 2) (bar = 100 μm).

stellate-shaped particles that were 10–20 μm in diameter (Figure 1F).

Discussion

In this study, we have reported the ex vivo laser confocal microscopic findings of cultured *Acanthamoeba* trophozoites obtained from *Acanthamoeba* keratitis patients. As a result, we consistently observed highly reflective, high-contrast *Acanthamoeba* trophozoites with no walls (mean size, 25.4 μm ; range, 17.1–58.5 μm). This is consistent with the previously published trophozoite sizes evaluated

by in vivo confocal microscopy (25–50 μm in diameter).^{11,13} The shapes of the trophozoites were highly pleomorphic not only by light microscopy, but also by confocal microscopy. *Acanthamoeba* trophozoites were characterized by the presence of acanthopodia, which was also visible by ex vivo laser confocal microscopy.

In clinical settings, direct scraping is still the “gold standard” for definitive diagnosis of *Acanthamoeba* keratitis. Recently, HRT2-RCM was shown to provide high-resolution images of *Acanthamoeba* cysts with round or ovoid-shaped structures from 10–20 μm in diameter;^{11–13} this allows for the rapid and non-invasive diagnosis of early-stage *Acanthamoeba* keratitis. However, it was quite difficult to identify *Acanthamoeba* trophozoites in keratitis patients. Shiraishi et al reported putative *Acanthamoeba* trophozoites in stromal images obtained by HRT2-RCM in only one case out of nine keratitis patients.¹³ They found that the *Acanthamoeba* trophozoites showed acanthopodia, nucleus, and karyosomes.¹³ We could not detect definitive *Acanthamoeba* trophozoites in the corneal epithelium and/or stroma, possibly because they cannot be distinguished from other highly reflective pathological structures.¹³

The limitation of this study is that the size and shape of the *Acanthamoeba* trophozoites obtained herein may not be readily applicable to images of in vivo conditions in the corneal tissue. However, considering the difficulty of visualizing *Acanthamoeba* trophozoites in corneal tissue by confocal microscopy, the ex vivo morphological data may be a first step to better recognizing *Acanthamoeba* trophozoite images in vivo by confocal microscopy.

In conclusion, ex vivo laser confocal microscopy was effective in demonstrating cultured *Acanthamoeba* trophozoites with various shapes and sizes. The observations of the current study may be helpful when similar structures are identified under in vivo conditions.

Disclosure

The authors report no conflicts of interest. They received no financial support for this study.

References

1. Nagington J, Watson PG, Playfair TJ, McGill J, Jones BR, Steele AD. Amoebic infection of the eye. *Lancet*. 1974;2(7896):1537–1540.
2. Ishibashi Y, Matsumoto Y, Watanabe R, et al. A case of *Acanthamoeba* keratitis: The first case report in Japan. *Acta Soc Ophthalmol Jpn*. 1988; 92(6):963–972. Japanese.
3. Hammersmith KM. Diagnosis and management of *Acanthamoeba* keratitis. *Curr Opin Ophthalmol*. 2006;17(4):327–331.
4. Winchester K, Mathers WD, Sutphin JE, Daley TE. Diagnosis of *Acanthamoeba* keratitis in vivo with confocal microscopy. *Cornea*. 1995; 14(1):10–17.

5. Cavanagh HD, Petroll WM, Alizadeh H, He YG, McCulley JP, Jester JV. Clinical and diagnostic use of in vivo confocal microscopy in patients with corneal disease. *Ophthalmology*. 1993;100(10):1444–1454.
6. Kaufman SC, Musch DC, Belin MW, et al. Confocal microscopy: a report by the American Academy of Ophthalmology. *Ophthalmology*. 2004;111(2):396–406.
7. Nakano E, Oliveira M, Portellinha W, de Freitas D, Nakano K. Confocal microscopy in early diagnosis of *Acanthamoeba* keratitis. *J Refract Surg*. 2004;20(Suppl 5):S737–S740.
8. Parmar DN, Awwad ST, Petroll WM, Bowman RW, McCulley JP, Cavanagh HD. Tandem scanning confocal corneal microscopy in the diagnosis of suspected *Acanthamoeba* keratitis. *Ophthalmology*. 2006;113(4):538–547.
9. Chew SJ, Beuerman RW, Assouline M, Kaufman HE, Barron BA, Hill JM. Early diagnosis of infectious keratitis with in vivo real time confocal microscopy. *CLAO J*. 1992;18(3):197–201.
10. Pfister DR, Cameron JD, Krachmer JH, Holland EJ. Confocal microscopy findings of *Acanthamoeba* keratitis. *Am J Ophthalmol*. 1996;121(2):119–128.
11. Matsumoto Y, Dogru M, Sato EA, et al. The application of in vivo confocal scanning laser microscopy in the management of *Acanthamoeba* keratitis. *Mol Vis*. 2007;13:1319–1326.
12. Kobayashi A, Ishibashi Y, Oikawa Y, Yokogawa H, Sugiyama K. In vivo and ex vivo laser confocal microscopy findings in patients with early-stage *Acanthamoeba* keratitis. *Cornea*. 2008;27(4):439–445.
13. Shiraishi A, Uno T, Oka N, Hara Y, Yamaguchi M, Ohashi Y. In vivo and in vitro laser confocal microscopy to diagnose *Acanthamoeba* keratitis. *Cornea*. 2010;29(8):861–865.
14. Oikawa Y, Kitagawa K, Yagita K, Endo T, Shima I, Ikeda T. Drug susceptibility of *Acanthamoeba* isolated from 13 Japanese patients with *Acanthamoeba* keratitis. *J Kanazawa Med Univ*. 2005;30(2):67–70.

Clinical Ophthalmology

Publish your work in this journal

Clinical Ophthalmology is an international, peer-reviewed journal covering all subspecialties within ophthalmology. Key topics include: Optometry; Visual science; Pharmacology and drug therapy in eye diseases; Basic Sciences; Primary and Secondary eye care; Patient Safety and Quality of Care Improvements. This journal is indexed on

Submit your manuscript here: <http://www.dovepress.com/clinical-ophthalmology-journal>

Dovepress

PubMed Central and CAS, and is the official journal of The Society of Clinical Ophthalmology (SCO). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.