

Economic evaluation of eribulin as second-line treatment for metastatic breast cancer in South Korea

Gabriel Tremblay¹
Unnati Majethia²
Janis L Breeze³
Ilias Kontoudis⁴
Jeongae Park⁵

¹Geneconomics Inc., Lévis, Quebec, Canada; ²Global Value and Access, Eisai Inc., Woodcliff Lake, NJ, ³Tufts Clinical and Translational Science Institute, Tufts University, and Institute for Clinical Research and Health Policy Studies, Tufts Medical Center, Boston, MA, USA; ⁴Global Value and Access, Eisai Limited, Hatfield, Hertfordshire, UK; ⁵Eisai Korea Inc., Gangnam-gu, Seoul, South Korea

Background: Metastatic breast cancer (MBC) is associated with poor prognosis, particularly for those patients with human epidermal growth factor receptor (HER2)-negative tumor. Similar to the rest of the world, treatment options are limited in South Korea following first-line chemotherapy with anthracyclines and/or taxanes. This study examined the cost-effectiveness and cost-utility of eribulin in South Korean patients with HER2-negative MBC who have progressed after usage of at least one chemotherapeutic regimen for advanced disease (second-line therapy).

Methods: A partition survival model was developed from the perspective of the South Korean health care system. The economic impact of introducing eribulin as second-line therapy for HER2-negative MBC was compared to that of capecitabine and vinorelbine. The analysis estimated incremental cost per life-year (LY), that is, cost-effectiveness, and cost per quality-adjusted life-year (QALY), that is, cost-utility, of eribulin for management of HER2-negative MBC in South Korea. The model accounted for overall survival, progression-free survival, drug costs, grade 3/4 adverse events, and health care utilization. Deterministic and probabilistic sensitivity analyses were performed to identify uncertainty in the results of the economic evaluation.

Results: Second-line eribulin was associated with greater benefits in terms of LY and QALY, compared to capecitabine and vinorelbine. The incremental cost-effectiveness ratio was ₩10.5M (approximately USD 9,200) per LY, and the incremental cost-utility ratio was ₩17M (approximately USD 14,800) per QALY in the basecase analysis. The incremental cost-utility ratio ranged from ₩12M (USD 10,461) to ₩27M (USD 23,538) per QALY in the deterministic sensitivity analysis. In the probabilistic sensitivity analysis, >99% of the simulations were below ₩50M (USD 42,300), and the lower and upper 95% confidence intervals were ₩3M (USD 2,600) and ₩24M (USD 20,900) per QALY, respectively.

Conclusion: There currently exist a limited number of treatment choices for women with HER2-negative MBC. Eribulin is a cost-effective option for second-line therapy in South Korea and should be added to the current indications for reimbursement.

Keywords: eribulin, metastatic breast cancer, cost-utility, economic analysis

Introduction

Women presenting with locally advanced or metastatic breast cancer (LABC/MBC) have a poor prognosis, with <25% surviving beyond 5 years.^{1,2} Though therapeutic options for patients with human epidermal growth factor receptor (HER2)-positive MBC generally consist of several lines of single-agent or combination chemotherapy, options are particularly limited for women with HER2-negative tumor, which is associated with a poor prognosis. Recent data from the Korean Breast Cancer Registry show that ~75% of women who received surgery for breast cancer had the HER2-negative subtype.³

Correspondence: Unnati Majethia
Eisai Inc., 155 Tice Boulevard, Woodcliff
Lake, NJ 07677, USA
Tel +1 551 999 0514
Fax +1 201 949 4011
Email unnati_majethia@eisai.com

Eribulin is a chemotherapy drug belonging to the halichondrin class of drugs that has a different mode of action compared to the existing chemotherapies. Eribulin is licensed as a third-line MBC drug and its label is recently expanded to include second line HER-2 negative MBC as well.^{4,5}

Study 301 established the safety and efficacy of eribulin as second-line chemotherapy for patients with MBC who previously received treatment with an anthracycline and a taxane.⁶ This was a multicenter, Phase III, open-label, randomized, two-arm study conducted in 1,102 patients (554 eribulin, 548 capecitabine) with LABC/MBC. Patients were prestratified according to their geographical region and HER2 status and then randomized in a 1:1 ratio to receive either eribulin or capecitabine. Eribulin was administered as an intravenous infusion of 1.23 mg/m² over 2–5 minutes on days 1 and 8 of a 21-day cycle. Exactly 1,250 mg/m² of capecitabine was administered orally twice daily in two equal doses on days 1–14, every 21 days. In a prespecified subgroup analysis, HER2-negative patients (~70% of randomized patients) treated with eribulin had significantly longer overall survival (OS) (15.9 months) than those who received capecitabine (13.5 months) ($P=0.030$).⁷

Halaven is currently reimbursed for patients with LABC/MBC who have previously received at least two chemotherapies in Korea. In line with the new approval label, this study examines the cost-utility of introducing eribulin following one prior chemotherapy (FOPC) for HER2-negative MBC in Korea, focusing on the patient population where eribulin was observed to provide the greatest clinical benefit. The primary economic endpoints were incremental cost per quality-adjusted life-year (QALY) supported by incremental cost per life-year (LY).

Methods

Economic perspective and patient population

A partition survival model was constructed from the perspective of the South Korean health care system to examine the cost-utility of introducing eribulin as second-line chemotherapy for HER2-negative LABC/MBC.

In the economic model, the eribulin comparator mix consisted of capecitabine and vinorelbine (cape/vin), with 50% of patients assumed to be treated with each agent. In comparison to the Study 301 design, vinorelbine was added to the treatment mix in order to reflect as accurately as possible the current clinical practice, since capecitabine and vinorelbine are the most widely used monotherapies in

South Korea based on the expert opinion (Nielsen Korea, unpublished data, 2014). A 50%/50% split in the utilization between treatments was assumed, since no local market share data were available. In the absence of comparative clinical data for vinorelbine versus eribulin or capecitabine in the specific patient population, vinorelbine was assumed to have equal efficacy to capecitabine based on Study 301. The hazard ratios for eribulin relative to capecitabine in Study 301 (HER2-negative patients, second-line treatment only) were 0.75 for OS (95% confidence interval [CI]: 0.60, 0.92) and 0.86 for progression-free survival (PFS) (0.69, 1.08). The hazard ratios for eribulin versus vinorelbine were assumed to be the same.

The economic endpoints were incremental cost per LY and incremental cost per QALY. Ethical approval for this study was not sought as this was an economic evaluation. It was however, previously obtained for the clinical trial which has been published separately.

Health states and treatments

The partition survival model included three health states: stable disease, progressive disease, and death. Patients entered the model in stable disease and switched to progressive disease or death. Primary therapy was assumed to be administered until tumor progression and secondary therapy after progression for a total treatment duration of maximum up to 8 months. This average duration of chemotherapy (7.35 months in Western Europe, rounded to 8 months – Western European data used as a proxy for South Korea, since no local data were available) was based on CancerMpact MBC data from Kantar Health,⁸ and included second-line treatment and beyond. The transition of patients between health states was based on data from Study 301. The partition in the current model was directly based on the Kaplan–Meier survivor function from patient-level data for the subgroup of HER2-negative patients in Study 301 who received second-line therapy.

Drug dose calculations were based on individual drug summary product characteristics. Wastage based on body surface area distribution was included in the analysis, and 10% dose rounding was employed for the smallest dose. Treatment cycles of 21 days⁹ were converted to 30.42-day (1-month) cycles for the ease of calculations (hereafter referred to as months).

Efficacy measures and survival extrapolation

The model considered OS and PFS data from Study 301. In the HER2-negative subgroup of patients in Study 301 who received eribulin or capecitabine as second-line therapies, survival was ~12% for eribulin and 7% for

Table 2 Health care resource utilization in MBC patients FOPC (costs in ₩)

Preprogression costs			Postprogression costs			End-of-life costs	
Type of cost	Monthly utilization	Cost per month	Type of cost	Monthly utilization	Cost per month	Type of cost	Cost for 2 weeks
Medical oncologist – first appointment	1.00	17,910	Medical oncologist-follow-up	1.96	27,205	Hospital/medical institution	4,213,038
Medical oncologist – follow-up	1.62	22,486	GP home visit	1.00	44,560	Total	4,213,038
Community nurse home visit	1.00	44,560	Psychotherapist	1.00	14,092		
Complete blood count	1.85	10,606	Community nurse home visit	1.00	44,560		
Blood chemistry panel, liver function test, CT scan	1.71	51,018	Complete blood count	1.83	10,491		
Urea and electrolytes	0.53	49,303	Blood chemistry panel, liver function test, CT scan, urea and electrolytes	1.78	53,106		
Total		195,883	Total		194,014		

Note: Data from National Health Insurance Services: South Korea.¹¹

Abbreviations: CT, computed tomography; FOPC, following one prior chemotherapy; GP, general practitioner; MBC, metastatic breast cancer.

of AEs. Health care utilization for the treatment of AEs was based on physician input. The costs associated with these treatments were obtained from the NHI cost database, 2014. Hospitalization costs were applied for AEs using a length of stay of 7.15 days and cost per hospitalization of ₩253,163 (USD 221), based on the World Health Organization CHOosing Interventions that are Cost-Effective (CHOICE) database (2008 data),¹⁴ resulting in a unit cost of ₩300,931 (USD 262) after applying inflation at 2.5% for 7 years.¹⁵

Utilities and quality of life

Health-related quality of life (HRQOL) data were collected in Study 301 and have been presented in Cortes et al.¹³ HRQOL data obtained using Quality of Life Questionnaire Cancer 30 in Study 301 were mapped to EuroQoL 5 Dimension Questionnaire (EQ-5D)-derived utility scores using a previously published and validated regression algorithm.¹⁶ The elicited utilities are presented in Hudgens et al.¹⁷ The resulting EQ-5D scores were used to infer utilities for the following states: baseline, tumor response, and progression.

Since the health states of the HRQOL analysis did not reflect the states considered in the economic evaluation, further post hoc calculations needed to be made. To determine the utility for the stable disease state, the incremental utility of tumor response was multiplied by the objective response rate obtained in Study 301. To determine the utility of the progression state, the EQ-5D utilities of the total study population were used to avoid a potential selection bias, since the observed Quality of Life Questionnaire Cancer 30 scores in Study 301 were only for eribulin and capecitabine arms (Table 3). Furthermore, the annual disutility of AEs,

Table 3 EQ-5D utilities

Health states	Eribulin utility scores (SD)	Capecitabine utility scores (SD)	Total study population scores (SD)
Baseline	0.704 (0.228)	0.691 (0.238)	0.697 (0.233)
Tumor response	0.780 (0.194)	0.783 (0.185)	0.782 (0.189)
Progression	0.705 (0.211)	0.651 (0.250)	0.679 (0.232)

(per treatment arm)

Note: Data from Hudgens et al.^{17,18}

Abbreviation: SD, standard deviation.

Table 4 Health state utilities used in the model

Health states	Eribulin	Capecitabine/vinorelbine arm
PFS	0.717	0.715
Progression	0.695	0.695

Abbreviation: PFS, progression-free survival.

estimated using independent linear mixed-effects models, was subtracted from the product of tumor response and objective response rate. The utility levels thus calculated and used in the model are presented in Table 4.^{17,18}

Sensitivity analyses

Deterministic SA and probabilistic SA (PSA) were conducted to identify uncertainty in the results of the economic analysis. The variables tested in the univariate deterministic SA included discounting rate, dose intensity, administration costs, direct health care costs for stable and progressive disease, secondary treatment costs, and utility level (Table 5). As the price of eribulin is a key variable, it was not incorporated into the SA but instead evaluated using a price acceptability curve.

A PSA was also developed using Monte Carlo simulations to create an incremental cost-effectiveness plane and a cost-effectiveness acceptability curve, in addition to determining the probability of cost-effectiveness, given a range of incremental cost-effectiveness ratio per QALY threshold. The PSA assessed first-order stochastic uncertainty related to the following variables: utility for each health state, costs (primary and secondary drug costs, administration costs),

preprogression survival, and OS (generating postprogression survival) for both arms (Table 6).

Results

Efficacy

The monthly partition is presented in Figure 1. The mean OS benefit over the 5-year horizon was of 21.75 months for eribulin and 17.13 months for cape/vin, with a mean difference of 4.61 months. The mean PFS benefit over the 5-year horizon was of 4.56 months for eribulin and 3.99 months for capecitabine, with a difference of mean 0.57 months.

Drug costs

Drug costs were ₩7,092,981 (USD 6,183) for eribulin versus ₩3,622,884 (USD 3,158) for cape/vin, with a difference of ₩3,470,098 (USD 3,025); 98% of this difference was due to primary drug costs. Other medical costs totaled ₩7,377,117 (USD 6,431) for eribulin and ₩6,823,025 (USD 5,948) for cape/vin, with a difference of ₩554,091 that was mainly due to improved survival in the eribulin arm. Overall, the analyses found a difference of ₩4,062,052 (USD 6,541) between eribulin and the cape/vin comparator. The total treatment costs are presented in Table 7.

Cost-effectiveness results

The benefits associated with eribulin compared to the cape/vin comparator were 1.81 LY versus 1.43 LY, with a

Table 5 Deterministic sensitivity analysis: scenario presentation

Scenario presentation	Low	Basecase	High
Scenario 1: Benefits discounting rate	0.0%	5.0%	7.0%
Scenario 2: Costs discounting rate	7.0%	5.0%	0.0%
Scenario 3: Costs and benefits discounting rates	0.0%	5.0%	7.0%
Scenario 4: Dose intensity	70.0%	85.4%	100.0%
Scenario 5: Administration costs	20.0%	0.0%	-20.0%
Scenario 6: Direct health care costs of stable state	-20.0%	0.0%	20.0%
Scenario 7: Direct health care costs of progression state	-20.0%	0.0%	20.0%
Scenario 8: New line of treatment costs after progression	-20.0%	0.0%	20.0%
Scenario 9: Utility of stable state of eribulin	0.788	0.717	0.645
Scenario 10: Utility of progression state of eribulin	0.765	0.695	0.626
Scenario 11: Utility of stable state of capecitabine	0.643	0.715	0.786
Scenario 12: Utility of progression state of capecitabine	0.626	0.695	0.765

Table 6 Parameters evaluated in probabilistic sensitivity analysis

Parameters	Variables	Point estimate	Standard error	Distribution	Source
Utility	Baseline – eribulin	0.713	0.23	Beta	Trial QOL data
	Tumor response – eribulin	0.801	0.19	Beta	Trial QOL data
	Progression – eribulin	0.695	0.21	Beta	Trial QOL data
	Baseline – TPC	0.713	0.24	Beta	Trial QOL data
	Tumor response – TPC	0.808	0.19	Beta	Trial QOL data
	Progression – TPC	0.695	0.25	Beta	Trial QOL data
Costs	Primary, secondary therapy and administration drug cost		SD = ±20%	Normal	Assumption
Survival	Preprogression – eribulin	4.56	0.42	Normal	Mean based on the clinical trial Kaplan–Meier and SD based on clinical trial (based on restricted OS mean)
	Postprogression – eribulin	17.19	1.05	Normal	Mean based on the clinical trial Kaplan–Meier and SD based on clinical trial (based on restricted OS mean)
	Preprogression – TPC	3.99	0.45	Normal	Mean based on the clinical trial Kaplan–Meier and SD based on clinical trial (based on restricted PFS mean)
	Postprogression – TPC	13.15	0.95	Normal	Mean based on the clinical trial Kaplan–Meier and SD based on clinical trial (based on restricted OS mean)
Dose intensity		0.854	70%–100%	Beta	Assumption

Abbreviations: OS, overall survival; PFS, progression-free survival; QOL, quality of life; SD, standard deviation; TPC, treatment of physician's choice.




Figure 1 Monthly partition analysis.

Abbreviations: OS, overall survival; PFS, progression-free survival.

difference of 0.38 LY. The incremental cost-effectiveness ratio was calculated as ₩10,564,275 (USD 9,200) for eribulin versus cape/vin.

Cost-utility results

The benefits associated with eribulin compared to the cape/vin comparator were 1.18 QALY versus 0.94 QALY, with a difference of 0.24 QALY. The incremental cost-utility ratio was calculated and found to be ₩16,898,483 (approximately USD 14,800) for eribulin versus cape/vin.

Since eribulin is indicated for life-threatening MBC, at a threshold of ₩50,000,000 (USD 42,300) per QALY,¹⁷ eribulin is likely to be a cost-effective treatment FOPC for HER2-negative MBC patients. The price acceptability curve is presented in Figure 2 and shows that the treatment is still cost-effective at different price levels.

Sensitivity analyses

Results of the deterministic SA are presented with a tornado graph in Figure 3. The univariate scenario analyses

demonstrated that the cost-utility of eribulin is sensitive to the utility level per health state and the dose intensity, but less sensitive to other variables.

The PSA found that the cost per LY ranged from ₩9M (USD 7,800) to ₩10M (USD 8,700), with an average of ₩9.1M (USD 7,900). The cost per QALY ranged from ₩13M (USD 11,300) to ₩14M (USD 12,200), with an average at ₩13.5M (USD 11,700). When looking at the probability of being under a certain threshold, 50% of the cost per QALY estimates were below ₩12.6M (USD 11,000), 75% were below ₩16.9M (USD 14,700), 90% were below ₩21.2M (USD 18,500), and 99% were below ₩35M (USD 30,500).

Table 7 Total treatment costs

Treatment	Eribulin	Capecitabine/ vinorelbine arm	Difference
Main therapy costs	4,897,727	1,502,572	3,395,155
Main therapy administration costs	58,357	48,695	9,662
Post-therapy – TPC costs	2,074,833	2,011,448	63,385
Post-therapy administration costs	62,065	60,169	1,896
Total drug costs	7,092,981	3,622,884	3,470,098
Direct medical costs			
Stable state costs	869,156	763,613	105,543
Progression state costs	3,060,246	2,364,651	695,595
End-of-life costs	3,447,715	3,694,761	-247,046
Total direct medical costs	7,377,117	6,823,025	554,091
Adverse events costs	57,626	19,764	37,863
Total costs	14,527,724	10,465,673	4,062,052

Abbreviation: TPC, treatment of physician’s choice.

One per 1,000 (0.1%) was above ₩50M (USD 42,300), the threshold used for this model. The lower and upper 95% CIs were ₩3M (USD 2,600) and ₩24M (USD 20,900) per QALY, respectively. The cost-effectiveness plane is shown in Figure 4.

Discussion

MBC remains a devastating disease with poor prognosis and limited treatment options. Eribulin has emerged as a safe and effective treatment for women with the disease, particularly those with the HER2-negative subtype.

The clinical data for the economic evaluation was obtained from the Phase III clinical trial data of eribulin against capecitabine. Given the maturity of the clinical data, no survival extrapolation was necessary. Hence, the partition in the current model was based on the Kaplan–Meier survivor function from patient-level data for the subgroup of HER2-negative patients in Study 301 who received eribulin or capecitabine FOPC.

Most of the costs used in the model (ie, drug acquisition costs, administration costs, AE management costs) were obtained from local data sources like the NHI Service and World Health Organization databases. The analysis excluded patients’ out-of-pocket expenses, carers’ costs, and lost productivity derived costs, since the analysis was conducted from the South Korean payer perspective.

The incremental cost-utility ratio of eribulin compared with capecitabine and vinorelbine was calculated as the ratio of the difference in cost to the difference in QALYs. Consistent with the economic evaluations of eribulin for second- and third-line therapy performed for other payer




Figure 2 Cost acceptability curve.

Abbreviations: ICER, incremental cost-effectiveness ratio; LY, life-year; QALY, quality-adjusted life-year.




Figure 3 Deterministic sensitivity analysis results: tornado diagram.




Figure 4 Cost-effectiveness plane. Abbreviation: QALY, quality-adjusted life-year.

perspectives,¹⁹ this analysis shows that eribulin as second-line therapy would be a cost-effective option in South Korea even against generic and less-expensive treatments such as capecitabine and vinorelbine.

The SA were highly consistent with the basecase analysis. The results of the PSA showed that the risk of introducing eribulin FOPC at the current price is low, as over 99% of the simulations were below ₩50,000,000 (USD 42,300) per QALY.

Limitations

Several limitations of this analysis should be noted. Most estimates used in the model were derived from a subgroup of patients from Study 301. However, stratified analysis

of HER2-negative patients was prespecified as part of the trial design and almost 70% of the participants had HER2-negative disease; the CIs associated with the effect estimates in this subgroup, therefore, reflect good precision.⁷

Conclusion

The time horizon in the model was bounded at 5 years, and therefore, the analyses do not reflect the costs associated with survival beyond that period, which may apply to ~10% of the patients in our model. However, we believe the 5-year model horizon appropriately balances the risk of longer extrapolation with greater uncertainty while capturing the majority of survival benefits for most patients.

Most importantly, vinorelbine was not included in Study 301, and thus, estimates for the efficacy of vinorelbine as second-line therapy were not available. However, because this drug is now commonly used as second-line therapy, it was felt important to include it in our analysis, and therefore, a conservative estimate of its efficacy was used based on the 301 trial. By adding vinorelbine to the comparator mix, we were able to make comparisons that reflect the current treatment patterns, which is an important goal of cost-effectiveness analysis.

There currently exist only a limited number of second-line treatment choices for women with HER2-negative MBC. Second-line eribulin is an important and cost-effective addition to the treatment mix in South Korea.

Acknowledgments

This study was funded by Eisai. Eisai developed the analysis with validation from external consultants.

Disclosure

Tremblay G was working for Eisai Inc. at the time of development of analysis. The authors report no other conflicts of interest in this work.

References

- Twelves C, Cortes J, Vahadat L, Wanders J, Kaufman P. Phase III trials of eribulin mesylate (E7389) in extensively pretreated patients with locally recurrent metastatic breast cancer. *Clin Breast Cancer*. 2010; 10(2):160–163.
- National Cancer Institute. *SEER Stat Fact Sheets: Breast*. Available from: <http://seer.cancer.gov/statfacts/html/breast.html>. Accessed January 14, 2016.
- You J, Kim Y, Moon H, Nam S, et al. Survival improvement in Korean breast cancer patients due to increases in early-stage cancers and hormone receptor positive/HER2 negative subtypes: a nationwide registry-based study. *J Breast Cancer*. 2015;18(1):8–15.
- Ezdrug* (2016) [prescribing information]. Available from: <https://ezdrug.mfds.go.kr:https://ezdrug.mfds.go.kr/#!/CCBAA03F010>. Accessed July 26, 2016.
- Halaven® 0.5mg/mL solution for injection Prescription Drug: Code of drug 04210 [prescribing information]. 2014.
- Kaufman P, Awada A, Twelves C, et al. Phase III open-label randomized study of eribulin mesylate versus capecitabine in patients with locally advanced or metastatic breast cancer previously treated with an anthracycline and a taxane. *J Clin Oncol*. 2015;33(6):594–601.
- Kaufman P, Awada A, Twelves C, et al. A phase III, open-label, randomized, multicenter study of eribulin mesylate versus capecitabine in patients with locally advanced or metastatic breast cancer previously treated with anthracycline. *Cancer Res*. 2012;72(24 Suppl):S6–S6.
- Kantar Health. Treatment architecture: Western Europe breast cancer. Kantar health. Available from: <https://authentication.khapps.com/>. Published March 2014. Accessed January 14, 2016.
- Eisai Australia Pty Ltd. (2015). *Halaven Product Information*. Available from: <https://www.ebs.tga.gov.au/ebs/picmi/picmirepository.nsf/pdf?OpenAgent&id=CP-2012-PI-02654-1&d=2015070116114622412>. Published June 2015. Accessed January 14, 2016.
- Majethia U, Tremblay G, He Y, Faria C, McCutcheon S, Koppenhafer L, Forsythe A. Economic burden of chemotherapy related toxicities in third line metastatic breast cancer patients. *Value Health*. 2014;17(7):A628.
- National Health Insurance Services: South Korea. Available from: <http://www.nhic.or.kr/static/html/wbd/g/a/wbdga0101.html>. Published 2016. Accessed January 14, 2016.
- ISPOR. (2016). *Pharmacoeconomic Guidelines Around The World: South Korea*. Available from: http://www.ispor.org/peguidelines/source/Korean_PE_Guidelines_Korean_Version.pdf. Accessed January 14, 2016.
- Cortes J, Hudgens S, Twelves C, et al. Health-related quality of life in patients with locally advanced or metastatic breast cancer treated with eribulin mesylate or capecitabine in an open-label randomized phase 3 trial. *Breast Cancer Res Treat*. 2015;154(3):509–520.
- World Health Organization. Cost effectiveness and strategic planning (WHO-CHOICE). Available from: <http://www.who.int/choice/costs/en/>. Published 2016. Accessed January 14, 2016.
- World Inflation Data. Historic inflation South Korea - CPI inflation. Available from: <http://www.inflation.eu/inflation-rates/south-korea/historic-inflation/cpi-inflation-south-korea.aspx>. Published 2016. Accessed January 14, 2016.
- Crott R, Versteegh M, Uyl-de-Groot C. An assessment of the external validity of mapping QLQ-C30 to EQ-5D preferences. *Qual Life Res*. 2013; 22:1045–1054.
- Hudgens S, Briggs A, Velikova G, Forsythe A, McCutcheon S, Kind P. Impact of treatment with eribulin (ERI) or capecitabine (CAP) for metastatic breast cancer (MBC) on EQ-5D utility derived from EORTC QLQ-C30. *Ann Oncol*. 2014;25(suppl 4):iv357–iv360.
- Hudgens S, Briggs A, Tremblay G, Forsythe A, Lloyd A. Comparison of methods to estimate health state utilities in metastatic breast cancer (MBC). *Value Health*. 2015;17:A557.
- Jang J. Drug reimbursement review in Korea – overview & case study. Presented at: ISPOR Asia-Pacific Conference; September 6–9, 2014; China, International Convention Center.

ClinicoEconomics and Outcomes Research

Publish your work in this journal

ClinicoEconomics and Outcomes Research is an international, peer-reviewed open-access journal focusing on health technology assessment, pharmacoeconomics and outcomes research in the areas of diagnosis, medical devices, and clinical, surgical and pharmacological intervention. The economic impact of health policy and health systems

Submit your manuscript here: <https://www.dovepress.com/clinicoeconomics-and-outcomes-research-journal>

organization also constitute important areas of coverage. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress