

# Clinical significance of quantitative *HER2* gene amplification as related to its predictive value in breast cancer patients in neoadjuvant setting

Ziping Wu<sup>1,\*</sup>  
 Shuguang Xu<sup>1,\*</sup>  
 Liheng Zhou<sup>1</sup>  
 Wenjin Yin<sup>2</sup>  
 Yanpin Lin<sup>1</sup>  
 Yueyao Du<sup>1</sup>  
 Yaohui Wang<sup>1</sup>  
 Yiwei Jiang<sup>1</sup>  
 Kai Yin<sup>1</sup>  
 Jie Zhang<sup>1</sup>  
 Jinsong Lu<sup>1</sup>

<sup>1</sup>Department of Breast Surgery, Renji Hospital, School of Medicine, Shanghai Jiao Tong University, Shanghai, People's Republic of China; <sup>2</sup>Cancer Institute, Fudan University Shanghai Cancer Center, Fudan University, Shanghai, People's Republic of China

\*These authors contributed equally to this work

**Background:** The aims of this study were to determine whether the quantitative *HER2* gene amplification level is related to the key clinicopathological features that represent the aggressiveness of breast cancer (BC) and to determine whether the quantitative *HER2* gene amplification level could predict the treatment response in the subset of *HER2*-positive patients who received neoadjuvant targeted therapy.

**Materials and methods:** Patients treated with weekly cisplatin- and paclitaxel-based neoadjuvant chemotherapy, who had undergone both immunohistochemistry and the fluorescence in situ hybridization test for *HER2*, were included in the study (n=103). For *HER2*-positive patients, defined as immunohistochemistry score 3+ or fluorescence in situ hybridization ratio  $\geq 2.0$ , trastuzumab was recommended with neoadjuvant chemotherapy (n=45). Pathological complete response was defined as complete pathological remission of tumor cells both in breast and axillary lymph nodes postoperation.

**Results:** In all patients enrolled in the study, a higher *HER2* amplification level was significantly correlated with larger tumor size and the absence of ER and PR expression. In *HER2*-positive patients treated with neoadjuvant trastuzumab concurrent with chemotherapy, both univariate and multivariate logistic regression showed that a higher *HER2/CEP17* ratio and *HER2* gene copy number were associated with a higher pathological complete response rate. When calculated by receiver operating characteristics analysis, an optimal cutoff of 4.5 for the *HER2/CEP17* ratio was expected to distinguish the most sensitive candidate for treatment with a combination of trastuzumab and neoadjuvant chemotherapy.

**Conclusion:** A higher *HER2* amplification level was correlated with larger tumor size and reduced ER and PR expression, which may indicate more aggressive tumor behavior. For *HER2*-positive patients, the *HER2/CEP17* ratio and *HER2* gene copy number may be good predictive factors for concurrent neoadjuvant trastuzumab and chemotherapy.

**Keywords:** breast cancer, neoadjuvant, quantitative *HER2* gene amplification, pathological complete response, predictive

## Introduction

The *HER2* gene encodes a 185 kDa monomeric protein, which is a receptor tyrosine kinase that belongs to the human epidermal growth factor receptor family.<sup>1</sup> Approximately 25% of breast cancer (BC) patients present with *HER2* gene amplification,<sup>1</sup> which is an independent prognostic factor of poor outcome. Anti-*HER2* therapy is now a standard treatment for *HER2*-positive patients both in adjuvant and metastatic settings.<sup>2</sup> When *HER2*-positive patients are treated with adjuvant trastuzumab therapy, the 10-year survival rate could be increased from 75.2% to 84%.

Correspondence: Jinsong Lu  
 Department of Breast Surgery, Renji Hospital, School of Medicine, Shanghai Jiao Tong University, No 160 Pujian Road, Shanghai 200127, People's Republic of China  
 Email lujss@163.com

Still, approximately 10% of these patients will inevitably develop recurrent or metastatic disease despite anti-HER2 treatment, reflecting the heterogeneous tumor biology of HER2-positive BC.<sup>3</sup>

In addition to the well-acknowledged fact that HER2 expression status represents a poor prognosis and good response to trastuzumab, effective methods are still required to distinguish the tumor biology of HER2-positive BC. As a routinely conducted test, fluorescence in situ hybridization (FISH) provides a linear measurement for the *HER2* gene amplification level, which seems to be a promising figure in reflecting the nature of HER2-positive BC. Several studies were carried out referring to this issue in adjuvant, neoadjuvant, and metastatic settings. In adjuvant settings, a large sample study by Xuan et al<sup>4</sup> observed an increased disease-free survival in trastuzumab-treated patients with a lower *HER2* amplification level. In addition, a study from Borley et al<sup>5</sup> observed an increased progression-free survival and overall survival in patients with lower *HER2* gene copy number. However, research that was correlated to the HERA trial showed no obvious influence of the *HER2* amplification degree on patient prognosis.<sup>6</sup> The latest meta-analysis from Xu et al<sup>7</sup> denied any correlations between this figure and survival.

In late-stage BC, a study from Fuchs et al<sup>8</sup> showed both higher *HER2* gene copy number (GCN) and that *HER2/CEP17* ratio is associated with shorter time to first metastasis in patients treated with trastuzumab and chemotherapy. Similarly another study also observed a shorter time to progression in patients with high *HER2/CEP17* ratio.<sup>9</sup> However, there was another study that observed an opposite trend: patients with high-level *HER2/CEP17* ratio showed a longer progression-free survival.<sup>10</sup>

In regard to neoadjuvant settings, results from different institutions were even more controversial.<sup>11–13</sup> It is still unclear whether *HER2* amplification level is correlated with patients' response to concurrent neoadjuvant therapy.

Disparity in the results may derive from the heterogeneity of patients and treatment in each study. Thus, we started to investigate this issue in the population of our prospective trials, in which patients received weekly paclitaxel in combination with cisplatin for neoadjuvant chemotherapy concurrent with trastuzumab in cases where the tumor was HER2-positive. In this study, we hypothesize that a higher *HER2/CEP17* ratio or *HER2* GCN is associated with more aggressive tumor biology and better pathological complete response (pCR) outcome in this highly uniform subset of patients.

## Materials and methods

### Patients and study design

BC patients from two paclitaxel- and cisplatin-based neoadjuvant clinical trials were included. The two trials were separately registered in [ClinicalTrials.gov](http://ClinicalTrials.gov) as SHPD001 (NCT02199418) and SHPD002 (NCT02221999). SHPD001 was reviewed and approved by two independent ethical committees and institutional review boards of RenJi Hospital, Shanghai Jiao Tong University and Fudan University Shanghai Cancer Center. SHPD002 was reviewed and approved by the independent ethical committee and institutional review board of RenJi Hospital, Shanghai Jiao Tong University. All patients provided written informed consent.

Women aged  $\geq 18$  years old with histologically confirmed locally advanced invasive BC were included. For all patients, paclitaxel 80 mg/m<sup>2</sup> was given weekly starting on day 1 for 16 weeks; cisplatin 25 mg/m<sup>2</sup> was given weekly on day 1, 8, and 15 every 28 days for 4 cycles. For HER2-positive patients in SHPD001, trastuzumab was recommended concurrently. All HER2-positive patients in SHPD002 received concurrent trastuzumab. Trastuzumab was given at a loading dose of 4 mg/kg, followed by a maintenance dose of 2 mg/kg, on day 1 for 16 weeks. For hormone receptor-positive patients in SHPD002, endocrine therapy of aromatase inhibitor or gonadotropin-releasing hormone agonist was randomized together with chemotherapy according to the patient's menstrual status. Planned surgery was given sequentially after neoadjuvant chemotherapy. All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

The study's primary outcome with regard to the two groups SHPD001 and SHPD002 was pCR, which was defined as the absence of tumor in the breast and axillary lymph nodes samples taken at the time of surgery.

Up until December 2016, 103 postoperative patients who went through the FISH test and 45 HER2-positive trastuzumab-treated patients were available for the analysis.

### IHC and FISH

ER, PR, and ki67 levels were assessed by immunohistochemistry (IHC) in paraffin-embedded tumor samples from biopsy. All patients enrolled in the clinical trial were tested by IHC to determine their HER2 status. Patients who tested 1+ to 3+ by IHC and some of the IHC 0 patients (n=103) were examined

by FISH. FISH was carried out on paraffinized tissue sections using the *HER2* DNA Dual Probe Kit (Linked-Biotech Pathology, Guangzhou, People's Republic of China). *HER2* positivity is defined according to American Society of Clinical Oncology (ASCO)/College of American Pathologists (CAP) recommendations from 2013.<sup>14</sup> In detail, 60 cells were routinely counted for FISH testing. Each cell was counted for *HER2* signal (red) and *CEP17* signal (green). Then, an average *HER2* signal per cell and an average *CEP17* signal per cell were used to obtain the final FISH ratio.

Both IHC and FISH assessment were conducted and confirmed by two qualified pathologists from the Department of Pathology, Shanghai Renji Hospital or Fudan University Shanghai Cancer Center.

## Statistical methods

Spearman's correlation test was used to determine the correlations between the *HER2* GCN, *HER2/CEP17* ratio, and other clinicopathological characteristics in patients who underwent FISH (n=103). In patients treated with concurrent trastuzumab (n=45), correlations between *HER2* GCN, the *HER2/CEP17* ratio, and pathological response to neoadjuvant therapy were calculated by both univariate and multivariate logistic regression analysis. Receiver operating characteristic (ROC) analysis was performed to see if *HER2* GCN and *HER2/CEP17* could independently predict patients' response to neoadjuvant therapy. Sensitivity and specificity were calculated to determine the optimal cutoff points. All statistical analyses were carried out using STATA Statistics SE 14 (Stata Corp LP, College Station, TX, USA). The results were considered significant when  $p < 0.05$ .

## Results

### Correlations of *HER2* gene amplification level and other clinicopathological characteristics

The baseline characteristics of all 103 patients who were available for FISH results to be obtained are shown in Table 1. Among these patients, *HER2* GCN ranged from 1.45 to 65 and *HER2/CEP17* ratios ranged from 1.09 to 50. No polysomy 17 cases were reported. *HER2* GCN was inversely related to ER expression level ( $p < 0.001$ ) and PR expression level ( $p < 0.001$ ) and positively related to ki67 level ( $p = 0.042$ ); the *HER2/CEP17* ratio was also inversely related with ER expression level ( $p < 0.001$ ) and PR expression level ( $p < 0.001$ ) and positively related with tumor size ( $p = 0.004$ ). No significant correlations were found between age and the *HER2* amplification level in either the ratio or GCN group (Table 2).

**Table 1** Baseline clinical characteristics of all patients

Characteristics	Number (n)	Percentage (%)
Age (years)		
$\geq 50$	59	57
$< 50$	44	43
Tumor size		
$> 5$ cm	58	56
$\leq 5$ cm	44	44
ER status		
ER +	78	76
ER -	25	24
PR status		
PR +	85	83
PR -	18	17
ki67 status <sup>a</sup>		
$> 30\%$	59	57
$\leq 30\%$	44	43
Menopause status		
Pre	51	50
Post	52	50

**Note:** <sup>a</sup>Most of the patients had a ki67 level greater than 14%, and thus 30% was used to separate the groups.

### Baseline clinical characteristics of trastuzumab-treated patients and their response to chemotherapy concurrent with target therapy

Among the 45 patients who were *HER2* positive and treated with trastuzumab, 22 patients reached pCR after surgery; the pCR rate was 48.9%. The baseline clinical characteristics of these 45 patients are summarized in Table 3. The detailed relationship between the *HER2/CEP17* ratio, *HER2* GCN, and neoadjuvant outcome for each patient is displayed in Figures 1 and 2. An obvious increase in pCR outcome was observed when the *HER2/CEP17* ratio or *HER2* GCN was elevated.

### Correlations of quantitative *HER2* gene amplification and pCR outcome

In univariate analysis, both a higher *HER2/CEP17* ratio (odds ratio [OR] = 1.827,  $p = 0.008$ ; 95% confidence interval [CI]:

**Table 2** Correlations between *HER2* amplification level and clinicopathological characteristics of all patients

Characteristics	N <sup>a</sup>	p-value (HER2 GCN) <sup>b</sup>	N	p-value (HER2/CEP17 ratio)
Age	101	0.051	103	0.465
Tumor size	100	0.194	102	<b>0.004</b>
ER status	101	<b>0.000</b>	103	<b>0.000</b>
PR status	101	<b>0.000</b>	103	<b>0.000</b>
Menopausal status	101	0.057	103	0.372
ki67	101	<b>0.042</b>	103	0.154

**Notes:** <sup>a</sup>101 patients had values for *HER2* GCN; <sup>b</sup>Bold facing indicates statistical significance.

**Abbreviation:** GCN, gene copy number.

**Table 3** Baseline clinical characteristics of HER2-positive trastuzumab-treated patients

Characteristics	Number (n)	Percentage (%)
Age (years)		
≥50	26	58
<50	19	42
Tumor size		
>5 cm	30	67
≤5 cm	15	33
ER status		
ER +	30	67
ER –	15	33
PR status		
PR +	33	73
PR –	12	7
ki67 status		
>30%	30	67
≤30%	15	33
Menopause status		
Pre	21	47
Post	24	53

1.176–2.851) and HER2 GCN amplification level (OR = 1.139,  $p=0.029$ ; 95% CI: 1.013–1.230) were associated with a better pCR rate, while ER expression was associated with a poorer pCR rate (OR = 0.211,  $p=0.025$ ; 95% CI: 0.053–0.824).

Similarly, in multivariate analysis, a higher HER2/CEP17 ratio (OR = 2.110,  $p=0.005$ ; 95% CI: 1.266–3.861) was also associated with a better pCR rate, while ER expression was associated with a poorer pCR rate (OR = 0.101,  $p=0.038$ ; 95% CI: 0.012–0.884). HER2 GCN was also positively related to pCR (OR = 1.147,  $p=0.050$ ; 95% CI: 1.000–1.317) when


adjusted. No significant correlations between other potential factors including patient age, PR status, and tumor size were observed (Tables 4–6).

## Optimal predictive cutoff points decided by ROC curve

The optimal cutoff point determined by ROC analysis was 4.5 for the HER2/CEP17 ratio and 14 for HER2 GCN, which were expected to distinguish the most sensitive and specific candidates for cisplatin- and paclitaxel-based weekly neoadjuvant chemotherapy combined with trastuzumab. The area under the curve value of the HER2/CEP17 ratio was 0.786 compared with 0.656 of HER2 GCN (Figure 3).

## Discussion


HER2 gene amplification was associated with poor survival outcome in BC patients. It is also an indication of trastuzumab treatment in HER2-positive BC patients.<sup>15</sup> However, the dosage effect of the HER2 gene amplification level on BC patients is not quite clear. Our study demonstrated that both HER2 GCN and HER2/CEP17 ratios are correlated with unfavorable prognostic factors. As for the predictive value, we demonstrated that both the HER2/CEP17 ratio and HER GCN were associated with better pCR outcome in trastuzumab-treated patients. This was the first time, at least to our knowledge, that correlations between tumor biology features, neoadjuvant treatment sensitivity, and both the HER2/CEP17 ratio and HER2 GCN were demonstrated in data from prospective clinical trials in the People's Republic of China.



**Figure 1** HER2/CEP17 ratio and neoadjuvant outcome of 44 patients treated with trastuzumab.

**Notes:** Every column represents an individual patient with an HER2/CEP17 ratio value (x-axis). The blue column represents pCR outcome, yellow column represents non-pCR outcome. An increasing tendency for pCR outcome was observed when the HER2/CEP17 ratio was elevated. One patient was not shown in the figure because the HER2/CEP17 value was too big (50).

**Abbreviation:** pCR, pathological complete response.



**Figure 2** *HER2* GCN and neoadjuvant outcome of 44 patients treated with trastuzumab.

**Notes:** Every column represents an individual patient with an *HER2* GCN value (x-axis). The blue column represents pCR outcome, yellow column represents non-pCR outcome. An increasing tendency for pCR outcome was observed when *HER2* GCN was elevated.

**Abbreviations:** GCN, gene copy number; pCR, pathological complete response.

We did FISH tests in patients with an IHC score equal to or greater than 0 at biopsy. This provided a more diverse population compared to studies conducted in patients who were mostly IHC 3+. In this set of patients, a higher *HER2* amplification level seemed to be accompanied by unfavorable clinicopathological factors. For instance, ER and PR status were inversely related to both *HER2* GCN and *HER2/CEP17* ratios. This result was supported by the work from Arnould et al.<sup>11</sup> In that research, high amplification of *HER2* was correlated with hormone receptor-negative tumors.

With regard to predictive value, existing data from clinical trials and observations have already shown that tumors with higher proliferation and invasion potential were more sensitive to chemotherapy. For instance, triple-negative BC often exhibits an increased pCR rate but a decreased overall

survival compared with other BC subtypes.<sup>16,17</sup> Similar to triple-negative BC, a study from Yu et al<sup>12</sup> demonstrated that *HER2*-positive BC was sensitive to neoadjuvant chemotherapy.<sup>12</sup> All the patients in that study were *HER2* positive and received neoadjuvant chemotherapy without targeted therapy. Yu et al observed a better pCR rate in patients with both higher *HER2/CEP17* ratio and *HER2* GCN.

Yu et al's study might indicate the sensitivity of patients with higher *HER2* amplification level to chemotherapy. Our study demonstrated, as well, that these patients are more sensitive to targeted therapy.

Our results are partially supported by the studies from Arnould et al<sup>11</sup> and Guiu et al,<sup>15</sup> where patients received concurrent trastuzumab and neoadjuvant chemotherapy of different regimens. Those patients were originally tested IHC 3+ or 2+ for *HER2* status. FISH assays were retrospectively conducted using the ASCO/CAP guideline version 2007.<sup>23</sup>

**Table 4** Univariate analyses for predictive factors of *HER2*-positive trastuzumab-treated patients

Characteristics	Univariate analysis				
	N	OR	95% CI	p-value <sup>b</sup>	
<i>HER2</i> GCN	44 <sup>a</sup>	1.139	1.013	1.230	<b>0.029</b>
<i>HER2/CEP17</i> ratio	45	1.827	1.170	2.851	<b>0.008</b>
Age (years) ≥50 vs <50	45	2.338	0.694	7.872	0.170
ER positive vs negative	45	0.211	0.053	0.824	<b>0.025</b>
PR positive vs negative	45	0.368	0.092	1.471	0.158
Tumor size (cm) >5 vs ≤5	45	0.680	0.219	2.108	0.504
Menopause status pre vs post	45	1.576	0.484	5.126	0.450
ki67	45	1.028	0.994	1.063	0.105

**Notes:** <sup>a</sup>44 patients had values for *HER2* GCN; <sup>b</sup>Bold facing indicates statistical significance.

**Abbreviations:** OR, odds ratio; CI, confidence interval; GCN, gene copy number.

**Table 5** Multivariate analyses for predictive factors of pCR using *HER2/CEP17* ratio

Characteristics	Multivariate analysis				
	N	OR	95% CI	p-value <sup>a</sup>	
<i>HER2/CEP17</i> ratio	45	2.210	1.266	3.861	<b>0.005</b>
Age (years) ≥50 vs <50	45	0.962	0.167	5.560	0.966
ER positive vs negative	45	0.078	0.009	0.673	<b>0.020</b>
PR positive vs negative	45	1.192	0.187	7.585	0.852
Tumor size (cm) >5 vs ≤5	45	0.719	0.480	1.077	0.110

**Note:** <sup>a</sup>Bold facing indicates statistical significance.

**Abbreviations:** pCR, pathological complete response; OR, odds ratio; CI, confidence interval.

**Table 6** Multivariate analysis for predictive factors of pCR using *HER2* GCN

Characteristics	Multivariate analysis			
	N <sup>a</sup>	OR	95% CI	p-value <sup>b</sup>
<i>HER2</i> GCN	44	1.147	1.000 1.317	<b>0.050</b>
Age (years) $\geq 50$ vs $< 50$	44	0.609	0.117 3.176	0.556
ER positive vs negative	44	0.190	0.343 1.046	0.056
PR positive vs negative	44	0.560	0.125 2.723	0.473
Tumor size (cm) $> 5$ vs $\leq 5$	44	0.881	0.654 1.186	0.404

**Notes:** <sup>a</sup>44 patients had values for *HER2* GCN; <sup>b</sup>Bold facing indicates statistical significance.

**Abbreviations:** pCR, pathological complete response; GCN, gene copy number; OR, odds ratio; CI, confidence interval.


Only *HER2* GCN was analyzed. When *HER2* GCN = 10 was used as the cutoff, the authors observed a better pCR rate in the cohort with high GCN. Our work is also supported by the work from Kogawa et al. Patients who received concurrent trastuzumab and neoadjuvant chemotherapy in that study showed a positive correlations between *HER2/CEP17* ratio and pCR.<sup>18</sup>

On the other hand, the study from Buzdar et al<sup>13</sup> showed no correlation between *HER2* amplification level and pCR outcome. However, only a brief conclusion was mentioned in the study. A detailed statistical method was not described.

Different from the neoadjuvant setting, results in the adjuvant settings are quite inconsistent. Briefly, we summarize some relevant studies in Table 7. In adjuvant settings, studies from Xuan et al<sup>4</sup> and Borley et al<sup>5</sup> found a positive correlation between *HER2* GCN and survival, but studies from Perez et al<sup>3</sup> and Dowsett et al<sup>6</sup> found no correlation. The difference between settings might derive from the schedule of trastuzumab. In neoadjuvant settings, trastuzumab is often used concurrently with chemotherapy. But in the adjuvant setting,

trastuzumab is usually used sequentially with chemotherapy. Concurrent chemotherapy and trastuzumab could increase the treatment effect, which might explain the major discrepancies between settings. Besides, the method of *HER2* signal counting might also influence the result. In the study from Perez et al, the maximum *HER2* signal count for the form of clouds was assigned as 20 per cell. This approach would not change the decision for trastuzumab treatment, but could underestimate the amplification level of some patients.

The pharmacological mechanism of trastuzumab was reported mainly from two aspects. One was described as anti-*HER2* dimerization and blockage of downstream cell signaling. The other was considered to be related to immune- and antibody-dependent cell-mediated cytotoxicity (ADCC).<sup>19</sup> When trastuzumab binds to the FC $\gamma$ R of natural killer cells, it would trigger ADCC and activate cell lysis. Some studies have reported that single-nucleotide polymorphisms of the FC $\gamma$ R might predict the efficacy of trastuzumab.<sup>20,21</sup> One of the studies focused on the FC $\gamma$ R gene *FCGR3A* and determined that single-nucleotide polymorphism rs396991 changed the amino acid on position 158 of *FCGR3A*. Patients with the low-affinity allele received less benefit from trastuzumab. Theoretically, a higher *HER2* gene amplification level may lead to higher *HER2* protein expression on the tumor cell surface. This is probably true considering the general consistency of the IHC and FISH results. At the same time, the tumor cell tends to combine with more antibodies, thus causing a greater extent of ADCC. Preclinical studies did determine that the ADCC effect of trastuzumab was partially relevant to *HER2* expression levels on tumor target cells in BC patients.<sup>22</sup> This was consistent with our result from the clinical samples.



**Figure 3** Optimal cutoff points of *HER2/CEP17* ratio and *HER2* GCN decided by ROC curves.

**Notes:** ROC curve using *HER2/CEP17* ratio (A) and *HER2* GCN (B) to select the best responder to concurrent neoadjuvant chemotherapy and trastuzumab. The AUC for *HER2/CEP17* ratio was 0.786, which was better than the AUC for *HER2* GCN (0.656).

**Abbreviations:** GCN, gene copy number; ROC, receiver operating characteristic; AUC, area under the curve.

**Table 7** Summary of the relevant research

Study	Settings	Treatment	Testing method	Conclusion
Borley et al, <sup>5</sup> 2014	Adjuvant	Adjuvant therapy (details not provided)	FISH (dual probe)	High <i>HER2</i> GCN associated with better survival
Xuan et al, <sup>4</sup> 2015		Trastuzumab-based chemotherapy	FISH (dual probe)	Both GCN and <i>HER2/CEP17</i> ratio (RATIO) associated with better survival
Perez et al, <sup>3</sup> 2014		Doxorubicin and cyclophosphamide followed by weekly paclitaxel with trastuzumab	FISH (dual probe)	No correlations between survival and RATIO/GCN
Dowsett et al, <sup>6</sup> 2009		Chemotherapy followed by trastuzumab	FISH (dual probe)	No correlations between survival and RATIO/GCN
Arnould et al, <sup>11</sup> 2007	Neoadjuvant	Trastuzumab-based chemotherapy with either docetaxel or docetaxel plus carboplatin	FISH (single probe)	High GCN associated with pCR
Buzdar et al, <sup>13</sup> 2007		Trastuzumab-based chemotherapy with paclitaxel followed by 5-fluorouracil, epirubicin and cyclophosphamide	FISH (dual probe)	No correlations between pCR and RATIO/GCN
Kogawa et al, <sup>18</sup> 2016		Chemotherapy with an anthracycline and/or a taxane given in combination or sequential regimens with or without trastuzumab	FISH (dual probe)	High RATIO associated with pCR
Yu et al, <sup>12</sup> 2012		Chemotherapy without trastuzumab	FISH (dual probe)	High GCN associated with pCR
Fuchs et al, <sup>8</sup> 2014	Metastatic	Trastuzumab-based treatment	FISH (dual probe)	High GCN associated with shorter time to first metastasis
Gullo et al, <sup>9</sup> 2009		Trastuzumab-based treatment	FISH (dual probe)	High RATIO associated with shorter TTP
Kim et al, <sup>10</sup> 2013		Trastuzumab plus taxane chemotherapy	FISH (dual probe)	High RATIO associated with longer PFS

**Abbreviations:** FISH, fluorescence in situ hybridization; GCN, gene copy number; pCR, pathological complete response; TTP, time to progression; PFS, progression-free survival.

Though the FISH result seemed simple and superficial, we should not neglect its value as a method for *HER2* gene quantization and as a therapeutic predictor. A highlight of our study is that we could use this routinely conducted test as a predictive and prognosis marker in *HER2*-positive patients, if validated.

Trastuzumab and pertuzumab are dual-targeted drugs for *HER2*-positive BC in the neoadjuvant setting. However, pertuzumab is not approved in the People's Republic of China currently. Thus, only trastuzumab was used in all the patients. Another major limitation of our study is the relatively small sample size. However, we did observe an obvious trend between the *HER2* amplification level and pCR results, despite the small sample size. Studies with larger sample sizes and assessing longer-term survival data are currently ongoing to validate our results in future studies.

## Conclusion

In summary, our study demonstrated that while using the latest ASCO/CAP guideline for *HER2* testing, a higher *HER2* amplification level was correlated with larger tumor size and

less ER and PR expression. In addition, both a higher *HER2/CEP17* ratio and *HER2* GCN could predict a better response to neoadjuvant therapy in trastuzumab-treated *HER2*-positive patients from neoadjuvant trials.

## Acknowledgment

This work is supported by grants from the National Natural Science Foundation of the People's Republic of China (grant numbers 81172505 and 81302302), the Doctoral Programs Foundation of the Ministry of Education of People's Republic of China (grant number 20120071120105), the Shanghai Natural Science Foundation (grant number 13ZR1452800), the Shanghai Municipal Commission of Health and Family Planning (grant numbers 20144Y0218, 201640006), the Science and Technology Commission of Shanghai Municipality (grant number 14411950202), the Clinical Research Plan of SHDC (grant number SHDC 12016231), and the Nurturing Fund of Renji Hospital 2015 (grant number RJZZ15-023). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Harris JR. HER2/ERBB2 Testing: Assessment of Status for Targeted Therapies. In: Harris JR, Lippman ME, Morrow M, Osborne CK, editors. *Diseases of the Breast*. 5th ed. Philadelphia: Lippincott Williams & Wilkins; 2014:324–438.
- National Comprehensive Cancer Network. NCCN Guidelines Version 2.2016: Breast Cancer.
- Perez EA, Romond EH, Suman VJ, et al. Trastuzumab plus adjuvant chemotherapy for human epidermal growth factor receptor 2-positive breast cancer: planned joint analysis of overall survival from NSABP B-31 and NCCTG N9831. *J Clin Oncol*. 2014;32(33):3744–3752.
- Xuan Q, Ji H, Tao X, et al. Quantitative assessment of HER2 amplification in HER2-positive breast cancer: its association with clinical outcomes. *Breast Cancer Res Treat*. 2015;150(3):581–588.
- Borley A, Mercer T, Morgan M, et al. Impact of HER2 copy number in IHC2+/FISH-amplified breast cancer on outcome of adjuvant trastuzumab treatment in a large UK cancer network. *Br J Cancer*. 2014;110(8):2139–2143.
- Dowsett M, Procter M, McCaskill-Stevens W, et al. Disease-free survival according to degree of HER2 amplification for patients treated with adjuvant chemotherapy with or without 1 year of trastuzumab: the HERA Trial. *J Clin Oncol*. 2009;27(18):2962–2969.
- Xu QQ, Pan B, Wang CJ, et al. HER2 amplification level is not a prognostic factor for HER2-positive breast cancer with trastuzumab-based adjuvant treatment: a systematic review and meta-analysis. *Oncotarget*. 2016;7(39):63571–63582.
- Fuchs EM, Kostler WJ, Horvat R, et al. High-level ERBB2 gene amplification is associated with a particularly short time-to-metastasis, but results in a high rate of complete response once trastuzumab-based therapy is offered in the metastatic setting. *Int J Cancer*. 2014;135(1):224–231.
- Gullo G, Bettio D, Torri V, Masci G, Salvini P, Santoro A. Level of HER2/neu gene amplification as a predictive factor of response to trastuzumab-based therapy in patients with HER2-positive metastatic breast cancer. *Invest New Drugs*. 2009;27(2):179–183.
- Kim JW, Kim JH, Im SA, et al. HER2/CEP17 ratio and HER2 immunohistochemistry predict clinical outcome after first-line trastuzumab plus taxane chemotherapy in patients with HER2 fluorescence in situ hybridization-positive metastatic breast cancer. *Cancer Chemother Pharmacol*. 2013;72(1):109–115.
- Arnould L, Arveux P, Couturier J, et al. Pathologic complete response to trastuzumab-based neoadjuvant therapy is related to the level of HER-2 amplification. *Clin Cancer Res*. 2007;13(21):6404–6409.
- Yu KD, Liu GY, Zhou XY, et al. Association of HER-2 copy number and HER-2/CEP-17 ratio with neoadjuvant taxane-containing chemotherapy sensitivity in locally advanced breast cancer. *Oncologist*. 2012;17(6):792–800.
- Buzdar AU, Valero V, Ibrahim NK, et al. Neoadjuvant therapy with paclitaxel followed by 5-fluorouracil, epirubicin, and cyclophosphamide chemotherapy and concurrent trastuzumab in human epidermal growth factor receptor 2-positive operable breast cancer: an update of the initial randomized study population and data of additional patients treated with the same regimen. *Clin Cancer Res*. 2007;13(1):228–233.
- Wolff AC, Hammond ME, Hicks DG, et al. Recommendations for human epidermal growth factor receptor 2 testing in breast cancer: American Society of Clinical Oncology/College of American Pathologists clinical practice guideline update. *Arch Pathol Lab Med*. 2014;138(2):241–256.
- Guiu S, Gauthier M, Coudert B, et al. Pathological complete response and survival according to the level of HER-2 amplification after trastuzumab-based neoadjuvant therapy for breast cancer. *Br J Cancer*. 2010;103(9):1335–1342.
- Wu K, Yang Q, Liu Y, Wu A, Yang Z. Meta-analysis on the association between pathologic complete response and triple-negative breast cancer after neoadjuvant chemotherapy. *World J Surg Oncol*. 2014;12:95.
- Liedtke C, Mazouni C, Hess KR, et al. Response to neoadjuvant therapy and long-term survival in patients with triple-negative breast cancer. *J Clin Oncol*. 2008;26(8):1275–1281.
- Kogawa T, Fouad TM, Liu DD, et al. High HER2/centromeric probe for chromosome 17 fluorescence in situ hybridization ratio predicts pathologic complete response and survival outcome in patients receiving neoadjuvant systemic therapy with trastuzumab for HER2-overexpressing locally advanced breast cancer. *Oncologist*. 2016;21(1):21–27.
- Musulino A, Naldi N, Bortesi B, et al. Immunoglobulin G fragment C receptor polymorphisms and clinical efficacy of trastuzumab-based therapy in patients with HER-2/neu-positive metastatic breast cancer. *J Clin Oncol*. 2008;26(11):1789–1796.
- Gavin PG, Song N, Kim SR, et al. Association of polymorphisms in FCGR2A and FCGR3A with degree of trastuzumab benefit in the adjuvant treatment of ERBB2/HER2-positive breast cancer: analysis of the NSABP B-31 trial. *JAMA Oncol*. 2017;3(3):335–341.
- Norton N, Olson RM, Pegram M, et al. Association studies of Fcγ receptor polymorphisms with outcome in HER2+ breast cancer patients treated with trastuzumab in NCCTG (Alliance) trial N9831. *Cancer Immunol Res*. 2014;2(10):962–969.
- Boero S, Morabito A, Banelli B, et al. Analysis of in vitro ADCC and clinical response to trastuzumab: possible relevance of FcγRIIIA/FcγRIIA gene polymorphisms and HER-2 expression levels on breast cancer cell lines. *J Transl Med*. 2015;13:324.
- Wolff AC, Hammond ME, Schwartz JN, et al. American Society of Clinical Oncology/College of American Pathologists guideline recommendations for human epidermal growth factor receptor 2 testing in breast cancer. *Arch Pathol Lab Med*. 2007;131(1):18–43.

### OncoTargets and Therapy

### Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

Dovepress

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.