

Diagnostic and prognostic value of microRNAs in cholangiocarcinoma: a systematic review and meta-analysis

Chao Sun¹
 Jie Zhu¹
 Bin Wu¹
 Jianlei Chen¹
 Zhenwei Zhu¹
 Peng Cai¹
 Wanliang Guo²
 Zhicheng Gu¹
 Jian Wang¹
 Shungen Huang¹

¹General Surgery Department, Children's Hospital of Soochow University, Suzhou, 215003, People's Republic of China; ²Radiology Department, Children's Hospital of Soochow University, Suzhou, 215003, People's Republic of China

Correspondence: Shungen Huang
 Department of General Surgery,
 Children's Hospital of Soochow
 University, Number 92 Zhongnan Street,
 Suzhou, 215003, People's Republic of
 China
 Tel +86 137 7608 8680
 Email drhuang01@163.com

Background and aim: Several dysregulated microRNAs (miRNAs) have been implicated in the pathogenesis of cholangiocarcinoma (CCA); however, small sample sizes and invariable research designs are limitations, hindering a thorough analysis of miRNAs as diagnostic and prognostic tools for CCA. This study aimed to systematically summarize the clinical value of miRNAs in human CCA both for all available miRNAs and single miRNA with multiple researches.

Methods: Pooled parameters included the area under the curve (AUC), sensitivity, specificity, and hazard ratios (HRs) to separately determine overall diagnostic and prognostic performance. Subgroup and sensitivity analyses were performed only in the event of heterogeneity. Thirty-four studies including 12 diagnostic studies and 22 prognostic studies were eligible for inclusion in this meta-analysis.

Results: We observed that miR-21, miR-26, miR-483, miR-106a, miR-150, miR-192, and miR-194 were employed for distinguishing patients with CCA from healthy controls. Pooled sensitivity, specificity, and AUC were 0.82 (95% confidence interval [CI] 0.77–0.86), 0.83 (95% CI 0.75–0.89), and 0.88 (95% CI 0.85–0.91), respectively. Abnormal expression of miR-21, miR-26a, miR-192, miR-200c, miR-221, miR-29a, miR-191, miR-181c, miR-34a, miR-106a, miR-203, and miR-373 in patients was confirmed to associate with poor survival rate. Pooled HRs and 95% CIs were calculated using STATA, resulting in the pooled HR of 1.47 (95% CI 0.91–2.37) for overall survival (OS), 0.67 (95% CI 0.16–2.81) for disease-free survival (DFS), 2.31 (95% CI 1.59–3.36) for progression-free survival (PFS), and 2.68 (95% CI 0.88–8.15) for relapse-free survival (RFS). Thus, CCA patients with dysregulated miRNA expression were confirmed to have shorter OS, DFS, PFS, and RFS. Data regarding the diagnostic and prognostic roles of miR-21 suggested pooled diagnostic results of miR-21 for sensitivity, specificity, and AUC were 0.85 (95% CI 0.76–0.91), 0.92 (95% CI 0.81–0.97), and 0.93 (95% CI 0.91–0.95), respectively, suggesting better diagnostic performance of miR-21 compared with other miRNAs. Meanwhile, pooled prognostic result of miR-21 for HR was 1.88 (95% CI 1.41–2.51), indicating miR-21 could more appropriately predict shorter OS in patients with CCA.

Conclusion: miRNAs may provide a new approach for clinical application, and miR-21 may be a promising biomarker for diagnosis and prognosis of CCA.

Keywords: microRNAs, miR-21, diagnosis, prognosis, cholangiocarcinoma, meta-analysis

Introduction

Cholangiocarcinoma (CCA) is a rare biliary malignancy prone to lymphatic metastasis. CCA is increasingly common, and currently, it is the most frequent primary hepatic malignancy.^{1,2} In some Asian countries, the elevated incidence of CCA appears to correlate with liver fluke *Opisthorchis viverrini* (*Ov*) infections. Liver fluke infections

subsequently result in chronic inflammation of the biliary tree and malignant transformation.^{3,4} Although the accuracy of current diagnostic methods for cancer has greatly improved, most patients with CCA are diagnosed at an unresectable stage of the disease, leading to a 5-year overall survival (OS) of <30%.⁵ There are currently only few diagnostic techniques for detecting early-stage CCA, and also, biomarker studies available are limited;^{6,7} as such, novel biomarkers for diagnosis and prognosis are needed.⁸

MicroRNAs (miRNAs) are endogenous small (~22 nt) noncoding RNAs that contribute to cell fate determination, proliferation, and cell death.^{9–11} Several studies indicate that miRNAs can be stably expressed in human plasma and serum, and miRNAs have also been demonstrated to be abnormally expressed in the circulatory system during tumorigenesis and inflammation.^{12,13} Kishimoto et al¹⁴ found that miR-21 was overexpressed in bile duct cancer (accuracy, 0.89; sensitivity, 84%; and specificity, 98%), indicating that it may be used as a biomarker for distinguishing bile duct cancer patients from normal subjects. Silakit et al¹⁵ confirmed high expression of miR-192 in sera of CCA patients, which predicted worse OS compared with those with low miR-192 expression. CA19-9 has been frequently used to diagnose and predict CCA patient outcomes.¹⁶ However, low sensitivity or low specificity prevents CA19-9 from being adopted globally.¹⁷ Therefore, identifying miRNAs unique to CCA may provide suitable biomarkers for detection and prognosis.

Methods

Search strategy

We searched PubMed, EMBASE, and the Web of Science for studies of associations between miRNAs and CCA patients. We used these search terms: (microRNA OR miRNA OR miR) and (cholangiocarcinoma OR bile duct cancer) and (diagnosis OR prognosis). We screened titles, abstracts, author information, and results. Studies were included if they focused on miRNAs in CCA patients identified by diagnosis

of histopathological confirmation, provided information about the relationship between miRNAs and diagnosis (summary receiver operating characteristic [SROC] curves or sensitivity/specificity) or prognosis (hazard ratio [HR] with 95% confidence interval [CI] or Kaplan–Meier curves), included healthy individuals as controls, and had clearly defined miRNA thresholds. Studies were excluded if they met the following criteria: reviews, letters, commentaries, or meeting records; duplicate publications; used cell lines; combined two or more miRNAs to calculate overall sensitivity/specificity or HR; and lacked sufficient data to evaluate HR and 95% CIs. Three individual investigators screened and evaluated retrieved articles. Disagreement was resolved with detailed discussion.

Quality assessment

This meta-analysis had diagnostic and prognostic components, so the Quality Assessment of Diagnostic Accuracy Studies 2 (QUADAS-2) and the Meta-analysis of Observational Studies in Epidemiology group (MOOSE) were used to evaluate diagnostic and prognostic accuracy of eligible studies.^{18,19} If studies did not meet these criteria, they were excluded. Bias risk and applicability concerns are detailed in Figure 1.

Data extraction

Two investigators independently examined each eligible study and its data using a standard protocol. We assessed first author, year of publication, study design, subject number, tumor stage, method of quantifying miRNA expression, and miRNA thresholds to stratify high- and low-subject groups. For diagnostic studies, sensitivity, specificity, true positives (TPs), false positives (FPs), false negatives (FN), and true negatives (TN) were included. For prognostic studies, HRs for OS or disease-free survival (DFS) or progression-free survival (PFS) or relapse-free survival (RFS), and 95% CIs and *p* values were extracted. If HR and relevant parameters




Figure 1 A graph of risk of bias and applicability concerns: a review of authors' judgments about each domain is presented as percentages across included studies.

were not available, we estimated HRs and 95% CIs using the methods of Parmar et al.²⁰

Statistical analysis

For diagnostic analysis, TPs, FPs, FNs, and TNs were extracted from included studies or recalculated based on the prevalence and sample size of each study. A bivariate model was used to calculate pooled sensitivity, specificity, positive and negative likelihood ratios (LRs), diagnostic odds ratios (DORs), and their 95% CIs.²¹ SROC curves were plotted by pooling sensitivity and specificity of each study to evaluate diagnostic effects, along with auto-generation of AUCs of SROC curves and the maximum point of intersection between sensitivity and specificity (Q value). Heterogeneity was examined using an I^2 test ($p < 0.1$ or $I^2 > 50\%$ indicated significant heterogeneity).²² Subgroup analysis was performed to identify potential heterogeneity sources, and Deeks' funnel plot was used to inspect publication bias ($p < 0.1$ indicated statistically significant publication bias).

For prognostic analysis, multivariate analysis was employed to avoid confounding of exposure effects.²³ Heterogeneity assessment of combined HRs was performed using Cochran's Q test and Higgins I^2 statistic ($p < 0.10$ and/or $I^2 > 50\%$ meant significant heterogeneity and the need for a fixed model). Otherwise, a random-effects model

was used, and subgroup analysis was conducted to identify the source of heterogeneity. An observed HR > 1 indicated elevations of miRNA were detrimental and correlated with worse survival, but HR < 1 suggested lows of miRNA were harmful and related to poor survival. Furthermore, Begg's funnel plots and Egger's linear regression tests were used to evaluate publication bias.²⁴ All analyses were performed with Stata 12.0 (Stata Corporation, College Station, TX, USA).

Results

As shown in Figure 2, a comprehensive search was performed on 30/04/2017, yielding a total of 373 results from the PubMed, EMBASE, and Web of Science. Among these results, 275 articles were identified as duplicates and irrelevant studies including reviews, textbooks, letters, or animal research after the screening of titles and abstract. The remaining 98 articles meeting the inclusion criteria were further thoroughly assessed. Finally, we found that 64 articles lacked sufficient data to allow for a meta-analysis or were not relevant to diagnoses or prognoses. Therefore, 28 articles containing 34 studies (including 12 for diagnostic analysis and 22 for prognostic analysis) focusing on the relationship between miRNAs and CCA were utilized for the final analysis. The main characteristics of each article are summarized in Tables 1 and 2.




Figure 2 A flowchart of literature search and study selection. Twenty-three case-control studies including 9 studies for diagnoses and 14 studies for prognoses were included in this meta-analysis.

Table 1 Main characteristics of the diagnostic studies included in the meta-analysis

Study	Year	miR	Country	Race	Specimen	Sample size		Sensitivity	Specificity	AUC	TP	FP	FN	TN	RNA extraction	Measurements
						Case	Control									
Wang et al ¹⁹	2015	26a	People's Republic of China	Asian	Serum	66	66	84.80%	81.80%	0.899	56	12	10	54	Qiagen miRNeasy kit	TaqMan
Wang et al ²⁰	2015	150	People's Republic of China	Asian	Plasma	15	15	80.60%	58.10%	0.764	13	3	2	12	Qiagen miRNeasy kit	SYBR
Silakit et al ²⁷	2015	192	Thailand	Asian	Urine	22	21	63.60%	66.70%	0.682	14	7	8	14	Qiagen miRNeasy kit	TaqMan
Silakit et al ²⁷	2015	21	Thailand	Asian	Urine	22	21	63.60%	71.40%	0.682	14	6	8	15	Qiagen miRNeasy kit	TaqMan
Wang et al ²⁸	2015	21	People's Republic of China	Asian	Serum	74	74	87.80%	90.50%	0.908	65	7	9	67	Qiagen miRNeasy kit	TaqMan
Kishimoto et al ¹⁴	2013	21	Japan	Asian	Plasma	94	50	85.10%	100.00%	0.930	80	1	14	49	miRVana PARIS kit	TaqMan
Cheng et al ¹⁷	2015	106a	People's Republic of China	Asian	Serum	103	20	81.60%	85.00%	0.890	84	3	19	17	Qiagen miRNeasy kit	TaqMan
Silakit et al ¹⁵	2014	192	Thailand	Asian	Serum	51	32	74.00%	72.00%	0.803	38	9	13	23	Qiagen miRNeasy kit	TaqMan
Selaru et al ³⁶	2009	21	USA	Caucasian	Tissue	20	14	95.00%	92.90%	0.995	19	1	1	13	TRIzol	TaqMan
Bernuzzi et al ³⁵	2016	483	Italy	Caucasian	Serum	30	30	93.33%	66.67%	0.770	28	10	2	20	miRVana PARIS kit	TaqMan
Bernuzzi et al ³⁵	2016	194	Italy	Caucasian	Serum	30	30	76.67%	73.33%	0.740	23	8	7	22	miRVana PARIS kit	TaqMan
Correa-Gallego et al ⁵¹	2016	21	USA	Caucasian	Plasma	25	7	88.00%	100.00%	0.940	22	0	3	7	miRVana PARIS kit	TaqMan

Abbreviations: AUC, area under the curve; FN, false negative; FP, false positive; TN, true negative; TP, true positive.

Table 2 Main characteristics of the prognostic studies included in the meta-analysis

Study	Year	miR	Population	Patient	Method	Cut-off	Sample	Survival	Follow-up	HR	95% CI	RNA extraction	Measurements	p-value
Silakit et al ¹⁵	2014	192	Thailand	51	qRT-PCR	2.0-fold	Serum	OS	78	2.520	1.306–4.85	Qiagen miRNeasy kit	TaqMan	0.006
Huang et al ³⁹	2013	21	People's Republic of China	41	qRT-PCR	Mean	Tissue	OS	42	1.998	1.156–3.456	DIG tailing kit	TaqMan	0.013
Huang et al ³⁹	2013	21	People's Republic of China	41	qRT-PCR	Mean	Tissue	RFS	38	1.825	1.041–3.202	DIG tailing kit	TaqMan	0.036
Qiao et al ⁴⁸	2015	34a	People's Republic of China	27	qRT-PCR	Median	Tissue	RFS	43	0.272	0.132–0.562	TRIZOL	SYBR	0.000
Wang et al ³⁸	2015	21	People's Republic of China	57	qRT-PCR	Quartile	Tissue	OS	60	1.588	1.064–2.371	Qiagen miRNeasy kit	TaqMan	0.024
Wang et al ³⁸	2015	21	People's Republic of China	57	qRT-PCR	Quartile	Tissue	PFS	60	1.888	1.117–3.191	Qiagen miRNeasy kit	TaqMan	0.018
Chu et al ⁵²	2015	200c	People's Republic of China	254	qRT-PCR	7.2-fold	Tissue	OS	107	1.926	1.156–3.209	miRVana PARIS kit	TaqMan	0.012
Cheng et al ¹⁷	2015	106a	People's Republic of China	103	qRT-PCR	1.0-fold	Serum	OS	71	0.200	0.106–0.38	TRIZOL	TaqMan	0.000
Chen et al ⁵³	2010	373	People's Republic of China	48	qRT-PCR	2.94-fold	Tissue	OS	43	0.205	0.091–0.463	miRVana PARIS kit	TaqMan	0.000
Chen et al ⁵³	2010	373	People's Republic of China	48	qRT-PCR	2.94-fold	Tissue	DFS	43	0.125	0.047–0.337	miRVana PARIS kit	TaqMan	0.000
Li et al ⁴⁷	2015	221	People's Republic of China	25	qRT-PCR	Median	Tissue	DFS	42	2.540	0.960–6.740	TRIZOL	SYBR	0.000
Li et al ⁴⁷	2015	221	People's Republic of China	25	qRT-PCR	Median	Tissue	PFS	43	2.419	1.290–4.536	TRIZOL	SYBR	0.006
Li et al ⁵⁴	2015	203	People's Republic of China	148	qRT-PCR	Mean	Tissue	OS	84	0.532	0.313–0.905	TRIZOL	MJ Mini PCR	0.02
Wang et al ⁴⁹	2015	26a	People's Republic of China	66	qRT-PCR	Mean	Serum	OS	60	3.461	1.331–5.364	Qiagen miRNeasy kit	TaqMan	0.013
Wang et al ⁴⁹	2015	26a	People's Republic of China	66	qRT-PCR	Mean	Serum	PFS	60	4.226	1.415–10.321	Qiagen miRNeasy kit	TaqMan	0.000
Chusorn et al ⁵⁵	2013	21	Thailand	23	qRT-PCR	Mean	Tissue	OS	60	2.332	1.166–4.664	TRIZOL	TaqMan	0.017
Deng and Chen ⁵⁶	2017	29a	People's Republic of China	125	qRT-PCR	Mean	Tissue	OS	60	3.508	1.772–6.947	TRIZOL	SYBR	0.000
Li et al ⁵⁷	2017	191	People's Republic of China	84	qRT-PCR	Mean	Tissue	OS	92	2.398	1.562–3.683	TruSeq RNA kit	TruSeq kit	0.000
Li et al ⁵⁷	2017	191	People's Republic of China	84	qRT-PCR	Mean	Tissue	DFS	92	2.267	1.124–4.571	TruSeq RNA kit	TruSeq kit	0.022
Wang et al ⁵⁸	2016	181c	People's Republic of China	62	RT-PCR	Mean	Tissue	OS	80	1.797	1.053–3.064	Qiagen miRNeasy kit	TaqMan	0.031
Garajova et al ⁵⁹	2015	21	Italy	41	RT-PCR	Mean	Tissue	OS	32	4.950	1.070–22.980	Qiagen miRNeasy kit	TaqMan	0.041
Garajova et al ⁵⁹	2015	21	Italy	41	RT-PCR	Mean	Tissue	RFS	32	6.210	1.290–29.890	Qiagen miRNeasy kit	TaqMan	0.023

Abbreviations: CI, confidence interval; DFS, disease-free survival; HR, hazard ratio; OS, overall survival; PFS, progression-free survival; RFS, relapse-free survival; RT-PCR, reverse transcription PCR; qRT-PCR, quantitative reverse transcription PCR.

Diagnostic and prognostic value of miRNAs for CCA

Twelve studies containing 552 patients and 380 healthy controls evaluated the diagnostic value of miRNAs for CCA. Various miRNAs were detected and shown to be associated with the diagnosis of CCA, including miR-21 (n = 5), miR-192 (n = 2), miR-26 (n = 1), miR-194 (n = 1), miR-106a (n = 1), miR-150 (n = 1), and miR-482 (n = 1). Out of all the investigations, eight studies were conducted in Asian populations, and the remaining four studies were performed in Caucasian populations. Different samples including serum (n = 6), plasma (n = 3), urine (n = 2), and tissue (n = 1) were used in all these different studies. All studies used quantitative reverse transcription PCR (qRT-PCR) to measure the expression of miRNAs. Sensitivity, specificity, and DOR data are shown in Figure 3A, indicating that miRNAs have relatively high diagnostic value for CCA. I^2 data are shown in Figure 4A. Meta-regression analysis revealed that plasma may be a source of heterogeneity, while other covariates, such as RNA extraction, measurements, specimen, and different races, may not contribute to heterogeneity (all $p > 0.05$) (Table 3). Pooled positive and negative LR data are shown in Figure 5A. Studies that included 1,155 patients from the People's Republic of China, Italy, and Thailand were collected, and 12 miRNAs were identified in CCA patients. Tissue and sera were sample sources, and 13 reports assessed the correlations between abnormal miRNA expression and OS, while four studies assessed the relationship between miRNAs expression and DFS. Limited studies were included for PFS (n = 3) and RFS (n = 2); thus, pooled HRs were respectively analyzed in the systemic review but without further heterogeneity analysis.

Increased expression of miR-21, miR-26a, miR-192, miR-200c, miR-191, miR-181c, miR-29a, and miR-221 was associated with a poor prognosis, as was decreased expression of miR-34a, miR-106a, miR-203, and miR-373 (Table 2). Heterogeneity data are shown in Figure 6A. A random-effects model was applied for the OS, DFS, and RFS subgroups, and the data are given in Table 4. Through subgroup analysis of OS determined by different elements, we found that the up-downregulation of miRNAs might be the source of heterogeneity ($p = 0.001$), while other factors (such as sample type, sample size, RNA extraction, and different measurements) may not lead to significant heterogeneity (all $p > 0.05$) (Table 4). The pooled results indicated that miRNAs were significant prognostic biomarkers for CCA patients (OS: HR = 1.47, 95% CI 0.91–2.37, n = 13; DFS: HR = 0.67, 95% CI 0.16–2.81, n = 4; PFS: HR = 2.31, 95%

CI 1.59–3.36, n = 3; RFS: HR = 1.46, 95% CI 1.00–2.14, n = 2). Sensitivity analysis data are shown in Figure 7. No obvious publication bias was found in the quantitative synthesis when Begg's funnel plot and Egger's test were applied to assess publication bias ($p = 0.326$) (Figure 8A).

Diagnostic and prognostic values of miR-21 for CCA

Notably, we found that abnormal expression of miR-21 was identified by five studies for diagnostic analysis and four studies for prognostic analysis of all included studies. Thus, we had to conduct a meta-analysis for evaluating the potential diagnostic and prognostic value of miR-21 for CCA.

Five studies containing 235 patients assessed the diagnostic value of miR-21 for CCA. The pooled sensitivity, specificity, positive LR (PLR), negative LR (NLR), and DOR were 0.85 (95% CI 0.76–0.91), 0.92 (95% CI 0.81–0.97), 10.9 (95% CI 4.0–29.5), 0.17 (95% CI 0.10–0.28), and 66 (95% CI 16–272), respectively, indicating that miR-21 has a higher diagnostic accuracy than the pooled miRNAs for CCA patients (Figure 4B). Additionally, the AUC was 0.93 (95% CI 0.91–0.95), indicating miR-21 has a stronger diagnostic value for CCA when compared with the pooled miRNAs (Figure 3B). Furthermore, there was no significant publication bias in this meta-analysis (Figure 5B).

Meanwhile, we performed the prognostic analysis on the relationship of miR-21 and the OS of CCA patients (n = 4) (Figure 6B). As heterogeneity among all studies was not observed (OS: $I^2 = 0.0\%$; $p = 0.452$), a fixed-effects model was used. We found significant association between abnormal miR-21 expression and poor OS (HR = 1.88, 95% CI 1.41–2.51). Publication bias was evaluated with funnel plots and Begg's tests. No significant publication biases were found (Figure 8B).

Discussion

CCA is the second most common primary hepatic malignancy arising from the bile duct epithelium,^{7,25} but it has few distinguishable symptoms, and patients are mostly diagnosed at the advanced tumor phase, resulting in the stubbornly high mortality in the recent years.^{26–28} Although the OS of patients with CCA is dismal, there is a large discrepancy between patients diagnosed at early and late stages, indicating an urgent need for novel diagnostic and prognostic biomarkers.²⁹

Current biomarkers for CCA have low sensitivity and specificity. Ince et al³⁰ constructed a receiver operating characteristic (ROC) curve and collected optimal thresholds for serum and biliary CA19-9 (serum: sensitivity, 49%; specific-




Figure 3 SROC curves describing the diagnostic performance of miRNAs in discriminating cancer patients from healthy subjects: **(A)** miRNAs and **(B)** miR-21. **Abbreviations:** AUC, area under the curve; miRNA, microRNA; SENS, sensitivity; SPEC, specificity; SROC, summary receiver operating characteristic.





Figure 4 Forest plots of sensitivity and specificity of each included publication: **(A)** miRNAs and **(B)** miR-21. **Abbreviations:** CI, confidence interval; miRNA, microRNA.

Table 3 Multivariate meta-regression analysis for the associations of miRNAs with susceptibility to various cancers

Results of subgroup and meta-regression analysis in the diagnosis meta-analysis										
	Sensitivity	95% CI		Specificity	95% CI		AUC	95% CI		Regression
Year										
Before 2010	0.95	–	–	0.93	–	–	0.99	–	–	0.2
After 2010	0.81	0.77	0.85	0.82	0.74	0.88	0.87	0.84	0.9	
Race										
Asian	0.8	0.74	0.85	0.83	0.74	0.9	0.87	0.84	0.9	0.13
Caucasian	0.88	0.83	0.94	0.8	0.58	0.92	0.92	0.89	0.94	
Sample size										
<100	0.81	0.72	0.89	0.75	0.67	0.81	0.81	0.78	0.85	0.07
≥100	0.85	0.8	0.88	0.9	0.81	0.95	0.87	0.83	0.89	
Specimen										
Serum	0.83	0.78	0.87	0.8	0.72	0.86	0.87	0.84	0.9	0.23
Plasma	0.86	0.78	0.91	0.95	0.76	0.99	0.87	0.84	0.90	0.05
Tissue	0.95	–	–	0.93	–	–	0.99	–	–	0.2
RNA extraction										
Qiagen miRNeasy kit	0.81	0.76	0.86	0.81	0.73	0.90	0.88	0.85	0.91	0.69
Others	0.85	0.78	0.91	0.85	0.75	0.96	0.86	0.83	0.89	
Measurements										
TaqMan	0.82	0.78	0.86	0.83	0.76	0.90	0.88	0.85	0.91	0.30
SYBR	0.87	0.69	1.00	0.81	0.53	1.00				

Note: The en-dashes indicate the 95% CI could not be calculated due to limited studies.

Abbreviations: AUC, area under the curve; CI, confidence interval; miRNA, microRNA.

ity, 84.5%; accuracy, 64%; biliary: sensitivity, 74%; specificity, 34%; accuracy, 69%) and noted the low sensitivity and specificity of these markers. They also reported sensitivity and specificity of biliary carcinoembryonic antigen (CEA) of 57% and 68%, respectively. Thus, CEA may be not ideal for distinguishing CCA from normal people. Huang et al³¹ identified elevated soluble fragments of cytokeratin 19 (CYFRA 21-1) from CCA patients sera and reported an AUC of 0.879 and a high specificity (96.2%) but a low sensitivity (75.6%). Therefore, traditional biomarkers are not suitable for improving the accuracy of diagnosis and prognosis.

miRNAs contribute to tumorigenesis by affecting cellular processes, including the cell cycle, angiogenesis, invasion, and metastasis.^{32–34} Because miRNAs are protected from RNases and remain stable in plasma and serum, they may have a potential role as biomarkers for early cancer detection.³⁵ However, studies focusing on diagnostic or prognostic values of miRNAs in CCA are inconsistent. For example, Selaru et al³⁶ performed qRT-PCR on 18 primary CCAs and 12 normal liver specimens and found that miR-21 was 95% sensitive and 100% specific in distinguishing between CCA and normal tissues (AUC 0.995). Silakit's report³⁷ revealed that the sensitivity and specificity of urine-derived miR-21 were 63.6% and 71.4%, respectively, for differentiating CCA patients from healthy controls (AUC 0.682). Wang et al³⁸ suggested that high tissue-derived miR-21 expression was associated with a higher risk of intrahepatic cholangiocarci-

noma (ICC) death compared to low tissue miR-21 expression and was an independent predictor of poor OS of ICC patients (HR = 3.519, 95% CI 1.411–5.702, $p = 0.021$). Huang et al³⁹ reported that CCA patients with high miR-21 expression had a mean 3-year OS of 15%, whereas patients with low miR-21 expression had a 33% 3-year OS (HR = 1.620, 95% CI 0.440–5.960, $p = 0.046$). Differences in specimen samples (ie, tissue or urine), different inclusion criteria, and different microarray techniques may explain discrepancies among these studies, but a lack of systematic evaluation complicates this conclusion. Knowledge on the diverse value of miRNAs is vague. Thus, we carried out this comprehensive and up-to-date research in a clinical context to precisely evaluate the diagnostic and prognostic accuracy of miRNAs.

In this systematic review, 12 diagnostic and 22 prognostic studies were included to study whether miRNAs are useful biomarkers for CCA. We noted that miRNAs had a relatively high diagnostic accuracy, and yielded a combined AUC of 0.88 (95% CI 0.85–0.91), with a pooled sensitivity of 0.82 and a pooled specificity of 0.83 for discriminating CCA cases. DOR, which combines the strength of sensitivity and specificity, is a useful indicator of diagnostic accuracy. In this meta-analysis, the DOR value for miRNAs was 22 (95% CI 11–43), indicating moderate diagnostic accuracy. However, the PLR and NLR were 4.8 (95% CI 3.1–7.3) and 0.22 (95% CI 0.16–0.29), respectively, suggesting that miRNAs may be insufficient in distinguishing patients with CCA because




Figure 5 Graph of Deeks' funnel plot asymmetry test: **(A)** miRNAs ($p = 0.30$) and **(B)** miR-21 ($p = 0.64$).
Abbreviations: ESS, effective sample size; miRNA, microRNA.

PLR > 10 and NLR < 0.1 are thresholds representing high accuracy. Subgroup and meta-regression analyses were applied if heterogeneity occurred in the diagnostic meta-analysis. We compared factors that may influence the heterogeneity and suggest that plasma specimens may be the chief sources of heterogeneity ($p = 0.05$). Tumor-derived circulating miRNA may have utility as a noninvasive blood biomarker, and both the sensitivity and specificity of blood-based assays (serum and plasma) exceeded 0.8, so they may have a high diagnostic value.

Elevated expression of miRNAs (miR-21, miR-26a, miR-29a, miR-181c, miR-191, miR-192, miR-200c, and miR-221) was associated with poor survival of CCA patients, and

decreased expression of miRNAs (miR-34a, miR-106a, miR-203, and miR-373) was associated with a worse prognosis. Pooled HR values of OS correlated with miRNA expression for CCA patients, and this suggested that specific miRNAs are independent risk factors for prognosis and may have use for clinical decision-making. The forest plot revealed heterogeneity in this meta-analysis ($I^2 = 88.1\%$; $p < 0.001$), so we performed meta-regression analysis to explore the source. Results showed that the up-downregulation of miRNAs was significantly related to heterogeneity ($p = 0.001$) and may partly explain the heterogeneity of the OS analysis. We also evaluated the relationship between miRNA expression and RFS, DFS, and PFS. The pooled HR of PFS from the

A




Figure 6 (Continued)

B




Figure 6 Forest plots of studies evaluating miRNAs expression level and cancer prognosis: (A) miRNAs and (B) miR-21.

Note: “↑” and “↓” indicate that elevated and decreased expression of microRNAs correlate with poor survival rate.

Abbreviations: CI, confidence interval; DFS, disease-free survival; HR, hazard ratio; miRNA, microRNA; OS, overall survival; PFS, progression-free survival; RFS, relapse-free survival.

Table 4 Subgroup analysis for association of miRNAs with OS in CCA

Variables	Coefficient	Standard error	t	p value	95% CI	
Up-down	-2.055098	0.344408	-5.97	0.001	-2.869494	-1.240703
Sample type	-0.0456237	0.3268338	-0.14	0.893	-0.8184629	0.7272155
Year	0.115651	0.3840489	0.3	0.772	-0.7924804	1.023782
Sample size	0.0126298	0.3844345	0.03	0.975	-0.8964135	0.921673
RNA extraction	0.7983387	0.4680503	1.71	0.12	-0.2445424	1.84122
Measurements	-0.4936747	0.5209893	-0.95	0.37	-1.654511	0.6671618

Abbreviations: CCA, cholangiocarcinoma; CI, confidence interval; miRNA, microRNA; OS, overall survival.




Figure 7 Sensitivity analyses assessing the influence of individual studies on the pooled abnormal miRNAs expression in OS subgroup.

Abbreviations: CI, confidence interval; miRNA, microRNA; OS, overall survival.




Figure 8 Begg's funnel plots for the assessment of publication bias in the meta-analysis for prognosis: **(A)** miRNAs ($p = 0.326$) and **(B)** miR-21 ($p = 0.174$).
Abbreviations: HR, hazard ratio; miRNA, microRNA; SE, standard error.

included studies was 2.31 (95% CI 1.59–3.36), indicating high expression of miRNAs might predict poor PFS for cancer patients. The included studies had low heterogeneity ($I^2 = 0.4\%$; $p = 0.366$), and the pooled HRs of DFS and RFS were 0.67 (0.16–2.81) and 2.68 (0.88–8.15), respectively, which showed that miRNAs can be used to monitor the therapeutic effects of radical resection or chemotherapy. However, due to limited studies, meta-regression and subgroup analyses for DFS and RFS subgroups were not performed, although there was significant heterogeneity.

Several miRNAs were confirmed to be associated with CCA patient variables, but most were assessed in one study, and only miR-21 was reported in at least three studies, suggesting it requires more investigation. For the diagnostic meta-analysis, four studies involving 210 patients and 159 healthy controls were included, and the pooled sensitivity and specificity of miR-21 were 0.85 (95% CI 0.76–0.91) and 0.92 (95% CI 0.81–0.97), respectively, with an ROC of 0.93 (95% CI 0.91–0.95). Compared with the pooled parameters of miRNAs, miR-21 had a higher diagnostic accuracy than other miRNAs and a better DOR (53, 95% CI 12–239), NLR (0.18, 95% CI 0.10–0.32), and PLR (9.4, 95% CI 3.5–25.4), indicating miR-21 was adequate to distinguish CCA patients from normal subjects. Data showed that patients with increased miR-21 had a higher risk of poor survival compared to those with low miR-21 (HR 1.88, 95% CI 1.41–2.51). There was no statistical heterogeneity among the included studies, so miR-21 may be an ideal prognostic marker for clinical decision-making. In conclusion, these findings proved that miR-21 could be a more suitable clinical biomarker with a higher capacity for discriminating CCA patients and healthy people. Exploring the sources of heterogeneity is necessary in a meta-analysis.

In the present study, different measures such as sample size, ethnicity, RNA extraction, and measurement methods were used to extract miRNAs in different studies. However, there is no evidence that these variables may influence the heterogeneity. In addition, high expression of miR-21 has been found to relate with hepatocellular carcinoma (HCC).⁴⁰ Our study further concluded that miR-21 correlate with diagnosis and prognosis of CCA with numerous miRNAs. It may not be possible for miR-21 to distinguish HCC and CCA. Thus, clinical features and techniques such as physical examination and CT/MRI scanning may need to be combined to differentiate CCA from HCC.⁴¹

miRNAs have been implicated in the pathogenesis of CCA^{42–45} because upregulated miRNAs suggest an oncogenic role and downregulation of miRNAs may be suppressive, so dysregulated miRNAs are correlated with the adverse clinical features, diminished survival, and poor prognosis of CCA patients.⁴⁶ Selaru et al³⁶ proved that miR-21 was oncogenic by inhibiting PDCD4 and TIMP3 suppressor genes in the biliary tree. Wang et al³⁸ confirmed the functional and mechanistic links between miR-21 and tumor suppressor genes (such as PTPN14 and PTEN) in the pathogenesis of ICC, which influenced the proliferation and tumor progression in ICC. Li et al⁴⁷ reported that miR-221 promoted extrahepatic CCA (EHCC) invasion and metastasis by targeting PTEN and formed a positive feedback loop with the β -catenin/c-Jun signaling pathways. Moreover, results from Qiao's study⁴⁸ suggested that miR-34a inhibited invasion and migration by targeting Smad4 to suppress the epithelial–mesenchymal transition via the TGF- β /Smad signaling pathway in human EHCC. In conclusion, altered miRNA expression is a key to tumorigenesis and can be used to identify clinicopathologic features of the disease.

Our study was limited by finding miR-21 in four studies, but up to 17 miRNAs were identified to be valuable for diagnosis or prognosis, and study heterogeneity occurred. Subgroup analysis was performed to find the source of heterogeneity, but we could not fully explain it. Likely differences in patients' baseline characteristics (ethnicity, gender, age, and tumor stage and grade), different thresholds for miRNA expression, and the way samples were prepared and preserved (ie, paraffin-fixed, formalin-fixed, freshly frozen tumors, or blood) made a difference. In addition, some HRs were calculated based on data extracted from survival curves, which might be less powerful than data obtained from articles directly.

Despite these limitations, our review had several important strengths. First, a relatively thorough systematic search was performed, and diagnostic and prognostic values of miRNAs in CCA were independently evaluated and verified. Our methods were rigorous and followed guidelines for conducting and reporting systematic reviews. Besides, further analysis and research was carried out to attest the diagnostic and prognostic roles of miR-21 expression in CCA patients.

Conclusion

miRNAs, especially miR-21, were identified to be highly correlated with CCA and could be potential and promising biomarkers in distinguishing CCA patients from healthy people; they could also contribute to predicting the progression of CCA. Furthermore, to better understand and use miRNAs as biomarkers in clinical detection, more large-scale and high-quality investigations are needed to confirm our results.

Acknowledgments

We thank the authors of all the included studies. This work was supported by a grant from the Science and Technology Project of Suzhou (SYS201764).

Disclosure

The authors report no conflicts of interest in this work.

References

- Siegel R, Ma J, Zou Z, Jemal A. Cancer statistics, 2014. *CA Cancer J Clin*. 2014;64(1):9–29.
- Zhang GW, Lin JH, Qian JP, Zhou J. Identification of risk and prognostic factors for patients with clonorchiasis-associated intrahepatic cholangiocarcinoma. *Ann Surg Oncol*. 2014;21(11):3628–3637.
- Sriamporn S, Pisani P, Pipitgool V, Suwanrungruang K, Kamsa-ard S, Parkin DM. Prevalence of *Opisthorchis viverrini* infection and incidence of cholangiocarcinoma in Khon Kaen, Northeast Thailand. *Trop Med Int Health*. 2004;9(5):588–594.
- Chamadol N, Pairojkul C, Khuntikeo N, et al. Histological confirmation of periductal fibrosis from ultrasound diagnosis in cholangiocarcinoma patients. *J Hepatobiliary Pancreat Sci*. 2014;21(5):316–322.
- Maithel SK, Gamblin TC, Kamel I, Corona-Villalobos CP, Thomas M, Pawlik TM. Multidisciplinary approaches to intrahepatic cholangiocarcinoma. *Cancer*. 2013;119(22):3929–3942.
- Andersen JB, Thorgeirsson SS. Genetic profiling of intrahepatic cholangiocarcinoma. *Curr Opin Gastroenterol*. 2012;28(3):266–272.
- Blechacz B, Gores GJ. Cholangiocarcinoma: advances in pathogenesis, diagnosis, and treatment. *Hepatology*. 2008;48(1):308–321.
- Brandi G, Venturi M, Pantaleo MA, Ercolani G; GICO. Cholangiocarcinoma: current opinion on clinical practice diagnostic and therapeutic algorithms: a review of the literature and a long-standing experience of a referral center. *Dig Liver Dis*. 2016;48(3):231–241.
- Lee YS, Dutta A. MicroRNAs in cancer. *Annu Rev Pathol*. 2009;4:199–227.
- Croce CM, Calin GA. miRNAs, cancer, and stem cell division. *Cell*. 2005;122(1):6–7.
- Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell*. 2004;116(2):281–297.
- Mitchell PS, Parkin RK, Kroh EM, et al. Circulating microRNAs as stable blood-based markers for cancer detection. *Proc Natl Acad Sci U S A*. 2008;105(30):10513–10518.
- Wang LG, Gu J. Serum microRNA-29a is a promising novel marker for early detection of colorectal liver metastasis. *Cancer Epidemiol*. 2012;36(1):e61–e67.
- Kishimoto T, Eguchi H, Nagano H, et al. Plasma miR-21 is a novel diagnostic biomarker for biliary tract cancer. *Cancer Sci*. 2013;104(12):1626–1631.
- Silakit R, Loilome W, Yongvanit P, et al. Circulating miR-192 in liver fluke-associated cholangiocarcinoma patients: a prospective prognostic indicator. *J Hepatobiliary Pancreat Sci*. 2014;21(12):864–872.
- Levy C, Lymp J, Angulo P, Gores GJ, Larusso N, Lindor KD. The value of serum CA 19-9 in predicting cholangiocarcinomas in patients with primary sclerosing cholangitis. *Dig Dis Sci*. 2005;50(9):1734–1740.
- Cheng Q, Feng F, Zhu L, et al. Circulating miR-106a is a novel prognostic and lymph node metastasis indicator for cholangiocarcinoma. *Sci Rep*. 2015;5:16103.
- Whiting P, Rutjes AW, Reitsma JB, Bossuyt PM, Kleijnen J. The development of QUADAS: a tool for the quality assessment of studies of diagnostic accuracy included in systematic reviews. *BMC Med Res Methodol*. 2003;3:25.
- Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA*. 2000;283(15):2008–2012.
- Parmar MK, Torri V, Stewart L. Extracting summary statistics to perform meta-analyses of the published literature for survival endpoints. *Stat Med*. 1998;17(24):2815–2834.
- Reitsma JB, Glas AS, Rutjes AW, Scholten RJ, Bossuyt PM, Zwinderman AH. Bivariate analysis of sensitivity and specificity produces informative summary measures in diagnostic reviews. *J Clin Epidemiol*. 2005;58(10):982–990.
- Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003;327(7414):557–560.
- Wu Y, Fu X, Zhu X, et al. Prognostic role of systemic inflammatory response in renal cell carcinoma: a systematic review and meta-analysis. *J Cancer Res Clin Oncol*. 2011;137(5):887–896.
- Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997;315(7109):629–634.
- Sirica AE. Cholangiocarcinoma: molecular targeting strategies for chemoprevention and therapy. *Hepatology*. 2005;41(1):5–15.
- Chu X, Zhao P, Lv Y, Liu L. Decreased expression of TFPI-2 correlated with increased expression of CD133 in cholangiocarcinoma. *Int J Clin Exp Pathol*. 2015;8(1):328–336.
- Wu T. Cyclooxygenase-2 and prostaglandin signaling in cholangiocarcinoma. *Biochim Biophys Acta*. 2005;1755(2):135–150.
- Nitta T, Sato Y, Ren XS, et al. Autophagy may promote carcinoma cell invasion and correlate with poor prognosis in cholangiocarcinoma. *Int J Clin Exp Pathol*. 2014;7(8):4913–4921.
- Patel T. Cholangiocarcinoma—controversies and challenges. *Nat Rev Gastroenterol Hepatol*. 2011;8(4):189–200.

30. Ince AT, Yıldız K, Baysal B, et al. Roles of serum and biliary CEA, CA19-9, VEGFR3, and TAC in differentiating between malignant and benign biliary obstructions. *Turk J Gastroenterol*. 2014;25(2):162–169.
31. Huang L, Chen W, Liang P, et al. Serum CYFRA 21-1 in biliary tract cancers: a reliable biomarker for gallbladder carcinoma and intrahepatic cholangiocarcinoma. *Dig Dis Sci*. 2015;60(5):1273–1283.
32. Iorio MV, Ferracin M, Liu CG, et al. MicroRNA gene expression deregulation in human breast cancer. *Cancer Res*. 2005;65(16):7065–7070.
33. Gramantieri L, Ferracin M, Fornari F, et al. Cyclin G1 is a target of miR-122a, a microRNA frequently down-regulated in human hepatocellular carcinoma. *Cancer Res*. 2007;67(13):6092–6099.
34. Yanaihara N, Caplen N, Bowman E, et al. Unique microRNA molecular profiles in lung cancer diagnosis and prognosis. *Cancer Cell*. 2006;9(3):189–198.
35. Bernuzzi F, Marabita F, Lleo A, et al. Serum microRNAs as novel biomarkers for primary sclerosing cholangitis and cholangiocarcinoma. *Clin Exp Immunol*. 2016;185(1):61–71.
36. Selaru FM, Olaru AV, Kan T, et al. MicroRNA-21 is overexpressed in human cholangiocarcinoma and regulates programmed cell death 4 and tissue inhibitor of metalloproteinase 3. *Hepatology*. 2009;49(5):1595–1601.
37. Silakit R, Loilome W, Yongvanit P, et al. Urinary microRNA-192 and microRNA-21 as potential indicators for liver fluke-associated cholangiocarcinoma risk group. *Parasitol Int*. 2017;66(4):479–485.
38. Wang LJ, He CC, Sui X, et al. MiR-21 promotes intrahepatic cholangiocarcinoma proliferation and growth in vitro and in vivo by targeting PTPN14 and PTEN. *Oncotarget*. 2015;6(8):5932–5946.
39. Huang Q, Liu L, Liu CH, et al. MicroRNA-21 regulates the invasion and metastasis in cholangiocarcinoma and may be a potential biomarker for cancer prognosis. *Asian Pac J Cancer Prev*. 2013;14(2):829–834.
40. Ali HEA, Abdel Hameed R, Effat H, et al. Circulating microRNAs panel as a diagnostic tool for discrimination of HCV-associated hepatocellular carcinoma. *Clin Res Hepatol Gastroenterol*. 2017;41(4):e51–e62.
41. Alexander LF, Harri P, Little B, Moreno CC, Mittal PK. Magnetic resonance imaging of primary hepatic malignancies in patients with and without chronic liver disease: a pictorial review. *Cureus*. 2017;9(8):e1539.
42. Peraldo Neia C, Cavalloni G, Chiorino G, Ostano P, Aglietta M, Leone F. Gene and microRNA modulation upon trabectedin treatment in a human intrahepatic cholangiocarcinoma paired patient derived xenograft and cell line. *Oncotarget*. 2016;7(52):86766–86780.
43. Loosen SH, Schueller F, Trautwein C, Roy S, Roderburg C. Role of circulating microRNAs in liver diseases. *World J Hepatol*. 2017;9(12):586–594.
44. Olaizola P, Lee-Law PY, Arbelaz A, et al. MicroRNAs and extracellular vesicles in cholangiopathies. *Biochim Biophys Acta*. 2018;1864(4 Pt B):1293–1307.
45. Li Z, Shen J, Chan MTV, Wu WKK. The role of microRNAs in intrahepatic cholangiocarcinoma. *J Cell Mol Med*. 2017;21(1):177–184.
46. Chen Y, Liu D, Liu P, Chen Y, Yu H, Zhang Q. Identification of biomarkers of intrahepatic cholangiocarcinoma via integrated analysis of mRNA and miRNA microarray data. *Mol Med Rep*. 2017;15(3):1051–1056.
47. Li J, Yao L, Li G, et al. miR-221 promotes epithelial-mesenchymal transition through targeting PTEN and forms a positive feedback loop with β -catenin/c-Jun signaling pathway in extra-hepatic cholangiocarcinoma. *PLoS One*. 2015;10(10):e0141168.
48. Qiao P, Li G, Bi W, Yang L, Yao L, Wu D. microRNA-34a inhibits epithelial mesenchymal transition in human cholangiocarcinoma by targeting Smad4 through transforming growth factor-beta/Smad pathway. *BMC Cancer*. 2015;15:469.
49. Wang LJ, Zhang KL, Zhang N, et al. Serum miR-26a as a diagnostic and prognostic biomarker in cholangiocarcinoma. *Oncotarget*. 2015;6(21):18631–18640.
50. Wang S, Yin J, Li T, et al. Upregulated circulating miR-150 is associated with the risk of intrahepatic cholangiocarcinoma. *Oncol Rep*. 2015;33(2):819–825.
51. Correa-Gallego C, Maddalo D, Doussot A, et al. Circulating plasma levels of microRNA-21 and microRNA-221 are potential diagnostic markers for primary intrahepatic cholangiocarcinoma. *PLoS One*. 2016;11(9):e0163699.
52. Chu CH, Chou W, Wang F, Yeh CN, Chen TC, Yeh TS. Expression profile of microRNA-200 family in cholangiocarcinoma arising from choledochal cyst. *J Gastroenterol Hepatol*. 2016;31(5):1052–1059.
53. Chen Y, Luo J, Tian R, Sun H, Zou S. miR-373 negatively regulates methyl-CpG-binding domain protein 2 (MBD2) in hilar cholangiocarcinoma. *Dig Dis Sci*. 2011;56(6):1693–1701.
54. Li J, Gao B, Huang Z, et al. Prognostic significance of microRNA-203 in cholangiocarcinoma. *Int J Clin Exp Pathol*. 2015;8(8):9512–9516.
55. Chusorn P, Namwat N, Loilome W, et al. Overexpression of microRNA-21 regulating PDCD4 during tumorigenesis of liver fluke-associated cholangiocarcinoma contributes to tumor growth and metastasis. *Tumour Biol*. 2013;34(3):1579–1588.
56. Deng Y, Chen Y. Increased expression of miR-29a and its prognostic significance in patients with cholangiocarcinoma. *Oncol Res Treat*. 2017;40(3):128–132.
57. Li H, Zhou ZQ, Yang ZR, et al. MicroRNA-191 acts as a tumor promoter by modulating the TET1-p53 pathway in intrahepatic cholangiocarcinoma. *Hepatology*. 2017;66(1):136–151.
58. Wang J, Xie C, Pan S, et al. N-myc downstream-regulated gene 2 inhibits human cholangiocarcinoma progression and is regulated by leukemia inhibitory factor/MicroRNA-181c negative feedback pathway. *Hepatology*. 2016;64(5):1606–1622.
59. Garajova I, Brandi G, Biasco G, et al. MiR-21 expression correlates with prognosis in extrahepatic radically resected cholangiocarcinomas treated with gemcitabine-based adjuvant chemotherapy. *Eur J Cancer*. 2015;51:S33–S33.

Cancer Management and Research

Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient. The manuscript management system is completely online and includes

Submit your manuscript here: <https://www.dovepress.com/cancer-management-and-research-journal>

a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress