Infection and Drug Resistance

Open Access Full Text Article

ORIGINAL RESEARCH

Association of MBL2 gene polymorphisms with pulmonary tuberculosis susceptibility: trial sequence meta-analysis as evidence

Raju K Mandal,^{1,*} Munawwar Ali Khan,^{2,*} Arif Hussain,³ Sajad A Dar,¹ Sultan Aloufi,⁴ Arshad Jawed,¹ Mohd Wahid,¹ Aditya K Panda,⁵ Mohtashim Lohani,⁶ Naseem Akhter,⁷ Saif Khan,⁸Bhartendu Nath Mishra,⁹ Shafiul Haque¹

Research and Scientific Studies Unit, College of Nursing and Allied Health Sciences, Jazan University, Jazan, Saudi Arabia; ²Department of Life and Environmental Sciences, College of Natural and Health Sciences. Zaved University. Dubai, United Arab Emirates; 3School of Life Sciences, Manipal Academy of Higher Education, Dubai, United Arab Emirates; ⁴Department of Clinical Laboratory Sciences, College of Applied Medical Sciences, University of Ha'il, Ha'il, Saudi Arabia; 5Centre for Life Sciences, Central University of Jharkhand, Ranchi, Jharkhand, India; ⁶Department of Emergency Medical Services, College of Applied Medical Sciences, Jazan University, Jazan, Saudi Arabia; ⁷Department of Laboratory Medicine, Faculty of Applied Medical Sciences, Al Baha University, Al Baha, Saudi Arabia; ⁸Department of Basic Sciences, College of Dentistry, University of Ha'il, Ha'il, Saudi Arabia; ⁹Department of Biotechnology, Institute of Engineering and Technology, Lucknow, Uttar Pradesh, India

*These authors contributed equally to this work

Correspondence: Shafiul Haque Research and Scientific Studies Unit, College of Nursing and Allied Health Sciences, Jazan University, Jazan 45142, Saudi Arabia Tel/Fax +966 17 317 4383 Email shafiul.haque@hotmail.com



Background: Mannose-binding lectin (MBL) or mannose-binding protein (MBP), encoded by MBL2 gene and secreted by the liver, activates complement system through lectin pathway in innate immunity against the host's infection. Conflictingly, a number of MBL2 variants, rs1800450 (A>B), rs1800451 (A>C), rs5030737 (A>D), rs7096206 (Y>X), rs11003125 (H>L), and rs7095891 (P>Q) allele, have been found to be associated with compromised serum levels and pulmonary tuberculosis (PTB) susceptibility. The present meta-analysis study was performed to evaluate the potential association of these MBL2 gene variants with PTB susceptibility.

Materials and methods: A quantitative synthesis was performed on PubMed (Medline), EMBASE, and Google Scholar web database searches. A meta-analysis was performed to calculate the pooled odds ratios and 95% CIs for all the genetic models.

Results: A total of 14 eligible studies were included to analyze their pooled data for associations between alleles, genotypes, and minor allele carriers. The statistical analysis revealed the significant reduced PTB risk with homozygous variant genotype of rs1800451 polymorphism (CC vs AA: P=0.043; OR =0.828, 95% CI =0.689–0.994). Contrary to this, the variant allele of rs5030737 polymorphism showed association with increased PTB risk (D vs A: P=0.026; OR =1.563, 95% CI =1.054–2.317). However, the other genetic models of rs1800450 (A>B), rs7096206 (Y>X), and rs11003125 (H>L) MBL2 gene polymorphisms did not divulge any association with PTB susceptibility.

Conclusion: The current meta-analysis concludes that rs1800451 (A>C) and rs5030737 (A>D) polymorphisms of MBL2 gene play a significant role in PTB susceptibility. Further, well-designed epidemiological studies with larger sample size including consideration of environmental factors are warranted for the future.

Keywords: meta-analysis, mannose-binding lectin, MBL2, pulmonary tuberculosis, PTB, polymorphism

Introduction

Tuberculosis (TB), caused by *Mycobacterium tuberculosis*, is one of the most prevalent contagious airborne disease/infection, which is very harmful to the body tissues. In spite of good advancements in the treatment strategies, it is still a major health problem among the public. Lately, many reputed health reports reveal 10.4 million new cases and about 1.4 million deaths caused by TB, globally, in the year 2015. Among the two commonly found TB types, pulmonary tuberculosis (PTB) is one of the most devastating form and remains a leading cause of morbidity and mortality, globally.¹ PTB is a two-stage process including infection of *M. tuberculosis* and progression to the disease. It is a complex disease where both environmental and genetic factors play a role in

Construction of the set of the se

185

infection and spread of the disease. The bacteria M. tuberculosis affects approximately one-third of the world's population but only 5%-10% of those infected/affected develop clinical TB during their lifetime.² It is suggestive of how host genetic factors largely influence the development of clinical disease following infection with *M. tuberculosis*,³ which is consistent with animal models of TB as well.⁴ Identifying the important susceptibility and resistance genes and their variations in the host will lead to better understanding of PTB pathogenesis and newer approaches for the treatment and prophylaxis. The ability of the host to prevent the proliferation of the dormant M. tuberculosis pathogen into a full-fledged disease depends on the competence of the immune system. Thus, understanding the molecular mechanism responsible for protective immunity is a necessary step for the development of improved therapies and PTB control.5

The first line of protective immunity to control the initial infection of M. tuberculosis is the innate immunity. Variation at genetic level in components of innate immunity, influenced by a variety of factors, leads to PTB susceptibility.^{6,7} The human mannose-binding lectin (MBL), or mannose-binding protein (MBP), encoded by the MBL2 gene is mapped on 10q11.2-q21 chromosome. It is a polypeptide of 248 amino acids with 32 kDa size8 and considered to be an important part of human innate immunity.9 MBL is a liver-derived collagen-like serum protein. It plays an important role in recognizing the pathogen by its carbohydrate domain present on the surface of a variety of pathogens including M. tuberculosis.^{10,11} MBL targets the invading microorganisms for phagocytosis and complement-mediated killing. It binds to surface carbohydrates and activate the complement cascade via the lectin pathway.12 The role of MBL in preventing and controlling infectious diseases has been widely investigated recently.13

A lot of variability exists in MBL serum levels in different individuals. Low serum levels of MBL2 or individuals producing low MBL2 genotypes have higher infectious disease risk including PTB.^{14,15} Alteration in MBL serum levels in different individuals is attributed to MBL2 gene polymorphisms, which result in the creation of MBL2 genotypes responsible for producing low MBL in ~5% population worldwide.¹⁶ The three commonly known MBL2 gene polymorphisms are present in the structural gene-coding region. They are found at codons: 52 (rs5030737, C>T transition, resulting Arg52Cys substitution, variant allele is known as "D"); 54 (rs1800450, G>A transition, resulting Gly54Asp substitution, variant allele is known as "B"); and 57 (rs1800451, G>A transition, resulting in Gly57Glu substitution, variant allele is known as "C"). The above said mutations lead to the structural changes in MBL protein, which cause a functional deficiency and a significant reduction in the circulating MBL.¹⁷ These polymorphisms are collectively known as "AO" and reveal the alteration in MBL serum levels. The allele "A" designates the wild-type allele, whereas the allele "O" is indicative of the presence of mutant allele(s) in either of the three polymorphisms.13 In addition, two other relevant single nucleotide polymorphisms (SNPs) at positions -550 (rs11003125, C>G, "HL" variant, where L is the wild-type allele) and -221 (rs7096206, G>C, "XY" variant, where Y is the wild-type allele) in the promoter region have been identified as protein expression regulators.¹⁸ These polymorphisms may change the transcription rate, which can further lead to change in the concentration of serum MBL. Genotypes containing structural polymorphisms and promoter variants have shown variations in MBL concentrations in different individuals which can be up to 1,000 folds.13 These polymorphisms affect MBL2 gene which reduces MBL protein level leading to decrease in innate immunity against pathogens. Analysis of the genetic variants of MBL2 gene in PTB appears to be helpful in identifying individuals at higher risk of PTB susceptibility.

A plethora of studies have been conducted in the past to evaluate the effect of MBL2 gene polymorphisms on the progress of infection in PTB.^{15,19–31} The results of these studies are contradictory. The reason for their contradiction is probably their small sample size which lacks adequate power to detect the effects of MBL2 gene polymorphisms on PTB. This weakness can be overcome by the use of meta-analysis, a statistical tool that explores the risk factors associated with different genetic diseases. A meta-analysis employs a quantitative method to combine the data drawn from individual studies with small sample sizes to provide a reliable conclusion. We, therefore, performed the current meta-analysis to identify the effect of the abovementioned MBL2 polymorphisms on overall PTB risk.

Materials and methods Study search and online databases

A systematic search was performed on various scholarly databases like PubMed, Medline, EMBASE, and Google Scholar. The combination of the keywords used for the search was "MBL OR MBL2 OR MBP OR mannose-binding protein OR mannose-binding lectin" gene (polymorphism OR mutation OR variant) AND tuberculosis OR PTB Susceptibility/Risk (last updated on December 2017). The titles and abstracts of the hits obtained by the above search criteria were evaluated. The most relevant publications were obtained after applying the eligibility criteria for thorough examination. In addition to the online database search, the reference lists of the selected research articles were also screened to find any missed relevant article during the initial search.

Criteria for inclusion and exclusion of the studies

To facilitate the proper interpretation of this study, heterogeneity was minimized. The following criteria were applied to select and include the published studies in this meta-analysis: 1) the study must have a case-control design investigating MBL2 gene polymorphism and PTB risk, 2) the patients included in the study should be confirmed PTB cases and controls should be TB free, 3) the study should have genotypic frequency available for both cases and controls, 4) the study should be published in English language only, and 5) the study should have used statistically acceptable data collection and analysis strategy. In addition to these criteria, in cases where the same case series was included in multiple articles, the most recent study with highest number of cases was included in this study. The main reasons followed for exclusion of the studies from this meta-analysis were as follow: 1) duplicate studies or overlapping publications, 2) the study designs based only on PTB cases, 3) the articles with no genotype frequency reported, and 4) the data of reviews and abstracts.

The information related to the identification and selection of the pertinent studies, by following the inclusion/exclusion criteria mentioned above, for the present meta-analysis has been given as PRISMA 2009 flow diagram (Figure 1).

Strategy for literature search

Two researchers (RKM and SK) individually examined all the titles and abstracts of the retrieved published articles using the designated online web database search in a sequential order. The full texts of all the articles, possibly qualified, were also retrieved. To evaluate the appropriateness of the study for the inclusion in this pooled analysis, one researcher (RKM) systematically examined all the full-text retrieved publications. Afterward, the second researcher (SK) performed the same procedure of text evaluation independently by selecting randomly 10% of the full-text articles. During the selection process, full agreement was found between the two researchers regarding study exclusion and selection criteria. After the selection of the final set of the eligible studies, another researcher (SAD) extrapolated the pertinent data from all the included studies. This step of data extrapolation was cross-checked by another researcher (MAK), independently, by collecting the information from all the selected articles. Discrepancies and discords occurred during the study selection were resolved amicably with thorough discussion in the presence of an adjudicator (SH).



Figure I PRISMA 2009 flow diagram depicting identification and selection process (inclusion/exclusion) of the relevant published articles dealing with MBL2 gene polymorphisms and pulmonary tuberculosis risk for the present meta-analysis.

Screening of the studies

The electronic searches performed on PubMed (Medline), EMBASE, and Google Scholar web databases resulted in short-listing of 36 research articles in the preliminary stage. All the articles were examined thoroughly for their relevance by checking their titles, abstracts, and full texts for this meta-analysis. The research articles predicting survival in MBL2 polymorphic PTB patients or considering MBL2 variants as indicators for therapeutic response were excluded from this study. Also excluded were studies reporting expression of MBL2 at mRNA or protein level, and the review articles. Only studies with a case-control or cohort design with genotypic frequencies available for all the three genotypes were included in this meta-analysis. Beside the selected database search, the reference articles cited in the potential retrieved studies were also checked carefully for other potentially relevant articles, but no germane study was found. After many rounds of careful screening (using the sequential approach) by applying the pre-set inclusion and exclusion criteria as shown in PRISMA 2009 flow diagram (Figure 1), 14 eligible original studies were considered worth to be included for this meta-analysis. These studies dealt with MBL2 rs1800450 (A>B), rs1800451 (A>C), rs5030737 (A>D), rs7096206 (Y>X), rs11003125 (H>L), and rs7095891 (P>Q) gene polymorphisms and PTB risk (Table 1).

Newcastle–Ottawa Scale (NOS) criteria for quality evaluation

For NOS evaluation, two independent investigators (RKM and SAD) separately performed the methodological quality assessment of the retrieved studies. The established NOS criteria for quality score analysis was used for the methodological quality assessment of the included studies.³² The NOS criteria focuses on three main aspects: 1) subject selection: 0–4 points/star; 2) comparability of subject: 0–2 points/star; 3) and clinical outcome: 0–3 points/star. The studies scoring five or more stars were considered of medium to high quality.^{32,33} In case of difference in the number of stars assigned to a particular study by the two investigators, the issue was fully debated and resolved by a thorough discussion with a third investigator, SH.

Extraction and evaluation of the data quality

Two investigators (RKM and SK) independently summarized the methodological quality assessment and data extracted from all the eligible and retrieved articles by following the standard procedure. To confirm the accuracy of the summarized data, a data-collection form was used while strictly following the criteria stated above. The major summarized characteristics from the retrieved studies were as follows: the

First author and year	Country	Ethnicity	Control	Cases	Methods	SNP	Association
Wu et al 2015 ¹⁹	China	Asian	453	151	PCR-RFLP	P>Q, Y>X, H>L, A>B	No
da Cruz et al 2013 ²⁰	Brazil	Mixed	148	119	Sequencing	H>L, Y>X, A>O, A>D, A>B, A>C	A>O, A>C risk
Araújo et al 2013 ²¹	Brazil	Mixed	159	133	PCR	A>B, A>C, A>D, A>O	No
Singla et al 2012 ²²	India	Asian	392	286	PCR-RFLP	A>B	Yes
Thye et al 2011 ³¹	Ghana	African	2,236	1,953	Pyrosequencing	H>L, Y>X, P>Q, A>C	A>C
Capparelli et al 2009 ¹⁵	Italy	Caucasian	288	277	PCR	A>O	Protective
Cosar et al 2008 ²³	Turkey	Caucasian	99	27	PCR-RFLP	A>B	Protective
Alagarasu et al 2007 ²⁴	India	Asian	146	225	PCR-SSP	A>B, A>D, A>C, A>O, Y>X	Y>X, A>O
Søborg et al 2007 ²⁵	Tanzania	African	426	424	PCR-SSP	A>B, A>D, A>C, A>O	No
Liu et al 2006 ²⁶	China	Asian	293	152	PCR-SSP	H>L, P>Q, Y>X, A>B	No
Selvaraj et al 2006 ²⁷	India	Asian	58	48	PCR-SSOP	A>O	No
Ozbaş-Gerçeker et al 2003 ²⁸	Turkey	Caucasian	100	49	PCR	A>B	No
Selvaraj et al 1999 ²⁹	India	Asian	109	202	PCR	A>B, A>C, A>D, A>O	A>C
Bellamy et al 1998 ³⁰	Gambia	African	422	397	PCR	A>B, A>C	Protective A>C

 Table I Main characteristics of all the studies included in this meta-analysis

Notes: MBL2 gene variants designated as rs1800450 (A>B), rs1800451 (A>C), rs5030737 (A>D), rs7096206 (Y>X), rs11003125 (H>L), rs7095891 (P>Q), combined rs1800450, rs1800451, rs5030737 (A>O).

Abbreviations: PCR-RFLP, polymerase chain reaction-restriction fragment length polymorphism; PCR-SSOP, polymerase chain reaction-sequence-specific oligonucleotide probes; PCR-SSP, polymerase chain reaction-sequence-specific amplification; SNP, single nucleotide polymorphism.

first author, the year of publication, the country of origin, the ethnicity of cases and controls and their number, the type of study, and the genotypic frequencies. The disagreements/ discrepancies, on any data item of the selected studies, between RKM and SK were resolved by open scientific discussion with the third arbitrator (SH) in order to reach a final consensus.

Statistical analyses

The pooled ORs and their corresponding 95% CIs were estimated in order to assess the association between MBL2 gene polymorphism and PTB risk. The chi-square-based Q-test was used to examine heterogeneity assumption.³⁴ A P-value <0.05 was the threshold for heterogeneity significance. In case of no heterogeneity, the data from single comparison was combined using a fixed-effects model;³⁵ otherwise, a random-effects model was used for pooling the data.³⁶ Additionally, for inter-study variability quantification, I^2 statistics was used. Higher values suggested an increasing degree of heterogeneity.37 The chi-squared test was used to calculate the Hardy-Weinberg equilibrium (HWE) among the controls. The funnel plot asymmetry was calculated using Egger's regression test (linear regression approach) on the natural logarithm scale of the OR. The significance of the intercept was measured by *t*-test, wherein the *P*-value <0.05 was the representation of statistically significant publication bias. A uniform resource locator http://www.meta-analysis.com/ pages/comparisons.html was used for comparative evaluation of the "meta-analysis" programs. The Comprehensive Meta-Analysis V2 software program (Biostat, Englewood, NJ, USA) was used to analyze all the statistical parameters in this meta-analysis.

Trial sequential analysis

According to the Cochrane handbook, the best meta-analysis includes all the eligible trials in the analysis. Though it may not be sufficient evidence, there are chances that sometimes meta-analysis may lead to bias (systematic errors) or chance (random errors). In order to reduce these errors, Trial sequential analysis (TSA) tool from Copenhagen Trial Unit, Center for Clinical Intervention Research, Denmark, was used. TSA helps in calculating the required information size, adjusts threshold for statistical significance, and estimates the power of current conclusion.^{32,38–40} When the TSA monitoring boundary crosses with the *Z*-curve before the required information size is reached, it confirms the robust evidence indicating the need for no further trials. When the case is reverse, further trials are necessarily required to be done in order to cross the monitoring boundaries. The Trial Sequential Analysis software (version 0.9, <u>http://www.ctu.</u> <u>dk/tsa/</u>) was used for this analysis. We performed two-sided hypothesis testing for the analysis and fixed type I error rate (α) at 5%. The results were combined by a DerSimonian–Laird random effects model. Sequential efficacy and harm monitoring boundaries that control type I error were constructed using Lan–DeMets version of the O'Brien–Fleming α -spending function, and the futility monitoring boundaries (for type II error) were constructed with an O'Brien–Fleming β -spending function.

Results

Quality assessment and characteristics of the selected studies

The major characteristics like genotypic distribution, minor allele frequency (MAF) in the controls/cases, and PTB susceptibility for all the 14 studies included in the present meta-analysis have been presented in Tables 2 and 3. All the 14 studies were evaluated for their quality according to the scoring system of NOS, and the majority of the studies (approximately 80%) were found to score five stars or more. The quality NOS scoring revealed moderate to good quality of all the studies included in this pooled analysis (Table 4).

Publication bias diagnosis, heterogeneity evaluation, and quantitative synthesis

The presence or absence of the publication bias in the included studies was checked by funnel plot asymmetry and Egger's regression statistics (P-value <0.05 considered as significant publication bias level). The shape of the funnel plots obtained and the outcomes of Egger's test showed absence/presence of publication bias in all the five comparative genetic models of MBL2 gene polymorphisms. Likewise, the presence/absence of heterogeneity in the included MBL2 gene polymorphism studies for all the five genetic models of this meta-analysis was evaluated by using Q-test and I^2 statistics. For each MBL2 polymorphism, random-effects model was applied to synthesize the data in the presence of heterogeneity, whereas fixed-effects model was adopted in the absence of heterogeneity. The outcomes of publication bias and heterogeneity evaluation and quantitative synthesis for each MBL2 polymorphism have been given in the following sections.

MBL2 rs1800450 polymorphism

A total of eleven case–control studies provided the sufficient data (including 2,258 controls and 1,789 PTB cases)

189

First author and year Controls				Cases				HWE	
	Genotype	e rs1800	450	Minor allele	Genot	ype rsl 8	800450	Minor allele	
	AA	AB	BB	MAF	AA	AB	BB	MAF	P-value
Wu et al 2015 ¹⁹	348	97	8	0.124	112	37	2	0.135	0.459
da Cruz et al 2013 ²⁰	124	21	3	0.091	95	22	2	0.109	0.079
Araújo et al 2013 ²¹	101	48	0	0.161	84	35	0	0.147	0.019
Singla et al 2012 ²²	207	155	30	0.274	175	100	11	0.213	0.895
Cosar et al 2008 ²³	71	27	1	0.146	23	4	0	0.074	0.366
Alagarasu et al 2007 ²⁴	86	33	4	0.166	120	42	12	0.189	0.704
Søborg et al 2007 ²⁵	271	13	1	0.026	289	9	0	0.015	0.063
Liu et al 2006 ²⁶	166	42	4	0.117	103	34	4	0.148	0.487
Ozbaş-Gerçeker et al 2003 ²⁸	76	20	4	0.14	40	9	0	0.091	0.090
Selvaraj et al 1999 ²⁹	84	24	1	0.119	137	51	14	0.195	0.615
Bellamy et al 1998 ³⁰	183	5	0	0.013	198	7	0	0.017	0.853
	Genotype	e rs 800	45 I		Genot	ype rsl8	80045 I		
	AA	AC	СС		AA	AC	СС		
da Cruz et al 2013 ²⁰	140	8	0	0.027	133	22	0	0.070	0.735
Araújo et al 2013 ²¹	101	2	0	0.009	84	4	0	0.022	0.920
Thye et al 2011 ³¹	1,002	977	257	0.333	885	815	194	0.317	0.421
Alagarasu et al 2007 ²⁴	86	12	1	0.070	120	11	1	0.049	0.439
Søborg et al 2007 ²⁵	271	112	20	0.188	289	115	20	0.182	0.065
Selvaraj et al 1999 ²⁹	103	5	1	0.032	176	25	1	0.066	0.006
Bellamy et al 1998 ³⁰	183	192	42	0.330	198	159	29	0.281	0.417
	Genotype	e rs5030	737		Genot	ype rs50	30737		
	AA	AD	DD		AA	AD	DD		
da Cruz et al 2013 ²⁰	138	9	1	0.037	108	10	1	0.050	0.067
Araújo et al 2013 ²¹	101	6	0	0.028	84	8	0	0.043	0.765
Alagarasu et al 2007 ²⁴	86	8	0	0.042	120	26	2	0.101	0.666
Søborg et al 2007 ²⁵	271	6	0	0.010	289	8	0	0.013	0.855
Selvaraj et al 1999 ²⁹	99	10	0	0.045	186	9	7	0.056	0.615

 Table 2 Genotypic distribution of MBL2 gene polymorphism included in meta-analysis: first

Abbreviations: HWE, Hardy–Weinberg equilibrium; MAF, minor allele frequency.

Table 3 Genotypic distribution	of MBL2 gene polymor	phisms included in this m	eta-analysis: second
---------------------------------------	----------------------	---------------------------	----------------------

First author and	Control	5			Cases				HWE
year	Combin 1800451	ed rs18004 , 5030737	450,	Minor allele	Combin 1800451	Combined rs1800450, 1800451, 5030737			
	AA	AO	00	MAF	AA	AO	00	MAF	P-value
da Cruz et al 2013 ²⁰	108	34	6	0.155	71	41	7	0.231	0.128
Araújo et al 2013 ²¹	101	56	2	0.188	84	47	2	0.191	0.057
Capparelli et al 2009 ¹⁵	166	112	10	0.229	55	158	61	0.510	0.087
Alagarasu et al 2007 ²⁴	86	53	7	0.229	120	82	24	0.287	0.747
Søborg et al 2007 ²⁵	271	131	30	0.221	289	132	22	0.198	0.013
Selvaraj et al 2006 ²⁷	37	18	3	0.206	24	19	5	0.302	0.678
Selvaraj et al 1999 ²⁹	68	39	2	0.197	107	73	22	0.289	0.175
	Genotype rs7096206			Genoty	be rs70962	06			
	YY	YX	XX		YY	YX	XX		
Wu et al 2015 ¹⁹	318	120	15	0.165	97	47	7	0.201	0.379
da Cruz et al 2013 ²⁰	110	32	6	0.148	75	40	4	0.201	0.076
Thye et al 2011 ³¹	1,663	486	31	0.125	1,437	396	26	0.120	0.502
Alagarasu et al 2007 ²⁴	72	61	13	0.297	136	79	10	0.22	0.987
Liu et al 2006 ²⁶	151	54	7	0.160	91	44	6	0.198	0.430
	Genotyp	e rsl I 003	125		Genoty	be rsl1003	125		
	HH	HL	LL		HH	HL	LL		
Wu et al 2015 ¹⁹	46	64	41	0.483	101	248	104	0.503	0.062
da Cruz et al 2013 ²⁰	19	61	68	0.665	8	45	66	0.743	0.366
Thye et al 2011 ³¹	9	287	1,878	0.929	7	266	1,570	0.924	0.576
Liu et al 2006 ²⁶	49	105	58	0.521	44	66	31	0.453	0.911

Abbreviations: HWE, Hardy–Weinberg equilibrium; MAF, minor allele frequency.

to calculate the ORs. Publication bias did not exist and heterogeneity was found in one genetic model (Table 5; Figure SI1). The pooled ORs revealed that MBL2 A>B gene polymorphism demonstrated no association with PTB risk in allelic contrast (B vs A: P=0.999; OR =1.000, 95% CI =0.799–1.251), homozygous (BB vs AA: P=0.495; OR =0.848, 95% CI =0.527–1.363), heterozygous (BA vs AA: P=0.717; OR =0.969, 95% CI=0.820–1.146), dominant (BB+ BA vs AA: P=0.704; OR =0.969, 95% CI =0.824–1.140), and recessive (BB vs BA+ AA: P=0.581; OR =0.876, 95% CI =0.547–1.402) genetic models, respectively (Figure 2).

In the stratification analysis based on the ethnicity, five studies were included for the analysis in the Asian population. No publication bias existed and heterogeneity was found in four genetic models (Table 6; Figure SI2). No significant relationship was observed with PTB in allelic contrast (B vs A: P=0.461; OR =1.131, 95% CI =0.815–1.571), homozygous (BB vs AA: P=0.582; OR =1.313, 95% CI =0.498–3.465), heterozygous (BA vs AA: P=0.947; OR =0.993, 95% CI =0.815–1.211), dominant (BB+ BA vs AA: P=0.605; OR

 Table 4 Quality assessment conducted according to the NOS for all the included studies

First author	Quality inc	dicators	
and year	Selection	Comparability	Exposure
Wu et al 2015 ¹⁹	* * *	*	**
da Cruz et al 2013 ²⁰	***	*	**
Araújo et al 2013 ²¹	* * *	*	**
Singla et al 2012 ²²	***	*	***
Thye et al 2011 ³¹	***	*	***
Capparelli et al 2009 ¹⁵	***	*	**
Cosar et al 2008 ²³	**	*	**
Alagarasu et al 2007 ²⁴	***	*	*
Søborg et al 2007 ²⁵	* * *	*	*
Liu et al 2006 ²⁶	* * *	*	***
Selvaraj et al 2006 ²⁷	* * *	*	*
Ozbaş-Gerçeker et al	*	*	**
200328			
Selvaraj et al 1999 ²⁹	***	*	**
Bellamy et al 1998 ³⁰	* *	*	**

Abbreviation: NOS, Newcastle-Ottawa Scale.

=1.083, 95% CI =0.800–1.467), and recessive (BB vs BA+ AA: *P*=0.576; OR =1.298, 95% CI =0.521–3.231), respectively (Figure 3).

MBL2 rs1800451 polymorphism

A total of seven case–control studies were included for this analysis accumulating 3,515 controls and 3,281 PTB cases. No publication bias was observed, whereas heterogeneity was found in three genetic models (Table 7; Figure SI3). The pooled ORs revealed that homozygous (CC vs AA: P=0.043; OR =0.828, 95% CI =0.689–0.994) genetic model was associated with reduced risk to PTB. However, other genetic models, allelic contrast (C vs A: P=0.921; OR =0.990, 95% CI =0.804–1.217), heterozygous (CA vs AA: P=0.724; OR =1.050, 95% CI =0.800–1.378), dominant (CC+ CA vs AA: P=0.856; OR =1.025, 95% CI =0.788–1.332), and recessive (CC vs CA + AA: P=0.093; OR =0.861, 95% CI =0.722–1.026), did not show any association with PTB (Figure 4).

The stratification analysis based on the ethnicity included three studies for the analysis of the African population. Publication bias and heterogeneity were absent (Table 8; Figure SI4). A reduced risk was observed with allelic (C vs A: P=0.025; OR =0.912, 95% CI =0.842–0.989) and homozygous (CC vs AA: P=0.047; OR =0.830, 95% CI =0.690–0.997) genetic models. On the other hand, heterozygous (CA vs AA: P=0.132; OR =0.918, 95% CI =0.822–1.026), dominant (CC+ CA vs AA: P=0.052; OR =0.901, 95% CI =0.812–1.001), and recessive (CC vs CA + AA: P=0.100; OR =0.863, 95% CI =0.723–1.029) genetic models did not show any risk with PTB (Figure 5).

MBL2 rs5030737 polymorphism

A total of five case–control studies were included, which comprises a sum of 735 controls and 876 confirmed PTB cases. There was no publication bias and heterogeneity (Table 9; <u>Figure SI5</u>). The pooled ORs revealed that variant allele was found to be associated with the increased risk (D vs A:

Table 5 Statistics to test the publication bias and heterogeneity (rs1800450 Overall)

Comparisons	Egger's regression analysis			Heteroger	eity analysis		Model used for the
	Intercept	95% CI	P-value	Q-value	P _{heterogeneity}	l² (%)	meta-analysis
B vs A	0.21	-2.33 to 2.74	0.86	20.21	0.03	50.52	Random
BB vs AA	0.70	–1.52 to 2.91	0.48	12.76	0.12	37.29	Fixed
BA vs AA	0.05	-1.90 to 1.99	0.96	9.34	0.50	0.00	Fixed
BB + BA vs AA	0.08	-2.28 to 2.45	0.94	14.27	0.16	29.90	Fixed
BB vs BA + AA	0.60	-1.50 to 2.71	0.52	11.61	0.17	31.11	Fixed

Study name	Statistics for each study	OR and 95% CI	
Due A		Of and 35% Of	Deletive
DVSA	Lower Upper OP limit limit Zuelus Durlur		Relative
	OR limit limit 2-value P-value		weight
Wu et al 2015	1.102 0.751 1.618 0.498 0.619	🖶	12.79
da Cruz et al 2013	1.222 0.693 2.156 0.692 0.489		8.85
Araujo et al 2013	0.898 0.559 1.442 -0.446 0.656	1 🕂 1	10.70
Singla et al 2012	0.717 0.556 0.925 -2.559 0.010		16.18
Cossar et al 2008	0.466 0.156 1.389 -1.370 0.171		3.50
Alagarasu et al 2007	1.170 0.762 1.798 0.718 0.473		11.69
Søborg et al 2007	0.567 0.246 1.307 -1.331 0.183	│─■┼│││	5.34
Liu et al 2006	1.309 0.842 2.035 1.196 0.232		11.42
Ozbas-Gerçeker et al 2003	0.621 0.281 1.373 -1.176 0.240		5.75
Selvarai et al 1999	1.795 1.113 2.894 2.400 0.016		10.61
Bellamy et al 1998	1.289 0.406 4.096 0.430 0.667		3.18
Combined	1.000 0.799 1.251 -0.001 0.999		
	0.01	0.1 1 10 100	
	0.01	0.1 1 10 100	
Study name	Statistics for each study	OR and 95% CI	
BB vs AA	Lower Upper		Relative
	OR limit limit Z-value P-value		weight
10045	0.777 0.400 0.740 0.047 0.750 I	I -I I I	0.00
Wu et al 2015	0.777 0.163 3.712 -0.317 0.752		9.22
da Cruz et al 2013	0.870 0.143 5.312 -0.151 0.880		6.89
Singla et al 2012	0.434 0.211 0.891 -2.275 0.023		43.56
Cossar et al 2008	1.014 0.040 25.750 0.009 0.993		2.16
Alagarasu et al 2007	2.150 0.671 6.893 1.288 0.198	│ ┼╋──│ │	16.62
Søborg et al 2007	0.313 0.013 7.707 -0.711 0.477	— • 	2.20
Liu et al 2006	1.612 0.394 6.585 0.665 0.506	╎─┼═──│ │	11.38
Ozbaş-Gerçeker et al 2003	0.210 0.011 3.996 -1.039 0.299		2.60
Selvaraj et al 1999	8.584 1.109 66.470 2.059 0.040	│	5.38
Combined	0.848 0.527 1.363 -0.682 0.495		
	0.01	0.1 1 10 100	
Study name	Statistics for each study	<u>OR and 95% CI</u>	
BA vs AA	Lower Upper		Relative
	OR limit limit Z-value P-value		weight
Wu et al 2015	1 185 0 768 1 830 0 766 0 443 I	1 1	14.90
do Cruz ot al 2012	1 267 0 710 2 622 0 026 0 240		14.89
Arouio et al 2013			6.55
Single et al 2013	0.877 0.520 1.479 -0.495 0.022		10.27
Singla et al 2012	0.765 0.555 1.055 -1.045 0.100		27.10
Alagarasu at al 2007	0.457 0.145 1.445 -1.555 0.165		2.12
Alagarasu et al 2007	0.912 0.555 1.555 -0.556 0.755		9.88
Cabora at al 2007	0.640 0.070 1.640 0.070 0.000 1		
Søborg et al 2007	0.649 0.273 1.543 -0.978 0.328		3.75
Søborg et al 2007 Liu et al 2006	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311		3.75 10.61
Søborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726		3.75 10.61 3.67
Søborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351		3.75 10.61 3.67
Søborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998	$ 0.649 0.273 1.543 -0.978 0.328 \\ 1.305 0.780 2.183 1.013 0.311 \\ 0.855 0.356 2.051 -0.351 0.726 \\ 1.303 0.747 2.272 0.933 0.351 \\ 1.294 0.404 4.149 0.433 0.665 \\ 1.294 0.404 4.149 0.433 0.665 \\ 1.294 0.404 4.149 0.433 0.665 \\ 1.294 0.404 4.149 0.433 0.665 \\ 1.294 0.404 0.414 0.433 0.665 \\ 1.294 0.404 0.414 0.433 0.665 \\ 1.294 0.404 0.414 0.433 0.665 \\ 1.294 0.404 0.414 0.433 0.665 \\ 1.294 0.404 0.414 0.433 0.665 \\ 1.294 0.404 0.414 0.433 0.665 \\ 1.294 0.404 0.414 0.433 0.665 \\ 1.294 0.404 0.414 0.433 0.665 \\ 1.294 0.404 0.414 0.433 0.665 \\ 1.294 0.404 0.414 0.433 0.665 \\ 1.294 0.404 0.414 0.433 0.665 \\ 1.294 0.404 0.414 0.433 0.665 \\ 1.294 0.404 0.433 0.665 \\ 1.294 0.404 0.414 0.433 0.665 \\ 1.294 0.404 0.414 0.433 0.665 \\ 1.294 0.404 0.414 0.433 0.665 \\ 1.294 0.404 0.414 0.433 0.665 \\ 1.294 0.404 0.414 0.433 0.665 \\ 1.294 0.404 0.414 0.414 0.414 0.433 0.665 \\ 1.294 0.414 $		3.75 10.61 3.67 9.09
Søborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined	$\begin{array}{cccccc} 0.649 & 0.273 & 1.543 & -0.978 & 0.328 \\ 1.305 & 0.780 & 2.183 & 1.013 & 0.311 \\ 0.855 & 0.356 & 2.051 & -0.351 & 0.726 \\ 1.303 & 0.747 & 2.272 & 0.933 & 0.351 \\ 1.294 & 0.404 & 4.149 & 0.433 & 0.665 \\ 0.969 & 0.820 & 1.146 & -0.362 & 0.717 \\ \end{array}$		3.75 10.61 3.67 9.09 2.07
Søborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined			3.75 10.61 3.67 9.09 2.07
Søborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01		3.75 10.61 3.67 9.09 2.07
Søborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01	0.1 1 10 100	3.75 10.61 3.67 9.09 2.07
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u>	0.1 1 10 100	3.75 10.61 3.67 9.09 2.07
Soborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper 0.8	0.1 1 10 100	3.75 10.61 3.67 9.09 2.07 Relative
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined <u>Study name</u> BB+BA vs AA	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value	0.1 1 10 100 OR and 95% Cl	3.75 10.61 3.67 9.09 2.07 Relative weight
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.326 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508	0.1 1 10 100 OR and 95% Cl	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60
Soborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined <u>Study name</u> BB+BA vs AA Wu et al 2015 da Cruz et al 2013	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404	0.1 1 10 100 OR and 95% CI	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622	0.1 1 10 100	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Singla et al 2012	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030	OR and 95% Cl	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48
Soborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined <u>Study name</u> BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Singla et al 2012 Cossar et al 2008	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162	0.1 1 10 100 OR and 95% CI	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Cossar et al 2013 Cossar et al 2008 Alagarasu et al 2007	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861	OR and 95% Cl	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Singla et al 2012 Cossar et al 2008 Alagarasu et al 2007 Seborg et al 2007	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245	OR and 95% CI	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 2.62
Soborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined <u>Study name</u> BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Singla et al 2012 Cossar et al 2007 Alagarasu et al 2007 Liu et al 2006	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.331 0.812 2.184 1.133 0.257	0.1 1 10 100 OR and 95% CI	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Cossar et al 2007 Cossar et al 2007 Seborg et al 2007 Liu et al 2007	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.669 0.820 1.146 -0.362 0.717 0.01	OR and 95% Cl	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75
Soborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined <u>Study name</u> BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Cossar et al 2013 Alagarasu et al 2007 Søborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.31 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.594 0.933 2.723 1.708 0.088	OR and 95% CI	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59
Soborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Cossar et al 2012 Cossar et al 2007 Alagarasu et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.366 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value P-value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.331 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.594 0.933 2.723 1.708 0.088 1.294 0.404 4.149 0.433 0.665	0.1 1 10 100 OR and 95% CI	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Singla et al 2012 Cossar et al 2008 Alagarasu et al 2007 Søborg et al 2007 Søborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 Statistics for each study Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.31 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.294 0.404 4.149 0.433 0.665 1.294 0.404 4.140 -0.381 0.704	OR and 95% Cl	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19
Soborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined <u>Study name</u> BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Cossar et al 2013 Alagarasu et al 2007 Saborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.063 0.257 1.416 -1.162 0.245 1.331 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.594 0.933 2.723 1.708 0.088 1.294 0.404 4.149 0.433 0.665 0.969 0.824 1.100 -0.381 0.704	OR and 95% CI	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19 1.94
Soborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Singla et al 2012 Cossar et al 2007 Saborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	0.1 1 10 100 OR and 95% CI OR and 95% CI 0.1 1 10 100	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19 1.94
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Cossar et al 2007 Seborg et al 2007 Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 Statistics for each study Lower Upper OR limit limit Z-value P-value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.331 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.294 0.404 4.149 0.433 0.665 0.969 0.824 1.140 -0.381 0.704 0.01	OR and 95% Cl	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19 1.94
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Cossar et al 2012 Cossar et al 2010 Alagarasu et al 2007 Søborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.31 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.294 0.404 4.149 0.433 0.665 0.969 0.824 1.140 -0.381 0.704 0.01 <u>Statistics for each study</u>	OR and 95% CI	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 2.7.48 2.00 10.47 3.62 10.75 3.59 9.19 1.94
Soborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined <u>Study name</u> BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Singla et al 2012 Cossar et al 2007 Saborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1999 Bellamy et al 1998 Combined	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.366 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.681 0.603 0.257 1.416 -1.162 0.245 1.331 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.594 0.933 2.723 1.708 0.088 1.294 0.404 4.149 0.433 0.665 0.969 0.824 1.140 -0.381 0.704 <u>0.01</u>	OR and 95% CI	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19 1.94 Relative
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Araujo et al 2013 Cossar et al 2012 Cossar et al 2007 Seborg et al 2007 Seborg et al 2007 Seborg et al 2007 Seborg et al 2007 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB vs BA+AA	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 Statistics for each study Lower Upper OR limit limit Z-value P-value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.331 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.594 0.933 2.723 1.708 0.088 1.294 0.404 4.149 0.433 0.665 0.969 0.824 1.140 -0.381 0.704 UDUE COR limit limit Z-value P-value	OR and 95% Cl OR and 95% Cl OR and 95% Cl OR and 95% Cl OR and 95% Cl	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19 1.94 Relative weight
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Cossar et al 2013 Alagarasu et al 2007 Saborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB vs BA+AA Wu et al 2015	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.331 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.294 0.404 4.149 0.433 0.665 0.969 0.824 1.140 -0.381 0.704 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 0.01	OR and 95% Cl	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19 1.94 Relative weight 9.09
Soborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined <u>Study name</u> BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Singla et al 2012 Cossar et al 2007 Saborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1999 Bellamy et al 1998 Combined <u>Study name</u> BB vs BA+AA Wu et al 2015 da Cruz et al 2013	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.366 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.661 0.603 0.257 1.416 -0.176 0.681 0.603 0.257 1.416 -0.176 0.438 1.391 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.594 0.933 2.723 1.708 0.088 1.294 0.404 4.149 0.433 0.665 0.969 0.824 1.140 -0.381 0.704 <u>0.01</u>	OR and 95% CI	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19 1.94 Relative weight
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Cossar et al 2013 Cossar et al 2013 Cossar et al 2010 Søborg et al 2007 Liu et al 2008 Alagarasu et al 2007 Søborg et al 2008 Corbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB vs BA+AA Wu et al 2015 da Cruz et al 2013 Sinda et al 2012	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.31 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.594 0.933 2.723 1.708 0.088 1.294 0.404 4.149 0.433 0.665 0.969 0.824 1.140 -0.381 0.704 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 0.01	OR and 95% Cl OR and 95% Cl	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19 1.94 Relative weight 9.09 6.79
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2012 Cossar et al 2007 Saborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1999 Bellamy et al 1998 Combined Study name BB vs BA+AA Wu et al 2015 da Cruz et al 2013 Singla et al 2012 Cossar et al 2008	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.331 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.594 0.933 2.723 1.708 0.088 1.294 0.404 4.149 0.433 0.665 0.969 0.824 1.140 -0.381 0.704 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 0.747 0.157 3.555 -0.367 0.714 0.826 0.136 5.027 -0.207 0.836 0.440 0.473 0.136 0.088 1.294 0.404 3.068 0.714 0.482 0.136 5.027 -0.207 0.836 0.443 0.238 0.980 -2.015 0.044 1.194 0.047 30 134 0.108 0.180	OR and 95% CI	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19 1.94 Relative weight 9.09 6.79 44.13
Soborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined <u>Study name</u> BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Singla et al 2012 Cossar et al 2007 Liu et al 2007 Diu et al 2008 Selvaraj et al 1999 Bellamy et al 1999 Bellamy et al 1998 Combined <u>Study name</u> BB vs BA+AA Wu et al 2015 da Cruz et al 2012 Cossar et al 2012 Cossar et al 2012 Cossar et al 2012	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.366 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value P-value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.331 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.594 0.933 2.723 1.708 0.088 1.294 0.404 4.149 0.433 0.665 0.969 0.824 1.140 -0.381 0.704 <u>0.01</u>	OR and 95% CI	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19 1.94 Relative weight 9.09 6.79 4.13 2.13
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Singla et al 2012 Cossar et al 2008 Alagarasu et al 2007 Selvaraj et al 2007 Selvaraj et al 2008 Selvaraj et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB vs BA+AA Wu et al 2015 da Cruz et al 2013 Singla et al 2012 Cossar et al 2012 Cossar et al 2007 Alagarasu et al 2007 Determet da 2007	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value P-value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.334 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.31 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.766 0.438 1.594 0.933 2.723 1.708 0.088 1.294 0.404 4.149 0.433 0.665 0.969 0.824 1.140 -0.381 0.704 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value P-value 0.01	OR and 95% Cl OR and 95% Cl	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19 1.94 Relative weight 9.09 6.79 44.13 2.13 2.13 16.57
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Cossar et al 2008 Alagarasu et al 2007 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB vs BA+AA Wu et al 2015 da Cruz et al 2013 Singla et al 2012 Cossar et al 2008 Alagarasu et al 2007 Selvaraj et al 1998 Combined	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.31 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.594 0.933 2.723 1.708 0.088 1.294 0.404 4.149 0.433 0.665 0.969 0.824 1.140 -0.381 0.704 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 0.01	OR and 95% CI	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19 1.94 Relative weight 9.09 6.79 44.13 2.13 16.57 2.16
Soborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined <u>Study name</u> BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Singla et al 2012 Cossar et al 2007 Saborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1999 Bellamy et al 1999 Bellamy et al 1999 Bellamy et al 1998 Combined <u>Study name</u> BB vs BA+AA Wu et al 2015 da Cruz et al 2012 Cossar et al 2007 Singla et al 2012 Cossar et al 2007 Singla et al 2012 Cossar et al 2007 Liu et al 2013 Singla et al 2012 Cossar et al 2007 Liu et al 2007 Liu et al 2007	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 -0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value P-value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.331 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.594 0.933 2.723 1.708 0.088 1.294 0.404 4.149 0.433 0.665 1.294 0.404 4.149 0.433 0.665 1.294 0.404 4.149 0.433 0.665 0.869 0.824 1.140 -0.381 0.704 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value P-value 0.714 0.157 3.555 -0.367 0.714 0.201 0.01	OR and 95% Cl	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19 1.94 Relative weight 9.09 6.79 44.13 2.13 16.57 2.16
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Cossar et al 2010 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB vs BA+AA Wu et al 2015 da Cruz et al 2013 Singla et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Singla et al 2012 Cossar et al 2013 Singla et al 2012 Cossar et al 2013 Singla et al 2012 Cossar et al 2008 Alagarasu et al 2007 Seborg et al 2007 Seborg et al 2007 Sudy name BB vs BA+AA	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value P-value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.31 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.594 0.933 2.723 1.708 0.088 1.294 0.404 4.149 0.433 0.665 0.969 0.824 1.140 -0.381 0.704 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value P-value 0.01	OR and 95% CI	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.09 10.47 3.62 10.75 3.59 9.19 1.94 Relative weight 9.09 6.79 44.13 2.16 11.26
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined <u>Study name</u> BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Cossar et al 2008 Alagarasu et al 2007 Selvaraj et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined <u>Study name</u> BB vs BA+AA Wu et al 2015 da Cruz et al 2013 Singla et al 2012 Cossar et al 2008 Alagarasu et al 2007 Seborg et al 2007 Liu et al 2015 da Cruz et al 2013 Singla et al 2017 Seborg et al 2007 Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.31 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.594 0.933 2.723 1.708 0.088 1.294 0.404 4.149 0.433 0.665 0.969 0.824 1.140 -0.381 0.704 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 0.747 0.157 3.555 -0.367 0.714 0.826 0.136 5.027 -0.207 0.836 0.483 0.238 0.980 -2.015 0.044 1.194 0.047 30.134 0.108 0.180 0.2204 0.694 7.002 1.340 0.180 0.318 0.013 7.831 -0.701 0.483 1.518 0.373 6.173 0.583 0.560 0.318 0.013 7.831 -0.701 0.483 1.518 0.373 6.173 0.583 0.560 0.317 0.011 4.105 -1.019 0.308 8.043 1.043 62.008 2.000 0.045	OR and 95% CI	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19 1.94 Relative weight 9.09 6.79 44.13 2.13 16.57 2.16 11.26 2.56
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Araujo et al 2013 Cossar et al 2008 Alagarasu et al 2007 Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB vs BA+AA Wu et al 2015 da Cruz et al 2012 Cossar et al 2007 Alagarasu et al 2007 Singla et al 2015 da Cruz et al 2013 Singla et al 2012 Cossar et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Combined	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 Statistics for each study Lower Upper OR limit limit Z-value P-value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.331 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.294 0.404 4.149 0.433 0.665 0.969 0.824 1.140 -0.381 0.704 UDUE COME UDUE COME UDUE COME UDUE 0.01	OR and 95% Cl	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19 1.94 Relative weight 9.09 6.79 44.13 2.13 16.57 2.16 11.26 5.31
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Cossar et al 2013 Cossar et al 2010 Carbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB vs BA+AA Wu et al 2015 da Cruz et al 2013 Singla et al 2007 Liu et al 2015 da Cruz et al 2013 Singla et al 2012 Cossar et al 2020 Alagarasu et al 2007 Seborg et al 2007 Liu et al 2015 da Cruz et al 2013 Singla et al 2012 Cossar et al 2008 Alagarasu et al 2007 Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Combined	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value P-value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.334 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.31 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.594 0.933 2.723 1.708 0.088 1.294 0.404 4.149 0.433 0.665 0.969 0.824 1.140 -0.381 0.704 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value P-value 0.01	OR and 95% CI OR and 95% CI	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19 1.94 Relative weight 9.09 6.79 44.13 2.16 11.26 2.56 5.31
Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Bellamy et al 1998 Combined Study name BB+BA vs AA Wu et al 2015 da Cruz et al 2013 Araujo et al 2013 Araujo et al 2013 Cossar et al 2008 Alagarasu et al 2007 Selvaraj et al 1999 Bellamy et al 1999 Bellamy et al 1998 Combined Study name BB vs BA+AA Wu et al 2015 da Cruz et al 2013 Singla et al 2012 Cossar et al 2008 Alagarasu et al 2007 Seborg et al 2017 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Singla et al 2017 Seborg et al 2007 Liu et al 2006 Ozbaş-Gerçeker et al 2003 Selvaraj et al 1999 Combined	0.649 0.273 1.543 -0.978 0.328 1.305 0.780 2.183 1.013 0.311 0.855 0.356 2.051 -0.351 0.726 1.303 0.747 2.272 0.933 0.351 1.294 0.404 4.149 0.433 0.665 0.969 0.820 1.146 -0.362 0.717 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 1.154 0.755 1.765 0.661 0.508 1.305 0.698 2.440 0.834 0.404 0.877 0.520 1.479 -0.493 0.622 0.710 0.521 0.967 -2.170 0.030 0.441 0.140 1.390 -1.397 0.162 1.046 0.633 1.727 -0.176 0.861 0.603 0.257 1.416 -1.162 0.245 1.31 0.812 2.184 1.133 0.257 0.713 0.303 1.678 -0.776 0.438 1.594 0.933 2.723 1.708 0.088 1.294 0.404 4.149 0.433 0.665 0.969 0.824 1.140 -0.381 0.704 0.01 <u>Statistics for each study</u> Lower Upper OR limit limit Z-value <i>P</i> -value 0.747 0.157 3.555 -0.367 0.714 0.826 0.136 5.027 -0.207 0.836 0.483 0.238 0.980 -2.015 0.044 1.194 0.047 30.134 0.108 0.180 0.2204 0.694 7.002 1.340 0.180 0.318 0.013 7.831 -0.701 0.483 1.518 0.373 6.173 0.553 0.560 0.431 0.013 7.831 -0.701 0.483 1.518 0.373 6.173 0.553 0.560 0.437 0.157 1.505 -0.367 0.714 0.826 0.136 5.027 -0.207 0.836 0.483 0.238 0.980 -2.015 0.044 1.194 0.047 30.134 0.108 0.180 0.210 0.11 4.105 -1.019 0.308 8.043 1.043 62.008 2.000 0.045 0.876 0.547 1.402 -0.552 0.581	0.1 1 10 100 OR and 95% CI 0.1 1 10 100 OR and 95% CI OR and 95% CI OR and 95% CI	3.75 10.61 3.67 9.09 2.07 Relative weight 14.60 6.73 9.63 27.48 2.00 10.47 3.62 10.75 3.59 9.19 1.94 Relative weight 9.09 6.79 44.13 2.13 16.57 2.16 11.26 2.56 5.31

C Decreases | Increases

Figure 2 Forest plot of ORs with 95% CI of PTB risk associated with the MBL2 rs1800450 (A>B) gene polymorphism for the overall population.

Notes: Black squares represent the value of OR and the size of the square indicates the inverse proportion relative to its variance. Horizontal lines represent the 95% CI of OR.

Comparisons	Egger's regre	ssion analysis	Heterogen	eity analysis	Model used for the		
	Intercept	95% CI	P-value	Q-value	P _{heterogeneity}	l² (%)	meta-analysis
B vs A	6.83	4.07 to 9.58	0.00	14.49	0.01	72.40	Random
BB vs AA	3.58	-0.66 to 7.84	0.07	11.46	0.02	65.08	Random
BA vs AA	3.82	-1.56 to 9.21	0.11	5.30	0.26	24.53	Fixed
BB + BA vs AA	6.06	1.82 to 10.30	0.02	9.37	0.05	57.33	Random
BB vs BA + AA	3.24	-1.01 to 7.49	0.09	10.26	0.04	61.02	Random

 Table 6 Statistics to test the publication bias and heterogeneity (rs1800450 Asian)

 Table 7 Statistics to test the publication bias and heterogeneity (rs1800451 overall)

Comparisons	Egger's regression analysis			Heterogene	eity analysis		Model used for the
	Intercept	95% CI	P-value	Q-value	P _{heterogeneity}	l² (%)	meta-analysis
C vs A	1.07	-0.93 to 3.07	0.22	14.49	0.02	58.58	Random
CC vs AA	-0.29	-1.57 to 0.99	0.52	1.29	0.86	0.001	Fixed
CA vs AA	1.19	-1.00 to 3.38	0.22	15.61	0.02	61.57	Random
CC + CA vs AA	1.18	-1.02 to 3.39	0.23	15.91	0.01	62.28	Random
CC vs CA + AA	-0.26	-1.16 to 0.64	0.42	0.71	0.95	0.001	Fixed

 Table 8 Statistics to test the publication bias and heterogeneity (rs1800451 African)

Comparisons	Egger's regression analysis			Heterogene	ity analysis		Model used for the
	Intercept	95% CI	P-value	Q-value	P _{heterogeneity}	l² (%)	meta-analysis
C vs A	-0.89	-23.14 to 21.35	0.70	2.09	0.35	4.29	Fixed
CC vs AA	-0.53	-16.60 to 15.54	0.75	1.22	0.54	0.00	Fixed
CA vs AA	-1.08	-21.67 to 19.50	0.62	1.78	0.41	0.00	Fixed
CC + CA vs AA	-1.10	-24.87 to 22.66	0.66	2.22	0.33	10.05	Fixed
CC vs CA + AA	–0.3 I	-11.52 to 10.69	0.78	0.59	0.74	0.00	Fixed

P=0.026; OR =1.563, 95% CI =1.054–2.317) of PTB in comparison to wild allele A. Other genetic models, homozygous (DD vs AA: *P*=0.168; OR =3.244, 95% CI =0.610–17.263), heterozygous (DA vs AA: *P*=0.258; OR =1.281, 95% CI =0.834–1.969), dominant (DD+ DA vs AA: *P*=0.088; OR =1.432, 95% CI =0.947–2.164), and recessive (DD vs DA + AA: *P*=0.177; OR =3.164, 95% CI =0.595–16.829), were not associated with any risk to PTB (Figure 6).

MBL2 combined rs1800450, rs1800451, rs5030737 polymorphism

A total of seven case–control studies provided enough data to calculate ORs, including 1,340 controls and 1,445 PTB cases. Publication bias was not found, but heterogeneity was present in all the genetic models (Table 10; Figure SI6). The pooled ORs revealed that combined gene polymorphism was not associated with PTB in allelic contrast (O vs A: P=0.063; OR =1.524, 95% CI =0.978–2.375), homozygous (OO vs AA: P=0.070; OR =2.797, 95% CI =0.920–8.508), heterozygous (OA vs AA: P=0.095; OR =1.481, 95% CI =0.934–2.347),

dominant (OO+ OA vs AA: *P*=0.075; OR =1.619, 95% CI =0.953–2.749), and recessive (OO vs OA+ AA: *P*=0.061; OR =2.297, 95% CI =0.962–5.485) genetic models (Figure 7).

The stratification analysis based on the ethnicity included three studies from the Asian population. There was no publication bias and heterogeneity (Table 11; Figure SI7). Increased risk was observed with allelic (O vs A: P=0.001; OR =1.502, 95% CI =1.183–1.907), homozygous (OO vs AA: P=0.001; OR =3.092, 95% CI =1.567–6.104) and recessive (OO vs OA+ AA: P=0.002; OR =2.855, 95% CI =1.464–5.569) genetic models. Other genetic models, heterozygous (OA vs AA: P=0.241; OR =1.201, 95% CI =0.884–1.631) and dominant (OO+ OA vs AA: P=0.023; OR =1.404, 95% CI =1.048–1.880), did not reveal any risk with PTB (Figure 8).

MBL2 rs7096206 polymorphism

A total of five case–control studies were included which provided 3,139 controls and 1,201 PTB cases. There was no publication bias, but heterogeneity was found in three genetic models (Table 12; Figure SI8). The pooled ORs

Study name		Statistics for each study	,		<u>OR</u>	and 95% CI			
B vs A		Lower Upper							Relative
	OR	limit limit Z-value	P-value						weight
Wu et al 2015 Singla et al 2012	0 717	0.556 0.925 -2.559	0.619						20.39
Alagarasu et al 2007	1.170	0.762 1.798 0.718	0.473			╽╼╆			19.05
Liu et al 2006	1.309	0.842 2.035 1.196	0.232			│ ┼╋╌	4		18.71
Selvaraj et al 1999	1.795	1.113 2.894 2.400	0.016						17.68
Combined	1.131	0.815 1.571 0.737	0.461		I		I I	I	
				0.1	0.2	0.5 1	2 5	10	
Study name		Statistics for each study	<u>_</u>		<u>0R</u>	and 95% CI			Deletive
BB VS AA	OR	limit limit Z-value	P-value						weight
Wu et al 2015	0.777	0.163 3.172 -0.317	0.752		- I ·		-	1	17.63
Singla et al 2012	0.434	0.211 0.891 –2.275	0.023						27.60
Alagarasu et al 2007	2.150	0.671 6.893 1.288	0.198			╶┼╋	-		22.15
Liu et al 2006 Selvarai et al 1999	8 584	1 109 66 470 2 059	0.506					_	19.32 13.20
Combined	1.131	0.498 3.465 0.550	0.582				. 1		13.23
				0.01	0.1	1	1 10	100	
				0.01	0.1	•	10	100	
Study name		Statistics for each study	<u>'</u>		<u>OR</u>	and 95% CI			
BA vs AA	OR	Lower Upper limit limit Z-value	<i>P</i> -value						Relative weight
Wu et al 2015	1 185	0.768 1.830 0.766	0 443	I	1	- ia -	I	I.	20.80
Singla et al 2012	0.763	0.553 1.053 -1.645	0.100			∎			37.87
Alagarasu et al 2007	0.912	0.535 1.555 -0.338	0.735						13.80
Liu et al 2006	1.305	0.780 2.183 1.013	0.311						14.83
Selvaraj et al 1999	1.303	0.747 2.272 0.933	0.351						12.70
Combined	0.995	0.013 1.211 -0.000	0.347	I 0.01	I	▼ 1	1 10	100	
				0.01	0.1	I	10	100	
Study name		Statistics for each study	<u>'</u>		<u>OR</u>	and 95% CI			
BB+BA vs AA	0.0	Lower Upper							Relative
		limit limit Z-value	P-value						weight
Wu et al 2015 Singla et al 2012	0.710	0.755 1.765 0.661	0.508						20.94
Alagarasu et al 2007	1.046	0.633 1.727 0.176	0.861						25.94 18.01
Liu et al 2006	1.331	0.812 2.184 1.133	0.257			│ -∓∎-	+		18.25
Selvaraj et al 1999	1.594	0.933 2.723 1.708	0.088						16.86
Combined	1.083	0.800 1.467 0.517	0.605		I		I	I	
				0.1	0.2	0.5 1	2 5	10	
Study name		Statistics for each at the				and 95% CI			
BB vs BA+AA		Lower Lipper							Relative
	OR	limit limit Z-value	P-value						weight
Wu et al 2015	0.747	0.157 3.555 -0.367	0.714		-		1	1	17.23
Singla et al 2012	0.483	0.238 0.980 -2.015	0.044			╶╋┥			28.72
Alagarasu et al 2007	2.204	0.694 7.002 1.340	0.180			┼╋	-1		22.30
Liu et al 2006 Selvarai et al 1999	1.518 8.043	1.043 62.008 2.000	0.045					_	19.08
Combined	1.298	0.521 3.231 0.560	0.576				٦		.2.01
				. 0.01	0.1	1	10	100	
						eases Incre	ases L		

Figure 3 Forest plot of ORs with 95% CI of PTB risk associated with the MBL2 rs1800450 (A>B) gene polymorphism for the Asian population. Notes: Black squares represent the value of OR and the size of the square indicates the inverse proportion relative to its variance. Horizontal lines represent the 95% CI of OR.

Study name	Statistics for each study	OR and 95% CI	
C vs A	Lower Upper		Relative
	OR limit limit Z-value P-value		weight
da Cruz et al 2013 Araujo et al 2013	2.750 1.205 6.278 2.402 0.016 2.372 0.429 13.108 0.990 0.322		5.35
Thye et al 2011	0.930 0.848 1.021 -1.529 0.126		33.04
Alagarasu et al 2007	0.681 0.313 1.483 -0.968 0.333		5.90
Søborg et al 2007	0.962 0.751 1.233 -0.303 0.762		23.47
Selvaraj et al 1999 Bellamy et al 1998	2.159 0.924 5.042 1.778 0.075		25.73
Combined	0.990 0.804 1.217 -0.099 0.921		
		0.01 0.1 1 10 100	
Chudu name	Obsticution for each shade		
	Statistics for each study	<u>OR and 95% CI</u>	Delether
	OR limit limit Z-value <i>P</i> -value		weight
Thye et al 2011	0.855 0.695 1.051 -1.486 0.137	🖬	78.26
Alagarasu et al 2007	0.717 0.044 11.617 -0.234 0.815		0.43
Søborg et al 2007	0.938 0.494 1.781 -0.196 0.844		8.16
Selvaraj et al 1999 Bellamy et al 1998	0.585 0.036 9.456 -0.377 0.706		0.43 12 71
Combined	0.828 0.689 0.994 -2.021 0.043		12.71
		0.01 0.1 1 10 100	
Study name	Statistics for each study	OR and 95% CI	Relative
CAVSAA	OR limit limit Z-value P-value		weight
da Cruz et al 2013	2.895 1.246 6.727 2.470 0.013		7.85
Araujo et al 2013	2.405 0.430 13.455 0.999 0.318		2.31
Thye et al 2011 Alagarasu et al 2007	0.944 0.830 1.075 -0.863 0.388		29.93 7.57
Søborg et al 2007	0.963 0.707 1.311 -0.241 0.810		22.73
Selvaraj et al 1999	2.962 1.087 7.879 2.125 0.034		6.10
Bellamy et al 1998	0.765 0.572 1.024 -1.802 0.072		23.49
Combined	1.000 0.000 1.070 0.000 0.724		
	0.01	3	
Study name	Statistics for each study	OR and 95% CI	
CC+CA vs AA	Lower Upper		Relative
1- 0	OR limit limit 2-value P-value	e <u> </u>	7 40
da Cruz et al 2013 Araujo et al 2013	2.895 1.246 6.727 2.470 0.013 2.405 0.430 13.455 0.999 0.318		2.16
Thye et al 2011	0.926 0.819 0.047 -1.231 0.218		29.88
Alagarasu et al 2007	0.662 0.288 1.520 -0.973 0.330	│ │ ─┼● <u>→</u> ─│ │ │	7.59
Søborg et al 2007 Selvarai et al 1999	0.959 0.716 1.284 -0.281 0.779		22.92
Bellamy et al 1998	0.743 0.562 0.980 -2.099 0.036		23.53
Combined	1.025 0.788 1.332 0.181 0.856		
		0.1 0.2 0.5 1 2 5 10	
Study name	Statistics for each study	OB and 95% CI	
CC vs CA+AA	Lower Upper		Relative
	OR limit limit Z-value P-value	9	weight
Thye et al 2011	0.879 0.721 1.070 -1.284 0.199		79.02
Alagarasu et al 2007	0.748 0.046 12.109 -0.204 0.838		0.40
Søborg et al 2007 Selvaraj et al 1999	0.537 0.033 8.675 -0.438 0.662		0.40
Bellamy et al 1998	0.725 0.442 1.190 -1.272 0.203		12.57
Combined	0.861 0.722 1.026 -1.678 0.093	Ⅰ Ⅰ ◀ Ⅰ Ⅰ	
		0.01 0.1 1 10 100	
		C Decreases Increases	

Figure 4 Forest plot of ORs with 95% CI of PTB risk associated with the rs1800451 (A>C) gene polymorphism for the overall population.

Notes: Black squares represent the value of OR and the size of the square indicates the inverse proportion relative to its variance. Horizontal lines represent the 95% CI of OR.

<u>Study name</u> C vs A		<u>Statistic</u> Lower	<u>cs for ea</u> Upper	ich study				OR and	<u>95% C</u>	<u>:I</u>			Relative
Thye et al 2011 Søborg et al 2007 Bellamy et al 1998 Combined	OR 0.930 0.962 0.790 0.912	limit 0.848 0.751 0.639 0.842	limit 1.021 1.233 0.978 0.989	Z-value -1.529 -0.303 -2.162 -2.239	<i>P</i> -value 0.126 0.762 0.031 0.025	0.5	-						weight 75.35 10.47 14.17
						0.5						2	
Study name		<u>Statistic</u>	<u>cs for ea</u>	<u>ich study</u>				OR and	<u>95% C</u>	<u>: </u>			
CC vs AA	OR	Lower limit	Upper limit	Z-value	<i>P</i> -value								Relative weight
Thye et al 2011	0.855	0.695	1.051	-1.486	0.137			- I ·	-				78.95
Søborg et al 2007	0.938	0.494	1.781	-0.196	.0.844				-	-1			8.23
Bellamy et al 1998	0.638	0.382	1.067	-1.712	0.087			_+-	-				12.82
Combined	0.830	0.690	0.997	-1.990	0.047			·					
						0.1	0.2	0.5	1	2	5	10	
Study name		Statistic	cs for ea	ich studv				OR and	95% C	; <u> </u>			
CA vs AA		Lower	Upper	<u></u>									Relative
	OR	limit	limit	Z-value	P-value			_					weight
Thye et al 2011	0.944	0.830	1.075	-0.863	0.388			-					72.69
Søborg et al 2007	0.963	0.707	1.311	-0.241	0.810				╼┼──				12.85
Bellamy et al 1998	0.765	0.572	1.024	-1.802	0.072			-					14.46
Combined	0.918	0.822	1.026	-1.508	0.132							I	
						0.5			1			2	
Study name		<u>Statistic</u>	s for ea	ch study				OR and	95% C	<u>:I</u>			
CC+CA vs AA	OR	Lower l limit	Upper limit	Z-value	<i>P</i> -value								Relative weight
Thye et al 2011	0.926	0.819	1.047	-1.231	0.218			-	+-				72.84
Søborg et al 2007	0.959	0.716	1.284	-0.281	0.779				-				12.93
Bellamy et al 1998	0.743	0.562	0.980	-2.099	0.036			-	-				14.23
Combined	0.901	0.812	1.001	-1.943	0.052								
						0.5			1			2	
Study name		Statistic	s for ea	ch study				OR and	95% C	<u>:</u>			
CC vs CA+AA	OR	Lower l	Upper limit	Z-value	<i>P</i> -value								Relative weight
Thve et al 2011	0.879	0 721	1 070	-1 284	0 799	I	I	1		Т	I	I.	79.65
Søborg et al 2007	0.948	0.502	1.790	-0.165	0.869					_			7.68
Bellamy et al 1998	0.725	0.442	1.190	-1.272	0.203				┛				12.67
Combined	0.863	0.723	1.029	-1.644	0.100								
						0 .1	0.2	∎ 0.5	• 1	2	5	1 0	
								Decrease	s Inc	reases	\Box		

Figure 5 Forest plot of ORs with 95% CI of PTB risk associated with the rs1800451 (A>C) gene polymorphism for the African population.

Notes: Black squares represent the value of OR and the size of the square indicates the inverse proportion relative to its variance. Horizontal lines represent the 95% CI of OR.

Comparisons	Egger's regre	ession analysis		Heteroger	neity analysis	Model used for the	
	Intercept	95% CI	P-value	Q-value	P _{heterogeneity}	l² (%)	meta-analysis
D vs A	-0.81	-8.87to 7.25	0.77	1.98	0.74	0.00	Fixed
DD vs AA	5.91	-153.15 to 165.67	0.72	0.81	0.67	0.00	Fixed
DA vs AA	-2.07	-22.86 to 18.71	0.77	6.43	0.17	37.77	Fixed
DD + DA vs AA	-0.0002	-12.68 to 12.68	0.99	3.36	0.50	0.00	Fixed
DD vs DA + AA	5.22	-165.83 to 176.28	0.76	0.87	0.65	0.00	Fixed

Table 9 Statistics to test the publication bias and heterogeneity (rs5030737 overall)

Table 10 Statistics to test the publication bias and heterogeneity (combined overall)

Comparisons	Egger's regression analysis			Heterogen	eity analysis	Model used for the	
	Intercept	95% CI	P-value	Q-value	P _{heterogeneity}	l² (%)	meta-analysis
O vs A	0.14	-11.91 to 12.18	0.98	67.58	0.00	91.12	Random
OO vs AA	1.29	-6.66 to 9.24	0.69	50.46	0.00	88.11	Random
OA vs AA	1.46	-8.71 to 11.63	0.73	42.54	0.00	85.89	Random
OO + OA vs AA	2.34	-9.59 to 14.27	0.64	61.81	0.00	90.29	Random
OO vs OA + AA	1.16	-5.02 to 7.36	0.65	31.66	0.00	81.05	Random

 Table II Statistics to test the publication bias and heterogeneity (combined Asian)

Comparisons	Egger's regression analysis			Heterogene	eity analysis	Model used for the	
	Intercept	95% CI	P-value	Q-value	P _{heterogeneity}	l² (%)	meta-analysis
O vs A	1.38	-18.90 to 21.54	0.56	0.69	0.71	0.00	Fixed
OO vs AA	1.69	-25.74 to 29.13	0.58	1.48	0.48	0.00	Fixed
OA vs AA	1.95	0.11 to 3.78	0.05	0.65	0.72	0.00	Fixed
OO + OA vs AA	1.72	-8.37 to 11.82	0.27	0.60	0.74	0.00	Fixed
OO vs OA + AA	1.46	-28.82 to 31.73	0.65	0.56	0.46	0.00	Fixed

revealed that Y>X polymorphism was not associated with PTB in allelic contrast (X vs Y: P=0.657; OR =1.056, 95% CI =0.830–1.344), homozygous (XX vs YY: P=0.707; OR =0.932, 95% CI =0.647–1.344), heterozygous (XY vs YY: P=0.476; OR =1.107, 95% CI =0.838–1.462), dominant (XX+ YX vs YY: P=0.548; OR =1.091, 95% CI =0.821–1.449), and recessive (XX vs XY+YY: P=0.685; OR =0.928, 95% CI =0.646–1.333) genetic models (Figure 9).

The ethnicity-based stratification analysis included three studies from the Asian population. Publication bias was not found, whereas heterogeneity was detected in two genetic models (Table 13; Figure SI9). Likewise, no risk was observed with allelic (X vs Y: P=0.905; OR =1.027, 95% CI =0.661–1.598), homozygous (XX vs YY: P=0.662; OR =0.884, 95% CI =0.509–1.536), heterozygous (XY vs YY: P=0.659; OR =1.059, 95% CI =0.822–1.363), dominant (XX+ YX vs YY: P=0.866; OR =1.042, 95% CI =0.642–1.691), and recessive (XX vs XY+YY: P=0.676; OR =0.890, 95% CI =0.517–1.534) genetic models (Figure 10).

MBL2 rs11003125 polymorphism

Four case–control studies were included to calculate the pooled ORs, including 2,685 controls and 2,556 PTB cases, wherein no publication bias was found but three genetic models showed heterogeneity (Table 14; Figure SI10). The pooled ORs did not demonstrate any risk with PTB in allelic contrast (L vs H: P=0.995; OR =0.999, 95% CI =0.805–1.241), homozygous (LL vs HH: P=0.878; OR =1.026, 95% CI =0.736–1.431), heterozygous (LH vs HH: P=0.430; OR =1.240, 95% CI=0.727–2.113), dominant (LL+ LH vs HH: P=0.537; OR =1.182, 95% CI =0.695–2.012), and recessive (LL vs LH+ HH: P=0.256; OR =0.918, 95% CI =0.791–1.064) genetic models (Figure 11).

The results of meta-analysis of MBL2 gene polymorphisms, ie, rs1800450 A>B, rs1800451 A>C, rs5030737 A>D, combined rs1800450, rs1800451, rs5030737 A>O, rs7096206 Y>X, and rs11003125 H>L and their association with PTB susceptibility have been summarized in Table 15.

Study name		<u>Stat</u>	istics for	each stud	dy		<u>OF</u>	and 95% (<u>CI</u>		
D vs A	OR	Lowe	er Upper	r Z-valı	ue <i>P</i> -valu	e					Relative weight
da Cruz et al 2013	1.376	6 0.596	3.176	0.747	0.455	T	1			Т	22.15
Araujo et al 2013	1.576	6 0.537	4.628	0.827	0.408				_		13.36
Alagarasu et al 2007	2.538	3 1.137	5.662	2.274	0.023				⊢│		24.07
Søborg et al 2007	1.247	0.430	3.617	0.406	0.685			<u>_</u>	-		13.67
Selvaraj et al 1999	1.250	0.580	2.689	0.586	0.558						26.74
Combined	1.500	0 1.004	2.317	2.225	0.020	 0.01	 0.1	1	 10	 100	
Study name		Statis	stics for e	each stud	У		OR and	1 95% CI			
DD vs AA		Lower	· Upper								Relative
	OR	limit	limit	Z-value	P-value	e					weight
da Cruz et al 2013	1.278	0.079	20.663	0.173	0.863			 	\rightarrow		36.08
Alagarasu et al 2007	3.589	0.170	75.704	0.822	0.411		-		■		30.06
Selvaraj et al 1999	8.003	0.452	141.572	1.419	0.156						33.86
Combined	3.244	0.610	17.263	1.380	0.168					I	
						0.01	0.1	1	10	100	
Study name		Statis	stics for e	each stud	У		OR and	<u>1 95% CI</u>			
DA vs AA	OP	Lower	· Upper	7-value	P-value	2					Relative
da Cruz et al 2013	1 / 20	0 557	3 617	0 735	0.463	, I					21 13
Araujo et al 2013	1.420	0.537	4 803	0.735	0.403				_		15.24
Alagarasu et al 2007	2.329	1.006	5.392	1.974	0.048						26.22
Søborg et al 2007	1.250	0.428	3.650	0.409	0.683				-		16.09
Selvaraj et al 1999	0.479	0.188	1.218	-1.546	0.122		-	-∎-{			21.22
Combined	1.281	0.834	1.969	1.130	0.258			•			
						0.01	0.1	1	10	100	
Study name		<u>Statis</u>	tics for ea	ach study	L		OR and	<u>1 95% CI</u>			
DD+DA vs AA		Lower	Upper	7-value	<i>P</i> -value						Relative
do Cruz ot al 2012	1 406	0.576	2 422	0 749	0 455						21 42
Araujo et al 2013	1.400	0.570	4 803	0.740	0.400						1/ 17
Alagarasu et al 2007	2.508	1.090	5.770	2.164	0.031					.	24.59
Søborg et al 2007	1.250	0.428	3.650	0.409	0.683		1 4	─┼┳─	┍┻╴╽		14.87
Selvaraj et al 1999	0.852	0.373	1.947	-0.381	0.703		+	╶╋╧──	4		24.96
Combined	1.432	0.947	2.164	1.703	0.088				⊦		
						0.1	0.2 0.5	1	2 5	10	
Study name		<u>Statisti</u>	cs for ea	ch study			OR and	<u>1 95% CI</u>			
DD vs DA+AA		Lower	Upper	7 '							Relative
	OR	limit	limit Z	2-value	P-value			-			weight
da Cruz et al 2013	1.246	0.077	20.128	0.155	0.877		+	╶╴╋╴╴			36.07
Alagarasu et al 2007 Selvarai et al 1000	3.225	0.153	67.923 148.512	0.753	0.451					-	30.07
Combined	3 164	0.595	16 829	1.452	0.146					\rightarrow	33.65
	0.104	2.000			5.111		I 0.1	1	10	100	
						0.01	U.1	1	UI	100	
						<	Uecrea	ses increa	ises 🖒	,	

Figure 6 Forest plot of ORs with 95% CI of PTB risk associated with the rs5030737 (A>D) gene polymorphism for the overall population. Notes: Black squares represent the value of OR and the size of the square indicates the inverse proportion relative to its variance. Horizontal lines represent the 95% CI of OR.

Study name		Statis	tics for	each stu	dy			OR	and 95%	CI		
O vs A	Le	ower	Upper		-							Relative
	OR	limit	limit	Z-value	P-value							weight
da Cruz et al 2013	1.633	1.057	2.525	2.208	0.027				⊥-■			13.95
Araujo et al 2013 Capparelli et al 2009	1.020	0.674	1.544	0.094	0.925			-	-			14.12
Alagarasu et al 2007	1.356	0.965	1.906	1.753	0.080				┼┳╌			15.29
Søborg et al 2007	0.873	0.694	1.100	-1.151	0.250			·	╉╴			15.44
Selvaraj et al 2006	1.659	0.887	3.102	1.586	0.113				-+-■	┣───		12.20
Selvaraj et al 1999	1.659	1.116	2.467	2.500	0.012							14.27
Combined	1.524	0.978	2.375	1.861	0.063	I	I	I		► I	I	
						0.1	0.2	0.5	1	2 5	10	
Study name	<u>S</u>	Statistic	s for ea	ch study				OR	and 95	<u>% CI</u>		
OO vs AA	L	ower	Upper									Relative
	OR	limit	limit	Z-value I	^p -value							weight
da Cruz et al 2013	1.775	0.573	5.498	0.994	0.320			-				14.62
Capparelli et al 2013	1.202	8 829	38 393	0.182 7 769	0.855							11.14
Alagarasu et al 2007	2.457	1.013	5.961	1.998	0.047						- 1	15.53
Søborg et al 2007	0.688	0.387	1.222	-1.277	0.201			-+-	┣┼╴			16.47
Selvaraj et al 2006	2.569	0.562	11.757	1.216	0.224			-	-	╎╋╴┼		13.03
Selvaraj et al 1999 Combined	6.991	1.593	30.682	2.577	0.010							13.20
Combined	2.797	0.920	0.000	1.012	0.070	I	I	I				
						0.1	0.2	0.5	1	2 5	10	
Study name	5	Statistic	cs for ea	ach study				OF	and 95	<u>% CI</u>		
OA vs AA	L	ower	Upper									Relative
	OR	limit	limit	Z-value	^D -value							weight
da Cruz et al 2013	1.834	1.064	3.161	2.184	0.029				Ŀ₽			13.87
Araujo et al 2013	1.009	0.622	1.637	0.037	0.971				-	_		14.45
Capparelli et al 2009	4.258	2.885	0.285	7.293	0.000				·	■		15.31 14.84
Søborg et al 2007	0.945	0.712	i 1.720	-0.379	0.040				T			16.08
Selvaraj et al 2006	1.627	0.714	3.711	1.158	0.247				∓∎	-		11.09
Selvaraj et al 1999	1.190	0.726	1.948	0.689	0.491				+			14.36
Combined	1.481	0.934	2.347	1.670	0.095				•			
						0.01	(0.1	1	10	100	
Study name		Statistic	e for or	ach etudy	,			OF	and 05	% CI		
00+0A vs AA	<u> </u>	ower	Unnor	<u>ion study</u>				01		<u>/// 01</u>		Relative
00.0110111	OR	limit	limit	Z-value	P-value							weight
da Cruz et al 2013	1.825	1.090	3.057	2.288	0.022	1		1		1	1	14.06
Araujo et al 2013	1.016	0.630	1.638	0.064	0.949				-#-			14.33
Capparelli et al 2009	5.418	3.717	7.898	8.788	0.000					=		14.99
Alagarasu et al 2007	1.266	0.831	1.928	1.099	0.272				_=			14.72
Søborg et al 2007	0.897	0.680	1.182	-0.772	0.440				■_			15.54
Selvaraj et al 1999	1.762	0.808	2 370	1.425	0.134					-		14.35
Combined	1.619	0.953	2.749	1.782	0.075							11.00
						0.01	().1	1	10	100	
Study name	<u>S</u>	Statistic	cs for ea	ach study	[<u>OR</u>	and 95%	CI		
OO vs OA+AA	OR L	_ower limit	Upper limit	Z-value	P-value							Relative weight
da Cruz et al 2013	1.479	0.483	4,525	0 686	0.493	I		I I		- 1	I	14 58
Araujo et al 2013	1.198	0.167	8.625	0.180	0.857					<u> </u>		9.66
Capparelli et al 2009	7.962	3.985	15.906	5.875	0.000				Γ			17.13
Alagarasu et al 2007	2.359	0.989	5.627	1.936	0.053				_ -∎	-		16.12
Søborg et al 2007	0.700	0.397	1.234	-1.232	0.218			1 -				17.76
Selvaraj et al 2006			A 1									4
Column-1 -1 -1 1000	2.132	0.482	9.420	0.998	0.318				╤┼═			12.31
Selvaraj et al 1999 Combined	2.132 6.539 2.297	0.482	9.420 28.359 5.485	0.998 2.508	0.318 0.012 0.061				┨			12.31 12.43
Selvaraj et al 1999 Combined	2.132 6.539 2.297	0.482 1.508 0.962	9.420 28.359 5.485	0.998 2.508 1.873	0.318 0.012 0.061							12.31 12.43

 $\langle \Box$ Decreases | Increases $\Box \rangle$

Figure 7 Forest plot of ORs with 95% CI of PTB risk associated with MBL2 combined rs1800450, rs1800451, rs5030737 (A>O) gene polymorphism for the overall population.

Notes: Black squares represent the value of OR and the size of the square indicates the inverse proportion relative to its variance. Horizontal lines represent the 95% CI of OR.

Study name	Statistics for each study	OR and 95% Cl	
O vs A	Lower Upper		Relative
OR	limit limit Z-value P-value		weight
Alagarasu et al 2007 1.356	0.965 1.906 1.753 0.080		49.22
Selvaraj et al 2006 1.659	0.887 3.102 1.586 0.113		14.07
Combined 1 502	1.110 2.407 2.300 0.012		30.21
	1.103 1.307 3.333 0.001		
		0.1 0.2 0.5 1 2 5 10	
Study name	Statistics for each study	OR and 95% Cl	
OO vs AA	Lower Upper		Relative
OR	limit limit Z-value P-value		weight
Alagarasu et al 2007 2.457	1.013 5.961 1.988 0.047		58.87
Selvaraj et al 2006 2.569	0.562 11.757 1.216 0.224		20.00
Selvaraj et al 1999 6.991	1.593 30.682 2.577 0.010	│ │ │ │ │ <mark>┤──┤■</mark>┤	21.14
Combined 3.092	1.567 6.104 3.254 0.001		
		0.1 0.2 0.5 1 2 5 10	
Study name	Statistics for each study	OR and 95% Cl	
OA vs AA	Lower Upper		Relative
Alagarasu et al 2007 1.109	0.712 1.726 0.457 0.648		47.77
Selvaraj et al 2000 1.027	0.714 3.711 1.150 0.247		13.70
Combined 1 201	0.884 1.621 1.172 0.241		30.40
Combined 1.201	0.004 1.031 1.173 0.241		
		0.1 0.2 0.5 1 2 5 10	
Study name	Statistics for each study	OR and 95% Cl	
		OK and 35 % Cr	Polativo
OC+OA VS AA OR	limit limit Z-value P-value		weight
Alagarasu et al 2007 1.266	0.831 1.928 1.099 0.272	│ │ │ ↓∰ →│ │ │	48.25
Selvaraj et al 2006 1.762	0.808 3.840 1.425 0.154	│ │ │ │ ─ <mark>─</mark> <mark>─</mark> ─│ │	14.07
Selvaraj et al 1999 1.473	0.915 2.370 1.594 0.111	┤││┼╋┼│││	37.69
Combined 1.404	1.048 1.880 2.276 0.023		
		0.1 0.2 0.5 1 2 5 10	
Study name	Statistics for each study	OR and 95% Cl	
OO vs OA+AA	Lower Upper		Relative
OR	limit limit Z-value P-value		weight
Alagarasu et al 2007 2.359	0.989 5.627 1.936 0.053		59.06
Selvaraj et al 2006 2.132	0.482 9.420 0.998 0.318		20.21
Selvaraj et al 1999 6.539	1.508 28.359 2.508 0.012		20.73
Combined 2.855	1 464 5 569 3 078 0 002		
	1.101 0.000 0.010 0.002		
		0.1 0.2 0.5 1 2 5 10	

Figure 8 Forest plot of ORs with 95% CI of PTB risk associated with MBL2 combined rs1800450, rs1800451, rs5030737 (A>O) gene polymorphism for the Asian population. Notes: Black squares represent the value of OR and the size of the square indicates the inverse proportion relative to its variance. Horizontal lines represent the 95% CI of OR.

Comparisons	Egger's regr	ession analysis		Heteroger	neity analysis	Model used for the		
	Intercept	95% CI	P-value	Q-value	P _{heterogeneity}	l² (%)	meta-analysis	
X vs Y	1.37	-4.30 to 7.03	0.50	12.54	0.01	68.11	Random	
XX vs YY	0.28	-5.45 to 6.02	0.88	5.13	0.27	22.10	Fixed	
XY vs YY	1.56	–2.99 to 6.11	0.35	11.08	0.03	63.13	Random	
XX + XY vs YY	1.44	-3.74 to 6.62	0.44	12.55	0.01	68.13	Random	
XX vs XY + YY	0.01	-4.88 to 4.91	0.99	3.62	0.46	0.00	Fixed	

 Table 12 Statistics to test the publication bias and heterogeneity (rs7096206 overall)

 Table 13 Statistics to test the publication bias and heterogeneity (rs7096206 Asian)

Comparisons	Egger's regression analysis			Heteroger	neity analysis	Model used for the	
	Intercept	95% CI	P-value	Q-value	P _{heterogeneity}	l² (%)	meta-analysis
X vs Y	9.80	-272.33 to 291.93	0.73	9.46	0.009	78.85	Random
XX vs YY	7.87	-117.43 to 133.17	0.57	5.07	0.08	60.57	Fixed
XY vs YY	0.09	-273.75 to 273.95	0.99	5.63	0.06	64.50	Fixed
XX + XY vs YY	-0.39	-321.95 to -321.15	0.99	7.93	0.02	74.78	Random
XX vs XY + YY	6.45	-90.85 to 103.75	0.55	3.52	0.17	43.15	Fixed

Table 14 Statistics to test the publication bias and heterogeneity (rs11003125 overall)

Comparisons	Egger's regression analysis			Heterogene	ity analysis	Model used for the	
	Intercept	95% CI	P-value	Q-value	P _{heterogeneity}	l² (%)	meta-analysis
L vs H	2.21	-10.49 to 14.91	0.53	7.98	0.05	62.39	Random
LL vs HH	1.88	-11.54 to 15.31	0.61	6.58	0.09	54.42	Fixed
LH vs HH	0.14	-12.97 to 13.24	0.97	7.81	0.05	61.58	Random
LL + LH vs HH	0.67	-12.35 to 13.69	0.85	8.64	0.03	65.26	Random
LL vs LH + HH	0.39	-6.83 to 7.59	0.84	4.65	0.20	35.43	Fixed

Sensitivity analysis

To examine the influence of each individual study on the overall risk of PTB, sensitivity analysis was accomplished. Leave-one-out sensitivity analysis (excluding single casecontrol study at a time) was performed and the pooled ORs recomputed for each genetic variant of MBL2. The same was tested during the subgroup analysis by the ethnicity. The ORs estimated for each polymorphism of MBL2 gene after excluding a single study did not show any major difference from the primary values (rs1800450 [A>B] [Overall: Figure SI11, Asian: Figure SI12]; rs1800451 [A>C] [Overall: Figure SI13, African: Figure SI14]; rs5030737 [A>D] [Overall: Figure SI15]; combined rs1800450, rs1800451, rs5030737 [A>O] [Overall: Figure SI16, Asian: Figure SI17]; rs7096206 [Y>X] [Overall: Figure SI18, Asian: Figure SI19], rs11003125 [H>L] [Overall: Figure SI20]) of these genetic variants. This analysis guaranteed the stability of the overall results of this meta-analysis.

Trial sequential analysis

TSA was used to investigate the relevance of MBL2 rs1800450 A>B, rs1800451 A>C, rs5030737 A>D, combined rs1800450, rs1800451, rs5030737 A>O, rs7096206 Y>X, and rs11003125 H>L gene polymorphisms with PTB susceptibility. The dominant model was used to study all these polymorphisms. As shown in Figure 12, the required information size for MBL2 rs1800450, rs1800451, and rs5030737 polymorphisms was 5,876, 24,257 and 3,611 patients, respectively. However, the accrued information size remained 4,047, 6,796, and 1,611. Similarly, for the promoter variants rs7096206, rs11003125, and combined structural polymorphism (A>O) of MBL2, the accrued information size was 4,340, 5,241 and 2,785, but the required information size was 19,554, 12,897, and 40,517, respectively (Figure 13). As depicted in Figures 12 and 13, the cumulative Z-curve of all the studied polymorphisms did not cross the trial monitoring boundary before reaching the required information size,

Study name X vs Y Wu et al 2015 da Cruz et al 2013 Thye et al 2011 Alagarasu et al 2007 Liu et al 2006 Combined	Statistics for each study Lower Upper OR limit limit Z-value P-value 1.276 0.916 1.776 1.441 0.149 1.447 0.922 2.270 1.608 0.108 0.953 0.834 1.089 -0.708 0.479 0.665 0.475 0.930 -2.386 0.017 1.297 0.877 1.918 1.305 0.192 1.056 0.830 1.344 0.444 0.657	OR and 95% Cl 0.01 0.1	Relative weight 19.71 14.97 28.63 19.51 17.18
Study name XX vs YY Wu et al 2015 da Cruz et al 2013 Thye et al 2011 Alagarasu et al 2007 Liu et al 2006 Combined	Statistics for each study Lower Upper OR limit limit Z-value P-value 1.530 0.606 3.860 0.900 0.368 0.978 0.267 3.583 -0.034 0.973 0.971 0.574 1.642 -0.111 0.912 0.407 0.170 0.974 -2.018 0.044 1.422 0.464 4.364 0.616 0.538 0.932 0.647 1.344 -0.376 0.707	OR and 95% Cl	Relative weight 15.60 7.92 48.30 17.55 10.63
Study name XY vs YY Wu et al 2015 da Cruz et al 2013 Thye et al 2011 Alagarasu et al 2007 Liu et al 2006	Statistics for each study Lower Upper OR limit limit Z-value P-value 1.284 0.855 1.928 1.205 0.228 1.833 1.058 3.177 2.161 0.031 0.943 0.811 1.096 -0.766 0.444 0.686 0.442 1.064 -1.683 0.092 1.352 0.840 2.175 1.243 0.214 1.107 0.838 1.462 0.713 0.476	OR and 95% Cl	Relative weight 19.57 14.54 30.63 18.29 16.97
Study name XX+XY vs YY Wu et al 2015 da Cruz et al 2013 Thye et al 2011 Alagarasu et al 2007 Liu et al 2006 Combined	Statistics for each study Lower Upper OR limit limit Z-value P-value 1.311 0.889 1.935 1.366 0.172 1.698 1.006 2.868 1.981 0.048 0.945 0.816 1.094 -0.761 0.447 0.637 0.418 0.969 -2.105 0.035 1.360 0.863 2.145 1.324 0.186 1.091 0.821 1.449 0.600 0.548	OR and 95% Cl	Relative weight 19.78 15.19 29.03 18.62 17.38
Study name XX vs XY+YY Wu et al 2015 da Cruz et al 2013 Thye et al 2011 Alagarasu et al 2007 Liu et al 2006 Combined	Statistics for each study Lower Upper OR limit limit Z-value P-value 1.419 0.568 3.550 0.749 0.454 0.823 0.227 2.987 -0.296 0.767 0.983 0.582 1.662 -0.063 0.650 0.476 0.203 1.116 -1.708 0.088 1.302 0.428 3.957 0.465 0.642 0.928 0.646 1.333 -0.405 0.685	OR and 95% CI	Relative weight 15.64 7.91 47.71 18.10 10.63

Figure 9 Forest plot of ORs with 95% CI of PTB risk associated with MBL2 rs7096206 (Y>X) gene polymorphism for the overall population. Notes: Black squares represent the value of OR and the size of the square indicates the inverse proportion relative to its variance. Horizontal lines represent the 95% CI of OR.

Study name	Statistics for each study	OR and 95% Cl	
X vs Y	Lower Upper		Relative
			weight
Alegeracy et al 2007 0 665	0.916 1.776 1.441 0.149		34.20
Liu et al 2006 1 207	0.877 1.918 1.305 0.192		31 70
Combined 1.027	0.661 1.598 0.120 0.905		01.70
		0.1 0.2 0.3 1 2 3 10	
Study name	Statistics for each study	OR and 95% CI	
XX vs YY	Lower Upper		Relative
OR	limit limit Z-value P-value		weight
Wu et al 2015 1.530	0.606 3.860 0.900 0.368	│ │ _│─┼╋┼──│ │	35.63
Alagarasu et al 2007 0.407	0.170 0.974 -2.018 0.044	│ │ ॑─॑<mark>॑</mark>┤<u></u>│ │ │ │	40.09
Liu et al 2006 1.422	0.464 4.364 0.616 0.538		24.28
Combined 0.884	0.509 1.536 -0.437 0.662		
		0.1 0.2 0.5 1 2 5 10	
Study name	Statistics for each study	OR and 95% Cl	
XY vs YY	Lower Upper	<u></u>	Relative
OR	limit limit Z-value P-value		weight
Wu et al 2015 1.284	0.855 1.928 1.205 0.228		38.64
Alagarasu et al 2007 0.686	0.442 1.064 -1.683 0.092		33.09
Liu et al 2006 1.352	0.840 2.175 1.243 0.214	│ │ │ ┼╋┼ │ │	28.27
Combined 1.059	0.822 1.363 0.442 0.659		
		0.1 0.2 0.5 1 2 5 10	
Study name	Statistics for each study	<u>OR and 95% Cl</u>	
XX+XY vs YY OR	Lower Upper limit limit Z-value P-value		Relative weight
Wu at al 2015 1 311			
	0 880 1 035 1 366 0 172		34.64
Alagarasu et al 2007 0.637	0.889 1.935 1.366 0.172 0.418 0.969 -2.105 0.035		34.64 33.38
Alagarasu et al 2007 0.637 Liu et al 2006 1.360	0.889 1.935 1.366 0.172 0.418 0.969 -2.105 0.035 0.863 2.145 1.324 0.186	│ │ ┼ <mark>┲</mark> ┼╋┤ │ │	34.64 33.38 31.98
Alagarasu et al 2007 0.637 Liu et al 2006 1.360 Combined 1.042	0.889 1.935 1.366 0.172 0.418 0.969 -2.105 0.035 0.863 2.145 1.324 0.186 0.642 1.691 0.168 0.866		34.64 33.38 31.98
Alagarasu et al 20070.637Liu et al 20061.360Combined1.042	0.889 1.935 1.366 0.172 0.418 0.969 -2.105 0.035 0.863 2.145 1.324 0.186 0.642 1.691 0.168 0.866		34.64 33.38 31.98
Alagarasu et al 2007 0.637 Liu et al 2006 1.360 Combined 1.042	0.889 1.935 1.366 0.172 0.418 0.969 -2.105 0.035 0.863 2.145 1.324 0.186 0.642 1.691 0.168 0.866		34.64 33.38 31.98
Alagarasu et al 2007 0.637 Liu et al 2006 1.360 Combined 1.042 Study name	0.889 1.935 1.366 0.172 0.418 0.969 -2.105 0.035 0.863 2.145 1.324 0.186 0.642 1.691 0.168 0.866	0.1 0.2 0.5 1 2 5 10 OR and 95% Cl	34.64 33.38 31.98
Alagarasu et al 2007 0.637 Liu et al 2006 1.360 Combined 1.042 Study name XX vs XY+YY OR	0.889 1.935 1.366 0.172 0.418 0.969 -2.105 0.035 0.863 2.145 1.324 0.186 0.642 1.691 0.168 0.866 Statistics for each study Lower Upper limit Limit Lower P-value	0.1 0.2 0.5 1 2 5 10 <u>OR and 95% Cl</u>	34.64 33.38 31.98 Relative weight
Alagarasu et al 2007 0.637 Liu et al 2006 1.360 Combined 1.042 Study name XX vs XY+YY OR Wu et al 2015 1.419	0.889 1.935 1.366 0.172 0.418 0.969 -2.105 0.035 0.863 2.145 1.324 0.186 0.642 1.691 0.168 0.866 Statistics for each study Lower Upper limit limit Z-value P-value 0.568 3.550 0.749 0.454	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	34.64 33.38 31.98 Relative weight 35.25
Alagarasu et al 2007 0.637 Liu et al 2006 1.360 Combined 1.042 Study name XX vs XY+YY OR 0R Wu et al 2015 1.419 Alagarasu et al 2007 0.476	0.889 1.935 1.366 0.172 0.418 0.969 -2.105 0.035 0.863 2.145 1.324 0.186 0.642 1.691 0.168 0.866 Statistics for each study Lower Upper limit Lower Upper limit Z-value P-value 0.568 3.550 0.749 0.454 0.203 1.116 -1.708 0.088	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	34.64 33.38 31.98 Relative weight 35.25 40.78
Alagarasu et al 2007 0.637 Liu et al 2006 1.360 Combined 1.042 Study name XX vs XY+YY OR Wu et al 2015 1.419 Alagarasu et al 2007 0.476 Liu et al 2006 1.302	0.889 1.935 1.366 0.172 0.418 0.969 -2.105 0.035 0.863 2.145 1.324 0.186 0.642 1.691 0.168 0.866 Statistics for each study Lower Upper limit 0.568 3.550 0.749 0.454 0.203 1.116 -1.708 0.088 0.428 3.957 0.465 0.642	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	34.64 33.38 31.98 Relative weight 35.25 40.78 23.96
Alagarasu et al 2007 0.637 Liu et al 2006 1.360 Combined 1.042 Study name XX vs XY+YY OR Wu et al 2015 1.419 Alagarasu et al 2007 0.476 Liu et al 2006 1.302 Combined 0.890	0.889 1.935 1.366 0.172 0.418 0.969 -2.105 0.035 0.863 2.145 1.324 0.186 0.642 1.691 0.168 0.866 Statistics for each study Lower Upper limit Z-value P-value 0.568 3.550 0.749 0.454 0.203 1.116 -1.708 0.088 0.428 3.957 0.465 0.642 0.517 1.534 -0.419 0.676	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	34.64 33.38 31.98 Relative weight 35.25 40.78 23.96
Alagarasu et al 20070.637Liu et al 20061.360Combined1.042Study nameXX vs XY+YYORWu et al 20151.419Alagarasu et al 20070.476Liu et al 20061.302Combined0.890	0.889 1.935 1.366 0.172 0.418 0.969 -2.105 0.035 0.863 2.145 1.324 0.186 0.642 1.691 0.168 0.866 Statistics for each study Lower Upper limit Z-value P-value 0.568 3.550 0.749 0.454 0.203 1.116 -1.708 0.088 0.428 3.957 0.465 0.642 0.517 1.534 -0.419 0.676	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	34.64 33.38 31.98 Relative weight 35.25 40.78 23.96

Figure 10 Forest plot of ORs with 95% CI of PTB risk associated with MBL2 rs7096206 (Y>X) gene polymorphism for the Asian population. Notes: Black squares represent the value of OR and the size of the square indicates the inverse proportion relative to its variance. Horizontal lines represent the 95% CI of OR.

Study name		Statis	tics for ea	ach study				<u>OR a</u>	nd 95% Cl	l		
L vs H	OR	Lower limit	Upper limit	Z-value	<i>P</i> -value							Relative weight
Wu et al 2015	1.083	0.834	1.405	0.598	0.550		Т	I	.	1	1	25.83
da Cruz et al 2013	1.458	0.999	2.128	1.955	0.051				⋤ _∎ ↓			18.26
Thye et al 2011	0.918	0.775	1.086	-0.999	0.318							33.03
Liu et al 2006	0.763	0.564	1.033	-1.751	0.080				.			22.88
Combined	0.999	0.805	1.241	-0.007	0.995				• I			
						0.1	0.2	0.5	• • 1 2	5	10	
						0.1	0.2	0.0			10	
Study name		Statis	stics for e	ach study				<u>OR a</u>	nd 95% C	l		
LL vs HH	OR	Lower limit	Upper limit	Z-value	<i>P</i> -value							Relative weight
Wu et al 2015	1.155	0.699	1.909	0.563	0.573			-				43.81
da Cruz et al 2013	2.305	0.944	5.629	1.833	0.067				-	∎──┼		13.86
Thye et al 2011	1.075	0.399	2.893	0.143	0.886				┢	-		11.27
Liu et al 2006	0.595	0.328	1.081	-1.705	0.088				-+			31.05
Combined	1.026	0.736	1.431	0.154	0.878			•	♦			
						0.1	0.2	0.5	1 2	5	10	
Study name		Statistics for each study						<u>OR a</u>	nd 95% C	l		
LH vs HH		Lower	Upper	7	D							Relative
	OR	limit	limit	Z-value	P-value							weight
Wu et al 2015	1.765	1.132	2.751	2.508	0.012					-		33.08
da Cruz et al 2013	1.752	1.704	4.359	1.206	0.228			-	┤_╺			19.05
Thye et al 2011	1.192	1.438	3.245	0.343	0.732				┤┛╌┤	_		17.07
Liu et al 2006	0.700	0.420	1.100	-1.370	0.171							30.80
Combined	1.240	0.727	2.113	0.789	0.430	I	I	1		I	I	
						0.1	0.2	0.5	1 2	5	10	
Study name		Statistics for each study						<u>OR a</u>	nd 95% C	l		
LL+LH vs HH		Lower	Upper	-								Relative
	OR	limit	limit	Z-value	P-value							weight
Wu et al 2015	1.527	1.012	2.303	2.017	0.044			1	┝╼═┹┼	-		32.80
da Cruz et al 2013	2.044	0.861	4.849	1.621	0.105				+			19.63
Thye et al 2011	1.090	0.405	2.933	0.171	0.864				╞	-		16.90
Liu et al 2006	0.663	0.411	1.069	-1.685	0.092			╶┼═	-			30.67
Combined	1.182	0.695	2.012	0.618	0.537			-				
						0.1	0.2	0.5	1 2	5	10	
Study name Statistics for each study						OR and 95% CI						
LL vs LH+HH		Lower	Upper	7-value	Pualua							Relative weight
M/	0.700			2-value		I	ı.	Ι.		I	I	12 42
vvu et al 2015	1 465	0.525	1.21/	-1.044	0.297				T_			12.42
da Cruz et al 2013	1.465	0.902	2.379	1.543	0.123				<u>,</u> • †	-		9.32
inye et al 2011	0.906	0.759	1.083	-1.084	0.278							09.00 9.77
Liu et al 2006	0.740	0.404	1.234	-1.137	0.200							0.77
Complined	0.918	0.791	1.004	-1.137	0.200	I	I	I	▼	1	I	
						0.1	0.2	0.5	1 2	5	10	
						<	$\langle \neg \rangle$	Decreases	s Incre	ases 🗌	\geq	

Figure 11 Forest plot of ORs with 95% CI of PTB risk associated with MBL2 rs I 1003125 (H>L) gene polymorphism for the overall population. Notes: Black squares represent the value of OR and the size of the square indicates the inverse proportion relative to its variance. Horizontal lines represent the 95% CI of OR.

MBL2 gene polymorphisms	Association	Ethnicity			
rs1800450 A>B	Overall	Asian			
B vs A	No risk	No risk			
BB vs AA	No risk	No risk			
BA vs AA	No risk	No risk			
BB + BA vs AA	No risk	No risk			
BB vs BA + AA	No risk	No risk			
rs1800451 A>C	Overall	African			
C vs A	No risk	Reduced risk			
CC vs AA	Reduced risk	Reduced risk			
CA vs AA	No risk	No risk			
CC + CA vs AA	No risk	No risk			
CC vs CA + AA	No risk	No risk			
rs5030737 A>D	Overall	Not studied			
D vs A	Increased risk	Not studied (due to inadequate studies)			
DD vs AA	No risk				
DA vs AA	No risk	_			
DD + DA vs AA	No risk				
DD vs DA + AA	No risk				
Combined rs1800450, 1800451, rs5030737 O>A	Overall	Asian			
O vs A	No risk	Increased risk			
OO vs AA	No risk	Increased risk			
OA vs AA	No risk	No risk			
OO + OA vs AA	No risk	No risk			
OO vs OA + AA	No risk	Increased risk			
rs7096206 Y>X	Overall	Asian			
X vs Y	No risk	No risk			
XX vs YY	No risk	No risk			
XY vs YY	No risk	No risk			
XX + YX vs YY	No risk	No risk			
XX vs XY + YY	No risk	No risk			
rs11003125 H>L	Overall	Not studied			
L vs H	No risk	Not studied (due to inadequate studies)			
LL vs HH	No risk				
LH vs HH	No risk				
LL + LH vs HH	No risk				
LL vs LH + HH	No risk				

Table 15 Summary of the results of association of MBL2 gene polymorphisms and PTB risk (overall and ethnicity-wise analysis)

Abbreviation: PTB, pulmonary tuberculosis.

indicating the cumulative evidence was not sufficient and further trials are needed (Figures 12 and 13).

Discussion

The passing years have proved the worth of SNPs as the most common and effective type of genetic variations studied in correlation with susceptibility to various diseases and have the potential to be considered as the markers for many complex diseases.⁴¹ The differences in susceptibility to infections are determined by the host genetic factors and the first line of the host defense is innate immunity. Therefore, genetic variations in components of innate immunity may influence susceptibility to infections.⁶

The humoral immunity against PTB is mediated by the innate immune components with the help of interference by many complement proteins. The diverse pattern-recognizing receptors or soluble pathogen-recognizing molecules (PRMs) enable innate immune system to recognize molecular motifs on pathogenic surfaces.⁶ One of the C-type soluble PRMs is MBL, which is secreted as a part of the acute-phase response by the liver. The MBL plays an important role in recognizing, initiating, regulating, and amplifying the innate immune defense.⁴²

The MBL triggers the lectin complement pathway after it recognizes the target microorganisms. The multiple opsonin C3b fragments are developed by this activation of



Figure 12 Trial sequence analysis of all the included studies dealing with MBL2 gene polymorphisms, based on dominant genetic model, and PTB risk: (A) rs1800450 (A>B), (B) rs1800451 (A>C), and (C) rs5030737 (A>D).

Abbreviations: PTB, pulmonary tuberculosis; RRR, relative risk reduction.



Figure 13 Trial sequence analysis of all the included studies dealing with MBL2 gene polymorphisms, based on dominant genetic model, and PTB risk: (A) combined rs1800450, rs1800451, rs5030737 (A>O), (B) rs7096206 (Y>X), and (C) rs11003125 (H>L). Abbreviation: PTB, pulmonary tuberculosis.

complement cascade, which eventually causes phagocytosis of the pathogen primarily by receptors of CR1 (CD35). The MBL is also believed to promote opsonophagocytosis of organisms directly and the clearance of apoptotic cells through various putative receptors.⁴³

Since MBL plays an important role in microorganism recognition during different stages of immunity, it possibly plays a significant role in the interaction of *M. tuberculosis* with the host. The MBL/MBP, a product of MBL2 gene, binds to the mannose groups present on several types of bacteria. MBP forms complexes with Mycobacteria and during diffusion in the blood enables their enhanced macrophage uptake. Thus, MBP binds and opsonizes Mycobacteria in order to boost their phagocytosis.⁴⁴ Earlier studies suggest that it is the phosphatidylinositol mannoside present on M. tuberculosis that binds with human MBP.45 Lipoarabinomannan present in M. tuberculosis cell wall acts as a binding protein, thereby enabling their uptake by the host macrophages. The variations in MBL protein serum levels have been reported to be influenced by SNPs located within the MBL2 gene promoter region and exon 1 by several studies.¹⁹⁻³¹ Inherited deficiency of MBL may impair the innate immune function, which may further lead to increased risk for the development of various infectious diseases including PTB.18

Previously, several case–control studies have reported the association of MBL2 gene polymorphisms and PTB susceptibility with inconsistency and lack of clarity. This may be because of their small sample sizes and ethnic variations leading to inadequate statistical power. Therefore, the current study was conducted to solve this conflict to provide a strong statistical basis and evidence for association of MBL2 gene polymorphisms and PTB susceptibility. The cumulative large sample size generates pooled ORs with sufficient statistical power and reduced random errors advantage.⁴⁶

This is the first dedicated meta-analytic study to evaluate the association between MBL2 gene polymorphisms and overall PTB susceptibility. This analysis included 14 studies considering all the eligible criteria. Most of the included studies have clearly described sample sizes, genotypes, case and healthy control descriptions. We found a significant reduced PTB risk associated with variant allele and homozygous genotype of rs1800451 (A>C) polymorphism. This suggests that individuals carrying homozygous genotype for MBL2 polymorphism may have a reduced risk for PTB in general. The homozygous variant may have a protective effect and helps the host in resisting infection and thereby PTB susceptibility. The enhanced resistance to mycobacterial infection might lead to high frequency of MBP variant alleles in diverse populations.⁴⁷ Conversely, variant allele of rs5030737 (A>D) showed increased risk to PTB. This indicates the dual role played by MBL2 polymorphism as a modifying gene in PTB development. Our data reiterate the support to the findings reporting involvement of the effects of combined host gene polymorphisms in the progression of *M. tuberculosis* infection. However, further large sample studies involving diverse populations are required for the confirmation.

Other MBL2 genetic polymorphisms, rs1800450 (A>B); combined polymorphisms rs1800450, rs1800451, rs5030737 (A>O); rs7096206 (Y>X); rs11003125 (H>L) did not show association with any increased or decreased risk of PTB in this analysis. Further investigations based on subgroup ethnicity found that rs1800451 (A>C) was associated with decreased PTB risk in the African population. Also, the combined polymorphisms (A>O) showed an increased PTB risk in the Asian population. However, the conclusions for the subgroup ethnicity analysis were suffered by the small sample size limitation which further warrants the studies to be conducted on a larger scale.

PTB is a multifactorial disease, which suggests that its susceptibility depends not only on the host genetic factors but also on the characteristics of infecting *Mycobacteria* and its interactions with other environmental factors.⁴⁸ The host and *M. tuberculosis* have evolved simultaneously over time, and the individual strains have adapted to their respective populations.⁴⁹ Hence, MBL2 gene polymorphisms alone could not be attributed for PTB predisposition.

As this study showed significant heterogeneity, we tried to minimize the likelihood of heterogeneity problem by performing the data analysis using random-effects model to obtain wider 95% CI. In addition, when studies were stratified by ethnicity, low heterogeneity was found in both, Asian and African population. Hence, we considered that the racial differences and ethnic origin of the study population, inadequate selection criteria of the subjects, and small sample size of each included study might be responsible for the foremost source of heterogeneity. Though, there are some limitations warranting to be addressed: firstly, the studies published in the English language and available from the selected electronic databases were included for this analysis, which makes it probable that some studies published in other languages or indexed in other electronic databases may have been missed; secondly, our results are based on unadjusted ORs; thirdly, our study is deficient in testing of gene-environment interactions because of insufficiency of the data available in the primary selected studies.

In opposition to the above stated limitations, the current study was complemented with several advantages as well including the testing of the publication bias and sensitivity analysis. No obvious publication bias was observed, and the results were robust as no single study yielded an obvious effect on the pooled ORs and the corresponding CIs during sensitivity analysis.

Conclusion

A meta-analysis is a powerful technique that provides a consensus using available data from different individual studies. The results of this study demonstrated that MBL2 rs1800451 (A>C) and rs5030737 (A>D) gene polymorphisms play a significant role in the risk of PTB disease. PTB has a significant public health impact. Hence, studies with a definite concept and character for MBL2 gene and infection or replication of M. tuberculosis are needed. The findings of this pooled study will help in augmenting the understanding associated with the role of MBL2 rs1800451 (A>C) and rs5030737 (A>D) gene polymorphisms and also aid in identifying the individuals prone to PTB infection. This eventually could be advantageous in achieving optimal therapeutic interventions to individual PTB patients. This study will also prove helpful for complete genotype profiling of MBL2 gene for PTB susceptibility and thereby would greatly assist in global control and outcome of this infectious disease. Moreover, this study also instigates the execution of studies with large sample sizes to decipher the potential molecular mechanism underlying PTB infection, ultimately aiding in development of molecular approaches for the detection of susceptible genotypes.

Acknowledgment

The authors are grateful to the Deanship of Scientific Research, Jazan University, Jazan-45142, Saudi Arabia, for the infrastructural (especially the Saudi Digital Library facility) and software related help for this research study.

Author contributions

Conceived and designed the study and experiments: RKM, MAK, AH, SAD, AJ, MW, AKP, ML, NA, SK, BNM, and SH. Performed the experiments: RKM, MAK, AH, SAD, AJ, AKP, SK, and SH. Analyzed the data: RKM, SAD, AKP, SK, and SH. Contributed reagents/materials/analysis tools: MAK, AH, MW, ML, NA, and BNM. Wrote the paper: RKM, SAD, AJ, SK, and SH. All authors reviewed the manuscript. All authors contributed toward data analysis, drafting and revising the paper, gave final approval of the version to be published and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

- World Health Organization (WHO). Global Tuberculosis Report; 2016. Available from: https://www.who.int/tb/publications/global_report/ gtbr2016_executive_summary.pdf. Accessed August 1, 2018.
- Frieden TR, Sterling TR, Munsiff SS, Watt CJ, Dye C. Tuberculosis. Lancet. 2003;362(9387):887–899.
- 3. Schluger NW. Changing approaches to the diagnosis of tuberculosis. *Am J Respir Crit Care Med.* 2001;164(11):2020–2024.
- Brown DH, Miles BA, Zwilling BS. Growth of *Mycobacterium tuber-culosis* in BCG-resistant and -susceptible mice: establishment of latency and reactivation. *Infect Immun.* 1995;63(6):2243–2247.
- Malik S, Schurr E. Genetic susceptibility to tuberculosis. *Clin Chem Lab Med.* 2002;40(9):863–868.
- Thiel S, Frederiksen PD, Jensenius JC. Clinical manifestations of mannan-binding lectin deficiency. *Mol Immunol*. 2006;43(1–2):86–96.
- Möller M, de Wit E, Hoal EG. Past, present and future directions in human genetic susceptibility to tuberculosis. *FEMS Immunol Med Microbiol.* 2010;58(1):3–26.
- Ezekowitz RA, Stahl PD. The structure and function of vertebrate mannose lectin-like proteins. J Cell Sci Suppl. 1988;9:121–133.
- 9. Medzhitov R, Janeway C Jr. Innate immunity. N Engl J Med. 2000;343(5):338–344.
- Petersen SV, Thiel S, Jensenius JC. The mannan-binding lectin pathway of complement activation: biology and disease association. *Mol Immunol*. 2001;38(2–3):133–149.
- Jensenius JC. The mannan-binding lectin (MBL) pathway of complement activation: biochemistry, biology and clinical implications. *Adv Exp Med Biol.* 2005;564:21–22.
- Neth O, Jack DL, Dodds AW, Holzel H, Klein NJ, Turner MW. Mannosebinding lectin binds to a range of clinically relevant microorganisms and promotes complement deposition. *Infect Immun.* 2000;68(2):688–693.
- Garred P, Larsen F, Madsen HO, Koch C. Mannose-binding lectin deficiency--revisited. *Mol Immunol*. 2003;40(2–4):73–84.
- Rantala A, Lajunen T, Juvonen R, et al. Mannose-binding lectin concentrations, MBL2 polymorphisms, and susceptibility to respiratory tract infections in young men. *J Infect Dis.* 2008;198(8):1247–1253.
- Capparelli R, Iannaccone M, Palumbo D, et al. Role played by human mannose-binding lectin polymorphisms in pulmonary tuberculosis. *J Infect Dis.* 2009;199(5):666–672.
- Garred P, Larsen F, Seyfarth J, Fujita R, Madsen HO. Mannose-binding lectin and its genetic variants. *Genes Immun.* 2006;7(2):85–94.
- Jensenius H, Klein DC, van Hecke M, Oosterkamp TH, Schmidt T, Jensenius JC. Mannan-binding lectin: structure, oligomerization, and flexibility studied by atomic force microscopy. *J Mol Biol.* 2009;391(1):246–259.
- Eisen DP, Minchinton RM. Impact of mannose-binding lectin on susceptibility to infectious diseases. *Clin Infect Dis.* 2003;37(11):1496–1505.
- Wu L, Deng H, Zheng Y, et al. An association study of NRAMP1, VDR, MBL and their interaction with the susceptibility to tuberculosis in a Chinese population. *Int J Infect Dis.* 2015;38:129–135.
- da Cruz HL, da Silva RC, Segat L, et al. MBL2 gene polymorphisms and susceptibility to tuberculosis in a northeastern Brazilian population. *Infect Genet Evol.* 2013;19:323–329.
- Araújo MS, Graça ES, Azevedo VN, et al. No evidence of association between MBL2A/O polymorphisms and *Mycobacterium tuberculosis* infection in populations from the Brazilian Amazon region. *Hum Immunol.* 2013;74(1):82–84.
- Singla N, Gupta D, Joshi A, Batra N, Singh J, Birbian N. Association of mannose-binding lectin gene polymorphism with tuberculosis susceptibility and sputum conversion time. *Int J Immunogenet*. 2012;39(1):10–14.
- Cosar H, Ozkinay F, Onay H, et al. Low levels of mannose-binding lectin confers protection against tuberculosis in Turkish children. *Eur J Clin Microbiol Infect Dis.* 2008;27(12):1165–1169.

- Alagarasu K, Selvaraj P, Swaminathan S, Raghavan S, Narendran G, Narayanan PR. Mannose binding lectin gene variants and susceptibility to tuberculosis in HIV-1 infected patients of South India. *Tuberculosis*. 2007;87(6):535–543.
- Søborg C, Andersen AB, Range N, et al. Influence of candidate susceptibility genes on tuberculosis in a high endemic region. *Mol Immunol.* 2007;44(9):2213–2220.
- Liu W, Zhang F, Xin ZT, et al. Sequence variations in the MBL gene and their relationship to pulmonary tuberculosis in the Chinese Han population. *Int J Tuberc Lung Dis.* 2006;10(10):1098–1103.
- Selvaraj P, Jawahar MS, Rajeswari DN, Alagarasu K, Vidyarani M, Narayanan PR. Role of mannose binding lectin gene variants on its protein levels and macrophage phagocytosis with live *Mycobacterium tuberculosis* in pulmonary tuberculosis. *FEMS Immunol Med Microbiol*. 2006;46(3):433–437.
- Ozbaş-Gerçeker F, Tezcan I, Berkel AI, et al. The effect of mannosebinding protein gene polymorphisms in recurrent respiratory system infections in children and lung tuberculosis. *Turk J Pediatr.* 2003;45(2):95–98.
- Selvaraj P, Narayanan PR, Reetha AM. Association of functional mutant homozygotes of the mannose binding protein gene with susceptibility to pulmonary tuberculosis in India. *Tuber Lung Dis.* 1999;79(4):221–227.
- Bellamy R, Ruwende C, McAdam KP, et al. Mannose binding protein deficiency is not associated with malaria, hepatitis B carriage nor tuberculosis in Africans. *QJM*. 1998;91(1):13–18.
- Thye T, Niemann S, Walter K, et al. Variant G57E of mannose binding lectin associated with protection against tuberculosis caused by *Mycobacterium africanum* but not by *M. tuberculosis. PLoS One.* 2011;6(6):e20908.
- Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol*. 2010;25(9):603–605.
- 33. Hu P, Huang MY, Hu XY, et al. Meta-analysis of C242T polymorphism in CYBA genes: risk of acute coronary syndrome is lower in Asians but not in Caucasians. *J Zhejiang Univ Sci B*. 2015;16(5):370–379.
- Wu R, Li B. A multiplicative-epistatic model for analyzing interspecific differences in outcrossing species. *Biometrics*. 1999;55(2):355–365.
- Mantel N, Haenszel W. Statistical aspects of the analysis of data from retrospective studies of disease. J Natl Cancer Inst. 1959;22(4):719–748.
- Dersimonian R, Laird N. Meta-analysis in clinical trials. Control Clin Trials. 1986;7(3):177–188.

- Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003;327(7414):557–560.
- Wetterslev J, Thorlund K, Brok J, Gluud C. Trial sequential analysis may establish when firm evidence is reached in cumulative meta-analysis. J Clin Epidemiol. 2008;61(1):64–75.
- Turner RM, Bird SM, Higgins JP. The impact of study size on metaanalyses: examination of underpowered studies in Cochrane reviews. *PLoS One.* 2013;8(3):e59202.
- Brok J, Thorlund K, Wetterslev J, Gluud C. Apparently conclusive meta-analyses may be inconclusive--trial sequential analysis adjustment of random error risk due to repetitive testing of accumulating data in apparently conclusive neonatal meta-analyses. *Int J Epidemiol.* 2009;38(1):287–298.
- 41. Shastry BS. SNP alleles in human disease and evolution. *J Hum Genet*. 2002;47(11):561–566.
- Boldt AB, Luty A, Grobusch MP, et al. Association of a new mannosebinding lectin variant with severe malaria in Gabonese children. *Genes Immun.* 2006;7(5):393–400.
- Ogden CA, Decathelineau A, Hoffmann PR, et al. C1q and mannose binding lectin engagement of cell surface calreticulin and CD91 initiates macropinocytosis and uptake of apoptotic cells. *J Exp Med.* 2001;194(6):781–795.
- Polotsky VY, Belisle JT, Mikusova K, Ezekowitz RA, Joiner KA. Interaction of human mannose-binding protein with *Mycobacterium avium. J Infect Dis.* 1997;175(5):1159–1168.
- 45. Hoppe HC, de Wet BJ, Cywes C, Daffé M, Ehlers MR. Identification of phosphatidylinositol mannoside as a mycobacterial adhesin mediating both direct and opsonic binding to nonphagocytic mammalian cells. *Infect Immun.* 1997;65(9):3896–3905.
- Ioannidis JP, Boffetta P, Little J, et al. Assessment of cumulative evidence on genetic associations: interim guidelines. *Int J Epidemiol.* 2008;37(1):120–132.
- Garred P, Harboe M, Oettinger T, Koch C, Svejgaard A. Dual role of mannan-binding protein in infections: another case of heterosis? *Eur J Immunogenet*. 1994;21(2):125–131.
- Kramnik I, Dietrich WF, Demant P, Bloom BR. Genetic control of resistance to experimental infection with virulent *Mycobacterium tuberculosis. Proc Natl Acad Sci U S A.* 2000;97(15):8560–8565.
- 49. Caws M, Thwaites G, Dunstan S, et al. The influence of host and bacterial genotype on the development of disseminated disease with *Mycobacterium tuberculosis*. *PLoS Pathog*. 2008;4(3):e1000034.

Infection and Drug Resistance

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed openaccess journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic

Submit your manuscript here: https://www.dovepress.com/infection-and-drug-resistance-journal

Dovepress

resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peerreview system, which is all easy to use. Visit http://www.dovepress.com/ testimonials.php to read real quotes from published authors.