

Apelin and chemerin as promising adipokines in children with type I diabetes mellitus

Asmaa Abd El Wakeel
Elsehmawy¹
Safinaz Ebrahim El-Toukhy²
Nora Mohamed Ahmed
Seliem³
Rehab Selim Moustafa⁴
Doaa Sayed Mohammed⁵

¹Department of Pediatrics, Faculty of Medicine, Al-Azhar University, Cairo, Egypt; ²Department of Medical Biochemistry, National Research Center, Cairo, Egypt; ³Biochemistry Department, Faculty of Medicine, Al-Azhar University, Cairo, Egypt; ⁴Child Health Department, National Research Center, Cairo, Egypt; ⁵Endocrine Department, Faculty of Medicine, Al-Azhar University, Cairo, Egypt

Background: Type 1 diabetes mellitus (T1DM) is one of the most common chronic diseases in children that may be due to micro or macrovascular complications. Diabetic renal disease or nephropathy is a common complication of DM, clinically silent and the only detectable abnormality due to the presence of microalbuminuria.

Subjects and methods: This study was a case–control study. Participants were classified into two groups. The first group included 40 children with T1DM and the second group included 30 matched healthy controls. Serum apelin (APLN), chemerin, cholesterol, and triglycerides (TG) levels were measured for each case. Also, albumin/creatinine ratio was analyzed in random urine sample.

Results: Comparison between T1DM patients and controls revealed that serum apelin, chemerin, cholesterol, TG levels, and albuminuria were significantly increased in cases compared to their controls. Significant positive correlations were found between HbA1c% and albuminuria for APLN and chemerin in the diabetic group. Whereas significant negative correlations were found between apelin and glomerular filtration rate (GFR).

Conclusion: Increased levels of serum apelin and chemerin in T1DM patients may be considered as promising adipokines for the development of diabetic complication.

Keywords: apelin, chemerin, children, diabetes mellitus

Introduction

Type 1 diabetes mellitus (T1DM) is a common, chronic and metabolic disease characterized by hyperglycemia as a cardinal metabolic feature.¹ Long-term damage, dysfunction and failure of various organs especially eyes, kidneys, nerves, heart and blood vessels are caused by chronic hyperglycemia of diabetes.² Diabetic nephropathy (DN) is an insidious and a common complication of diabetes mellitus (DM). At its onset, DN is clinically silent, and the only detectable abnormality due to the presence of microalbuminuria, which is defined as urine albumin of 30–300 g/day (or also known as albuminuria).³

DN is believed to be the consequence of poorly controlled hyperglycemia. The mechanisms leading to renal injury in diabetes mellitus is evolving, may be caused by accumulation of injurious metabolic products, such as glycosylated compounds, involvement of the renin–angiotensin system, and endothelial and podocyte injury being involved in the pathogenesis of DN.³

Glomerular hypertrophy, thickening of glomerular basement membrane and mesangial expansion with the formation of Kimmelstiel–Wilson nodules are the classical changes that are seen in the glomerular compartment that are usually accompanied with glomerular lesions including tubulo-interstitium undergoing fibrosis, as well

Correspondence: Asmaa Abd El Wakeel Elsehmawy
Department of Pediatrics, Faculty of Medicine, Al-Azhar University, Zhraa Elmadi, Nerco tower No.5, Cairo, Egypt
Email Asmaawakeel@yahoo.com

as arteriolar thickening and hyalinization in the vascular compartment of the kidney. There is an imbalance of extracellular matrix (ECM) synthesis and degradation, which is most likely responsible for the accumulation of excessive matrices in various compartments of the kidney. Thus, the genesis of Kimmelstiel–Wilson lesion, a diagnostic feature of DN, is not only due to the excessive synthesis of ECM glycoproteins but also due to a decreased degradation by matrix metalloproteinases.⁴

Hyperglycemia-induced hyperfiltration is an early observable abnormality in DM. At this stage, patients are asymptomatic, blood pressure is normal, there is no detectable microalbuminuria, and therefore it is considered as a latent stage of diabetic nephropathy. Presence of microalbuminuria (also known as albuminuria) in patients with DM1 or DM2 is recognized as an early marker of DN.⁵

Chemerin, a discovered adipocytokine, is a chemoattractant protein that acts as a ligand for the G-protein-coupled receptor CMKLR1 (also known as ChemR23). Chemerin is a 14 kDa protein secreted in an inactive form as prochemerin and is activated through cleavage of the C-terminus by inflammatory and coagulation serine proteases. It is involved in glucose and lipid metabolism. Elevated levels of this peptide have been associated with insulin resistance and systemic inflammation.⁶

Apelin (APLN), adipocytokine, is a peptide known as the endogenous ligand of the G-protein-coupled receptor Apelin. Apelin-13, the most active member of the APLN group, is expressed in a variety of tissues, including kidney and endothelial cells. In addition, increased body mass in type 2 diabetic patients might contribute to increase APLN levels in the blood.⁷

Aim of the study

The aim of the current study was to evaluate the level of serum chemerin and apelin in type 1 diabetic children and their correlation with glycated hemoglobin and proteinuria.

Materials and methods

Participants

Seventy children of both genders were enrolled for this study and divided into two groups.

Study group: included 40 patients with T1DM were randomly selected from the outpatient pediatric clinic.

Control group: included 30 apparently healthy children.

The study was carried out in Alzhraa University Hospital, Faculty of Medicine for Girls, Al-Azhar University, Cairo, Egypt: Pediatric and Biochemistry Departments.

Design

Participants were evaluated during a hospital visit to determine family medical history, current medical conditions, medication use, self-report of the presence of any smokers in the household and demographic information.

In addition, physical examinations, including height and weight measurements and laboratory testing using blood samples for lipid profile, were performed. Children with renal diseases or urological problems, presence of urinary tract infection or taking any medication rather than insulin or those with chronic devastating diseases were excluded from this study.

The body weight was measured using Seca scale to the nearest 0.1 kg barefoot and in light clothes, after emptying the urinary and gastrointestinal apparatuses (Seca Model 770, Hamburg, Germany).

Investigations

In the morning, after 12–14 hours fasting, 7 mL of blood was collected from all participants and divided into two: one half was taken in a plain tube to separate serum and another half was taken in a tube containing EDTA. Serum was separated to determine fasting blood glucose, urea, and creatinine levels and with the another half of blood sample following analysis were performed.

1. Glycosylated hemoglobin (HbA1c%)
HbA1c% was measured using a spectrophotometer (Clinical Chemistry Analyzer 7; Germany).
2. Lipid profile and serum creatinine
Lipid profile was measured using an autoanalyzer (Beckman Coulter Synchron CX 9; Beckman Coulter Inc., Brea, CA, USA). For cholesterol measurements, serum total cholesterol was determined by a commercial kit (Boehringer-Mannheim, Germany).⁸ The concentration of triglycerides (TG) was measured using a TechnoCon AutoAnalyzer II (TechnoCon Instruments, Tarrytown, NY, USA).
3. Albumin/creatinine ratio (ACR) screening for microalbuminuria was assessed for fresh morning urine samples by measuring ACR using ELISA kit provided by Orgentec Diagnostika, GmbH (Mainz, Germany).⁹
4. Serum APLN and chemerin

Serum apelin and chemerin were assayed using human ELISA kit (Cusabio, Wuhan, Hubei Province, China).

Serum chemerin

The human chemerin ELISA, standards and samples were incubated in microtitration wells pre-coated with polyclonal antihuman chemerin antibody. Incubation was followed

by washing, and then biotin-labeled polyclonal antihuman chemerin antibody was added and incubated with the captured chemerin. After another washing, streptavidin-HRP conjugate was added. After the last washing step, the remaining conjugate was allowed to react with the substrate solution. The reaction was stopped by addition of acidic solution, and absorbance of the resulting yellow product was measured. The absorbance was proportional to the concentration of chemerin. A standard curve was constructed by plotting absorbance values against chemerin.^{10,11}

Serum APLN

This assay employs the competitive inhibition enzyme immunoassay technique. A monoclonal antibody specific to apelin has been pre-coated into a microplate. A competitive inhibition reaction was launched between biotin-labeled APLN and biotin-unlabeled APLN (standards or samples) with the pre-coated antibody specific to APLN. After incubation the unbound conjugate was washed off, avidin conjugated to horseradish peroxidase (HRP) was added to each microplate well and

incubated. The amount of bound HRP conjugate was inversely proportional to the concentration of APLN in the sample. After addition of the substrate solution, the intensity of color developed was inversely proportional to the concentration of APLN in the sample. Detection range was 0.1–1000 ng/mL.^{12,13}

Statistics

Data were collected, revised, coded, and entered on the Statistical Package for Social Science (IBM SPSS) version 20 statistical analysis software (IBM Corporation, Armonk, NY, USA). The statistical tools used include mean, SD, and *t*-test.

Results

The descriptive statistics of the study groups are shown in Table 1. Patients and healthy volunteers were matched for the mean of both age and gender. Significant increase in blood level of urea, creatinine, fasting blood sugar, cholesterol, TG and HbA1c% was shown in diabetic children in comparison to the control. Increased ACR and GFR in diabetic children was shown in comparison to healthy controls.

Table 1 Demographic, anthropometric, and chemical data of the studied groups

		Control group	Diabetic group	Test value	P-value
Age/year	Mean ± SD	10.90±1.51	10.73±1.90	0.416	0.679
	Range	8.5–14	8–13.5		
BMI	Mean ± SD	20.24±0.98	26.63±4.02	–8.519	0.000
	Range	18.43–21.74	21.65–39.06		
Z-score (BMI)	Median (IQR)	–1.01 (–1.19–0.88)	0.36 (–0.22–0.76)	–7.026	0.000
	Range	–1.43 to –0.70	–0.72–3.12		
Duration of disease/year	Mean ± SD	–	4.18±0.74	NA	NA
	Range	–	3–5.5		
Urea (mg/dL)	Mean ± SD	19.45±3.69	28.28±7.86	–5.689	0.000
	Range	13.5–25	14.5–42		
Creatinine (mg/dL)	Mean ± SD	0.61±0.10	1.05±0.11	–17.585	0.000
	Range	0.44–0.83	0.82–1.25		
FBS (mg/dL)	Mean ± SD	77.87±5.99	147.73±40.82	–9.283	0.000
	Range	69–89	89–243		
Cholesterol (mg/dL)	Mean ± SD	112.00±10.47	174.20±32.22	–10.163	0.000
	Range	94–128	125–244		
Triglycerides (mg/dL)	Mean ± SD	72.80±6.05	94.05±21.40	–5.274	0.000
	Range	62.9–82.1	71.3–156		
HbA1c%	Mean ± SD	4.72±0.41	7.35±0.90	–14.826	0.000
	Range	3.79–5.43	5.69–8.87		
APLN (pg/mL)	Mean ± SD	114.18±7.05	767.31±95.42	–37.345	0.000
	Range	100.57–126.45	604.94–869.34		
Chemerin (pg/mL)	Mean ± SD	83.33±5.51	117.12±25.79	–7.044	0.000
	Range	72.04–92.18	83.16–180		
Albumin/creatinine	Mean ± SD	2.35±0.99	20.60±18.21	–5.475	0.000
	Range	1–4.5	0.7–114.2		
GFR, mL/min/1.73 m ²	Mean ± SD	184.20±51.89	95.50±21.47	9.770	0.000
	Range	102–295	63–148		

Abbreviations: BMI, body mass index; HbA1c%, glycosylated hemoglobin; APLN, apelin; GFR, glomerular filtration rate.

Both APLN and chemerin showed a statistically significant increase in diabetic children when compared to their healthy counterparts (Table 1). Both APLN and chemerin showed a statistically significant increase in non-controlled diabetic children (Table 2).

The cutoff point for APLN and chemerin was >126.45 and 92.18, respectively, with 100% sensitivity and specificity for APLN, and 74% and 100% for chemerin in the prediction of nephropathy in diabetic children (Table 3).

We found a positive correlation of APLN and chemerin with BMI, urea, fasting blood glucose, glycosylated hemoglobin and ACR. In addition, we also found a positive correlation of APLN with cholesterol and triglyceride as shown in Table 4.

Correlation analysis between GFR in diabetic children and the laboratory data revealed that there were negative correlations between GFR and fasting blood sugar, urea, creatinine, cholesterol, triglycerides, HbA1c%, and APLN (Table 5).

Discussion

The present study was a controlled study that was carried out on 70 children of both genders and divided into two groups. The diabetic group included 40 children with T1DM who were randomly selected from the outpatient pediatric clinic; there were 21 (52.5%) females and 19 (47.5%) males, with age ranging from 8 to 13.5 years. The control group included 30 apparently healthy children including 12 (40%) females and 18 (60%) males with age ranging from 8.5 to 14 years.

The aim of this study was to evaluate the level of serum chemerin and APLN in patients with type 1 diabetes and their correlation with proteinuria and HbA1c%.

In the present study there were significant increases in BMI, blood urea, creatinine, fasting blood sugar, cholesterol,

TG, and HbA1c% in diabetic children in comparison to the control. Increased ACR was shown in diabetic children in comparison to healthy controls.

Table 4 Correlation between demographic, anthropometric, and laboratory data with APLN and chemerin

	APLN		Chemerin	
	r	P-value		
Age	0.297	0.062	-0.023	0.890
BMI	0.563	0.000	0.507	0.001
Duration	-0.005	0.977	-0.052	0.752
Urea	0.600	0.000	0.409	0.009
Creatinine	0.003	0.987	-0.044	0.787
Fasting blood glucose	0.644	0.000	0.765	0.000
Cholesterol	0.406	0.009	0.205	0.205
Triglycerides	0.364	0.021	-0.138	0.397
HbA1c%	0.710	0.000	0.721	0.000
Albumin/creatinine	0.338	0.033	0.577	0.000
GFR (mL/min/1.73m ²)	-0.315	0.048	-0.123	0.449

Abbreviations: APLN, apelin; BMI, body mass index; HbA1c%, glycosylated hemoglobin; GFR, glomerular filtration rate.

Table 5 Correlation between GFR and the laboratory data

	GFR (mL/min/1.73 m ²)	
	r	P-value
FBS (mg/dL)	-0.314	0.048
Urea (mg/dL)	-0.633	0.000
Creatinine (mg/dL)	-0.630	0.000
Cholesterol (mg/dL)	-0.400	0.011
Triglycerides (mg/dL)	-0.319	0.045
HbA1c%	-0.392	0.012
APLN (pg/mL)	-0.315	0.048
Chemerin (pg/mL)	-0.123	0.449

Abbreviations: GFR, glomerular filtration rate; HbA1c%, glycosylated hemoglobin; APLN, apelin.

Table 2 Chemerin and APLN levels in both controlled and non-controlled diabetic children

		Controlled group	Uncontrolled group	Test value	P-value	Significance
HbA1c%	Mean ± SD	6.46±0.27	7.96±0.61	-9.248	0.000	HS
	Range	5.79-6.83	7.19-8.92			
APLN (pg/mL)	Mean ± SD	650.33±38.93	836.60±25.49	-18.328	0.000	HS
	Range	604-723.56	792-869.57			
Chemerin (pg/mL)	Mean ± SD	89.67±4.85	135.36±15.18	-11.606	0.000	HS
	Range	83.19-96	123.66-180			

Abbreviations: HbA1c%, glycosylated hemoglobin; APLN, apelin.

Table 3 The cutoff point, sensitivity, specificity of APLN and chemerin as an early marker in detection of diabetic nephropathy

	Cutoff point	AUC	Sensitivity	Specificity	Positive predictive value	Negative predictive value
APLN (pg/mL)	>126.45	1.000	100.00	100.00	100.0	100.0
Chemerin (pg/mL)	92.18	0.922	75.00	100.00	100.0	75.0

Abbreviations: APLN, apelin; AUC, area under the curve.

In our study there was significant increase in serum chemerin in diabetic children in comparison to the controls, and this was in agreement with previous studies on T2DM.^{14,15} Moreover, in adolescent type 1 diabetic patients, Abd El Dayema et al¹⁶ showed that chemerin was significantly higher in diabetic group than the controls.

In our study there was a significant increase in serum APLN in diabetic children in comparison to the controls and this is in agreement with previous studies of APLN in type 1 diabetic patients.¹⁷⁻²⁰

In the present study, we reported that serum cholesterol and TG were significantly increased in T1DM patients than in healthy controls. From the previous results, we considered patients with T1DM are at high risk of developing premature vascular affection because of hyperlipidemia and thus should be screened well for this serious complication; we found a positive correlation of APLN with cholesterol and TG. In support of these results, another study found statistically significant differences between diabetic and non-diabetic groups related to serum levels of TG and cholesterol.²¹ Few studies have described the effects of APLN on lipid metabolism; one stated that APLN was shown to inhibit lipolysis.²² This was ensured by Than et al²³ who found that APLN increases the stability of lipid vacuoles making them more resistant to lipases.

APLN has a role in energy metabolism: it improves sensitivity of insulin in insulin-resistant obese mice, and it is related to an increase in glucose uptake in skeletal muscle.²⁴ Synthesis of APLN is affected by insulin and plasma APLN levels is increased in obesity in association with hyperinsulinemia.²⁵ Furthermore, clinical studies demonstrated that chemerin modulates inflammation. Mechanisms linking chronic inflammation to the development of diabetes remain elusive, but it can be assumed that any effects may be exerted via insulin secretion, insulin resistance, or both.²⁶

In our study both apelin and chemerin levels are more increased in non controlled diabetic children compared the controlled one (table 2) and both chemerin and APLN were positively correlated with fasting blood glucose and glycosylated hemoglobin and this is in agreement with previous studies that showed a correlation between serum APLN and HbA1c% in children with T1DM.^{17,18,20}

ACR is used to screen people with chronic conditions, such as diabetes that put them at an increased risk of developing kidney disease. Studies have shown that identifying individuals in the very early stages of kidney disease helps people and healthcare providers adjust treatment. Controlling diabetes by maintaining tight glycemic control delays or prevents the progression of kidney disease.²⁷

Both APLN and chemerin, novel adipocytokines, were increased in response to inflammation. In vitro studies have revealed that APLN and the APLN receptor can induce the sprouting of endothelial cells in an autocrine or paracrine manner, thus suggesting a role for APLN in angiogenesis.²⁸ Moreover, it was found that chemerin induced functional angiogenesis in human endothelial cells, by promoting migration, capillary tube formation, activation of endothelial gelatinase (matrix metalloproteinases-2/matrix metalloproteinases-9) and activation of phosphatidylinositol 3-kinase/Akt and MAPKs pathways, which is a key mechanism for angiogenesis and cell survival. Therefore, the high level of serum chemerin possibly promotes the development of diabetic complication and diabetic nephropathy.²⁹

The activation of the immune system and chronic inflammation are both involved in pathogenesis of DM and its complications as a result of DN. Some studies have demonstrated that cytokines, chemokines, growth factors, adhesion molecules, nuclear factors as well as immune cells such as monocytes, lymphocytes and macrophages are all involved in DM pathogenesis and of course play an important role in DM complications.³⁰

In the present study there were increases in ACRs in diabetic children in comparison to the healthy controls and there were positive correlations between APLN and chemerin blood levels with alp/creatinine ratio. Correlation of serum APLN with microalbuminuria was confirmed by another study by Guo et al,³¹ which reported that APLN and chemerin may induce glomerular endothelial cell proliferation and then nephropathy, also there was a negative correlation between serum APLN and GFR, which was stated in the study on type 2 diabetes.³² The cutoff point of APLN and chemerin was >126.45 and 92.18, respectively, with 100% sensitivity and specificity for APLN and 74%, 100% for chemerin in the prediction of nephropathy in diabetic children (Table 3, Figure 1). To our knowledge this is the first study to mention the cutoff for both markers in diabetic children. We can state that increased serum APLN and chemerin in diabetic children may contribute to development of diabetic complication.

Conclusion

In conclusion, increased levels of serum APLN and chemerin were increased in T1DM patients, which may be considered as promising adipokines in diabetic complication development, and hence measuring serum APLN and chemerin in diabetic children is of benefit for detection of diabetic complications.




Figure 1 ROC curve demonstrates apelin and chemerin sensitivity and specificity in predicting nephropathy risk in T1DM children.

Limitations

One of the limitations of the study was small sample size as it was a single-center study and we included all children only referred to our hospital during the period from January to June 2018.

Acknowledgments

This work was supported by the Department of Pediatrics, Faculty of Medicine for Girls – Al-Aazhar University and Biochemistry Department of the national Research center. The authors thank the families of the children who agreed to participate in this work.

Author contributions

All authors contributed to data analysis, drafting and revising the article, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Svoren BM, Jospe N. Diabetes mellitus in children. In: Kliegman RM, Stanton BF, St Geme JW, Score NF, Behrman RE editors. *Nelson's Textbook of Pediatrics*. 20th ed. Elsevier, Inc. Chapter 589 2016:2760–2782.
2. American Diabetes Association (ADA). Diagnosis and classification of diabetes mellitus. *Diabetes Care*. 2013;35(1):64–71.
3. Kher K, Mietus-Snyder M. Kidney disease associated with diabetes mellitus and metabolic syndrome. In: Kher KK, Schnaper HW, Greenbaum LA, editors. *Clinical Pediatric and Nephrology*. 3rd ed. Boca Raton, FL: CRC Press; 2017, Part E:533–540.
4. Kolset SO, Reinholt FP, Jenssen T. Diabetic nephropathy and extracellular matrix. *J Histochem Cytochem*. 2012;60(12):976–986.
5. Demirel F, Tepe D, Kara O, Esen I. Microvascular complications in adolescents with type 1 diabetes mellitus. *J Clin Res Pediatr Endocrinol*. 2013;5(3):145–149.
6. Li Y, Shi B, Li S. Association between serum chemerin concentrations and clinical indices in obesity or metabolic syndrome: a meta-analysis. *PLoS One*. 2014;9(12):e113915.
7. Huang Z, Wu L, Chen L. Apelin/APJ system: a novel potential therapy target for kidney disease. *J Cell Physiol*. 2018;233(5):3892–3900.
8. Flegg HM. Ames Award Lecture 1972. An investigation of the determination of serum cholesterol by an enzymatic method. *Ann Clin Biochem*. 1973;10(1–6):79–84.
9. Mogensen CE. Microalbuminuria predicts clinical proteinuria and early mortality in maturity-onset diabetes. *N Engl J Med*. 1984;310(6):356–360.
10. Takahashi M, Takahashi Y, Takahashi K, et al. Chemerin enhances insulin signaling and potentiates insulin-stimulated glucose uptake in 3T3-L1 adipocytes. *FEBS Lett*. 2008;582(5):573–578.
11. Burtis CA, Ashwood ER. *Tietz Textbook of Clinical Chemistry*. 2nd ed. Philadelphia: WB Saunders; 1994:1017–1089.
12. Kaliora AC, Kanellos PT, Gioxari A, Karathanos VT. Regulation of GIP and ghrelin in healthy subjects fed on sun-dried raisins: a pilot study with a crossover trial design. *J Med Food*. 2017;20(3):301–308.
13. Vehapoglu A, Ustabas F, Ozgen TI, Terzioğlu S, Cermik BB, Ozen OF. Role of circulating adipocytokines vaspin, apelin, and visfatin in the loss of appetite in underweight children: a pilot trial. *J Pediatr Endocrinol Metab*. 2015;28(9–10):1065–1071.
14. Khaled Y, Rashed L. Serum chemerin levels and chemerin rs17173608 genotypes in the susceptibility of diabetic nephropathy in Egyptian diabetic patients. *Egypt J Obes Diabetes Endocrinol*. 2016;2(1): 18–22.
15. El-Gohary IE, Abedi-Karima, Hashad DI. Serum chemerin level: does it have a role in progression of diabetic nephropathy. *Am J Intern Med*. 2016;4(2–1):13–17.
16. El Dayem SMA, Battah AA, El Bohy AEM, El Shehaby A, El Ghaffar EA. Relationship of plasma level of chemerin and vaspin to early atherosclerotic changes and cardiac autonomic neuropathy in adolescent type 1 diabetic patients. *J Pediatr Endocrinol Metab*. 2015;28(3–4):265–273.
17. Dayem SMAE, Battah AA, Bohy AEME, Yousef RN, Ahmed AM, Talaat AA. Apelin, nitric oxide and vascular affection in adolescent type 1 diabetic patients. *Open Access Maced J Med Sci*. 2017;5(7): 934–939.
18. Meral C, Tascilar E, Karademir F, et al. Elevated plasma levels of apelin in children with type 1 diabetes mellitus. *J Pediatr Endocrinol Metab*. 2010;23(5):497–502.
19. Habchi M, Duvillard L, Cottet V, et al. Circulating apelin is increased in patients with type 1 or type 2 diabetes and is associated with better glycaemic control. *Clin Endocrinol*. 2014;81(5):696–701.

20. Al-Suhaimi EA, Al-Kulaifi FM, Ravinayagam V, Al-Qahtani MH. Serum adipocytokines, metabolic and immunological correlations in type 1 diabetes mellitus (T1DM) children. *Open Endocrinol J.* 2012;6(1):110–116.
21. Nascimento AMMAD, Sequeira IJ, Vasconcelos DF, Gandolfi L, Pratesi R, Nóbrega YKM. Endothelial dysfunction in children with type 1 diabetes mellitus. *Arch Endocrinol Metab.* 2017;61(5):476–483.
22. Yue P, Jin H, Xu S, et al. Apelin decreases lipolysis via G(q), G(i), and AMPK-dependent mechanisms. *Endocrinology.* 2011;152(1):59–68.
23. Than A, Cheng Y, Foh L-C, et al. Apelin inhibits adipogenesis and lipolysis through distinct molecular pathways. *Mol Cell Endocrinol.* 2012;362(1-2):227–241.
24. Yue P, Jin H, Aillaud M, et al. Apelin is necessary for the maintenance of insulin sensitivity. *Am J Physiol Endocrinol Metab.* 2010;298(1):E59–E67.
25. Soriguer F, Garrido-Sanchez L, Garcia-Serrano S, et al. Apelin levels are increased in morbidly obese subjects with type 2 diabetes mellitus. *Obes Surg.* 2009;19(11):1574–1580.
26. Weigert J, Neumeier M, Wanninger J, et al. Systemic chemerin is related to inflammation rather than obesity in type 2 diabetes. *Clin Endocrinol.* 2010;72(3):342–348.
27. American Diabetes Association Position Statement: Standards of Medical Care in Diabetes --2015. *Diabetes Care.* 2015;38(Suppl. 1):S1–S94.
28. Kidoya H, Ueno M, Yamada Y, et al. Spatial and temporal role of the apelin/APJ system in the caliber size regulation of blood vessels during angiogenesis. *Embo J.* 2008;27(3):522–534.
29. Bozaoglu K, Curran JE, Stocker CJ, et al. Chemerin, a novel adipokine in the regulation of angiogenesis. *J Clin Endocrinol Metab.* 2010;95(5):2476–2485.
30. Navarro-González JF, Mora-Fernández C. The role of inflammatory cytokines in diabetic nephropathy. *J Am Soc Nephrol.* 2008;19(3):433–442.
31. Guo C, Liu Y, Zhao W, et al. Apelin promotes diabetic nephropathy by inducing podocyte dysfunction via inhibiting proteasome activities. *J Cell Mol Med.* 2015;19(9):2273–2285.
32. Dawood A, Abdelraof M, El Ghobashy Y. The relationship between serum apelin level and different grades of diabetic nephropathy in type 2 diabetic patients. *Egypt J Obes Diabetes Endocrinol.* 2017;3(1):32–37.

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy

Dovepress

Publish your work in this journal

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert

opinion and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-targets-and-therapy-journal>