

LKB1 suppresses glioma cell invasion via NF- κ B/Snail signaling repression

This article was published in the following Dove Medical Press journal:
OncoTargets and Therapy

Ye Yuan¹
Shi-Lin Li²
Yu-Lin Cao³
Jun-Jun Li¹
Qiang-Ping Wang¹

¹Department of Neurosurgery, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430022, People's Republic of China; ²Department of Neurology, Qitaihe Qimei Hospital, Qitaihe 154600, People's Republic of China; ³Department of Hematology, Wuhan General Hospital of Guangzhou Military Area Command, Wuhan 430070, People's Republic of China

Background: Liver kinase B1 (LKB1) is involved in various human diseases. Aberrant expression of LKB1 expression is involved in glioma progression and associated with prognosis, however, the specific mechanism involving NF- κ B/Snail signaling pathways remain unknown.

Materials and methods: In the present study, quantitative real-time PCR analysis was used to investigate the expression of LKB1 tumor tissue samples and cell lines. In glioma cell lines, CCK-8 assay, transwell invasion and migration assays were used to investigate the effects of LKB1 on proliferation and invasion.

Results: We observed that LKB1 knockdown promoted glioma cell proliferation, migration and invasion. This effect was induced through NF- κ B/Snail signaling activation. Also, LKB1 overexpression suppressed proliferation, migration, and invasion, which could be rescued by Snail overexpression.

Conclusion: Taken together, our results show that LKB1 knockdown promotes remarkably glioma cell proliferation, migration and invasion by regulating Snail protein expression through activating the NF- κ B signaling. This may serve as a potential prognostic marker and therapeutic target for glioma.

Keywords: glioma, proliferation, migration, invasion, LKB1, NF- κ B, snail

Introduction

Gliomas are the most common primary malignant intracranial brain neoplasms in adults, accounting for >50% of the central nervous system tumor; it has the feature of rapid progression and poor prognosis.^{1,2} Glioma is characterized by rapid progression and poor prognosis due to its highly invasive characteristics. Therefore, elucidation of specific mechanisms underlying this invasive nature of glioma is imperative to explore novel diagnostic methods and treatments.

Liver kinase B1 (LKB1), also known as serine/threonine protein kinase 11, is located on chromosome 19p13.3 and encodes a 48-kDa serine/threonine protein kinase.³ Functionally, LKB1 plays pivotal roles in cellular processes such as apoptosis, metabolism, and cell cycle regulation. As a multifunctional molecular protein, LKB1 acts as a vital metabolic enzyme in the 5'-adenosine monophosphate-activated protein kinase (AMPK) signaling pathway,⁴ and LKB1 deactivation often mediates the mammalian target of rapamycin signaling pathway,⁵ which is critical for modulating cell proliferation, survival, and energy metabolism. Growing evidence has indicated that *LKB1* functions as a tumor-suppressor gene by modulating AMPK or AMPK-related kinases, which further modulate cell cycle, apoptosis, polarity, and metabolism.⁶ Although previous studies have reported the clinical importance of LKB1 expression in gliomas and role of LKB1-AMPK pathway in suppressing glioma cell proliferation, the specific mechanisms underlying NF- κ B/Snail signaling remain unknown.

Correspondence: Qiang-Ping Wang
Department of Neurosurgery, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430022, People's Republic of China
Email kouxiantang7595@163.com

NF- κ B signaling pathway is activated by several signaling molecules and it is a very crucial regulatory transcription factor for cell survival, growth, and differentiation.^{7,8} Previous studies have demonstrated that NF- κ B activation blocked the Snail degradation, subsequently promoting cancer cell migration and invasion.^{9,10}

In this study, we elucidated LKB1 expression and function in glioma. Our results indicated that LKB1 is down-regulated in glioma tissues. Moreover, cellular experiments revealed the pivotal role of LKB1 in inhibiting glioma cell growth, migration, and invasion, which is likely mediated through NF- κ B signaling.

Materials and methods

Cell lines and reagents

HA, HM, U-87MG, H4, U-251MG, and LN-18 cells (Cellcook Biotech Co., Ltd., Guangzhou, China), which were routinely checked for contamination and authenticated by the company, were purchased. Cells were grown in DMEM (Hyclone Laboratories, San Angelo, TX, USA) supplemented with 10% FBS (Gibco, Waltham, MA, USA). 4-Methyl-N1-(3-phenylpropyl)-1, 2-benzenediamine (JSH-23), and caffeic acid phenethyl ester (CAPE) were purchased from Sigma-Aldrich Co. (St Louis, MO, USA). Anti-LKB1 (3050), anti-Snail (3879), anti-Slug (9585), anti-P-IKK α/β (2697), anti-total IKK α (2682), anti-total IKK β (8943), anti-NF- κ B p65 (8242), and anti- β -actin (4970) antibodies were purchased from Cell Signaling Technology (Danvers, MA, USA).

Patients and sample preparation

A total of 49 samples of surgically resected gliomas were collected from patients who visited Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, from September 2017 to March 2018. All patients provided written informed consent, and the study was approved by the hospital authorities. Before brain surgery, no patient received radiotherapy or chemotherapy. After surgical resection, all samples were immediately snap-frozen in liquid nitrogen and fixed in formaldehyde solution. This research was approved by the Human Research Committee of Huazhong University of Science and Technology and China Anti-Cancer Association and was performed in accordance with the Helsinki Declaration.

Establishment of knockdown and overexpression cells

To establish stable knockdown and overexpression cell lines, full-length shRNA sequences that specifically targeted LKB1, Snail, or I κ B α were cloned into pLKO.1-MSCV-Puro

or pcDNA3.0 vectors and were sequenced bidirectionally. Plasmid construction and lentivirus packaging were completed by Shanghai GeneChem Co., Ltd. (Shanghai, China). Cells were infected by the lentivirus according to the manufacturer's protocol and selected by puromycin (Sigma-Aldrich Co.) after 2 weeks to obtain cell lines with stable expression. Empty vector pcDNA3.0 or pLKO.1 and scrambled shRNA were used as a negative control. The shLKB1 target sequence was 5'-CATCTACTCAGGACTTCAC-3', and the Snail target sequence was 5'-GCTGAGCTGTTACTAGGACAA-3'. The short hairpin RNA was designed based on the LKB1 sequence: 5'-CATTGTGCACAAGGACATCAA-3' and I κ B α sequence: 5'-GACGAGAAAGATCATTGAAAT-3'. Cells infected with empty vector or scrambled shRNA were used as controls.

Transwell invasion and migration assays

Cell migration and invasion experiments were performed using the Transwell system (Corning Inc., Corning, NY, USA) based on the manufacturer's instructions.¹¹ To assess invasion, filters were precoated with Matrigel (BD Biosciences, San Jose, CA, USA). Approximately 6×10^3 cells were added to the top chamber containing serum-free DMEM. The bottom chamber contained 600 μ L of DMEM supplemented with 20% FBS. Following 24 hours of incubation, cells on the upper surface were gently removed using a cotton swab, and then the membrane was fixed with 4% formaldehyde for 20 minutes and stained with 0.1% crystal violet solution (Servicebio, Wuhan, China) for 30 minutes. Cells that migrated to the lower surface of the membrane were photographed and counted under a microscope. The same experiment was performed for the migration assay except the filters were not precoated with Matrigel.

Cell counting Kit-8 (CCK-8) assay

For CCK-8 assay, cells were counted and seeded at a density of 5,000 cells/well in 96-well plates containing complete culture medium. After treatment with 10 μ L CCK-8 (Dojindo Laboratories, Kumamoto, Japan), absorbance was measured at 450 nm using a microplate reader (Spectra Max M2 reader; Molecular Devices LLC, Sunnyvale, CA, USA). Each experiment was performed in triplicate.

Western blotting

Total protein was lysed from glioma tissues or cells using RIPA lysis buffer (Boster Biological Technology, Pleasanton, CA, USA) with 1 mM phenylmethane sulfonyl fluoride (Boster Biological Technology) according to the manufacturers' instruction.¹² Samples were supplemented with phosphatase inhibitors (Boster Biological Technology) for phosphorylated

protein extraction. After clarification by centrifugation at 12,000×g for 20 minutes at 4°C, soluble protein concentration was measured using the bicinchoninic acid (Boster Biological Technology) method. For immunoblotting, total proteins/phosphorylated protein samples (60 µg) were incubated at 100°C for 10 minutes, separated by 12% SDS-PAGE (Boster Biological Technology) at 120 V for 2 hours, and transferred on a polyvinylidene difluoride membrane (EMD Millipore, Billerica, MA, USA), which was incubated with 5% fat-free milk in PBS for 2 hours and incubated with primary antibodies at 4°C overnight and with secondary antibodies at room temperature for 1 hour. Finally, enhanced chemiluminescence (Boster Biological Technology) was used to visualize samples exposed to X-ray film, and β-actin (1:1,000; Cell Signaling Technology) was used as an internal control. To analyze relative protein expression, ImageJ2X analysis software package (National Institute of Mental Health, Bethesda, MD, USA) was used to measure the intensity of specific bands. All experiments were repeated thrice.

Quantitative real-time PCR (qRT-PCR)

Total RNA was lysed from previously collected glioma specimens using TRIZOL reagent (Thermo Fisher Scientific, Waltham, MA, USA). cDNA synthesis and RT-PCR were performed using SYBR® Premix Ex Taq™ Kit (Takara Bio Inc., Shiga, Japan). 18S rRNA was used as an internal control. LKB1 primer sequences were as follows: forward 5'-AGGGATGCTTGAGTACGAACC-3' and reverse 5'-GTCCTCCAAGTACGGCACC-3'. Snail primer sequences were as follows: forward 5'-TCGGAAGCCTAACTACAGCGA-3' and reverse 5'-AGATGAGCATTGGCAGCGAG-3'. Relative mRNA expression level was standardized as described earlier.^{13,14}

Immunohistochemical (IHC) and immunofluorescence staining analyses

IHC analysis and semiquantitative scoring were performed as described previously.¹⁵ Cells were grown on 35 mm dishes and fixed with 4% paraformaldehyde. After fixation, cells were penetrated with Triton X-100, closed with 5% BSA in PBS for 1.5 hours at room temperature, and dyed with anti-LKB1 (1:500) with 0.3% BSA in PBS and Cy3-conjugated secondary antibodies (1:100; Cell signaling Technology, Danvers, MA, USA) with 0.3% BSA in PBS. Fluorescence was visualized under laser scanning confocal microscope (Olympus Corporation, Tokyo, Japan).

IHC analysis and semiquantitative scoring were performed as described previously.¹⁶ Scores were defined based on cell-positive dyeing intensity (0= negative; 1= weak;

2= moderate; and 3= strong) multiplied by the proportion of positively dyed cells (75%–100%=4; 50%–74%=3; 25%–49%=2; 1%–24%=1; and 0%=0), leading to scores ranging 0–12.

Dual luciferase reporter assays

U-87MG and U-251MG cells were routinely seeded in 24-well plates for 24 hours before transfection with NF-κB firefly luciferase reporter plasmid and phRL-TK (Shanghai GeneChem Co., Ltd.) using Lipofectamine™ 3000 (Thermo Fisher Scientific) according to the manufacturer's protocol. The Renilla luciferase expression plasmid was used as an internal control. At 24 hours after transfection, cells were harvested and lysed with 100 µL of 1× passive lysis buffer (Boster Biological Technology). Subsequently, luciferase activities were calculated using a Dual-Glo Luciferase kit (Promega Corporation, Fitchburg, WI, USA).

Statistical analysis

Data are shown as mean and SD. Statistical significance was tested using two-tailed Student's *t*-test and chi-squared test. Survival analyses were performed using log-rank tests and Kaplan–Meier plots. Multivariate survival analyses were performed using a Cox regression model. Statistical analysis was performed using the Empower Stats Statistical Software Program Version 2.16.1, with *P*<0.05 considered significant.

Ethics approval and consent to participate

The use of human tissues was approved by the Human Research Committee of Huazhong University of Science and Technology (S304). Written informed consent was obtained from each patient.

Results

LKB1 was often downregulated in human glioma tissues

RT-PCR showed remarkable differences in LKB1 expression among human glioma specimens of different grades. Compared to low-grade gliomas, high-grade gliomas showed significantly less *LKB1* gene expression (Figure 1A, *P*<0.001). Western blotting showed that LKB1 was often downregulated at a higher grade (Figures 1B and S1). We measured LKB1 expression levels by Western blotting in two normal brain tissues and glioma tissues of various grades (Figure S2) and checked the expression levels of LKB1 in glioma cell lines including U87MG and U251MG and in normal astrocytes (Figure S3). IHC analyses were performed to test LKB1 expression in 49 human glioma samples. As shown in Figure 1C (Figure S4), immunostaining




Figure 1 LKB1 is often downregulated in human gliomas.

Notes: (A) Relative LKB1 expression levels measured by RT-PCR in 31 patients with LGG and 18 patients with HGG. (B) Sixteen glioma tissues were assessed for LKB1 expression by Western blot, including four grade I, four grade II, four grade III, and four grade IV. (C) Representative images and (D) scores of the IHC analysis of LKB1 expression in the paraffin-embedded different grade glioma. (E) Kaplan–Meier curves for DFS of glioma patients with low vs high expression of LKB1. The median LKB1 expression was used as the cutoff value. Statistical significance was assessed using two-tailed Student's *t*-test. Scale bars: 50 μ m. ** $P<0.01$, and *** $P<0.001$.

Abbreviations: DFS, disease-free survival; HGG, high grade glioma; IHC, Immunohistochemical; LGG, low grade glioma; LKB1, liver kinase B1; RT, real-time.

intensity of LKB1 evidently differed among different grades of gliomas. Compared to low-grade gliomas, high-grade gliomas showed weaker LKB1 immunostaining intensity and significantly decreased LKB1 protein expression on quantification analyses (Figure 1D, $P<0.01$). Correlation analysis of LKB1 expression with clinicopathological characteristics of 49 glioma samples with informative IHC demonstrated that downregulated LKB1 expression was significantly correlated with the Karnofsky Performance Scale (KPS) score ($P=0.001$) and recurrence ($P=0.01$, Table 1); this was consistent with the results of a previous report that KPS was an independent predictor of survival.¹⁷ In addition, Kaplan–Meier analysis demonstrated that low LKB1 expres-

sion was correlated with shorter recurrence-free survival in malignant gliomas (Figure 1E, $P<0.05$). Multivariate Cox regression analysis further proved that disease-free survival was significantly higher in tumors with high LKB1 protein expression than in those with low LKB1 protein expression (HR: 2.097, 95% CI: 1.557–2.637, $P=0.013$).

LKB1 downregulation promoted glioma cell proliferation, migration, and invasion

To explore the role of LKB1 in glioma cell proliferation, we knocked down LKB1 in U-87 and U-251 cells. At 48 hours after transfection, the downregulation efficacy of LKB1 was measured by qRT-PCR and western blotting

Table I Association of LKB1 expression with clinicopathological characteristics in human gliomas

Features	No	LKB1		P-value
		Low	High	
Age (years)				0.79
<50	26	16	10	
≥50	23	15	8	
Gender				0.96
Male	27	17	10	
Female	22	14	8	
Tumor size, cm				0.005
<5	28	13	15	
≥5	21	18	3	
Tumor location				0.93
Supratentorial	35	22	13	
Subtentorial	14	9	5	
KPS				0.001
<90	26 ^a	22 ^a	4	
≥90	21	8	13	
WHO grade				0.0006
Low-grade (I+II)	31	14	17	
High-grade (III+IV)	18	17	1	
Tumor recurrence				0.01
No	32 ^a	16 ^a	16	
Yes	16	14	2	

Notes: ^aPartial data not available; statistics based on available data.

Abbreviations: KPS, Karnofsky Performance Scale; LKB1, liver kinase B1.

(Figure 2A and B). CCK-8 assay demonstrated that cell growth rates were markedly higher in the LKB1 knockdown group than in the scramble control group (Figure 2C, $P<0.001$). Colony-forming assay showed that compared to the control group, the LKB1 knockdown group yielded much greater number of colonies (Figure 2D, $P<0.01$). Transwell migration assay indicated that the number of migrated cells significantly increased after LKB1 knockdown (Figure 2E, $P<0.01$). Invasion assay showed that compared to the control

group, the LKB1 knockdown group showed more invasive cells (Figure 2F, $P<0.01$).

LKB1 upregulation suppressed glioma cell proliferation, migration, and invasion

LKB1 was overexpressed using lentiviruses containing shRNA to specifically target ELTD1 in the glioma cell lines U-87 and U-251. Transfection efficiency was verified using qRT-PCR and Western blotting (Figure 3A and B), and CCK-8, colony-forming, and Transwell assays were used to test effects of LKB1 on glioma cell proliferation, migration, and invasion. LKB1 overexpression significantly suppressed glioma cell proliferation, migration, and invasion (Figure 3C–F).

Snail is involved in LKB1-regulated glioma cell proliferation, migration, and invasion

Snail protein has been reported to regulate several cellular processes through a series of mechanisms. Previous studies have indicated that Snail overexpression promoted glioma cell proliferation, migration, and invasion. Therefore, LKB1 upregulation may repress Snail expression, thereby suppressing glioma cell proliferation, migration, and invasion. Our results showed obvious increase in Snail protein expression after LKB1 knockdown (Figure 4A). However, the Slug protein expression level remains unchanged (Figure S5). In addition, to verify whether Snail participated in LKB1-regulated glioma cell proliferation and invasion, we used lentiviruses containing shRNA to target Snail in U-87 and U-251 cells (Figure 3B). As expected, Snail overexpression abrogated the effects of LKB1 overexpression




Figure 2 (Continued)




Figure 2 Downregulation of LKB1 promotes glioma cell proliferation, migration, and invasion.

Notes: (A) The knockdown efficiency against LKB1 was verified by Western blot in U-87 and U-251 cells. (B) Growth curves between none, scramble, and LKB1(KD) by CCK-8 assay. The results are shown as the mean \pm SD of at least three independent experiments. (C–E) Representative images (left panels) and histogram quantification (right panels) of the (C) colony formation assay, (D) Transwell migration, and (E) invasion assays with U-87 and U-251 cells. Statistical significance was assessed using one-way ANOVA followed by Dunnett's tests for multiple comparisons. Scale bars: 50 μ m. ** P <0.01 and *** P <0.001.

Abbreviations: CCK-8, Cell Counting Kit-8; LKB1, liver kinase B1; KD, knockdown.

on inhibiting U-87MG and U-251MG cell proliferation (Figure 4C), colony formation (Figure 4D), migration (Figure 4E), and invasion (Figure 4F). Taken together, these results indicate that LKB1 inhibited glioma cell proliferation, migration, and invasion by regulating Snail protein expression.

LKB1 regulates glioma cell proliferation, migration, and invasion via the NF- κ B/Snail signaling pathway

We treated LKB1-knockdown cells with two chemical inhibitors of NF- κ B, CAPE, and JSH-23. First, we confirmed

the inhibition efficiency by p65 in U-87 and U-251 cells (Figure 5A). Both CAPE and JSH-23 attenuated the Snail expression induced by LKB1 knockdown in U-87 and U-251 cells (Figure 5B). Next, we confirmed whether the NF- κ B signaling pathway was activated after LKB1 knockdown. After LKB1 knockdown, I κ B α subunit decreased, the phosphorylation of I κ B kinase added, and the phosphorylation

of p65 increased (Figure 5C); all these effects were rescued by I κ B α silencing. Furthermore, p65 migrated to U-87 cell nuclei after LKB1 knockdown; this effect was also rescued by I κ B α overexpression (Figure 5D). Then, U-87 cells were infected with an NF- κ B luciferase reporter plasmid to confirm the transcriptional activation of NF- κ B. LKB1 knockdown induced NF- κ B activation and simultaneous I κ B α -guided




Figure 3 (Continued)




Figure 3 Upregulation of LKB1 suppresses glioma cell proliferation, migration, and invasion. **Notes:** (A) The overexpression efficiency against LKB1 was verified by Western blot in U-87 and U-251 cells. (B) Growth curves between none, vector, and LKB1 (OE) by CCK-8 assay. The results are shown as the mean \pm SD of at least three independent experiments. (C–E) Representative images (left panels) and histogram quantification (right panels) of the (C) colony formation assay, (D) Transwell migration, and (E) invasion assays with U-87 and U-251 cells. Statistical significance was assessed using one-way ANOVA followed by Dunnett’s tests for multiple comparisons. Scale bars: 50 μ m. ****** P <0.01. **Abbreviations:** CCK-8, Cell Counting Kit-8; LKB1, liver kinase B1; OE, overexpression.




Figure 4 (Continued)




Figure 4 Snail is involved in LKB1-regulated glioma cell proliferation, migration, and invasion.

Notes: (A) LKB1 knockdown decreased the protein level of Snail. Other proteins remain unchanged. (B) The overexpression efficiency against Snail was verified by Western blot in U-87 and U-251 cells. (C) Growth curves between none, vector, Snail(OE), LKB1(OE), and Snail(OE) + LKB1(OE) by CCK-8 assay. The results are shown as the mean \pm SD of at least three independent experiments. (D–F) Representative images (left panels) and histogram quantification (right panels) of the (D) colony formation assay, (E) Transwell migration, and (F) invasion assays with U-87 and U-251 cells. Statistical significance was assessed using one-way ANOVA followed by Dunnett's tests for multiple comparisons. Scale bars: 50 μ m. ** $P < 0.01$ and *** $P < 0.001$.

Abbreviations: CCK-8, Cell Counting Kit-8; LKB1, liver kinase B1; OE, overexpression.

luciferase overexpression toward the baseline (Figure 5E). Finally, we knocked down I κ B α and probed Snail increased expression (Figure 5F). Collectively, these results indicate that NF- κ B activation is involved in Snail expression.

Discussion

Gliomas account for ~40% of the diagnosed primary central nervous system tumors. Among gliomas, glioblastoma is the most common malignant type that is associated with very poor prognosis, with survival duration for most patients being only 15 months from diagnosis.^{18,19} Therefore, identification of novel and efficient molecular markers is urgent to diagnose and treat patients with glioma.

In this study, we measured LKB1 expression in 49 glioma samples using RT-PCR and in 16 glioma samples using Western blotting and found that LKB1 was often downregulated in

high-grade gliomas. Subsequently, we analyzed the clinical relevance of LKB1, which highlighted the potential of LKB1 as a novel clinical prognostic index. Huang et al have reported that LKB1 downregulation promoted tumor progression and predicted unfavorable prognosis in patients with glioma.²⁰ However, precise mechanisms underlying this role remain largely unknown. In this study, we observed that the LKB1 protein modulated tumor cell proliferation, migration, and invasion, thereby, playing an important role in tumor progression. Through gain and loss of function experiments, we found that LKB1 inhibited glioma cell proliferation, migration, and invasion. LKB1 knockdown markedly promoted tumor cell proliferation, whereas ELTD1 overexpression inhibited tumor cell growth.

Furthermore, by screening for possible signaling pathways using a dual luciferase reporter assay (Figure S6), we




Figure 5 LKB1 regulates glioma cell proliferation, migration, and invasion via the NF-κB/Snail signaling pathway. **Notes:** (A and B) U-87 and U-251 cells transfected with LKB1 plasmid and co-cultured with NF-κB signaling inhibitors were reaped, and the lysates were immune-blotted for (A) p-P65 Tot.P65, (B) Snail, Slug, and β-actin. (C) LKB1 knockdown decreased the protein level of IκBα and increased the protein level of p-IKK and p-P65. Other proteins remain unchanged. (D) The p65 migrated to the nuclei of U-87 cells after LKB1 knockdown and its effects were abrogated by IκBα overexpression. (E) LKB1 knockdown induced NF-κB activation and IκBα overexpression abrogated its effects. (F) IκBα knockdown increased the protein level of Snail. Slug remains unchanged. Statistical significance was assessed using one-way ANOVA followed by Dunnett's tests for multiple comparisons. Scale bars: 50 μm. **P<0.01. **Abbreviations:** LKB1, liver kinase B1; NS, no statistical significance.

found that the *LKB1* gene regulated NF- κ B signaling, which plays important roles in cancer growth and metastasis. P65 is a regulatory transcription factor particularly vital in cell survival, growth, and differentiation.^{21,22} NF- κ B activation blocked Snail degradation, subsequently promoting cancer cell migration and invasion.^{23,24} Moreover, LKB1 knockdown markedly enhanced NF- κ B reporter activity. In particular, the expression of tumor-related gene *Snail* was upregulated at the mRNA and protein levels when LKB1 expression was disrupted in glioma cells, suggesting that Snail is the downstream effector molecule of LKB1.

Conclusion

In summary, we demonstrated the function and expression patterns of LKB1 in glioma and frequently downregulated LKB1 expression in glioma tissues. Our findings indicated that the tumor suppressor role of LKB1 is possibly conferred via NF- κ B/Snail signaling; therefore, targeting NF- κ B/Snail signaling may inhibit tumor cell growth. Understanding the precise role of LKB1 in the NF- κ B/Snail axis will improve our knowledge of LKB1-induced tumor growth inhibition and enable the development of novel molecular therapeutic strategies against glioma.

Acknowledgments

The study was funded by the Funds for Creative Research of Union Hospital, Tongji Medical College, Huazhong University of Science and Technology (02.03.2017-65). No external funding was received for this study.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Wen PY, Kesari S. Malignant gliomas in adults. *N Engl J Med Overseas Ed*. 2008;359(5):492–507.
2. Louis DN, Perry A, Reifenberger G, et al. The 2016 World Health organization classification of tumors of the central nervous system: a summary. *Acta Neuropathol*. 2016;131(6):803–820.
3. Boudeau J, Sapkota G, Alessi DR. LKB1, a protein kinase regulating cell proliferation and polarity. *FEBS Lett*. 2003;546(1):159–165.
4. Shackelford DB, Shaw RJ. The LKB1-AMPK pathway: metabolism and growth control in tumour suppression. *Nat Rev Cancer*. 2009;9(8):563–575.
5. Green AS, Chapuis N, Lacombe C, Mayeux P, Bouscary D, Tamburini J. LKB1/AMPK/mTOR signaling pathway in hematological malignancies: from metabolism to cancer cell biology. *Cell Cycle*. 2011;10(13):2115–2120.
6. Bakula D, Müller AJ, Zuleger T, et al. WIPI3 and WIPI4 β -propellers are scaffolds for LKB1-AMPK-TSC signalling circuits in the control of autophagy. *Nat Commun*. 2017;8:15637.
7. Yu L, Li L, Medeiros LJ, Young KH. NF- κ B signaling pathway and its potential as a target for therapy in lymphoid neoplasms. *Blood Rev*. 2017;31(2):77–92.
8. Xie B, Nie S, Hu G, et al. The involvement of NF- κ B/Klotho signaling in colorectal cancer cell survival and invasion. *Pathol Oncol Res*. Epub 2019 Jan 5.
9. Wu Y, Deng J, Rychahou PG, Qiu S, Evers BM, Zhou BP. Stabilization of snail by NF- κ B is required for inflammation-induced cell migration and invasion. *Cancer Cell*. 2009;15(5):416–428.
10. Bloom MJ, Saksena SD, Swain GP, Behar MS, Yankeelov TE, Sorace AG. The effects of IKK-beta inhibition on early NF- κ B activation and transcription of downstream genes. *Cell Signal*. 2019;55:17–25.
11. Ni R, Huang Y, Wang J. miR-98 targets ITGB3 to inhibit proliferation, migration, and invasion of non-small-cell lung cancer. *Oncotargets Ther*. 2015;8:2689–2697.
12. Ye H, Liu K, Qian K. Overexpression of long noncoding RNA HOTTIP promotes tumor invasion and predicts poor prognosis in gastric cancer. *Oncotargets Ther*. 2016;9:2081–2088.
13. Pang Y, Mao H, Shen L, Zhao Z, Liu R, Liu P. MiR-519d represses ovarian cancer cell proliferation and enhances cisplatin-mediated cytotoxicity in vitro by targeting XIAP. *OTT*. 2014;7:587–597.
14. Xiao J, Lin H-Y, Zhu Y-Y, Zhu Y-P, Chen L-W. MiR-126 regulates proliferation and invasion in the bladder cancer BLS cell line by targeting the PIK3R2-mediated PI3K/Akt signaling pathway. *Oncotargets Ther*. 2016;9:5181–5193.
15. Yang L-H, Wang Y, Qiao S, et al. Liver-enriched activator protein 1 as an isoform of CCAAT/enhancer-binding protein beta suppresses stem cell features of hepatocellular carcinoma. *Cancer Manag Res*. 2018;10:873–885.
16. Dai X, Zhao L, Peng F. Primary clear cell carcinoma of the thymus and literature comparison of features. *Cancer Manag Res*. 2018;10:513–518.
17. Lacroix M, Abi-Said D, Fournay DR, et al. A multivariate analysis of 416 patients with glioblastoma multiforme: prognosis, extent of resection, and survival. *J Neurosurg*. 2001;34(2):190–198.
18. Thon N, Kreth S, Kreth F-W. Personalized treatment strategies in glioblastoma: MGMT promoter methylation status. *Oncotargets Ther*. 2013;6:1363–1372.
19. Thomas AA, Brennan CW, Deangelis LM, Omuro AM. Emerging therapies for glioblastoma. *JAMA Neurol*. 2014;71(11):1437–1444.
20. Huang J, Chen H, Wei Q, Zhang Z, Zhong Z, Xu Y. Downregulation of LKB1 promotes tumor progression and predicts unfavorable prognosis in patients with glioma. *Oncol Lett*. 2017;13(3):1688–1694.
21. Hai L, Liu P, Yu S, et al. Jagged1 is Clinically Prognostic and Promotes Invasion of Glioma-Initiating Cells by Activating NF- κ B(p65) Signaling. *Cell Physiol Biochem*. 2018;51(6):2925–2937.
22. Zhang P, Chen X-B, Ding B-Q, Liu H-L, He T. Down-regulation of ABCE1 inhibits temozolomide resistance in glioma through the PI3K/Akt/NF- κ B signaling pathway. *Biosci Rep*. 2018;38(6):BSR20181711.
23. Lu Z, Li Y, Wang J, et al. Long non-coding RNA NKILA inhibits migration and invasion of non-small cell lung cancer via NF- κ B/Snail pathway. *J Exp Clin Cancer Res*. 2017;36(1):54.
24. Zhang Y, Yang X, Wu H, Zhou W, Liu Z. MicroRNA-145 inhibits migration and invasion via inhibition of fascin 1 protein expression in non-small-cell lung cancer cells. *Mol Med Rep*. 2015;12(4):6193–6198.

Supplementary materials




Figure S1 LKB1 expression levels in two NBTs and glioma tissues of various grades.

Abbreviations: LKB1, liver kinase B1; NBT, normal brain tissue.




Figure S2 Expression levels of LKB1 in glioma cell lines including U87MG and U251MG and in normal astrocytes.

Abbreviation: LKB1, liver kinase B1.




Figure S3 The Slug protein expression level remains unchanged in U-87 and U-251.

Notes: *** $P < 0.001$.

Abbreviations: LKB1, liver kinase B1; KD, knockdown.




Figure S4 LKB1 level in low grade and high grade glioma.

Abbreviations: HGG, high grade glioma; LGG, low grade glioma; LKB1, liver kinase B1.




Figure S5 Immunostain for LKB1 in LGG and HGG.

Notes: Magnification $\times 20$.

Abbreviations: HGG, high grade glioma; LGG, low grade glioma; LKB1, liver kinase B1.




Figure S6 Luciferase reporter assay to screening signaling pathway.

Abbreviations: HGG, high grade glioma; LGG, low grade glioma; LKB1, liver kinase B1; KD, knockdown.

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.