


High incidence of multidrug-resistant *Escherichia coli* coharboring *mcr-1* and *bla*_{CTX-M-15} recovered from pigs

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Muhammad Shafiq¹ 
Jinhu Huang¹
Sadeeq Ur Rahman²
Jan Mohammad Shah¹
Li Chen¹
Yi Gao¹
Mengli Wang¹
Liping Wang¹

¹Moe Joint International Research Laboratory of Animal Health and Food Safety, Laboratory of Veterinary Pharmacology and Toxicology, College of Veterinary Medicine, Nanjing Agricultural University, Nanjing, 210095, People's Republic of China; ²College of Veterinary Sciences and Animal Husbandry, Section Microbiology, Abdul Wali Khan University, Mardan, KP, Pakistan

Purpose: The coexistence of mobile colistin (COL)-resistant gene *mcr-1* with extended-spectrum beta-lactamase (ESBL) gene in *Escherichia coli* has become a serious threat globally. The aim of this study was to investigate the increasing resistance to COL and in particular its coexistence with ESBL-producing *E. coli* recovered from pig farms in China.

Materials and methods: *E. coli* were isolated from 14 pig farms in Jiangsu China. Susceptibility testing was identified by micro-dilution method. PCR assay and nucleotide sequencing were used to detect COL-resistant genes, *mcr-1* to *-5*, as well as ESBL genes, *bla*_{CTX-M}, *bla*_{SHV} and *bla*_{TEM}. Conjugation experiment, plasmid replicon typing of the multidrug resistance (MDR), S1-PFGE and DNA southern hybridization were performed to study the transferability of these genes.

Results: Overall, 275 *E. coli* isolates were recovered from a total of 432 cloacal and nasal swabs. More than 90% of the isolates were MDR, of which 70.18% were resistant to COL. Of these 275 isolates, *mcr-1* was identified as the most predominant gene carried by 71.63% (197/275) of isolates, 39.59% (78/197) of the isolates were harboring both *mcr-1* and ESBL genes (*bla*_{CTX-M}, *bla*_{SHV} and *bla*_{TEM}). ESBL genotyping showed that *bla*_{CTX-M} was the most predominant ESBL (68.49%) followed by *bla*_{SHV} (16.4%) and *bla*_{TEM} (15%). Sequencing revealed that the most common variants of *bla*_{CTX-M} identified were, *bla*_{CTX-M-15} (69%), *bla*_{CTX-M-55} (29%) and *bla*_{CTX-M-1} (1.8%). IncHI2, IncFIB, IncFIC, IncN and IncX4 were found to be the most common Inc-types found both in donors and in transconjugants and were associated with the transfer of the *mcr-1* and ESBL encoding genes. Six strains carried a total of five different plasmids: approximately 97-, 130-, 160-, 227- and 242-kb plasmids.

Conclusion: The coexistence of the *mcr-1*- and *bla*_{CTX-M-15}-carrying isolates displaying high MDR, recovered from *E. coli* of pig origin, is a major concern for both humans and veterinary medicine.

Keywords: *E. coli*, colistin, *mcr-1*, ESBL, coexistence

Introduction

Antimicrobial resistance (AMR) has now been widely recognized as a crucial threat to human and animal health as the extensive use of antimicrobials in humans as well as in food-producing animals.¹ Global consumption of antimicrobials in animal settings may rise up to 67% by 2030, determined predominantly by BRICS (Brazil, Russia, India, China and South Africa) countries, as large-scale and intensive farming operations are greatly in demand with the upsurge in revenue and animal protein consumption.^{2,3} This heavy antimicrobial practice creates a selective pressure that contributes to the emergence and spread of bacterial resistance. One of the major concerns is the rapid increase

Correspondence: Liping Wang
Moe Joint International Research Laboratory of Animal Health and Food Safety, Laboratory of Veterinary Pharmacology and Toxicology, College of Veterinary Medicine, Nanjing Agricultural University, Yifu Building 2021, Nanjing 210095, People's Republic of China
Tel +86-025-2584395573
Fax +86-025-84398669
Email wlp71@163.com

of the multidrug-resistant (MDR) *Escherichia coli* in animal settings and clinical medicine.^{4–6} This is not only because of the lessened number of useful antimicrobials for curing MDR *E. coli* infections, but also due to the potential transfer of MDR *E. coli* strains from animals to humans, especially which producing extended-spectrum β -lactamases (ESBLs) and carbapenemases, and display resistance to colistin (COL).⁷

ESBLs are β -lactamases that confer resistance to oxyimino “second- and third-generation” cephalosporin’s (eg, cefotaxime (CTX), ceftriaxone and ceftazidime) and aztreonam.^{8,9} ESBL-producing bacteria were first reported in 1980, soon after the introduction of the third-generation cephalosporin’s (CTX and ceftiofur (CEF) into clinical settings.¹⁰ Currently, there are more than 350 ESBL genes that have been reported, and these genes are commonly developed through point mutations of the classical SHV-1 and TEM-1 β -lactamases and more increasingly prominent the CTX-M types.^{11–13} Among the CTX-M enzymes, *bla*_{CTX-M-55} has become the leading CTX-M type in ESBL-producing *E. coli* isolates of animal origin during the last decade.^{14,15} In contrast, *bla*_{CTX-M-15} seems to be the most extensive types in isolates of human origin.¹⁶ ESBL-producing *E. coli* are highly linked with multiple plasmids and studies have reported that ESBL genes are often carried on IncF, IncI1, IncN, IncHI1 and IncHI2 in food-producing animals worldwide.^{13,17–19} There is potential for ESBL genes/plasmid spreading between *E. coli* from animals, food and humans.^{20,21}

The co-occurrence of ESBL genes and *mcr-1* in *E. coli* was reported from China in 2016.²² Rhouma and Letellier assumed that a historic relation existed between ESBL genes, carbapenemase genes and *mcr-1*.²³ A recent study proposed that cephalosporin resistance is commonly spread in animals and humans through distinct plasmids.²⁴ It is highly expected that food-producing animals have become the most significant reservoirs in disseminating these resistance genes in the community through horizontal gene transfer. To assess the co-occurrence and emergence of *mcr* and ESBL genes in *E. coli* of pig source, we examined 14 pig farms of Jiangsu province in China to evaluate the current scenario of these resistant genes in pigs and further clarified the predominant genotype and plasmids diversity of *mcr* and ESBL genes.

Materials and methods

Collection of samples

A total of 432 samples (400 from healthy and 32 from dead pigs) were collected from 14 commercial pig farms in

Jiangsu, China (Figure S1), during the period of August 2016 until December 2017. From each farm, samples were randomly collected. The anal swabs were collected by inserting the swab into the rectum and being rotated. To collect nasal swabs from swine, the nose was wiped with a piece of paper and a sterile swab was inserted into the nasal cavity and rotated for 3 s at 90°. From 32 dead pigs, all samples were aseptically obtained from different organs. All collected samples were immediately transported at 4°C to the laboratory for microbial examination and processed within 4 hrs.

Isolation and screening of ESBL-producing and COL-resistant *E. coli*

All samples were directly streaked onto MacConkey agar (Binhe Microorganism Reagent Co. Ltd., Hangzhou, China) supplemented with CTX (1 μ g/mL) and COL (2 μ g/mL) for the screening of possible ESBL-producing and COL nonsensitive *E. coli* as previously described.²⁵ Plates were incubated at 37°C for 18–24 hrs. Presumptive *E. coli* colonies with dark pink to red colors were confirmed microscopically and further verified by species-specific PCR as described previously.²⁶ Confirmed *E. coli* strains were stored in Luria–Bertani medium (Oxoid, United Kingdom) containing 40% (vol/vol) glycerol in aliquots at –80°C until further use.

ESBL-producing *E. coli* were further confirmed by double-disk synergy (DDS) testing as recommended by the Clinical and Laboratory Standards Institute (CLSI) guideline, using antibiotic discs of ceftazidime (30 μ g), ceftazidime plus clavulanic acid (30/10 μ g), CTX (30 μ g) and CTX plus clavulanic acid (30/10 μ g). DDS test was performed for phenotypic detection of ESBLs. The test result is considered as positive if the zone of inhibition is ≥ 5 mm larger with clavulanic acid than without.²⁷

Antimicrobial susceptibility testing

Antimicrobial susceptibility was performed by minimum inhibitory concentration (MIC) determination using broth microdilution method against 11 antibiotics for all 275 isolates and 17 antibiotics for transconjugants. The MIC data was interpreted according to the CLSI recommendations.²⁷ Antibiotics used in this study, comprised of 5 β -lactams – ampicillin (AMP), CTX, ceftiofur (CFX), CEF and meropenem (MEM) – and 12 non- β -lactams – COL, ciprofloxacin (CIP), chloramphenicol (CHL), enrofloxacin (ENR), gentamycin (GEN), kanamycin, nalidixic acid, polymyxin-B (POL-B), tetracycline (TET), trimethoprim, streptomycin and sulfamethoxazole. The MIC of COL was determined by broth

micro-dilution method recommended by the joint CLSI-EUCAST polymyxin breakpoints working group (www.EUCAST.org), CLSI VET01-A4 is used for CEF and ENR which are missing in the human CLSI M100-S27. *E. coli* ATCC 25922 was used as a quality control in antimicrobial susceptibility testing. Isolates that exhibited resistance to more than 3 antimicrobial agents were classified as MDR.²⁸

PCR assays for detection of *mcr* and ESBL genes

PCR assay was used to detect COL-resistant genes *mcr-1* to *mcr-5* as well as ESBL genes (*bla*_{CTX-M}, *bla*_{SHV} and *bla*_{TEM}). Total DNA was isolated by conventional boiling method. All these resistant genes were screened via PCR-based diagnostics with specific primers, as previously described. All the primers and PCR conditions used in this study are listed in Table 1. All PCR positive amplicons of these targeted genes were sequenced by Sanger sequencing in TSINGKE Corporation (Nanjing, PR China).

Conjugation experiment

To determine the transferability of resistance genes, 15 COL-resistant *E. coli* isolates were selected as donors for conjugation. *E. coli* EC-600 (Nal^R, Rif^R) was used as recipient bacteria. Conjugation experiments were performed as previously described.²⁹ These putative transconjugants were further confirmed using antibiotic susceptibility testing, PCR detection and plasmid incompatibility (Inc) groups typing carried by the transconjugants.

Plasmid replicon typing

Plasmid DNA was extracted from both donors and transconjugants using the Wizard Genomic DNA Purification kit (Promega) and was characterized by PCR-based replicon typing method (PBRT). Eighteen pairs of primers were designed to perform 5 multiplex and 3 simple PCR targeting the FIA, FIB, FIC, HI1, HI2, IncI1, L/M, N, P, W, T, A/C, K, B/O, X, Y, F and FIIA replicons as previously mentioned.³⁰ While primers for two other plasmids IncI2 and IncX4 were designed

Table 1 PCR primers and conditions used in this study

Primer name	PCR target	Sequence (5'- 3')	Annealing temperature (°C)	Product size (bp)	Reference
<i>E. coli</i>-specific					
UAL UAR	<i>uidA</i>	TGGTAATTACCGACGAAAACG GC ACG CGT GGT TAC AGT CTT GCG	62	147	58
ESBL genes					
CTX-MA CTX-MB	<i>bla</i> _{CTX-M}	CGC TTT GCG ATG TGC AG ACC GCG ATA TCG TTG GT	54	550	58
SHV-F SHV-R	<i>bla</i> _{SHV}	GGG TTA TTC TTA TTT GTC GC TTA GCG TTG CCA GTG CTC	58	930	58
TEM-F TEM-R	<i>bla</i> _{TEM}	ATA AAA TTC TTG AAG ACG AAA GAC AGT TAC CAA TGC TTA ATC	56	1086	58
MCR genes					
CLR5-F CLR5-R	<i>mcr-1</i>	CGG TCA GTC CGT TTG TTC CTT GGT CGG TCT GTA GGG	58	309	41
MCR-2-F MCR-2-R	<i>mcr-2</i>	TGT TGC TTG TGC CGA TTG GA AGA TGG TAT TGT TGG TTG CTG	58	567	42
MCR-3-F MCR-3-R	<i>mcr-3</i>	TTG GCA CTG TAT TTT GCA TTT TTA ACG AAA TTG GCT GGA ACA	50	542	43
MCR-4F MCR-4R	<i>mcr-4</i>	ATT GGG ATA GTC GCC TTT TT TTA CAG CCA GAA TCA TTA TCA	58	487	44
MCR-5-F MCR-5-R	<i>mcr-5</i>	ATG CGG TTG TCT GCA TTT ATC TCA TTG TGG TTG TCC TTT TCT G	50	1644	45

separately which were missing in the previous replicon typing. All the PBRT primers and PCR conditions used in this study are listed in the Supplementary Table S1. PCR amplicons of plasmids were sequenced by Sanger sequencing (TSINGKE Corporation, Nanjing, PR China) and retrieved sequences were used to confirm replicon types by using BLAST tool available at NCBI web (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>).

Pulsed field gel electrophoresis (PFGE) and Southern hybridization

To determine the genetic relatedness and location of transmissible *mcr-1*-positive elements, the six conjugative *E. coli* strains were characterized by S1-PFGE and Southern hybridization using a probe specific for *mcr-1*. Genomic DNA from each of the isolate was digested with S1 nuclease (Thermo scientific) and was examined by PFGE as previously described.³¹

Southern hybridizations of plasmid DNA were performed with a digoxin-labeled *mcr-1*-specific probe according to the manufacturer's instructions (Roche Diagnostics, Mannheim, Germany) as previously described.³²

Statistical analysis

Differences in the AMR profiles of *E. coli* isolates with or without *mcr-1* were assessed by a two-tailed Chi-square test or Fisher's exact test using the Statistical Packages of Social Sciences software for Windows, version 20.0 (IBM Corp., Armonk, NY), with $P < 0.05$ set as the level of significant differences.

Results

Bacterial isolation and antimicrobial susceptibility

Overall, 275 *E. coli* isolates (243 from healthy and 32 from dead pigs) were recovered from 432 samples of 14 different pig farms. All isolates were observed to be ESBL-producing and COL nonsensitive as determined by phenotypic approaches, giving a carriage rate of 63.6% (275/432). Of these 275 *E. coli* isolates, 174 (63.27%) were from feces, 69 (25.09%) were from nasal and remaining 32 (11.63%) were from diseased and dead pigs.

The MICs were obtained from the antibiotic susceptibility testing for all isolates. To determine the resistance profiles of 275 representative *E. coli* strains, susceptibility of 11 antibiotics were used (Table 2). Of the 275 *E. coli* isolates, the resistant rate to TET was 97.81%, followed by AMP (96.72%), CHL

(94.54%), CFX (86.18%), CTX (78.18%), CEF (77.81%), CIP (73.81%), POL-B (71.27%), GEN (70.54%), and COL (70.18%). In contrast, the most effective antibiotic against these isolates was MEM with 99.6% susceptibility. Majority of the *E. coli* strains showed considerable MDR to β -lactams and several non- β -lactams groups, including polypeptides group, fluoroquinolones, aminoglycosides, amphenicol, quinolone group, sulfonamides and TET.

mcr-1 and ESBL genes are prevalent among *E. coli* strains

Although 70.18% (193 of 275) swine *E. coli* isolates conferred resistance to COL, the *mcr-1* carriage rate was 71.63% (197/275) (Table 3) and only *mcr-1* gene was detected in these COL-resistant *E. coli* isolates. No other colistin-resistant gene (*mcr-2* to *mcr-5*) could be detected in the study population of *E. coli* isolates. The *mcr-1* gene was detected in all farms, and the prevalence rate was enormously high 71.6%, ranging from 47.8% to 100% in different farms, while 40.6% in diseased isolates (Table 3).

We recovered a total of 146 (53.09%) ESBL-producing *E. coli* strains from 275 samples collected from 14 different farms of pigs (Figure S2 and Table 5). Among the 146 ESBL-producing isolates, 68.49% (100/146) harbored *bla*_{CTX-M} genes, 22.60% (33/146) harbored *bla*_{SHV}, while 18.49% (27/146) were carrying *bla*_{TEM} genes. Among them, 6 isolates carried *bla*_{CTX-M} and *bla*_{SHV}, 3 isolates contained *bla*_{CTX-M} and *bla*_{TEM}, and one isolate had three genes of *bla*_{CTX-M}, *bla*_{SHV}, and *bla*_{TEM}.

The resistance patterns of *E. coli* are different in *mcr-1*-positive and *mcr-1*-negative isolates

The *mcr-1*-positive *E. coli* isolates displayed more resistance to other antimicrobials than those of *mcr-1*-negative isolates (Table 4). For *mcr-1*-positive *E. coli*, all isolates, 100% (197/197), possessed not less than 3 antibiotics resistance pattern, while about 96.15% (75/78) of COL-negative isolates did ($P=0.006$), with 3 COL-sensitive isolates which displayed resistance to not more than 2 antibiotics. In addition, about two-thirds of *mcr-1*-positive *E. coli* isolates, 68.52% (135/197), showed resistance to at least 9 drugs, but only 3.84% (3/78) of *mcr-1*-negative isolates did ($P=0.000$). Interestingly, 68.02% (134/197) of *mcr-1*-positive strains presented resistance profiles to 10 drugs, but no COL-negative *E. coli* did ($P=0.000$).

Table 2 Distribution of MICs of 11 antibiotics for 275 MDR *Escherichia coli* isolates

Antibiotics	MIC (mg/L)											MIC ₅₀	MIC ₉₀	Resistance						
	0.008	0.016	0.032	0.063	0.125	0.25	0.5	1	2	4	8				16	32	64	128	256	>256
Ampicillin	-	-	-	-	-	-	0	1	1	1	1	6	4	12	23	38	189	>256	96.72%	
Cefotaxime	-	-	-	0	8	9	11	14	18	15	14	19	31	45	44	33	14	256	78.18%	
Cefoxitin	-	-	-	-	0	3	2	11	3	1	18	72	51	56	30	9	19	128	86.18%	
Ceftiofur	-	-	-	-	0	3	21	23	14	12	26	19	28	33	56	31	9	256	77.81%	
Chloramphenicol	-	-	-	-	-	-	0	0	2	1	7	5	17	34	35	61	113	>256	94.54%	
Ciprofloxacin	-	0	15	19	5	6	3	14	10	25	26	42	47	38	23	2	-	16	64	73.81%
Colistin	-	-	0	9	29	8	14	22	29	110	41	7	4	2	0	-	-	4	8	70.18%
Gentamycin	-	-	-	0	12	14	9	17	13	4	12	25	54	56	27	16	16	32	256	70.54%
Meropenem	0	16	158	75	10	4	3	2	3	1	0	0	0	-	-	-	0.032	0.0625	0.36%	
Polymyxin-B	-	-	-	3	12	12	15	37	41	87	49	16	2	1	0	-	-	4	8	71.27%
Tetracycline	-	-	-	-	-	-	0	3	0	3	0	1	16	41	47	69	96	256	97.81%	

Notes: Red vertical lines indicate the breakpoints between intermediate and resistant values. White areas indicate range of tested dilutions for each antibiotic; the MIC₅₀ and MIC₉₀ values are concentrations at which ≥50% and ≥90% of isolates are inhibited.

Abbreviations: MIC, minimum inhibitory concentration.

Table 3 Prevalence of *mcr-1* & or ESBL-producing *E. coli* in swine samples collected from different farms of Jiangsu China

Farm numbers	No. of positive <i>E. coli</i> samples	No. of <i>mcr-1</i> positive <i>E. coli</i> isolates (%)	No. of <i>mcr-1</i> & ESBL producing <i>E. coli</i> isolates (%)
Farm 1	32	22 (68.7)	10 (31.2)
Farm 2	11	7 (63.6)	3 (27.2)
Farm 3	10	8 (80.0)	4 (40.0)
Farm 4	26	18 (69.2)	4 (15.3)
Farm 5	14	14 (100)	2 (14.2)
Farm 6	14	13 (92.8)	2 (14.2)
Farm 7	7	5 (71.4)	N.D.
Farm 8	14	12 (85.7)	6 (42.8)
Farm 9	16	14 (87.5)	9 (56.2)
Farm 10	43	37 (86.0)	11 (25.5)
Farm 11	23	11 (47.8)	5 (21.7)
Farm 12	12	9 (75.0)	6 (50)
Farm 13	14	7 (50.0)	4 (28.5)
Farm 14	7	7 (100.)	5 (71.4)
Diseased	32	13 (40.6)	7 (21.8)
Total	275	197	78

Abbreviations: ESBL, extended-spectrum β -lactamase; N.D., not determined.

Coexistence of *mcr-1* and ESBL genes screened among the strains

Based on the results of this study, among 197 *mcr-1* positive isolates, 39.59% (78/197) were identified as carrying both *mcr-1* and ESBL genes. Distribution of all *mcr-1* positive *E. coli* isolates (n=78) harboring ESBL genes are analyzed and presented in Figure 1 and Table 5. Our findings indicate that the combination of *mcr-1* with *bla*_{CTX-M} was the most prevalent with the rate of 70.51% (55/78) followed by the combination of *mcr-1* and *bla*_{SHV} (14.10%, 11/78). Finally, 7.69% (6/78) isolates were identified carrying both *mcr-1* and *bla*_{TEM}. Furthermore, combination of *mcr-1* with two or

Table 4 The resistance patterns of *E. coli* with or without *mcr-1*

Resistance profiles	<i>mcr-1</i> positive (n=197)	<i>mcr-1</i> negative (n=78)	Chi-square (P-value)
≥ 11 drugs	0 (0)	0.0 (0)	-
≥ 10 drugs	68.02 (134)	0.0 (0)	0.000
≥ 9 drugs	68.52 (135)	3.84 (3)	0.000
≥ 8 drugs	81.21 (160)	58.97 (46)	0.000
≥ 7 drugs	89.84 (177)	76.92 (60)	0.005
≥ 6 drugs	93.90 (185)	88.46 (69)	0.125
≥ 5 drugs	99.49 (196)	91.02 (71)	0.000
≥ 4 drugs	100.0 (197)	93.58 (73)	0.000
≥ 3 drugs	100.0 (197)	96.15 (75)	0.006
≥ 2 drugs	100.0 (197)	98.71 (77)	0.111

more than two ESBL genes was also identified. Results showed that a total of 5.12% (4/78) of *mcr-1*-positive isolates also carried *bla*_{CTX-M} and *bla*_{SHV}, while 1.28% (1/78) of *mcr-1*-positive isolates carried *bla*_{CTX-M} and *bla*_{TEM}. Interestingly, a single isolate was carrying (*mcr-1*+ *bla*_{CTX-M} + *bla*_{SHV}+ *bla*_{TEM}). In this study, *mcr-1* and *bla*_{CTX-M} were identified as the dominant genes (Table 5).

As *mcr-1* in combination with *bla*_{CTX-M} were identified as the most prevalent (70.51%, 55/78), we further sequenced the *bla*_{CTX-M} genes to explore the subtypes. Sequencing analysis of these 55 *bla*_{CTX-M} isolates showed that all these *mcr-1*-positive isolates were harboring *bla*_{CTX-M-1} group. The most prevalent variants identified in these 55 isolates belonged to this group were *bla*_{CTX-M-15} in 38/55 (69%) isolates, followed by *bla*_{CTX-M-55} in 16/55 (29%) isolates and *bla*_{CTX-M-1} in one isolate (1.8%).

mcr-1 and ESBL genes could be conjugative transfer by plasmids with different replicon type

Conjugation experiments were performed on random 15 *mcr-1* positive isolates. Of the 15 *mcr-1* resistant isolates, 10 isolates were carrying additional *bla*_{CTX-M}, while a single isolate was harboring *bla*_{TEM}. Of these 15 isolates, 12 were successfully transferred to *E. coli* EC-600. The resistance profiles of the 12 transconjugants were identical to those of the *mcr-1* and *bla*_{CTX-M} carrying *E. coli* donor isolates, indicating the transfer of antibiotic resistance. In addition, resistant to several non- β -lactam antibiotics, such as aminoglycosides, fluoroquinolones, TET, macrolides and sulfonamides, were also co-transferred along with COL and β -lactam resistance. MICs of COL of these transconjugants revealed 4- to 8-fold increase as compared with the recipient EC-600 (0.125 μ g/mL).

PCR-based replicon typing (PBRT) showed that in the *E. coli* isolates carrying *mcr-1* and *bla*_{CTX-M}, the plasmids with different replicons, including IncHI2 (n=7), IncFIB (n=7), IncFIC (n=4), IncP (n=4), IncFrepB (n=4), IncN (n=3), IncX4 (n=2) IncY (n=2) and IncI1 (n=1), were detected in the donor strains (Figure 2 and Table 6). However, PBRT of the transconjugants confirmed only five replicons, IncHI2, IncFIB, IncFIC, IncN and IncX4, which were present in both donors and transconjugants and were associated with the transfer of the *mcr-1* and ESBL genes.

S1-PFGE analysis demonstrated that these six strains carried multiple plasmids varying in sizes ranging from

Table 5 Distribution of various resistance genes among 275 MDR *Escherichia coli* isolates from pigs

Colistin resistant gene	Extended-spectrum beta-lactamase genes			No. of isolates
	<i>bla</i> _{CTX}	<i>bla</i> _{SHV}	<i>bla</i> _{TEM}	
<i>mcr-1</i>				
+				119
	+			1
		+		34
			+	13
				15
+	+			55
+		+		11
+			+	6
+	+	+	+	1
+	+	+		4
+	+		+	1
	+	+	+	1
	+	+		2
	+		+	2
		+	+	1
N.D.	N.D.	N.D.	N.D.	9
197	100	33	27	275

Abbreviations: +, positive; N.D., not determined.

~97 kb to 242 kb (Figure 3A). Southern hybridization assay confirmed that the *mcr-1* gene recovered from these six strains was positioned on the following five different types of plasmids: with the size of approximately 97, 130, 160, 227 and 242 kb, respectively (Figure 3B).

Discussion

China alone produces and consumes roughly half the planet's pigs, about 500 million annually, and has been the leading consumption of antibiotics in the world.³³ The increased usage of antibiotics may trigger the emergence of AMR. Reports on emergence of AMR particularly resistance of β -lactam and COL are increasing all over the world.³⁴⁻³⁷ The prevalence of ESBL in animal origin has been rising since 2003, with slight variances amongst terrestrial regions and different animal species.^{4,14,38} We report on the high incidence of *mcr-1*-carrying ESBL-producing *E. coli* recovered from pigs in Jiangsu, China. Our results indicated that all the *mcr-1* and ESBL-producing *E. coli* isolates showed MDR. The majority of these isolates (77-86%) showed resistance to cephalosporin (Table 2). In addition, high resistance was also observed to common β -lactam and non- β -lactam antimicrobials such as AMP, fluoroquinolones, aminoglycosides, amphenicol, quinolones, sulfonamides and TET which are commonly using in human as well in veterinary practice. Many recent studies have reported MDR ESBL-producing *E. coli* isolated from poultry,³ pigs,³⁹ cattle³⁶ and humans.⁴⁰

Recently, plasmid-mediated COL-resistant genes *mcr-1* to *mcr-8* have been widely discovered around the world.^{34,41-47} Herein, we screened 275 MDR *E. coli* isolated from 14 pig farms from Jiangsu province for the presence of *mcr-1* to *mcr-5* genes. Only *mcr-1* gene was detected in isolates from every farm, and the carriage rate was extremely high in 71.6% (197/275). The high prevalence rate of the *mcr-1* found in this study




Figure 1 Distribution of various resistance genes in combination.




Figure 2 Detection of plasmid replicon types in multidrug-resistant *E. coli* using PCR assay.

from pigs in Jiangsu is consistent with very recent reports from China.^{48,49} These recent studies in pigs reported similarly high *mcr-1*-positive carriage (79.2% and 76.2%), ranging from 45% to 100% in different provinces, while the *mcr-1* rate in Jiangsu province reported by⁴⁸ was 71.9% which is very similar to our findings 71.6%. The present study and, together with the previous studies, confirmed a surprisingly high rate of *mcr-1* in swine farms and is likely associated with the prolonged and extensive practice of COL as a growth promoter in pigs.

The coexistence of *mcr-1* with other resistance genes in an *E. coli* was reported in China.⁵⁰ One recent study also assumed that a historic bridge existed between *mcr-1* and ESBL.²³ However, there is scarcity in the incidence of the coexistence of *mcr-1* and ESBL in the pig origin. Herein, we screened 275 MDR *E. coli*-resistant strains from pig farms in Jiangsu during 2016–2017 and found a high occurrence of *mcr-1*-positive strains with ESBL in the swine 39.5% (78/197), which was very high from the previous report.⁵¹ A very recent longitudinal study from China investigated the co-rising of *mcr-1* and ESBL in chicken isolates.¹⁴ Among ESBL-positive strains, we found that *bla*_{CTX-M} is the most predominant.

In this study, 39.5% *mcr-1*-positive *E. coli* strains were detected which coexist in different ESBL genes. Among them, *bla*_{CTX-M} was the predominant one which was found in 55 *mcr-1*-positive *E. coli* 70.5% (Figure 1). On

sequence-based analysis of these 55 *bla*_{CTX-M} isolates, interestingly the *bla*_{CTX-M-15} gene was found to be the most prevalent *bla*_{CTX-M} gene (69%) followed by *bla*_{CTX-M-55} (29%). The spread of the *bla*_{CTX-M-15} gene is a common *bla*_{CTX-M} enzyme and detected widely in Enterobacteriaceae of human origin.^{19,52} Very few studies have reported the co-occurrence of *mcr-1* and *bla*_{CTX-M-15} of human origin.^{22,53} From China, one recent study in dairy cows also found *bla*_{CTX-M-15} as the second prevalent ESBL gene 21.4% (62/275), but there was no *mcr-1* gene detected in those *bla*_{CTX-M-15} isolates.⁵⁴ Another study also reported the coexistence of *mcr-1* and *bla*_{CTX-M-15} in Turkey hen meat.⁵⁵ Herein, this is the first investigation that reveals the coexistence of *mcr-1* and *bla*_{CTX-M-15} in *E. coli* strains of pig origin in a very high proportion.

A significant increase in the *bla*_{CTX-M-55} was found in *E. coli* by over a period of ten years.⁴ Many previous studies reported that *bla*_{CTX-M-55} in human strains in China has become the second dominant *bla*_{CTX-M} type and even the occurrence of *bla*_{CTX-M-55} was higher than *bla*_{CTX-M-15}.⁵¹ A high rate of *mcr-1* and *bla*_{CTX-M-55} was recently detected from the chicken origin in China,¹⁴ which was consistent with our results. Thus, the concurrent dissemination of the *mcr-1* harboring *bla*_{CTX-M-15} and *bla*_{CTX-M-55} mediated by a single bacterial clone is existing which suggests that *mcr-1* is found in the diverse reservoirs.

Table 6 Conjugation experiments and plasmid replicon type detection for 15 *mcr-I*-positive *E. coli*

Strain	CTX MIC (mg/L)	COL MIC (mg/L)	Plasmid (Inc) types	Resistance genes	Resistance profile for non-beta-lactam antibiotics GEN ENR SUL CIP KEN CHL TET TRM STR POL-B
EC-9	128	16	H12,II,Y, FIC	MCR-1, CTX-M-15	GEN, ENR, NAL, SUL, CIP, KEN, CHL, TET, TRM, STR, POL-B
EC-9-T	128	4	H12, FIC	MCR-1, CTX-M-15	GEN, ENR, NAL, SUL, KEN, CHL, TET, TRM, POL-B
EC-37	256	8	N, P	MCR-1, CTX-M-55	NAL, SUL, CFX, CEF, KEN, CHL, TET, TRM, STR, POL-B
EC-37-T	32	4	N	MCR-1, CTX-M-55	NAL, SUL, CFX, CEF, KEN, CHL, TET, TRM, STR, POL-B
EC-48	256	4	FIB, P, FrepB, N	MCR-1, CTX-M-55	GEN, ENR, NAL, SUL, KEN, CHL, TET, TRM, STR, POL-B
EC-48-T	32	2	FIB, N	MCR-1, CTX-M-55	GEN, ENR, NAL, SUL, KEN, CHL, TET, TRM, STR, POL-B
EC-52	128	4	FIB,N, P	MCR-1, CTX-M-55	ENR, NAL, SUL, CEF, KEN, CHL, TET, TRM, STR, POL-B
EC-52-T	256	2	FIB,N	MCR-1, CTX-M-55	ENR, NAL, SUL, CEF, KEN, CHL, TET, TRM, STR
EC-29	256	8	H12, FIB	MCR-1, CTX-M-15	GEN, NAL, SUL, CFX, CEF, KEN, CHL, TET, TRM, STR, POL-B
EC-29-T	0.5	2	H12, FIB	MCR-1, CTX-M-15	GEN, NAL, SUL, KEN, CHL, TET, TRM, POL-B
EC-34	256	8	FIB, H12, FrepB	MCR-1, CTX-M-15	GEN, NAL, SUL, CEF, KEN, CHL, TET, TRM, STR, POL-B
EC-34-T	>256	8	H12,FIB	MCR-1, CTX-M-15	GEN, NAL, SUL, CEF, KEN, CHL, TET, TRM, POL-B
EC-40	128	4	H12, FIB, FrepB	MCR-1, CTX-M-15	ENR, NAL, SUL, CFX, CEF, CIP, KEN, CHL, TET, TRM, POL-B
EC-40-T	64	2	H12, FIB	MCR-1, CTX-M-15	ENR, NAL, SUL, CFX, CEF, KEN, CHL, TET, TRM, POL-B
EC-1	256	4	X4, FIC	MCR-1, TEM	GEN, ENR, NAL, SUL, CIP, KEN, CHL, TET, TRM, STR, POL-B
EC-1-T	256	2	X4, FIC	MCR-1, TEM	GEN, NAL, SUL, CEF, CIP, KEN, CHL, TET, TRM, STR, POL-B
EC-25F	256	2	H12,FIB, P	MCR-1, CTX-M-15	GEN, ENR, NAL, SUL, CIP, KEN, CHL, TET, TRM, STR, POL-B
EC-25F-T	>256	4	H12, FIB	MCR-1, CTX-M-15	ENR, NAL, SUL, CIP, KEN, CHL, TET, TRM, STR, POL-B
EC-55	64	8	H12,FIB, FIC,Y	MCR-1, CTX-M-55	ENR, NAL, SUL, CIP, KEN, CHL, TET, TRM, STR, POL-B
EC-55-T	64	4	H12,FIC	MCR-1, CTX-M-55	ENR, NAL, SUL, CIP, KEN, CHL, TET, TRM, POL-B, STR
EC-20	2	4	X4, FIC, FrepB	MCR-1	GEN, ENR, NAL, SUL, CIP, KEN, CHL, TET, TRM, POL-B
EC-20-T	0.25	4	X4, FIC	MCR-1	GEN, NAL, SUL, KEN, CHL, TET, TRM, STR, POL-B
EC-19	128	32	H12	MCR-1	GEN, NAL, SUL, CIP, KEN, CHL, TET, TRM, POL-B
EC-19-T	256	4	H12	MCR-1	GEN, NAL, SUL, KEN, CHL, TET, TRM, POL-B
EC-600	0.25	0.125	ND	ND	GEN, NAL, SUL, KEN, CHL, TET, TRM, POL-B

Abbreviations: COL, colistin; CTX, ceftaxime; GEN, gentamycin; ENR, enrofloxacin; NAL, nalidixic-acid; SUL, sulfamethoxazole; CIP, ciprofloxacin; KEN, kenamycin; CHL, chloramphenicol; TET, tetracycline; TRM, trimethoprim; STR, streptomycin; POL-B, polymyxin-B; ND, not determined.




Figure 3 (A) S1-nuclease pulsed-field gel electrophoresis profiles of six *E. coli* and (B) Southern Blot hybridization of six *E. coli* carrying the *mcr-1* plasmid. **Notes:** M – Salmonella H9812 (15.0–242.5 kb); 1: E-55; 2: E-01; 3: E-09; 4: E-02; 5: E-44; and 6: E-37. The arrows in the figure represent the position of the *mcr-1* plasmid.

In addition, *bla*_{CTX-M-15} and *bla*_{CTX-M-55} were previously reported on conjugative plasmids, ie, FIB, IncI1, IncHI2, IncK, IncP and IncN.^{35,56} Therefore, we also detected these incompatibility types by PCR typing. While *mcr-1* gene was often found on conjugative plasmids like IncF, IncI2, IncHI2, IncN, IncP and X4, which exhibit an unexpected diversity.⁵⁷ In conjugation experiment, IncHI2, IncFIB, IncFIC, IncN and IncX4, were found in both donors and transconjugants and were associated with the transfer of the *mcr-1* and ESBL encoding genes.

Genetic representation of *mcr-1*-carrying plasmids demonstrated that this gene is located on different conjugative elements of ~97 kb and 242 kb in size. The fact that *mcr-1*-carrying *E. coli* isolates display divergent PFGE profiles suggests that these elements may play a vital role in *mcr-1* transmission. The incidence of closely related plasmids that carry

mcr-1 and ESBL resistance genes among genotypically varied *E. coli* strains from various origins is a threat for alarm as it indicates that plasmids can easily disseminate from animals to humans and the spread of these plasmids may be remarkably challenging to control.

Considering that *bla*_{CTX-M} has become the most prevalent ESBL type of animal origin in the last few years, this situation may suggest that *mcr-1* and *bla*_{CTX-M} emerged and arose due to the extensive use of antimicrobial practice in animal farming in the last decade. Our results also suggested that *bla*_{CTX-M-15} and *bla*_{CTX-M-55} and other β -lactamase genes cohabiting with *mcr-1* positive isolates is a potential threat to public health as the pig carrying these genes may enter the food chain. It is recommended that we should pay high consideration in monitoring the incidence of ESBL-producing & COL-resistant *E. coli* in both clinical and food-producing animals.

Conclusion

Our study reported a high incidence of the *mcr-1*-carrying ESBL-producing *E. coli* recovered from pigs in Jiangsu, China. The coexistence of the *mcr-1*- and *bla*_{CTX-M-15}-carrying isolates displaying MDR, recovered from pig origin is a major concern for both humans and veterinary medicine. The presence of these genes on the conjugative plasmids with the ability to transfer between similar strains which contain other drug resistance genes emphasizes on urgent intervention.

Acknowledgments

This work was supported by the National Key Research and Development Program of China (2018YFD0500300), the National Natural Science Foundation of China (31702292, 31872517), the Natural Science Foundation of Jiangsu Province (BK20170710), the China Postdoctoral Science Foundation (2017M611841, 2018T110515) and the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD).

Disclosure

The authors report no conflicts of interest in this work.

References

- World Health Organization. *Global Action Plan On Antimicrobial Resistance*. World Health Organization; 2015. Available from: http://www.wpro.who.int/entity/drug_resistance/resources/global_action_plan_eng.pdf. Accessed July 2, 2019.
- Van Boeckel TP, Brower C, Gilbert M, et al. Global trends in antimicrobial use in food animals. *Proc Natl Acad Sci U S A*. 2015;112(18):5649–5654. doi:10.1073/pnas.1503141112
- Brower CH, Mandal S, Hayer S, et al. The prevalence of extended-spectrum beta-lactamase-producing multidrug-resistant *Escherichia coli* in poultry chickens and variation according to farming practices in Punjab, India. *Environ Health Perspect*. 2017;125(7):077015. doi:10.1289/EHP36
- Rao L, Lv L, Zeng Z, et al. Increasing prevalence of extended-spectrum cephalosporin-resistant *Escherichia coli* in food animals and the diversity of CTX-M genotypes during 2003–2012. *Vet Microbiol*. 2014;172(3–4):534–541. doi:10.1016/j.vetmic.2014.05.029
- Kong LH, Lei CW, Ma SZ, et al. Various sequence types of *Escherichia coli* isolates cohabiting bla_{NDM-5} and *mcr-1* genes from a commercial swine farm in China. *Antimicrob Agents Chemother*. 2017;61(3). doi:10.1128/AAC.02167-16
- Wang R, Liu Y, Zhang Q, et al. The prevalence of colistin resistance in *Escherichia coli* and *Klebsiella pneumoniae* isolated from food animals in China: coexistence of *mcr-1* and bla_{NDM} with low fitness cost. *Int J Antimicrob Agents*. 2018;51(5):739–744. doi:10.1016/j.ijantimicag.2018.01.027
- Jackson CR, Davis JA, Frye JG, Barrett JB, Hiott LM. Diversity of plasmids and antimicrobial resistance genes in multidrug-resistant *Escherichia coli* isolated from healthy companion animals. *Zoonoses Public Health*. 2015;62(6):479–488. doi:10.1111/zph.2015.62.issue-6
- Bush K, Jacoby GA. Updated functional classification of β -lactamases. *Antimicrob Agents Chemother*. 2010;54(3):969–976.
- Moawad AA, Hotzel H, Neubauer H, et al. Antimicrobial resistance in Enterobacteriaceae from healthy broilers in Egypt: emergence of colistin-resistant and extended-spectrum β -lactamase-producing *Escherichia coli*. *Gut Pathog*. 2018;10:39. doi:10.1186/s13099-018-0266-5
- Yamasaki S, Le TD, Vien MQ, Van Dang C, Yamamoto Y. Prevalence of extended-spectrum beta-lactamase-producing *Escherichia coli* and residual antimicrobials in the environment in Vietnam. *Anim Health Res Rev*. 2017;18(2):128–135. doi:10.1017/S1466252317000160
- Pitout JD, Nordmann P, Laupland KB, Poirel L. Emergence of Enterobacteriaceae producing extended-spectrum beta-lactamases (ESBLs) in the community. *J Antimicrob Chemother*. 2005;56(1):52–59. doi:10.1093/jac/dki166
- Cheng VC, Wong SC, Ho PL, Yuen KY. Strategic measures for the control of surging antimicrobial resistance in Hong Kong and mainland of China. *Emerg Microbes Infect*. 2015;4(2):e8. doi:10.1038/emi.2015.8
- Brolund A, Sandegren L. Characterization of ESBL disseminating plasmids. *Infect Dis (Lond)*. 2016;48(1):18–25. doi:10.3109/23744235.2015.1062536
- Feng S, Shen C, Chen H, et al. Co-production of MCR-1 and NDM-5 in *Escherichia coli* isolated from a colonization case of inpatient. *Infect Drug Resist*. 2018;11:1157–1161. doi:10.2147/IDR.S171164
- Abbas G, Khan I, Mohsin M, Sajjad Ur R, Younas T, Ali S. High rates of CTX-M group-1 extended-spectrum beta-lactamases producing *Escherichia coli* from pets and their owners in Faisalabad, Pakistan. *Infect Drug Resist*. 2019;12:571–578. doi:10.2147/IDR.S189884
- Woerther PL, Burdet C, Chachaty E, Andremont A. Trends in human fecal carriage of extended-spectrum beta-lactamases in the community: toward the globalization of CTX-M. *Clin Microbiol Rev*. 2013;26(4):744–758. doi:10.1128/CMR.00023-13
- Carattoli A. Plasmids in Gram negatives: molecular typing of resistance plasmids. *IJMM*. 2011;301(8):654–658. doi:10.1016/j.ijmm.2011.09.003
- Casella T, Nogueira MCL, Saras E, Haenni M, Madec JY. High prevalence of ESBLs in retail chicken meat despite reduced use of antimicrobials in chicken production, France. *Int J Food Microbiol*. 2017;257:271–275. doi:10.1016/j.ijfoodmicro.2017.07.005
- Day MJ, Rodriguez I, van Essen-Zandbergen A, et al. Diversity of STs, plasmids and ESBL genes among *Escherichia coli* from humans, animals and food in Germany, the Netherlands and the UK. *J Antimicrob Chemother*. 2016;71(5):1178–1182. doi:10.1093/jac/dkv485
- Dolejska M, Papagiannitsis CC. Plasmid-mediated resistance is going wild. *Plasmid*. 2018. doi:10.1016/j.plasmid.2018.09.010
- Bi Z, Berglund B, Sun Q, et al. Prevalence of the *mcr-1* colistin resistance gene in extended-spectrum beta-lactamase-producing *Escherichia coli* from human faecal samples collected in 2012 in rural villages in Shandong Province, China. *Int J Antimicrob Agents*. 2017;49(4):493–497. doi:10.1016/j.ijantimicag.2016.12.018
- Zhang H, Seward CH, Wu Z, Ye H, Feng Y. Genomic insights into the ESBL and MCR-1-producing ST648 *Escherichia coli* with multi-drug resistance. *Sci Bull*. 2016;61:875–878. doi:10.1007/s11434-016-1086-y
- Rhouma M, Letellier A. Extended-spectrum β -lactamases, carbapenemases and the *mcr-1* gene: is there a historical link? *Int J Antimicrob Agents*. 2017;49(3):269–271. doi:10.1016/j.ijantimicag.2016.11.026
- de Been M, Lanza VF, de Toro M, et al. Dissemination of cephalosporin resistance genes between *Escherichia coli* strains from farm animals and humans by specific plasmid lineages. *PLoS Genet*. 2014;10(12):e1004776. doi:10.1371/journal.pgen.1004541
- Vounba P, Rhouma M, Arsenaault J, Bada Alamedjji R, Fravallo P, Morris Fairbrother J. Prevalence of colistin resistance and *mcr-1*/*mcr-2* genes in ESBL/AmpC-producing *E. coli* isolated from chickens in Canada (Quebec), Senegal and Vietnam. *J Glob Antimicrob Resist*. 2019. doi:10.1016/j.jgar.2019.05.002

26. Tantawiwat S, Tansuphasiri U, Wongwit W, Wongchotigul V, Kitayaporn D. Development of multiplex PCR for the detection of total coliform bacteria for *Escherichia coli* and *Clostridium perfringens* in drinking water. *Southeast Asian J Trop Med Public Health*. 2005;36(1):162–169.
27. CLSI. *Performance Standards for Antimicrobial Susceptibility Testing, CLSI Supplement M100S*. Wayne (PA): Clinical and Laboratory Standards Institute. 2016.
28. Magiorakos AP, Srinivasan A, Carey RB, et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect*. 2012;18(3):268–281. doi:10.1111/j.1469-0691.2011.03570.x
29. Alba P, Leekitchaenphon P, Franco A, et al. Molecular epidemiology of mcr-encoded colistin resistance in enterobacteriaceae from food-producing animals in Italy revealed through the EU harmonized antimicrobial resistance monitoring. *Front Microbiol*. 2018;9(1217). doi:10.3389/fmicb.2018.01217
30. Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods*. 2005;63(3):219–228. doi:10.1016/j.mimet.2005.03.018
31. Barton BM, Harding GP, Zuccarelli AJ. A general method for detecting and sizing large plasmids. *Anal Biochem*. 1995;226(2):235–240. doi:10.1006/abio.1995.1220
32. Cao L, Li X, Xu Y, Shen J. Prevalence and molecular characteristics of mcr-1 colistin resistance in *Escherichia coli*: isolates of clinical infection from a Chinese University Hospital. *Infect Drug Resist*. 2018;11:1597–1603. doi:10.2147/IDR.S166726
33. Collignon P, Voss A. China, what antibiotics and what volumes are used in food production animals? *Antimicrob Resist Infect Control*. 2015;4:16
34. AbuOun M, Stubberfield EJ, Duggett NA, et al. mcr-1 and mcr-2 (mcr-6.1) variant genes identified in *Moraxella* species isolated from pigs in Great Britain from 2014 to 2015. *J Antimicrob Chemother*. 2018;73(10):2904. doi:10.1093/jac/dky272
35. Bai L, Hurlley D, Li J, et al. Characterisation of multidrug-resistant Shiga toxin-producing *Escherichia coli* cultured from pigs in China: co-occurrence of extended-spectrum beta-lactamase- and mcr-1-encoding genes on plasmids. *Int J Antimicrob Agents*. 2016;48(4):445–448. doi:10.1016/j.ijantimicag.2016.06.021
36. Ali T, Rahman SU, Zhang L, et al. Characteristics and genetic diversity of multi-drug resistant extended-spectrum beta-lactamase (ESBL)-producing *Escherichia coli* isolated from bovine mastitis. *Oncotarget*. 2017;8(52):90144–90163. doi:10.18632/oncotarget.21496
37. Nakayama T, Kumeda Y, Kawahara R, Yamaguchi T, Yamamoto Y. Carriage of colistin-resistant, extended-spectrum beta-lactamase-producing *Escherichia coli* harboring the mcr-1 resistance gene after short-term international travel to Vietnam. *Infect Drug Resist*. 2018;11:391–395. doi:10.2147/IDR.S153178
38. Kraemer JG, Pires J, Kueffer M, et al. Prevalence of extended-spectrum beta-lactamase-producing Enterobacteriaceae and Methicillin-Resistant *Staphylococcus aureus* in pig farms in Switzerland. *Sci Total Environ*. 2017;603–604:401–405
39. Liu X, Liu H, Wang L, et al. Molecular characterization of extended-spectrum β -Lactamase-Producing multidrug resistant *Escherichia coli* from swine in Northwest China. *Front Microbiol*. 2018;9:1756. doi:10.3389/fmicb.2018.01756
40. Huang IF, Lee WY, Wang JL, et al. Fecal carriage of multidrug-resistant *Escherichia coli* by community children in southern Taiwan. *BMC Gastroenterol*. 2018;18(1):86. doi:10.1186/s12876-018-0863-2
41. Liu YY, Wang Y, Walsh TR, et al. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis*. 2016;16(2):161–168. doi:10.1016/S1473-3099(16)30197-9
42. Xavier BB, Lammens C, Ruhel R, et al. Identification of a novel plasmid-mediated colistin-resistance gene, mcr-2, in *Escherichia coli*, Belgium, June 2016. *Euro Surveill*. 2016;21(27):30280.
43. Yin W, Li H, Shen Y, et al. Novel plasmid-mediated colistin resistance gene mcr-3 in *Escherichia coli*. *mBio*. 2017;8(3):pii:e00543-17. doi:10.1128/mBio.00543-17
44. Carattoli A, Villa L, Feudi C, et al. Novel plasmid-mediated colistin resistance mcr-4 gene in *Salmonella* and *Escherichia coli*, Italy 2013, Spain and Belgium, 2015 to 2016. *Euro Surveill*. 2017;22(31):pii:30589. doi:10.2807/1560-7917.ES.2017.22.31.30589
45. Borowiak M, Fischer J, Hammerl JA, Hendriksen RS, Szabo I, Malorny B. Identification of a novel transposon-associated phosphoethanolamine transferase gene, mcr-5, conferring colistin resistance in d-tartrate fermenting *Salmonella enterica* subsp. *enterica* serovar Paratyphi B. *J Antimicrob Chemother*. 2017;72(12):3317–3324. doi:10.1093/jac/dkx066
46. Yang YQ, Li YX, Lei CW, Zhang AY, Wang HN. Novel plasmid-mediated colistin resistance gene mcr-7.1 in *Klebsiella pneumoniae*. *J Antimicrob Chemother*. 2018. 73 1791–1795. doi:10.1093/jac/dky111
47. Wang X, Wang Y, Zhou Y, et al. Emergence of a novel mobile colistin resistance gene, mcr-8, in NDM-producing *Klebsiella pneumoniae*. *Emerg Microbes Infect*. 2018;7(1):122. doi:10.1038/s41426-018-0124-z
48. Zhang J, Chen L, Wang J, et al. Molecular detection of colistin resistance genes (mcr-1, mcr-2 and mcr-3) in nasal/oropharyngeal and anal/cloacal swabs from pigs and poultry. *Sci Rep*. 2018;8(1):3705. doi:10.1038/s41598-018-22084-4
49. Tong H, Liu J, Yao X, et al. High carriage rate of mcr-1 and antimicrobial resistance profiles of mcr-1-positive *Escherichia coli* isolates in swine faecal samples collected from eighteen provinces in China. *Vet Microbiol*. 2018;225:53–57. doi:10.1016/j.vetmic.2018.09.018
50. Li A, Yang Y, Miao M, et al. Complete sequences of mcr-1-harboring plasmids from extended-spectrum-beta-lactamase- and carbapenemase-producing enterobacteriaceae. *Antimicrob Agents Chemother*. 2016;60(7):4351–4354. doi:10.1128/AAC.00550-16
51. Hu YY, Cai JC, Zhou HW, et al. Molecular typing of CTX-M-producing *Escherichia coli* isolates from environmental water, swine feces, specimens from healthy humans, and human patients. *Appl Environ Microbiol*. 2013;79(19):5988–5996. doi:10.1128/AEM.01740-13
52. Fischer J, Rodriguez I, Baumann B, et al. blaCTX-M-(1)(5)-carrying *Escherichia coli* and *Salmonella* isolates from livestock and food in Germany. *J Antimicrob Chemother*. 2014;69(11):2951–2958. doi:10.1093/jac/dku270
53. Sanchez-Benito R, Iglesias MR, Quijada NM, et al. *Escherichia coli* ST167 carrying plasmid mobilisable mcr-1 and blaCTX-M-15 resistance determinants isolated from a human respiratory infection. *Int J Antimicrob Agents*. 2017;50(2):285–286. doi:10.1016/j.ijantimicag.2017.05.005
54. Zheng B, Feng C, Xu H, et al. Detection and characterization of ESBL-producing *Escherichia coli* expressing mcr-1 from dairy cows in China. *J Antimicrob Chemother*. 2018;74(2):321–325.
55. Falgenhauer L, Waezsada S-E, Gwozdinski K, et al. Chromosomal locations of mcr-1 and bla CTX-M-15 in fluoroquinolone-resistant *Escherichia coli* ST410. *Emerg Infect Dis*. 2016;22(9):1689–1691. doi:10.3201/eid2209.160692
56. Ceccarelli D, Kant A, van Essen-Zandbergen A, et al. Diversity of plasmids and genes encoding resistance to extended spectrum cephalosporins in commensal *Escherichia coli* from dutch livestock in 2007–2017. *Front Microbiol*. 2019;10:76. doi:10.3389/fmicb.2019.00076
57. Wang Q, Sun J, Li J, et al. Expanding landscapes of the diversified mcr-1-bearing plasmid reservoirs. *Microbiome*. 2017;5(1):70. doi:10.1186/s40168-017-0288-0
58. Ali T, Ur Rahman S, Zhang L, et al. ESBL-producing *Escherichia coli* from cows suffering mastitis in china contain clinical class 1 integrons with CTX-M linked to ISCR1. *Front Microbiol*. 2016;7:1931. doi:10.3389/fmicb.2016.01931

Supplementary materials



Figure S1 Map of sampling sites in Jiangsu province.




Figure S2 Screening of *mcr-1* and ESBL encoding genes in *E. coli*.

Notes: PCR product was separated on 1% agarose gel. Lane 1 and 6 shows 2000 bp molecular marker (Vazyme, Beijing, China); Lane 2, (*mcr-1*), Lane 3 (*blaCTX-M*) isolate; Lane 4, (*blaTEM*); and Lane 5 shows (*blaSHV*).

Table S1 List of primers used for plasmid replicons typing in this study

Primers	Sequence (5' to 3')	Target sites/ genes	Annealing temperature	Amplicons size	References
PBRT* primers					
H11-F H11-R	GGAGCGATGGATTACTTCAGTAC TGCCGTTTCACCTCGTGAGTA	parA-parB	58 °C	471-bp	1
H12 -F H12- R	GGCTCACTACCGTTGTCATCCT CGAAAGCCGGACGGCAGAA	RNAI	58 °C	644-bp	1
11-F 11-F	CGAAAGCCGGACGGCAGAA TCGTCGTTCCGCCAAGTTCGT	iterons	58 °C	139-bp	1
X -F X -R	AACCTTAGAGGCTATTTAAGTTGCTGAT TGAGAGTCAATTTTATCTCATGTTTTAGC	ori □	58 °C	376-bp	1
L/M-F L/M-F	GGATGAAAATCATCAGCATCTGAAG CTGCAGGGGCGATTCTTTAGG	repA, B, C	58 °C	785-bp	1
N-F N-R	GTCTAACGAGCTTACCGAAG GTTTCAACTCTGCCAAGTTC	repA	58 °C	559-bp	1
FIA- F FIA- R	CCATGCTGGTTCTAGAGAAGGTG GTATATCCTTACTGGCTTCCGCAG	iterons	58 °C	462-bp	1
FIB -F FIB -R	GGAGTTCTGACACACGATTTTCTG CTCCCGTCGCTTCAGGGCATT	repA	58 °C	702-bp	1
W-F W-R	CCTAAGAACAACAAAGCCCCCG GGTGCGCGGCATAGAACCGT	repA	58 °C	242-bp	1
Y-F Y-R	AATTCAAACAACACTGTGCAGCCTG GCGAGAATGGACGATTACAAAACCTT	repA	58 °C	765-bp	1
P-F P-R	CTATGGCCCTGCAAACGCGCCAGAAA TCACGCGCCAGGGCGCAGCC	iterons	58 °C	534-bp	1
FIC -F FIC -R	GTGAACTGGCAGATGAGGAAGG TTCTCCTCGTCGCCAAACTAGAT	repA2	58 °C	262-bp	1
A/C -F A/C -R	GAGAACCAAAGACAAAGACCTGGA ACGACAAACCTGAATTGCCTCCTT	repA	58 °C	465-bp	1
T-F T-R	TTGGCCTGTTTGTGCCTAAACCAT CGTTGATTACACTTAGCTTTGGAC	repA	58 °C	750-bp	1
FIIS -F FIIS -R	CTGTCGTAAGCTGATGGC CTCTGCCACAACTTCAGC	repA	58 °C	270-bp	1
FrepB-F FrepB-R	TGATCGTTTAAGGAATTTTG GAAGATCAGTCACACCATCC	RNAI/repA	58 °C	270-bp	1
K/B -F K/B-R	GCGGTCCGGAAAGCCAGAAAAC TCTTTCACGAGCCCCGCCAAA	RNAI	58 °C	160 bp	1

(Continued)

Table S1 (Continued).

Primers	Sequence (5' to 3')	Target sites/ genes	Annealing temperature	Amplicons size	References
B/O-F B/O-R	GCGGTCCGAAAGCCAGAAAAC TCTGCGTTCCGCCAAGTTCGA	RNAI	58 °C	159 bp	¹
X4- F X4-R	AGCAAACAGGGAAAGGAGAAGACT TACCCCAAATCGTAACCTG	-	62 °C	569 bp	²
Incl2-F Incl2-R	ATTGTTGCGTGGCTTCAT TGGAGAGGATTAAGGAGGAA	-	60 °C	353 bp	This study

References

1. Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods*. 2005;63(3):219–228.
2. Johnson TJ, Bielak EM, Fortini D, et al. Expansion of the IncX plasmid family for improved identification and typing of novel plasmids in drug-resistant Enterobacteriaceae. *Plasmid*. 2012;68(1):43–50.

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>