

eIF2 α promotes vascular remodeling via autophagy in monocrotaline-induced pulmonary arterial hypertension rats

This article was published in the following Dove Press journal:
Drug Design, Development and Therapy

Linya Guo,^{1,*} Yanbing Li,^{2,3,*}
Ying Tian,^{4,5,*} Shaoxin Gong,⁶
Xi Chen,¹ Tianhong Peng,¹
Aiping Wang,^{1,4,5,7}
Zhisheng Jiang⁷

¹Clinical Anatomy & Reproductive Medicine Application Institute, School of Medicine, University of South China, Hengyang 421001, People's Republic of China;

²National Key Discipline of Human Anatomy, Southern Medical University, Guangzhou 510000, Guangdong, People's Republic of China; ³Guangdong Engineering Research Center for Translation of Medical 3D Printing Application, Guangzhou, 510000, Guangdong, People's Republic of China; ⁴Institute of Clinical Research, Affiliated Nanhua Hospital, University of South China, Hengyang 421002, Hunan, People's Republic of China; ⁵Postdoctoral Research Institute on Basic Medicine, University of South China, Hengyang, 421001, Hunan, People's Republic of China; ⁶Department of Pathology, First Affiliated Hospital, University of South China, Hengyang 421001, Hunan, People's Republic of China; ⁷Key Laboratory for Arteriosclerosis of Hunan Province, Institute of Cardiovascular Disease, University of South China, Hengyang 421001, Hunan, People's Republic of China

*These authors contributed equally to this work

Correspondence: Aiping Wang
Institute of Clinical Research, Affiliated Nanhua Hospital, University of South China, Hengyang 421001, Hunan, People's Republic of China
Email waiping2011@163.com

Zhisheng Jiang
Key Laboratory for Arteriosclerosis of Hunan Province, Institute of Cardiovascular Disease, University of South China, Hengyang 421001, Hunan, People's Republic of China
Email zsjiang2005@163.com

Purpose: Eukaryotic initiation factor 2 α (eIF2 α) plays important roles in the proliferation and survival of pulmonary artery smooth muscle cells (PASMCs) in animal hypoxia-induced pulmonary hypertension models. However, the underlying mechanism remains unknown at large. Autophagy has been reported to play a key role in the vascular remodeling in pulmonary arterial hypertension (PAH). The purposes of this study are to determine the functions of eIF2 α and autophagy in the vascular remodeling of the monocrotaline-induced PAH rats and to clarify the correlation between eIF2 α and autophagy.

Methods: We established a rat model of monocrotaline-induced PAH, and we established a cell model of platelet derived growth factor (PDGF)-induced PASMCs proliferation. The vascular morphology and the expression of eIF2 α , LC3B, and p62 were assessed in the pulmonary arterial tissue of Sprague-Dawley rats and PDGF-induced PASMCs.

Results: Autophagy was significantly active in monocrotaline model group (MCT)-induced PAH rats, which obviously promotes vascular remodeling in MCT-induced PAH rats. Furthermore, the proliferation of PASMCs was induced by PDGF in vitro. The expression of LC3B, eIF2 α was increased in the PDGF-induced PASMCs proliferation, and the expression of p62 was reduced in the PDGF-induced PASMCs proliferation. Moreover, eIF2 α siRNA downregulated the expression of eIF2 α and LC3B, and upregulated the expression of p62 in PDGF-induced PASMCs proliferation. eIF2 α siRNA inhibited the PDGF-induced PASMCs proliferation. Finally, chloroquine can upregulate the protein expression of LC3B and p62, it also can inhibit proliferation in PDGF-induced PASMCs.

Conclusion: Based on these observations, we conclude that eIF2 α promotes the proliferation of PASMCs and vascular remodeling in monocrotaline-induced PAH rats through accelerating autophagy pathway.

Keywords: eIF2 α , autophagy, PASMCs, PAH, monocrotaline

Introduction

Pulmonary arterial hypertension (PAH) is an extremely malignant disease of the cardiovascular system. Its main clinical features are increased pulmonary vascular resistance and pulmonary artery pressure, which ultimately leads to right heart failure and even death.¹⁻³ The pathophysiological mechanism of PAH is very complicated, and pulmonary vascular remodeling is the main pathological change in PAH. The influencing factors of pulmonary vascular remodeling involve a variety of cells, importantly, the abnormal proliferation of pulmonary artery smooth muscle cells (PASMCs) is the most important cause of pulmonary vascular remodeling in PAH.^{4,5} However, the underlying mechanism remains unknown. Therefore, to further study the

pathophysiological mechanism of PAH and find new drugs to effectively treat PAH have become a top priority.

Recent studies have shown that translation initiation factors play a key role in regulating certain functional proteins to promote cell proliferation.⁶ However, eukaryotic initiation factor 2 α (eIF2 α) plays a key role in regulating cell proliferation and hypertrophy as the regulatory subunit of the translation initiation factor family, and participated in the regulation of smooth muscle cell proliferation and migration.^{7–10} Our previous study also confirmed that the expression of p-eIF2 α and eIF2 α was upregulated in both pulmonary artery tissue of HPH rats and hypoxic-treated PSMCs proliferation.¹¹ Meanwhile, after transfected with eIF2 α siRNA in hypoxia-promoted PSMCs proliferation, it can significantly inhibit the proliferation of PSMCs.¹¹ These results reveal for the first time the role of eIF2 α as a new pro-proliferating protein in pulmonary vascular remodeling in HPH. However, the underlying mechanism remains unknown.

Autophagy is an efficient subcellular degradation pathway of lysosomal-dependent in eukaryotic cells, which plays an important role in scavenging waste, cell growth, maintaining protein metabolism balance, and stabilization of intracellular environment.^{12–14} Abnormal autophagy breaks the balance of the body and leads to disease development in some pathological environments.¹⁵ Recent studies have shown that autophagy plays a vital role in various human diseases.^{16–19} More and more studies have reported that autophagy plays an important role in cardiovascular diseases, and recent studies have found that activation of autophagy promotes the development of monocrotaline-induced PAH; moreover, inhibition of autophagy can inhibit the development of monocrotaline-induced PAH.^{20,21} Furthermore, studies have shown that eIF2 α can activate endoplasmic reticulum autophagy after oxidative damage in the endoplasmic reticulum.²² Based on the above studies, we speculated that eIF2 α could regulate pulmonary vascular remodeling in monocrotaline-induced PAH via autophagy pathway. In this study, we will reveal a key role of eIF2 α in monocrotaline-induced PAH.

Materials and methods

Animals

Male Sprague-Dawley (SD) rats (weighing 150–200 g) were obtained from the Laboratory Animal Center, School of Medicine, University of South China (Hengyang, People's Republic of China). All

surviving animals were handled in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals. The experimental protocol was approved by the medicine animal welfare committee of Medicine School, University of South China (Hengyang, People's Republic of China).

Animals experiments

Animals experiments each SD rat is 150–200 g/head. After 1 week of adaptive feeding, each rat was weighed and labeled; rats were randomly divided into three groups: 1) control group (Control); 2) vehicle group (Vehicle), each rat was intraperitoneally injected with the same dose of vehicle; 3) monocrotaline model group (MCT), each rat was intraperitoneally injected with monocrotaline (60 mg/kg); pulmonary hypertension model was established for 4 weeks. Rats in each group were weighed and treated separately. Rats were intraperitoneally injected with chloral hydrate (30 mg/kg). Right ventricular systolic pressure (RVSP) and mean pulmonary arterial pressure (mPAP) were monitored for each rat. Subsequently, the right ventricle and left ventricle (RV, LV) and interventricular septum (S) of the rats were dissected separately. The length of the tibia was measured and weighed to calculate the ratio of RV to (LV+S) and tibia length to RV, which is a key indicator for assessing RV hypertrophy. Pulmonary tissue was isolated for analysis of mRNA and protein expression.

HE staining

The isolated rat lung tissue was stained with HE to assess the degree of pulmonary vascular remodeling. After paraffin sectioning, the sections are placed in distilled water and then placed in an aqueous solution of hematoxylin for staining for about 10 mins; then, the slices were placed into ammonia and acid water for several seconds; after the slice was rinsed in running water for 1 hr, it was placed in distilled water for several seconds; then, the sections were dehydrated in alcohol at concentrations of 90% and 70% for 10 mins each; subsequently, the sections were stained with eosin staining solution for about 2–3 mins; after sectioning, it is dehydrated with 100% alcohol and then placed in xylene; the sections were sealed and placed in an incubator for drying; finally, the sections were photographed with a microscope to observe the morphology of the blood vessels. Pulmonary arterial wall thickness (WT) was calculated by the following formula: $WT (\%) = \frac{\text{area}_{\text{ext}} - \text{area}_{\text{int}}}{\text{area}_{\text{ext}}} \times 100$, where area_{ext} and area_{int} are the area bounded by external and internal elastic lamina, respectively.

Western blot analysis

Protein concentration was measured after protein extraction from pulmonary artery tissue and primary cultured PSMCs. Extracted proteins were detected by Western blot analysis. Primary antibody was purchased from Cell Signaling Technology (CST), USA. Optical density analysis was performed using Image J 1.43 (National Institutes of Health). Primary antibodies against eIF2 α (1:1000 dilution, CST, Danvers, MA, USA, #5324s) were purchased from CST. Primary antibodies against p-eIF2 α (1:1000 dilution, CST, #3398s) were purchased from CST. Primary antibodies against LC3B (1:1000 dilution, CST, #12741s) were obtained from CST. Primary antibodies against p62 (1:1000 dilution, CST, #88588) were purchased from CST. Primary antibody against β -actin (BM0627, 1:250) was purchased from Boster. Goat anti-rabbit IgG secondary antibody was purchased from Beyotime (A0408, 1:2000 dilution).

Real-time PCR analysis

RNA concentration was measured by spectrophotometer. Total RNA was extracted by TRIzol. Reverse transcription reaction using PrimeScriptTM RT reagent (TaKaRa, Shiga, Japan). Fluorescence quantification using SYBR[®] PremixEx TaqTM II kit (TaKaRa, Shiga, Japan). Real-time PCR amplification using the ABI 7500 Real-Time PCR System (Applied Biosciences, Foster City, CA, USA). The data were quantified using the $\Delta\Delta$ CT method, and the β -actin gene was used as a reference.

Cell culture and experiments

SD rats were anesthetized with 10% chloral hydrate (0.3 mg/kg) by intraperitoneal injection. The chest and abdomen hair of rat were removed and disinfected, and then the chest of rat was dissected immediately on the ultra-clean operating table to take the pulmonary artery tissue. The endovascular cell and adventitia of the pulmonary artery were removed. The pulmonary artery tissue was cut into 1 \times 1 mm³ with ophthalmic scissor and inoculated in containing 20% FBS- DMEM high glycosylation. The third generation of cells was used for subsequent experiments after the PSMCs growing to 70–80%. After the cells were grown to 70–80% confluence, the medium was replaced with 10% FBS, serum was starved for 24 hrs, and the cells were synchronized to G0 phase. Platelet-derived growth factor-BB (PDGF-BB, R&D Systems, USA) (10 ng/mL) stimulated PSMCs at different times

(6, 12, 24, 48, 72 hrs), cells were collected at the peak of cell viability; to explore autophagy in the PDGF-induced PSMCs proliferation, we designed the following experiment: PSMCs (passage 2–3) were plated in 24-well plates grown to subconfluence, then quiesced in serum-free medium for 24 hrs before chloroquine (CLQ, Sigma, USA) (10 μ M) treatment in 10% FCS for 24 hrs. The cultured cells were preincubated with autophagy inhibitor chloroquine for 2 hrs, and PDGF-BB (10 ng/mL) was stimulated for 48 hrs.

Proliferation assay

Cell Counting Kit-8 (CCK-8, Yiyuan Biotechnologies, Guangzhou, People's Republic of China) specific steps are according to the manufacturer's instructions. Cell suspension (100 μ L) was cultured in a 96-well plate, and the plate was pre-incubated at 37°C and 5% CO₂ condition for 24 hrs. CCK-8 (10 μ L) solution was added to the plate for 1–4 hrs. Finally, the OD value was read at the absorbance of 450 nm.

Small interference RNA transfection

The eIF2 α siRNA were purchased from Ribobio (Guangzhou, People's Republic of China). The eIF2 α siRNA segment was transfected by transient transfection. The cells were planted in 24-well plates, the transfection was treated when the confluence of PSMCs grown at about 30%. Subsequently, according to the manufacturer's instructions, the irrelevant fragment transfection complex and the eIF2 α siRNA transfection complex were, respectively, transfected into PDGF-induced PSMCs according to manufacturer's instructions. After transfection, PSMCs were cultured with a concentration of 5% CO₂ and a temperature of 37°C for 4–6 hrs. Then exchange for fresh medium (20% fetal bovine serum), continue to culture cells for 48 hrs in PDGF treatment condition. The specific experiments were divided into four groups: control group (Control), PDGF group (PDGF), PDGF+negative group (+Ng), PDGF+siRNA group (+50 nM).

Statistical analysis

All data were statistically analyzed and the data were expressed as mean \pm standard error ($\bar{x}\pm$ s.e.m). The q-PCR results were processed by ABI 7500 real-time PCR instrument; the Western blot data were analyzed by Image J software. All the data were statistically analyzed by SPSS 20.0 statistical software. The bar graph was made by Graphpad Prism 6.0. Statistical analysis was performed

by unpaired Student's *t*-test for two groups or analysis of variance followed by Newman-Student-Keuls test for multiple groups. Bilateral $P < 0.05$ was considered statistically significant.

Results

Vascular remodeling in MCT-induced PAH rats

In order to clarify whether the establishment of the MCT-induced PAH rat model was successful, we performed hemodynamic analysis on SD rats in each group. The mPAP, RVSP, RV/(LV+S), and right ventricular/tibia length ratios were significantly increased in the MCT model group compared with the control group (Figure 1A-D). There was no change in the vehicle group compared with that of the control group. HE staining results showed that the pulmonary arteries of the lung tissue in the MCT-induced PAH rats showed significant thickening (Figure 2A-C) and the percentage of the medial thickness of the pulmonary arterioles (WT%) was significantly increased (Figure 2D), showing obvious characteristics of pulmonary arteriole remodeling. The above results

indicated that the MCT-induced PAH model in SD rats was successfully constructed.

The mRNA expression of Ki-67 and PCNA in MCT-induced PAH rats

The mRNA expression of proliferation markers Ki-67 and PCNA in pulmonary arteries tissue was detected by real-time PCR. The results showed that the mRNA expression levels of Ki-67 and PCNA were significantly increased in the pulmonary arteries of MCT-induced PAH rats (Figure 2E, F) which was significantly different from that of control group ($P < 0.05$). The results suggest that PASMCs significantly proliferate in MCT-induced PAH rats.

The expression of p-eIF2 α , eIF2 α , LC3B, and p62 in MCT-induced PAH rats

The mRNA expression of eIF2 α in pulmonary arteries was detected by real-time PCR, the results showed that the mRNA level of eIF2 α in the pulmonary arteries tissue of MCT-induced PAH rats was significantly upregulated (Figure 3A). To examine whether eIF2 α and autophagy were activated in MCT-induced PAH rats, the protein levels of eIF2 α , p-eIF2 α ,




Figure 1 Changes in hemodynamic parameters. (A) Mean pulmonary artery pressure (mPAP). (B) Right ventricular systolic pressure (RVSP). (C) Right ventricle/(left ventricle+ventricular septum) [RV/(LV+S)] (D) RV/tibia length. The data are expressed as mean \pm standard error; $n = 8$, $*P < 0.05$ vs Control.

Abbreviations: Control, control group; Vehicle, vehicle group; MCT, monocrotaline group.




Figure 2 Pulmonary vascular remodeling was observed by HE staining. (A–C) monocrotaline-induced pulmonary vascular remodeling was observed by HE staining: (A–C) monocrotaline-induced pulmonary artery remodeling in PAH rats (200 \times), scale bars=100 μ m. (D) Thickness of vascular media (WT=WT (%)= $\frac{\text{area}_{\text{ext}}-\text{area}_{\text{int}}}{\text{area}_{\text{ext}}}\times 100$, where area_{ext} and area_{int} are the area bounded by external and internal elastic lamina, respectively). (E) Ki-67 mRNA expression. (F) PCNA mRNA expression. The data are expressed as mean \pm standard error, n=8, * P <0.05 vs Control.

Abbreviations: Ki-67, nuclear protein, marker of cell proliferation; PCNA, cell proliferating nuclear antigen; Control, control group; Vehicle, vehicle group; MCT, monocrotaline group.

LC3B, and p62 were examined by Western blotting. In MCT-induced PAH rats, the levels of p-eIF2 α , eIF2 α , and LC3B protein were increased in the pulmonary arteries compared with that of control group (Figure 3B), which was significantly different from the control group (P <0.05); however, the protein

expression of p62 was reduced in the pulmonary arteries tissue of the MCT-induced PAH rats (Figure 3B). Statistical analysis of protein gray value scans showed statistical significance (Figure 3C–E); these results suggest that eIF2 α and autophagy both are activated in MCT-induced PAH rats.




Figure 3 Expression changes of p-eIF2 α , eIF2 α , and LC3B, P62 in pulmonary arterial tissue of MCT-induced PAH rats. (A) Expression of eIF2 α mRNA in pulmonary artery tissue. (B) Protein expression of p-eIF2 α , eIF2 α , LC3B, and p62 in pulmonary artery tissue in MCT-induced PAH rats. (C) The ratio of p-eIF2 α /total eIF2 α . (D) The ratio of LC3B-II/LC3B-I. (E) Statistical analysis of p62 gray value. The data are expressed as mean \pm standard error, n=8, * P <0.05 vs Control.

PDGF-induced PSMCs proliferation

PASMCs were treated with PDGF for stimulating proliferation, cells proliferation was tested by CCK-8. As depicted in **Figure 4A**, PDGF-induced PASMCs proliferation successfully, which was significantly different from the control group ($P<0.05$). The mRNA expression of Ki-67 and PCNA in PDGF-induced PASMCs was detected by real-time PCR, the results showed that the mRNA expression levels of Ki-67 and PCNA were significantly increased in the PDGF-induced PASMCs proliferation (**Figure 4B, 4C**); the result indicates that PDGF successfully induces the proliferation of PASMCs.

Effect of eIF2 α siRNA on PDGF-induced PASMCs proliferation

To confirm the role of eIF2 α in PDGF-induced PASMCs proliferation, we transfected eIF2 α siRNA (50 μ M) into PDGF-induced PASMCs to observe whether eIF2 α siRNA can inhibit the proliferation of PASMCs. The protein expression of eIF2 α after transfection with eIF2 α siRNA was detected by Western blotting. As depicted in **Figure 5A**, the protein expression of eIF2 α and p-eIF2 α were upregulated in the PDGF group, but eIF2 α siRNA significantly downregulated the protein expression of eIF2 α and p-eIF2 α compared with that of PDGF-induced PASMCs group (**Figure 5B**, $P<0.05$). The effect of eIF2 α siRNA on the proliferation of PASMCs was analyzed by CCK-8 to observe whether eIF2 α siRNA can inhibit the proliferation of PDGF-induced PASMCs. As shown in **Figure 6A**, PDGF obviously induced the PASMCs proliferation, whereas eIF2 α siRNA markedly inhibited PDGF-induced PASMC proliferation in vitro. Real-time PCR results show that the mRNA expression of Ki67 and PCNA was inhibited after transfection with eIF2 α siRNA in PDGF-induced PASMCs (**Figure 6B, C**), which was significantly different

from the PDGF group ($P<0.05$). The above results indicate that eIF2 α siRNA inhibits the expression of eIF2 α and the PDGF-induced PASMCs proliferation.

Effect of chloroquine on PDGF-induced PASMCs proliferation

To confirm that eIF2 α siRNA can inhibit autophagy activation, the protein expression of LC3B and p62 after transfection with eIF2 α siRNA was detected by Western blotting. As depicted in **Figure 5A**, the protein expression of LC3B was upregulated and the protein expression of p62 was downregulated, but eIF2 α siRNA significantly downregulated the protein expression of LC3B and upregulated the protein expression of p62 compared with that of PDGF-group (**Figure 5C, 5D**, $P<0.05$). To confirm the role of autophagy in PDGF-induced PASMCs proliferation, PASMCs were pretreated with chloroquine, a specific autophagy inhibitor, and then PDGF-induced proliferation. First, the protein expression of LC3B and p62 was detected by Western blotting after treatment with chloroquine in the PDGF-induced PASMCs proliferation. As depicted in **Figure 7A**, chloroquine significantly upregulated the protein expression of LC3B and p62 in CLQ-group compared with that of PDGF-group (chloroquine is an autophagic lysosome inhibitor) (**Figure 7B, C**, $P<0.05$). We examined the effects of chloroquine on the PDGF-induced PASMCs proliferation by using CCK-8. As shown in **Figure 7D**, chloroquine extremely inhibited the proliferation of PDGF-induced PASMCs. Real-time PCR results showed that the mRNA expression of Ki67 and PCNA was significantly inhibited after treatment with chloroquine (**Figure 7E, F**). These results indicate that chloroquine can inhibit PDGF-induced PASMCs proliferation. In conclusion, autophagy plays an important role in the pulmonary vascular remodeling of MCT-induced PAH,




Figure 4 PDGF-induced PASMCs proliferation. **(A)** CCK-8 detected proliferation. **(B)** Ki-67 mRNA expression. **(C)** PCNA mRNA expression. Ki-67: nuclear protein, marker of cell proliferation. The data are expressed as mean \pm standard error, $n=3$, $*P<0.05$ vs Control.

Abbreviations: PCNA, cell proliferating nuclear antigen; Control, control group; Vehicle, vehicle group; PDGF, PDGF group.




Figure 5 The expression changes of eIF2 α , p-eIF2 α , LC3B, and p62 in the PDGF-induced PSMCs proliferation and the effect of eIF2 α siRNA on autophagy. **(A)** Expression of eIF2 α , p-eIF2 α , LC3B, and p62 proteins in the different groups. **(B)** Statistical analysis of gray values of different groups of total eIF2 α . **(C)** The ratio of LC3B-II/LC3B-I. **(D)** Analysis of gray value of p62. Control: control group. PDGF: cell proliferation group. +Ng: PDGF+negative group. +eIF2 α siRNA (50 nM): after transfected with eIF2 α siRNA (50 nM) for 4–6 hrs, the medium was replaced and PDGF was induced for 48 hrs. The data are expressed as mean \pm standard error, n=3, *P<0.05 vs Control. #P<0.05 vs PDGF.




Figure 6 Effect of eIF2 α siRNA on proliferation of PDGF-induced PSMCs. **(A)** Effect of eIF2 α siRNA on PDGF-induced PSMCs proliferation. **(B)** Effect of eIF2 α siRNA on Ki-67 mRNA expression. **(C)** Effect of eIF2 α siRNA on PCNA mRNA expression. +Ng: PDGF+negative group. +eIF2 α siRNA (50 nM): after transfected with eIF2 α siRNA (50 nM) for 4–6 hrs, the medium was replaced and treated with PDGF for 48 hrs. The data are expressed as mean \pm standard error; n=3, *P<0.05 vs Control. #P<0.05 vs PDGF.

Abbreviations: Control, control group; PDGF, cell proliferation group.

importantly, eIF2 α plays a key role via autophagy pathway in MCT-induced PAH.

Discussion

The primary findings of the study were as follows: 1) pulmonary arterial remodeling was observed in rats treated with monocrotaline, which also exhibited elevated the protein expression of eIF2 α , p-eIF2 α , LC3B and reduced the protein expression of p62; 2) eIF2 α plays an important role via autophagy pathway in the pulmonary vascular

remodeling in MCT-induced PAH; and 3) hloroquine attenuated PDGF-induced PSMC proliferation and expression of autophagy. Autophagy plays an important role in the pulmonary vascular remodeling in MCT-induced PAH.

Increased pulmonary vascular resistance and pulmonary vascular remodeling are two basic pathological features of PAH.^{23–25} In this study, we have used monocrotaline to induce pulmonary vascular remodeling of PAH rats. Studies have proved that PSMCs play a crucial role in the pulmonary vasculature of PAH.^{26–28} In our study, primary




Figure 7 Effect of chloroquine on PDGF-induced PSMCs proliferation. **(A)** The proteins expression of LC3 and p62 in different groups. **(B)** The ratio of LC3B-II/LC3B-I. **(C)** Statistical analysis of gray value of p62. **(D)** Effect of chloroquine on the proliferation of PDGF-induced PSMCs. **(E)** The effect of chloroquine on Ki-67 mRNA expression. **(F)** The effect of chloroquine on PCNA mRNA expression (10 nM) for 2 hrs, then was replaced the culture media and treated it with PDGF for 48 hrs. The data are expressed as mean±standard error; n=3, * $P<0.05$ vs Control. # $P<0.05$ vs PDGF.

Abbreviations: Control, control group; PDGF, cell proliferation group; +Vehicle, PDGF+Vehicle group; +CLQ, cells treated with chloroquine.

PASMCs are treated with PDGF to induce the proliferation. The above results showed the animal models and cell models were successfully established.

eIF2 α is a regulatory subunit of eukaryotic translation initiation factor 2, which is a key protein that catalyzes the initiation of protein synthesis.^{29,30} eIF2 α , as an important member of the translation initiation factor family, plays a key role in the regulation of cell proliferation, in addition, it is involved in the regulation of smooth muscle cell proliferation and migration.⁷⁻¹⁰ Our previous studies also confirmed that the expression of p-eIF2 α and eIF2 α was upregulated in both HPH rats in pulmonary arterial tissue and hypoxic-treated PASMCs proliferation, and all of these effects were inhibited by eIF2 α siRNA.¹¹ But the underlying mechanism remains unknown. In our study, we established monocrotaline-induced PAH rats and PDGF-induced PASMCs proliferation. Our study has found the following results: 1) both p-eIF2 α and eIF2 α expression were upregulated in both PDGF-induced PASMCs proliferation and MCT-induced PAH rats; 2) the mRNA expression of Ki-67 and PCNA in pulmonary arterial tissue was significantly increased in monocrotaline-induced PAH rats and PDGF-induced PASMCs proliferation; 3) eIF2 α expression was significantly upregulated, and the above effects were inhibited by eIF2 α siRNA in PDGF-induced PASMCs proliferation; and 4) eIF2 α siRNA significantly inhibited the proliferation of the PDGF-induced PASMCs.

These results reveal the important role of eIF2 α as a new pro-proliferating protein in pulmonary vascular remodeling in MCT-induced PAH. These results indicate that eIF2 α can be used as a new target for the prevention and treatment of PAH.

The expression of eIF2 α is elevated in monocrotaline-induced PAH rats and PDGF-induced PASMCs proliferation, but what kind of downstream mechanism does eIF2 α play role by regulating? This is one of the main issues to be explored in this experiment. Autophagosomes bind to lysosomes to form autophagolysosome during autophagy, which can degrade damaged macromolecules and organelles in the cytoplasm. It can provide raw materials for the normal survival and metabolism of cells by degrading products.^{31,32} Many research show that autophagy is closely related to the occurrence and development of lung diseases.^{21,33} Recently, some studies have shown that autophagy is involved in the development of PAH induced by monocrotaline.^{21,33} Studies have shown that eIF2 α can active endoplasmic reticulum autophagy after oxidative damage in the endoplasmic reticulum. Therefore, we speculate that the proliferative effect of eIF2 α in MCT-induced PAH may be related to the activation of autophagy. We also found the following results: 1) the expression of LC3B was significantly upregulated in both PDGF-induced PASMCs proliferation and MCT-induced PAH rats, and the expression of p62 was down-regulated, consistent with the activation of




Figure 8 The proposed pathway of eIF2 α mediation contributes to pulmonary vascular remodeling. eIF2 α can promote PASMCS proliferation and pulmonary vascular remodeling by activating autophagy in MCT-induced PAH rats and PDGF-induced PASMCS proliferation; and eIF2 α siRNA can inhibit autophagy activation and inhibit the proliferation of PASMCS; chloroquine inhibits the proliferation of PASMCS by inhibiting autophagy.

autophagy; and (2) eIF2 α siRNA significantly inhibited the expression of LC3B and increased the expression of p62 in PDGF-induced PASMCS proliferation. To further confirm the effect of autophagy on the proliferation of PASMCS, next we use chloroquine to inhibit autophagy pathways in PDGF-induced PASMCS. The expression of LC3B protein was up-regulated in PDGF-induced PASMCS. Chloroquine treatment further increased LC3B expression in these cells consistent (Chloroquine is an autophagic lysosome inhibitor). We examined the effects of chloroquine on p62. The results show that the expression of p62 was markedly reduced in PDGF-induced PASMCS, but chloroquine treatment up-regulated p62 expression in PDGF-induced PASMCS, and treatment with chloroquine profoundly inhibited the proliferation of PDGF-induced PASMCS. These results show that autophagy is involved in the development of PAH and can be inhibited by eIF2 α siRNA.

Taken together, in this experimental study, we found that the eIF2 α expression was up-regulated both in the rat PAH model induced by monocrotaline and the PASMCS induced by PDGF. The high expression of eIF2 α leads to the activation of autophagy, which in turn promotes the proliferation of PASMCS, and leads to the pulmonary vascular remodeling in PAH. We further found that the pathophysiological process that the important role of eIF2 α promoting PASMCS proliferation and activating of autophagy can be blocked by eIF2 α siRNA. At the same time, chloroquine can inhibit the proliferation of PASMCS by inhibiting autophagy. These results indicate that eIF2 α is an important proliferation protein in the pulmonary

vascular remodeling in PAH. As a kind of protein closely related to protein translation, people's understanding of eIF2 α tends to be deeper. The relationship between eIF2 α and tumor provides new ideas for clinical tumor treatment. At present, there are some exploratory studies using eIF2 α as a therapeutic target.^{34,35} This suggests indicate that eIF2 α may be a very promising target for clinical treatment. Although there are still some problems to be solved so far, some existing research results have pointed out the direction for future research and provided new ideas for the treatment of PAH. We have reason to believe that eIF2 α protein can be used as a new target for the prevention and treatment of PAH in the further research, and blocking the function of eIF2 α may reduce and delay the development of PAH and improve the therapeutic effect of PAH. These research will provide a key regulatory molecule and a new target for the exploration of PAH.

In summary, our research found that eIF2 α promotes the proliferation of PASMCS and vascular remodeling in monocrotaline-induced PAH rats through accelerating autophagy pathway (Figure 8).

Acknowledgment

This project was financed by the National Natural Science Foundation of China (grant number 81600040 to Wang A; 31871169 to Tian Y), the Province Natural Science Foundation of Hu Nan (grant number 2017JJ3279 to Wang A), the Postdoctoral Science Fund of China (grant number 2017M622589 to Wang A), Key Project of Education Department of Hu Nan (grant number 18A243 to Wang A).

Disclosure

The authors declare that there are no conflicts of interest in this work.

References

- Miura Y, Fukumoto Y, Sugimura K, et al. Identification of new prognostic factors of pulmonary hypertension. *Circ J*. 2010;74(9):1965–1971.
- Lan NSH, Massam BD, Kulkarni SS, Lang CC. Pulmonary arterial hypertension: pathophysiology and Treatment. *Diseases*. 2018;6(2):pii:E38. doi:10.3390/diseases6020038
- Guiot J, Parzibut G, Weber T, et al. Pulmonary arterial hypertension. *Rev Med Liege*. 2019;74(3):139–145.
- Santos-Ribeiro D, Mendes-Ferreira P, Maia-Rocha C, Adão R, Leite-Moreira AF, Brás-Silva C. Pulmonary arterial hypertension: basic knowledge for clinicians. *Arch Cardiovasc Dis*. 2016;109(10):550–561. doi:10.1016/j.acvd.2016.03.004
- Yu L, Tu Y, Jia X, et al. Resveratrol protects against pulmonary arterial hypertension in rats via activation of silent information regulator 1. *Cell Physiol Biochem*. 2017;42(1):55–67. doi:10.1159/000477115
- Zheng Q, Ye J, Cao J. Translational regulator eIF2alpha in tumor. *Tumour Biol*. 2014;35(7):6255–6264. doi:10.1007/s13277-014-1789-0
- Jiang L, Zang D, Yi S, et al. A microRNA-mediated decrease in eukaryotic initiation factor 2alpha promotes cell survival during PS-341 treatment. *Sci Rep*. 2016;6:21565. doi:10.1038/srep21565
- Bentley JK, Hershenson MB. Airway smooth muscle growth in asthma: proliferation, hypertrophy, and migration. *Proc Am Thorac Soc*. 2008;5(1):89–96. doi:10.1513/pats.200705-063VS
- Liu X, Bennett RL, Cheng X, Byrne M, Reinhard MK, May WS Jr. PKR regulates proliferation, differentiation, and survival of murine hematopoietic stem/progenitor cells. *Blood*. 2013;121(17):3364–3374. doi:10.1182/blood-2012-09-456400
- Bennett RL, Pan Y, Christian J, Hui T, May WS Jr. The RAX/PACT-PKR stress response pathway promotes p53 sumoylation and activation, leading to G₁ arrest. *Cell Cycle*. 2012;11(2):407–417. doi:10.4161/cc.11.2.18999
- Wang AP, Li XH, Yang YMA, et al. Critical role of the mTOR/eIF2alpha pathway in hypoxia-induced pulmonary hypertension. *PLoS One*. 2015;10(6):e0130806. doi:10.1371/journal.pone.0130806
- Mizushima N, Levine B, Cuervo AM, Klionsky DJ. Autophagy fights disease through cellular self-digestion. *Nature*. 2008;451(7182):1069–1075. doi:10.1038/nature06639
- Yang Z, Klionsky DJ. Eaten alive: a history of macroautophagy. *Nat Cell Biol*. 2010;12(9):814–822. doi:10.1038/ncb0910-814
- Yang Z, Zeng B, Pan Y, Huang P, Wang C. Autophagy participates in isoliqurritigenin-induced melanin degradation in human epidermal keratinocytes through PI3K/AKT/mTOR signaling. *Biomed Pharmacother*. 2018;97:248–254. doi:10.1016/j.biopha.2017.10.070
- Yorimitsu T, Klionsky DJ. Eating the endoplasmic reticulum: quality control by autophagy. *Trends Cell Biol*. 2007;17(6):279–285. doi:10.1016/j.tcb.2007.04.005
- Levine B, Kroemer G. Autophagy in the pathogenesis of disease. *Intern Med*. 2013;132(1):27–42.
- Grootaert MOJ, Roth L, Schrijvers DM, De Meyer GRY, Martinet W. Defective autophagy in atherosclerosis: to die or to senesce? *Oxid Med Cell Longev*. 2018;2018:7687083. doi:10.1155/2018/7687083
- Roy S, Debnath J. Autophagy and Tumorigenesis. *Semin Immunopathol*. 2010;32(4):383–396. doi:10.1007/s00281-010-0213-0
- Yuan H, Jiang C, Zhao J, et al. Euxanthone attenuates Aβ₁₋₄₂-induced oxidative stress and apoptosis by triggering autophagy. *J Mol Neurosci*. 2018;66(4):512–523. doi:10.1007/s12031-018-1175-2
- Zhou Y, Wang Y, Wang X, et al. The protective effects of kappa-opioid receptor stimulation in hypoxic pulmonary hypertension involve inhibition of autophagy through the AMPK-MTOR pathway. *Cell Physiol Biochem*. 2017;44(5):1965–1979. doi:10.1159/000485886
- Long L, Yang X, Southwood M, et al. Chloroquine prevents progression of experimental pulmonary hypertension via inhibition of autophagy and lysosomal bone morphogenetic protein type II receptor degradation. *Circ Res*. 2013;112(8):1159–1170. doi:10.1161/CIRCRESAHA.111.300483
- Tsai TC, Lai KH, Su JH, Wu YJ, Sheu JH. 7-acetylsinimaximol B induces apoptosis and autophagy in human gastric carcinoma cells through mitochondrial dysfunction and activation of the PERK/eIF2alpha/ATF4/CHOP signaling pathway. *Mar Drugs*. 2018;16(4):pii:E104. doi:10.3390/md16040104
- Kim D, George MP. Pulmonary hypertension. *Med Clin North Am*. 2019;103(3):413–423. doi:10.1016/j.mcna.2018.12.002
- Humbert M, Sitbon O, Simonneau G. Treatment of pulmonary arterial hypertension. *N Engl J Med*. 2004;351(14):1425–1436. doi:10.1056/NEJMra040291
- McLaughlin VV, Archer SL, Badesch DB, et al. ACCF/AHA 2009 expert consensus document on pulmonary hypertension a report of the American college of cardiology foundation task force on expert consensus documents and the American heart association developed in collaboration with the American college of chest physicians; American thoracic society, inc.; and the pulmonary hypertension association. *J Am Coll Cardiol*. 2009;53(17):1573–1619. doi:10.1016/j.jacc.2009.01.004
- Li X, He Y, Xu Y, et al. KLF5 mediates vascular remodeling via HIF-1alpha in hypoxic pulmonary hypertension. *Am J Physiol Lung Cell Mol Physiol*. 2016;310(4):L299–L310. doi:10.1152/ajplung.00189.2015
- Mandegar M, Fung YC, Huang W, Remillard CV, Rubin LJ, Yuan JX. Cellular and molecular mechanisms of pulmonary vascular remodeling: role in the development of pulmonary hypertension. *Microvasc Res*. 2004;68(2):75–103. doi:10.1016/j.mvr.2004.06.001
- Morrell NW, Adnot S, Archer SL. Cellular and molecular basis of pulmonary arterial hypertension. *J Am Coll Cardiol*. 2009;54:S20–S31. doi:10.1016/j.jacc.2009.04.018
- Morrell NW, Adnot S, Archer SL. Eukaryotic initiation factor 2 phosphorylation and translational control in metabolism. *Adv Nutr*. 2012;3(1 Suppl):307–321. doi:10.3945/an.112.002113
- Page AB, Owen CR, Kumar R, et al. Persistent eIF2alpha(P) is colocalized with cytoplasmic cytochrome c in vulnerable hippocampal neurons after 4 hours of reperfusion following 10-minute complete brain ischemia. *Acta Neuropathol*. 2003;106(1):8–16. doi:10.1007/s00401-003-0693-2
- Lin NY, Beyer C, Giessel A, et al. Autophagy regulates TNFalpha-mediated joint destruction in experimental arthritis. *Ann Rheum Dis*. 2013;72(5):761–768. doi:10.1136/annrheumdis-2012-201671
- Kroemer G, Mariño G, Levine B. Autophagy and the integrated stress response. *Mol Cell*. 2010;40(2):280–293. doi:10.1016/j.molcel.2010.09.023
- Liu X, Cao H, Li J, et al. Autophagy induced by DAMPs facilitates the inflammation response in lungs undergoing ischemia-reperfusion injury through promoting TRAF6 ubiquitination. *Cell Death Differ*. 2017;24(4):683–693. doi:10.1038/cdd.2017.1
- Shi B, Shao B, Yang C, Guo Y, Fu X, Gan N. Upregulation of JHDM1D-AS1 protects PDLSCs from H₂O₂-induced apoptosis by decreasing DNJC10 via phosphorylation of eIF2α. *Biochimie*. 2019;9084(19):30191–30199.
- Sengupta S, Seigny CM, Bhattacharya P, Jordan VC, Clarke R. Estrogen-induced apoptosis in breast cancers is phenocopied by blocking dephosphorylation of eukaryotic initiation factor 2 alpha (eIF2α) protein. *Mol Cancer Res*. 2019;17(4):918–928. doi:10.1158/1541-7786.MCR-18-0481

Drug Design, Development and Therapy

Dovepress

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which has also

been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/drug-design-development-and-therapy-journal>