

Association of diabetes knowledge with glycemic control and self-care practices among Pakistani people with type 2 diabetes mellitus

This article was published in the following Dove Press journal:
Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy

Allah Bukhsh^{1,2}
Tahir Mehmood Khan^{1,2}
Muhammad Sarfraz Nawaz³
Hafiz Sajjad Ahmed⁴
Kok Gan Chan^{5,6}
Bey-Hing Goh^{1,2,7,8}

¹School of Pharmacy, Monash University, Bandar Sunway, Selangor, Malaysia;

²Institute of Pharmaceutical Sciences, University of Veterinary and Animal Sciences, Lahore, Pakistan; ³Department of Pharmacy, Quaid-i-Azam University Islamabad, Islamabad, Pakistan; ⁴Capital Hospital, Islamabad, Pakistan;

⁵International Genome Centre, Jiangsu University, Zhenjiang, People's Republic of China; ⁶Division of Genetics and Molecular Biology, Faculty of Science, Institute of Biological Sciences, University of Malaya, Kuala Lumpur, Malaysia;

⁷Biofunctional Molecule Exploratory (BMEX) Research Group, School of Pharmacy, Monash University Malaysia, Bandar Sunway, Selangor Darul Ehsan, Malaysia; ⁸Health and Well-being Cluster, Global Asia in the 21st Century Platform, Monash University Malaysia, Bandar Sunway, Malaysia

Objective: This study explored the relationship of disease knowledge with glycemic control and self-care practices in adult Pakistani people diabetes (PWD).

Methods: People diagnosed with type 2 diabetes (n=218) were selected from three health care centers, located in different cities of Pakistan. Disease knowledge and self-care practices were assessed by Urdu versions of Diabetes Knowledge Questionnaire (DKQ) and Diabetes Self-Management Questionnaire (DSMQ), using a cross-sectional design. Chi-square and correlation analysis were applied to explore the relationship of disease knowledge with glycemic control and self-care practices. Linear regression was used to explore the predictors for disease knowledge.

Results: Majority of the sample was >45–60 years old (48.8%), suffering from type 2 diabetes mellitus for <5 years (49.5%) and had poor glycemic control (HbA1c \geq 7%; n=181 participants). Disease knowledge was significantly associated ($p<0.05$) with patient's gender, level of education, family history of diabetes, nature of euglycemic therapy, and glycemic control. Correlation matrix showed strongly inverse correlations of DKQ with glycosylated hemoglobin levels ($r=-0.62$; $p<0.001$) and strongly positive with DSMQ sum scale ($r=0.63$; $p<0.001$). PWD having university-level education ($\beta=0.22$; 95% Confidence Interval (CI) 0.189, 0.872; $p<0.01$), doing job ($\beta=0.22$; 95% CI 0.009, 0.908]; $p=0.046$), and use of oral hypoglycemic agents in combination with insulin ($\beta=-0.16$; 95% CI [-1.224, -0.071]; $p=0.028$) were the significant predictors for disease knowledge.

Conclusion: Disease knowledge significantly correlated with glycosylated hemoglobin levels and self-care activities of PWD. These findings will help in designing patient-tailored diabetes educational interventions for yielding a higher probability of achieving target glycemic control.

Keywords: knowledge, glycosylated hemoglobin, self-management, glycemic control, HbA1c

Introduction

Type 2 diabetes mellitus (T2DM) is a chronic disease with rapidly increasing prevalence worldwide, but its rate is much higher in low- and middle-income countries such as Pakistan. With an overall prevalence rate of 6.9%, Pakistan has the 10th largest population of people with diabetes (PWD) in the world.¹ According to recent updates from the International Diabetes Federation (IDF), there are 7.5 million adult cases of diabetes in Pakistan. If appropriate interventional strategies are not adopted, this number is expected to reach 16.7 million by the year 2045.¹

Effective management of diabetes depends not only appropriate euglycemic medicine use but also on patients' knowledge about their medicines, healthy eating options,

Correspondence: Allah Bukhsh
School of Pharmacy, Monash University,
Jalan Lagoon Selatan, Bandar Sunway
47500, Selangor, Malaysia
Tel +60 113 362 3703
Fax +60 35 514 6364
Email allah.bukhsh@monash.edu

Kok Gan Chan
International Genome Centre, Jiangsu
University, Zhenjiang, People's Republic of
China
Tel +60 37 967 7748
Email kokgan@um.edu.my

exercise, and self-monitoring of blood glucose levels.²⁻⁴ Despite various pharmacological options available today, diabetes prevalence continues to increase. Poor knowledge about the disease and its associated complications among PWD has not only been reported from low- and middle-income countries but from high-income countries as well.⁵⁻⁷ Poor disease knowledge is one of the main reasons for inadequate self-care behaviors and glycemic control.⁸ Glycated hemoglobin (HbA1c) is a primary measure for glycemic control in PWD. Poor glycemic control (HbA1c \geq 7%) leads to increased mortality and serious complications such as kidney failure, myocardial infarction, stroke, retinopathy, hypertension, and many other associated micro- and macro-vascular complications.^{9,10}

Although disease knowledge alone does not ensure required behavior modifications, the assessment of patients' disease knowledge is a cornerstone in designing tailored health educational intervention.^{11,12} Diabetes education, with consequent improvements in patients' knowledge about disease and self-care activities, results in achieving optimal glycemic control and reduction in disease-related complications.^{8,13}

High prevalence and poor glycemic control among PWD become a challenging health issue of the country. In order to run effective diabetes educational interventions, patients' knowledge about disease must be known. In our recently published study, we have observed that self-care activities of PWD were significantly associated with their glycemic control.¹⁴ Additionally, we also reported that respondents' level of education and use of oral hypoglycemic agents were significant predictors of their self-care practices.¹⁴ So far, little is known about the disease knowledge and its association with glycemic control among PWD of Pakistan. Diabetes patients with good knowledge of disease are expected to maintain desired glycemic control. In this study, we attempted to assess the patients' diabetes knowledge and its association with their glycated hemoglobin and self-care activities by using the recently validated Urdu-version of Diabetes Knowledge Questionnaire.¹⁵ The findings of this study will be helpful in identifying the areas where patients' knowledge about the disease can be improved by designing a patient-tailored diabetes educational intervention in Pakistan.

Methodology

The cross-sectional study design was adopted using the validated Urdu versions of 24-items Diabetes Knowledge Questionnaire (DKQ)¹⁵ and 16-items Diabetes Self-Management Questionnaire (DSMQ)¹⁶ (English and

Urdu version of DKQ and DSMQ are freely available online, the web addresses for both are provided at the end of the manuscript). Data were collected from PWD between February 2017 to September 2017 in outpatient clinics of three health care facilities located in different cities of Pakistan, namely Capital Hospital Islamabad, Akhuwat Diabetes Clinic Lahore, and Awan Medical Complex Lahore.

Study population

In order to be included in the study, patients must have been diagnosed with T2DM (for at least 1 year at the time of interview), older than 30 years, had recent glycated hemoglobin (HbA1c) lab test reports (not older than 2 months at the time of interview), able to understand Urdu language, and willing to take part in the study. Patients suffering from other types of diabetes, cognitive impairment, and any terminal illness were excluded from the study.

Sample size

Daniel's (1999) formula was used for sample size calculation of this study.¹⁷

$$n = \frac{Z^2 P(1 - P)}{d^2}$$

where, n=population sample size, Z=statistic for a level of confidence, P expected prevalence or proportion, and d=precision. The minimum required sample size for the study was n=202, based on 6.9% diabetes prevalence in Pakistan,¹⁸ using Z=1.96 for 95% level of confidence and precision value d=0.035. Thus, data were collected from n=230 subjects, assuming a 10% attrition rate.

Ethical consideration

Monash University Human Research and Ethics approved the study (Ethics approval No. 7767) after seeking permission for data collection from the respective health care facilities in Pakistan. Participation in the study was on a voluntary basis. Informed written consent was obtained from each respondent after providing details of study purpose and procedure. In the case of an illiterate person, we took written consent from their accompanying family members. Participants were assured that their identity and information will be kept confidential.

Data and procedure

Data were collected from the consenting PWD, who met our study inclusion criteria. Participants' eligibility was determined after reviewing their medical records and with the consultation of the health care professionals at data collection facility (n=760). All adult PWD who fulfilled the eligibility criteria of the study (n=245) were invited to take part in the study. Demographic information of the respondents was collected from the willing participants in a private area of the health care facility. Participants were required to complete two previously validated self-administered questionnaires in the Urdu language. Urdu version of Diabetes Knowledge Questionnaire (DKQ)¹⁵ was used to measure participants' disease knowledge, whereas, Urdu version of Diabetes Self-management Questionnaire (DSMQ)¹⁵ was administered to assess their self-care practices. For those participants who were illiterate, questionnaires were read out and explained to them, by the investigators. Out of n=245 eligible patients who agreed to participate in our study, 230 PWD finally participated in the study (response rate 93.7%). Data were analyzed for n=218 patients, after excluding the incomplete questionnaires.

Study tools

Patient's socio-demographic and clinical information

Patients medical record files were used to collect information about their age (years), gender, weight, height, marital status, smoking status, level of education, employment status, family history of diabetes, duration of diabetes, nature of medication, and clinical lab data. Respondents' recent HbA1c value taken from their medical records was considered as a measure of glycemic control.

Diabetes Knowledge Questionnaire (DKQ)

The DKQ is a 24-item questionnaire, designed by Starr County Diabetes Education Study,¹¹ to elicit information about patients' understanding of the cause of their disease, its associated complications, blood glucose levels, diet, and physical activity. The DKQ has been recently translated and validated for Pakistani PWD in the Urdu language.¹⁵ The DKQ has three response options "yes", "no", and "don't know". One point is awarded for each correct option, whereas, no point or negative scoring for the incorrect option. Its scoring involves summing-up the

points obtained by each participant. A higher score represents better disease knowledge.

Diabetes Self-Management Questionnaire (DSMQ)

Self-care activities were assessed with 16-item DSMQ.¹⁹ Translated and validated Urdu-version DSMQ was used in this study.¹⁶ The DSMQ measures the patient's self-care activity for last the 60 days on a 4-point Likert scale. It has four sub-scales; namely, Glucose Management (GM; comprising 5 items), Dietary Control (DC; comprising 4 items), Physical Activity (PA; comprising 3 items), and 4) Health-Care Use (HU; comprising 3 items). There are seven positively keyed (item no. 1, 2, 3, 4, 6, 8, and 9) and nine reverse-coded items (item no. 5, 7, 10, 11, 12, 13, 14, 15, and 16). The 16th item of DSMQ, which measures overall self-care of the respondent, is only included in the sum-scale. The scoring of DSMQ involves summing-up scores of all items (after reversing nine negatively keyed items) and transforming into a scale ranging from 0 to 10. A higher score represents better self-care.

Statistical analysis

Data were analyzed by Statistical Package for Social Science (SPSS 24.0. Inc., Chicago, IL, USA). Demographic and clinical variables were presented by using descriptive statistics. Data distribution was analyzed by the Shapiro–Wilk test.²⁰ Mann–Whitney U-test and Kruskal–Wallis test were applied to compare groups with non-normal distribution. American Diabetes Association (ADA) guideline was used to define glycemic control in this study. Patients with values of HbA1c<7% were classified as "good glycemic control", whereas those with HbA1c≥7% were classified as "poor glycemic control".^{21–23} Spearman rank-order (two-tailed test) was applied to find the association of DKQ median scores with DSMQ sum scale and its subscales (GM, DC, PA, and HU). Multiple linear regression was applied to identify significant predictors of diabetes knowledge. Statistical tests were considered significant at $p<0.05$.

Results

The demographic details of the respondents have been presented in Table 1. A total of 230 PWD participated in this study. Whereas, after removing the incomplete questionnaire, data of 218 patients were analyzed. Both genders equally participated in this study (male=51.4% and female=48.6%). The majority of the participants, n=102,

Table 1 Characteristics of the study sample and their association with diabetes knowledge (N=218)

Parameter	n (%); median (IQR)	DKQ sum score median (IQR)	p-value
Gender			0.04 ^a
Male	112 (51.4)	14 (11.25–17)	
Female	106 (48.6)	13 (10–16)	
Age (years)	52 (45–60)		0.92 ^b
30–45 years	65 (29.8)	14 (11–16)	
>45–60 years	102 (48.8)	14 (10–17)	
>60 years	51 (23.4)	19 (11–17)	
BMI (kg/m ²)	28.53 (24.99–32.32)		0.51 ^b
Underweight (<18.5)	3 (1.4)	10 (9–10)	
Normal (18.5–<25)	51 (22.9)	13 (10–16)	
Overweight (25–<30)	85 (38.5)	14 (11–17)	
Obese (≥30)	79 (37.2)	13 (11–17)	
Smoking			0.66 ^a
No	205 (94.5)	14 (10.5–17)	
Yes	13 (5.5)	13 (10–15)	
Education			0.004 ^b
No formal education	48 (22)	13 (10–15)	
Primary level	37 (17)	13 (9.5–17)	
Secondary level	88 (40.4)	14 (11–17)	
University level	45 (20.6)	15 (12–19)	
Family history of diabetes			0.03 ^b
First degree relatives	129 (59.2)	13 (10–17)	
Second degree relatives	3 (1.4)	10 (9–10)	
Both first and second degree relatives	17 (7.8)	17 (13.5–19)	
No history	69 (31.7)	14 (11–16)	
Working status			0.02 ^b
Jobless	8 (3.7)	13 (10–15.25)	
Housewives/stay at home	104 (47.7)	13 (10–16)	
Business	33 (15.2)	13 (11–16)	
Doing Job	59 (27.1)	15 (12–18)	
Retired	14 (6.4)	15 (13–18)	
Diabetes duration (years)	5 (2–10)		0.18 ^b
<5 years	108 (49.5)	14 (12–17)	
5–9 years	44 (20.2)	13 (10–17.75)	
>9–<15 years	41 (18.8)	13 (11–17)	
≥15 years	25 (11.5)	11 (10–15.5)	
Anti-diabetic therapy			0.002 ^b
Exclusively insulin	36 (16.5)	13 (10–15.75)	
Combined with medication	67 (30.7)	13 (10–15)	
Oral hypoglycemic agents only	115 (52.8)	15 (12–18)	
HbA1c value (%)	8.9 (7.27–10.2)		<0.001 ^a
Good glycemic control (<7%)	37 (17)	18 (16–19)	
Poor glycemic control (≥7%)	181 (83)	13 (10–15)	

Notes: Data are n (%) or M±SD. ^aRegards a Association with DKQ Sum scale; Mann–Whitney U Test (two-tailed test). ^bAssociation with DKQ Sum scale; Kruskal–Wallis test. **Abbreviations:** HbA1c, glycated hemoglobin; BMI, body mass index; DKQ, Diabetes Knowledge Questionnaire.

were of age ranging from 45 to 60 years, non-smokers (94.5%), and married (97.7%). With regard to the level of education, 22% had no formal education, 17% had primary level education, whereas, rest 61% had secondary and university-level education. More than half of the respondents (59.2%) reported for a family history of diabetes among their first degree relatives only. Nearly half of the respondents (49.5%) were living with diabetes for less than 5 years. Only 22.9% were having a normal body mass index, whereas 38.5% were overweight and the rest 37.2% were obese.

Current pharmacological therapy consisted of oral hypoglycemic agents only in 52.5%; exclusively insulin in 16.5% and 30.7% used a combination of oral hypoglycemic agents and insulin. Good glycemic control ($HbA1c < 7\%$) was observed in only 17% of the study participants.

Details of respondents' demographic information and their association with disease knowledge are presented in Table 1.

Diabetes knowledge was significantly related to glycosylated hemoglobin ($r = -0.62$, $p < 0.001$). PWD having good glycemic control ($HbA1c < 7\%$) scored significantly higher ($p < 0.001$) for diabetes knowledge (DKQ sum-scale score 18; IQR 16–19), as compared to those with poor glycemic control (DKQ sum-scale score 13; IQR 10–15).

Patient's age, BMI, smoking status, and duration of diabetes were not found to have statistically significant ($p > 0.05$) association with their disease knowledge. Significantly ($p < 0.05$) higher scores for DKQ sum-scale were obtained by the study respondents with male gender (14; IQR 11.25–17), university-level education (15; IQR 12–19), having family history of diabetes in both primary and secondary degree relatives (17; IQR 13.5–19), and those who were using oral hypoglycemic agents only (15; IQR 12–18). (Table 1).

Significantly better ($p < 0.05$) knowledge of disease and blood glucose testing was observed in subjects with good glycemic control; details are presented in Table 2. Strongly positive correlation of DKQ sum-scale was observed with DSMQ sum-scale ($r = 0.63$, $p < 0.001$), and with three sub-scales of DSMQ; namely Glucose Management ($r = 0.61$, $p < 0.001$), Dietary Control ($r = 0.65$, $p < 0.001$), and Healthcare Use ($r = 0.55$, $p < 0.001$). Whereas, a moderately positive correlation was observed between the DKQ sum-scale and Physical activity sub-scale of DSMQ ($r = 0.44$, $p < 0.001$). Details are presented in Table 3.

Multiple linear regression analysis was run to predict the patients' disease knowledge (DKQ sum-scale score) based on the patients' gender, age, working status, family history of diabetes, duration of diabetes, level of education, and nature of euglycemic medicine. Significant regression equation was observed ($F(7, 210) = 4.591$, $p < 0.001$), with R^2 of 0.133. Respondents' predicted disease knowledge is equal to $12.226 - 1.037$ (gender) $+ 0.458$ (working status) $+ 0.230$ (family history of diabetes) $+ 0.530$ (level of education) $+ 0.005$ (age) $- 0.019$ (duration of diabetes) $- 0.648$ (nature of euglycemic medicine). Regression analysis revealed that respondents' working status ($\beta = 0.21$; [95% CI 0.008, 0.918]; $p = 0.04$), level of education ($\beta = 0.18$; [95% CI 0.108, 0.121]; $p = 0.01$) and nature of euglycemic therapy ($\beta = -0.15$; [95% CI -1.206 , -0.013]; $p = 0.04$), were the significant predictors for their disease knowledge. Whereas, patient's gender, duration of diabetes, age, and family history of diabetes were insignificant predictors for patients' disease knowledge. (Table 4)

Discussion

Health education for patients suffering from chronic diseases is an important source to improve their self-care practices. Existing literature about patients' disease knowledge plays a vital role in designing a patient-tailored and outcome-oriented educational intervention. Diabetes-related self-care practices, such as, healthy eating practices, being physically active, blood glucose monitoring, and adhering to the recommended euglycemic therapy, can benefit patients in achieving desired glycemic control.^{24,25} Limited literature is available on disease knowledge of people with diabetes in Pakistan. This study is first of its kind to appraise the association of disease knowledge with glycemic control and self-care practices of PWD in Pakistan.

Our study revealed that disease knowledge was higher among those with university-level education, male gender, doing job, or retired from a job, and the differences were statistically significant ($p < 0.05$). Furthermore, those respondents who had diabetes in their first- and second-degree relatives had better knowledge. Surprisingly, disease knowledge of respondents who were using only oral euglycemic agents was significantly better ($p < 0.05$) than those who were using either insulin alone or in combination with oral euglycemic agents. Improved self-care practices were also observed in our recent study among PWD using oral hypoglycemic agents only.¹⁴ However, statistically insignificant

Table 2 Disease knowledge of people with diabetes according to glycemic control (n=218)

Sr. No.	Question	Correct answer, n (%)			p-value
		Total (n=218)	Good glycemic control (<7%, (n=37)	Poor glycemic control (≥7%, (n=181)	
1	Eating too much sugar and other sweet foods is a cause of diabetes.	51 (23.4)	19 (51.4)	32 (17.7)	<0.001
2	The usual cause of diabetes is lack of effective insulin in the body.	157 (72)	35 (9.6)	122 (67.4)	0.001
3	Diabetes is caused by failure of the kidneys to keep sugar out of the urine.	63 (29.9)	15 (40.5)	48 (26.5)	0.086
4	Kidneys produce insulin.	132 (60.6)	29 (78.4)	103 (56.9)	0.015
5	In untreated diabetes, the amount of sugar in the blood usually increases.	203 (93.1)	36 (97.3)	167 (92.3)	0.271
6	If I am diabetic, my children have a higher chance of being diabetic.	192 (88.1)	36 (97.3)	156 (86.2)	0.057
7	Diabetes can be cured.	18 (8.3)	3 (8.1)	15 (8.3)	0.971
8	A fasting blood sugar level of 210 is too high.	147 (67.4)	36 (97.3)	111 (61.3)	<0.001
9	The best way to check my diabetes is by testing my urine.	108 (49.5)	31 (83.8)	77 (42.5)	<0.001
10	Regular exercise will increase the need for insulin or other diabetic medication.	106 (48.6)	23 (62.2)	83 (45.9)	0.071
11	There are two main types of diabetes: Type 1 (insulin-dependent) and Type 2 (noninsulin dependent).	153 (70.2)	34 (91.9)	119 (65.7)	0.002
12	An insulin reaction is caused by too much food.	70 (32.1)	19 (51.4)	51 (28.2)	0.006
13	Medication is more important than diet and exercise to control my diabetes.	70 (32.1)	15 (40.5)	55 (30.4)	0.228
14	Diabetes often causes poor circulation.	144 (66.1)	26 (70.3)	118 (65.2)	0.552
15	Cuts and abrasions on diabetes heal more slowly.	206 (94.5)	36 (97.3)	170 (93.9)	0.412
16	Diabetics should take extra care when cutting their toenails.	207 (95)	37 (100)	170 (93.9)	0.124
17	A person with diabetes should cleanse a cut with iodine and alcohol.	25 (11.5)	9 (24.3)	16 (8.8)	0.007
18	The way I prepare my food is as important as the foods I eat.	184 (84.4)	32 (86.5)	152 (84)	0.702
19	Diabetes can damage my kidneys.	188 (86.2)	37 (100)	151 (83.4)	0.008
20	Diabetes can cause loss of feeling in my hands, fingers and feet.	175 (80.3)	35 (94.6)	140 (77.3)	0.016
21	Shaking and sweating are signs of high blood sugar.	110 (50.5)	30 (81.1)	80 (44.2)	<0.001
22	Frequent urination and thirst are signs of low blood sugar.	115 (52.8)	31 (83.8)	84 (46.4)	<0.001
23	Tight elastic hose or socks are not bad for diabetics.	155 (71.1)	35 (94.6)	120 (66.3)	0.001
24	A diabetic diet consists mostly of special foods.	24 (11)	7 (18.9)	17 (9.4)	0.092

Table 3 Correlation between diabetes knowledge and self-management practices of type 2 diabetes patients (N=218)

Parameter	Median (IQR)	Correlation	p-value
DSMQ "Sum Scale"	3.96 (2.71–6.88)	0.63	<0.001
Subscale "Glucose Management"	4.67 (3.33–7.33)	0.61	<0.001
Subscale "Dietary Control"	4.17 (2.5–6.88)	0.65	<0.001
Subscale "Physical Activity"	3.33 (1.11–6.67)	0.44	<0.001
Subscale "Health-Care Use"	4.44 (2.22–6.67)	0.55	<0.001

Note: Spearman rank-order (two-tailed test) was applied to check correlation between DKQ and DSMQ scales. The DSMQ scale scores are transformed to a scale ranging from 0 to 10.
Abbreviations: DSMQ, Diabetes Self-Management Questionnaire; DKQ, Diabetes Knowledge Questionnaire; IQR, interquartile range.

association ($p>0.05$) was observed between respondents' age, disease knowledge, and duration of disease.

Good glycemic control was observed in only 17% ($n=37$) of the study participants. Like this study, a similar trend was reported in two other studies conducted in urban areas of Pakistan, where good glycemic control was achieved in less than one-third of the study participants.^{26,27} In our study, we found a statistically significant association of ($p<0.01$) higher diabetes knowledge with low HbA1c levels and better self-care practices. A similar trend was also reported in a study conducted in Ethiopia, where a high level of disease knowledge was significantly associated with patients' literacy and their good self-care practices.⁸ In contrast, Dussa et al reported that disease knowledge in Indian PWD was not correlated with their HbA1c levels.²⁸ Besides poor disease knowledge and inadequate self-care practices, poor glycemic control may be partly reflective of more severe disease and treatment not being suitably intensified. Diabetes is a progressive disease and most of the patients in our study (52.8%) were using OHA and were suffering from diabetes for a longer duration.

Further findings indicated that subjects with good glycemic control scored higher ($p<0.050$) for knowledge about the disease, its associated complications, normal

blood glucose levels, and its monitoring. However, as a whole, majority of the study participants had poor knowledge about diet and physical activity, which was also represented in the form of lower score for DSMQ's subscale "Physical activity" [3.33 (95% CI 1.11–6.67)] and DSMQ's sub-scale "Dietary control" [4.17 (95% CI 2.5–6.88)]. Similar findings were also reported from an urban city of Pakistan (Peshawar), where 75% of the participants were not performing any form of exercise to control their blood sugar levels.²⁹

Besides being physically active, healthy eating practices also play a vital role in diabetes self-care. Inadequate knowledge regarding diet and poor dietary practices could be other contributing factors for poor glycemic control among the study participants. Our results are also consistent with those from low- and middle-income countries, as well as high-income countries, which indicated that glycemic control is associated with disease knowledge and self-care practices of PWD.^{30,31}

This study identified several patients' demographic characteristics associated with their disease knowledge. The higher level of patients' education followed by the job status of the respondents was strong predictors for better disease knowledge, which is in line with the findings of previous studies conducted in low- and middle-income countries, where diabetes knowledge was related to patients' education levels.^{8,28} Relating to the present study, only one-fifth (21%) of the respondents who had university-level education scored maximum for knowledge about diabetes. Patients' level of education was also identified as a significant predictor for their self-care practices in one of our recently published studies.¹⁴

Besides pharmacological intervention, improving knowledge and awareness among PWD and their family members will lead to better health-related outcomes. The findings of our study recommend that the contents of an educational intervention should be designed keeping in view the educational status of the patients, as majority of

Table 4 Linear association of patients demographic characteristics on DKQ, Diabetes Knowledge Questionnaire; (N=218)

Predictors	β [95% CI]	p-value
Gender	-0.14 [-2.601, 0.527]	0.193
Working status	0.22 [0.009, 0.908]	0.046
Anti-diabetes medication	-0.16 [-1.224, -0.071]	0.028
Family history of diabetes	0.09 [-0.107, 0.567]	0.180
Level of education	0.22 [0.189, 0.872]	0.003
Duration of diabetes (years)	-0.03 [-0.103, 0.065]	0.651
Age (years)	0.02 [-0.032, 0.043]	0.783

Notes: Gender: reference category = female; working status: reference category = jobless; anti-diabetes medication: reference category = oral hypoglycemic agents; education: reference category = no formal education.

the patients with low level of education had poor disease knowledge. Second, while imparting diabetes education, job nature of the PWD must be taken into account, because irrespective of the education-level, the disease knowledge of the majority of our respondents (housewives) was comparable to that of illiterate respondents.

Strengths and limitations

Assessment of diabetes knowledge and self-care activities by using DKQ and DSMQ, which were recently validated for PWD in Pakistan, is one of the strengths of this study. The study is one of its kind to appraise the association of diabetes knowledge with glycemic control and self-care activities in PWD living in urban areas of Pakistan. One of the limitations of the current study is its cross-sectional design and location of the studied population, as most of the participants were belonging an urban area of Pakistan, so the findings may not be representative of all people living in rural areas. Further limitation includes the possibility of self-reporting bias, as patients might be unwilling to reveal deficiencies in the self-care practices and may not be accurate all the time.

Conclusion

Peer support approaches are endorsed by WHO as both effective and cost-effective in providing ongoing support to people with diabetes especially for low- and middle-income countries, such as Pakistan. Just as represented in our study findings, most of the PWD had low education levels, so providing education to family members/peers in addition to diabetes patients could prove to be more effective. Improved knowledge about diabetes and self-care practices will help patients to optimize their lifestyle and reduce the chances of diabetes-associated complications. Additionally, health education to PWD family members will also support patients in coping with recommended dietary and other lifestyle modifications. Our findings lend support to tailoring diabetes educational programs for PWD in Pakistan. So culturally sensitive and patient-tailored educational interventions by health care providers are likely to be more effective at achieving desired clinical outcomes.

Diabetes questionnaire links

Diabetes Self-management Questionnaire (Urdu Version) is available online at: https://static-content.springer.com/esm/art%3A10.1186%2Fs12955-017-0776-8/MediaObjects/12955_2017_776_MOESM2_ESM.pdf.

Diabetes Self-management Questionnaire (English version) is available online at: https://static-content.springer.com/esm/art%3A10.1186%2Fs12955-017-0776-8/MediaObjects/12955_2017_776_MOESM1_ESM.pdf. Diabetes Knowledge Questionnaire (Urdu version) is available online at: https://www.frontiersin.org/files/Articles/274955/fpubh-05-00139-HTML/image_m/fpubh-05-00139-at001.jpg.

Acknowledgments

AB gratefully acknowledges Monash University for providing a PhD Merit Scholarship. KGC thanks the University of Malaya for grantd (PG136-2016A and PG135-2016A). GBH thanks Monash Global Asia in the Twenty-First Century (GA21) for research grants (GA-HW-19-L01 and GA-HW-19-501) and the External Industry Grant from Biotek Abadi Sdn Bhd (vote no. GBA-81811A).

Disclosure

The authors report no conflicts of interest in this work.

References

- International Diabetes Federation. *IDF Diabetes Atlas - 8th Edition*; 2017. Available from: www.idf.org/diabetesatlas. Accessed March 13, 2019.
- Bukhsh A, Khan TM, Lee S, et al. Efficacy of pharmacist based diabetes educational interventions on clinical outcomes of adults with type 2 diabetes mellitus: a network meta-analysis. *Front Pharmacol*. 2018;9:339. doi:10.3389/fphar.2018.00339
- Bukhsh A, Tan XY, Chan KG, Lee L-H, Goh B-H, Khan TM. Effectiveness of pharmacist-led educational interventions on self-care activities and glycemic control of type 2 diabetes patients: a systematic review and meta-analysis. *Patient Prefer Adherence*. 2018;12:2457. doi:10.2147/PPA.S176067
- Shrivastava SR, Shrivastava PS, Ramasamy J. Role of self-care in management of diabetes mellitus. *J Diabetes Metab Disord*. 2013;12(1):14. doi:10.1186/2251-6581-12-14
- Al-Maskari F, El-Sadig M, Al-Kaabi JM, Afandi B, Nagelkerke N, Yeatts KB. Knowledge, attitude and practices of diabetic patients in the United Arab Emirates. *PLoS One*. 2013;8(1):e52857. doi:10.1371/journal.pone.0052857
- Deepa M, Bhansali A, Anjana R, et al. Knowledge and awareness of diabetes in urban and rural India: the Indian Council of Medical Research India diabetes study (phase I): Indian Council of Medical Research India diabetes 4. *Indian J Endocrinol Metab*. 2014;18(3):379. doi:10.4103/2230-8210.131191
- Islam SMS, Niessen LW, Seissler J, et al. Diabetes knowledge and glycemic control among patients with type 2 diabetes in Bangladesh. *Springerplus*. 2015;4(1):284. doi:10.1186/s40064-015-1103-7
- Kassahun T, Gesesew H, Mwanri L, Eshetie T. Diabetes related knowledge, self-care behaviours and adherence to medications among diabetic patients in Southwest Ethiopia: a cross-sectional survey. *BMC Endocr Disord*. 2016;16(1):28. doi:10.1186/s12902-016-0114-x
- Gibbons CH, Goebel-Fabbri A. Microvascular complications associated with rapid improvements in glycemic control in diabetes. *Curr Diab Rep*. 2017;17(7):48. doi:10.1007/s11892-017-0902-3

10. Association AD. 5. Glycemic targets. *Diabetes Care*. 2016;39 (Supplement 1):S39–S46. doi:10.2337/dc16-S008
11. Garcia AA, Villagomez ET, Brown SA, Kouzekanani K, Hanis CL. The starr county diabetes education study development of the Spanish-language diabetes knowledge questionnaire. *Diabetes Care*. 2001;24(1):16–21. doi:10.2337/diacare.24.1.16
12. Schapira MM, Swartz S, Ganschow PS, et al. Tailoring educational and behavioral interventions to level of health literacy: a systematic review. *MDM Policy Pract*. 2017;2(1):2381468317714474. doi:10.1177/2381468317714474
13. Chrvala CA, Sherr D, Lipman RD. Diabetes self-management education for adults with type 2 diabetes mellitus: a systematic review of the effect on glycemic control. *Patient Educ Couns*. 2016;99(6):926–943. doi:10.1016/j.pec.2015.11.003
14. Bukhsh A, Khan TM, Nawaz MS, et al. Association of diabetes-related self-care activities with glycemic control of patients with type 2 diabetes in Pakistan. *Patient Prefer Adherence*. 2018;12:2377. doi:10.2147/PPA.S176067
15. Bukhsh A, Lee SWH, Pusparajah P, Khan AH, Khan TM. Psychometric properties of the Urdu version of diabetes knowledge questionnaire. In: *Front Publ Health*. 2017;5:139.
16. Bukhsh A, Lee SWH, Pusparajah P, Schmitt A, Khan TM. Psychometric properties of the Diabetes Self-Management Questionnaire (DSMQ) in Urdu. *Health Qual Life Outcomes*. 2017;15(1):200. doi:10.1186/s12955-017-0776-8
17. Daniel W. *Biostatistics: A Foundation for Analysis in the Health Sciences*. 7th ed. New York: R Wiley; 1999.
18. International Diabetes Federation. *IDF Diabetes Atlas - 7th ed.*; 2016. Available from: <http://www.diabetesatlas.org/>. Accessed February 13, 2016.
19. Schmitt A, Gahr A, Hermanns N, Kulzer B, Huber J, Haak T. The Diabetes Self-Management Questionnaire (DSMQ): development and evaluation of an instrument to assess diabetes self-care activities associated with glycaemic control. *Health Qual Life Outcomes*. 2013;11(1):1. doi:10.1186/1477-7525-11-83
20. Ghasemi A, Zahediasl S. Normality tests for statistical analysis: a guide for non-statisticians. *Int J Endocrinol Metab*. 2012;10(2):486–489. doi:10.5812/ijem.3505
21. Ali MK, Bullard KM, Saaddine JB, Cowie CC, Imperatore G, Gregg EW. Achievement of goals in US diabetes care, 1999–2010. *N Engl J Med*. 2013;368(17):1613–1624. doi:10.1056/NEJMsa1213829
22. García-Pérez L-E, Álvarez M, Dilla T, Gil-Guillén V, Orozco-Beltrán D. Adherence to therapies in patients with type 2 diabetes. *Diabetes Ther*. 2013;4(2):175–194. doi:10.1007/s13300-013-0034-y
23. Lee SWH, Lee JY, San San Tan C, Wong CP. Strategies to make Ramadan fasting safer in type 2 diabetics: a systematic review and network meta-analysis of randomized controlled trials and observational studies. *Medicine*. 2016;95:2. doi:10.1097/MD.00000000000004864
24. Heisler M, Smith DM, Hayward RA, Krein SL, Kerr EA. How well do patients' assessments of their diabetes self-management correlate with actual glycemic control and receipt of recommended diabetes services? *Diabetes Care*. 2003;26(3):738–743. doi:10.2337/diacare.26.3.738
25. Chiu C-J, Wray LA. Peer reviewed: factors predicting glycemic control in middle-aged and older adults with type 2 diabetes. *Prev Chronic Dis*. 2010;7:1.
26. Ullah Khan A, Zafar Ali Khan M, Nadeem M, Yasmeen BR, Fakhr A. Status of glycemic control in patients of type 2 diabetes mellitus. *Pak Armed Forces Med J*. 2013;63:2.
27. Hassan MM, Zafar A, Akram S. Glycemic control and clinical outcomes of diabetic pulmonary tuberculosis patients. *Pak J Med Health Sci*. 2018;12(1):371-+.
28. Parimalakrishnan S, Dussa K, Sahay R. Assessment of diabetes knowledge using diabetes knowledge questionnaire among people with type 2 diabetes mellitus. *Asian J Pharm Clin Res*. 2015;8(2):254–256.
29. Gul N. Knowledge, attitudes and practices of type 2 diabetic patients. *J Ayub Med Coll Abbottabad*. 2010;22(3):128–131.
30. D'Souza MS, Karkada SN, Venkatesaperumal R, Natarajan J. Self-care behaviours and glycemic control among adults with type 2 diabetes. *GSTF J Nurs Health Care (JNHC)*. 2017;2:1.
31. Houle J, Beaulieu MD, Chiasson JL, et al. Glycaemic control and self-management behaviours in Type 2 diabetes: results from a 1-year longitudinal cohort study. *Diabetic Med*. 2015;32(9):1247–1254. doi:10.1111/dme.12686

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy

Dovepress

Publish your work in this journal

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert opinion

and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-targets-and-therapy-journal>