

# Co-Occurrence of the *bla*<sub>KPC-2</sub> and *Mcr-3.3* Gene in *Aeromonas caviae* SCAC2001 Isolated from Patients with Diarrheal Disease

This article was published in the following Dove Press journal:  
*Infection and Drug Resistance*

Lingtong Tang<sup>1,2,\*</sup>  
Jianglian Huang<sup>3,\*</sup>  
Junping She<sup>1</sup>  
Kelei Zhao<sup>4</sup>  
Yingshun Zhou<sup>1</sup>

<sup>1</sup>Department of Pathogenic Biology, School of Basic Medicine, Southwest Medical University, Luzhou 646000, Sichuan, People's Republic of China;

<sup>2</sup>Department of Clinical Laboratory, People's Hospital of Gao County, Yibing 644000, Sichuan, People's Republic of China;

<sup>3</sup>Department of Clinical Laboratory, The Second Affiliated Hospital of Xiamen Medical College, Xiamen 361600, People's Republic of China; <sup>4</sup>Antibiotics Research and Re-Evaluation Key Laboratory of Sichuan Province, Sichuan Industrial Institute of Antibiotics, Chengdu University, Chengdu 610052, Sichuan, People's Republic of China

\*These authors contributed equally to this work

**Purpose:** To characterize the genetic feature of a multi-drug-resistant *Aeromonas caviae* strain isolated from the diarrhea sample of a 45-year-old male patient with acute diarrhea.

**Materials and Methods:** Whole-genome of the *A. caviae* strain SCAC2001 was sequenced via the Illumina system, followed by a series of bioinformatic analyses to describe the genetic feature.

**Results:** The genome sequence of *A. caviae* SCAC2001 was assembled into 340 scaffolds (305 of them were > 1000 bp in length and 4,487,370 bp in total) with an average G+C content of 61.09%. Phylogenetic analysis showed that the *A. caviae* SCAC2001 strain was highly similar to the *A. caviae* strain R25-2 and T25-39. Resistome analysis identified that *A. caviae* SCAC2001 carried 13 antimicrobial resistance genes, including  $\beta$ -lactams (*bla*<sub>KPC</sub>, *bla*<sub>CTX-M-14</sub>, *bla*<sub>TEM-1</sub>, *bla*<sub>OXA-10</sub>, *bla*<sub>OXA-427</sub>, *bla*<sub>VEB-3</sub> and *bla*<sub>MOX-6</sub>), aminoglycosides (*aadA1*), fluoroquinolones (*aac(6)-Ib-cr*), phenicol resistance (*catB3*), sulfonamide (*sul1*), trimethoprim (*dfrA5*) and colistin resistance (*mcr-3.3*). And also, *A. caviae* SCAC2001 carried 54 putative virulence genes including the type IV pilus, fimbria, flagellarthe, and hemolysin A encoding genes, and 12 pathogen–host interactions (PHI) genes. There were also four genomic islands and eight prophages in the genome of *A. caviae* SCAC2001. In addition, *A. caviae* SCAC2001 also carried three secondary metabolism products coding clusters including nonribosomal peptide synthetases (nrps), hserlactone and bacteriocin.

**Conclusion:** *A. caviae* SCAC2001 carries many resistance genes, a variety of virulence factors, PHI genes and four genomic islands and eight prophages, which poses a severe threat to infectious diseases control strategies, diagnosis methods and clinical treatment.

**Keywords:** *Aeromonas caviae*, *bla*<sub>KPC-2</sub>, *mcr-3.3*, virulence factors, secondary metabolism products

Correspondence: Yingshun Zhou  
Department of Pathogenic Biology,  
School of Basic Medicine, Southwest  
Medical University, Luzhou, Sichuan  
646000, People's Republic of China, No.  
319, Zhongshan Road,  
Tel +86-0830-3160073  
Email yingshunzhou@swmu.edu.cn

Kelei Zhao  
Antibiotics Research and Re-Evaluation  
Key Laboratory of Sichuan Province,  
Sichuan Industrial Institute of Antibiotics,  
No. 168, Huaguan Road, Chengdu  
610052, Sichuan, People's Republic of  
China  
Tel +86-028-84216035  
Email zhaokelei@cdu.edu.cn

## Introduction

Colistin is the last resort for the treatment of infections caused by multidrug-resistant bacteria, particularly the carbapenem-resistant microorganisms.<sup>1,2</sup> However, the mobile colistin-resistant gene *mcr-1*, was first reported in the *Enterobacteriaceae* by Liu et al<sup>3</sup> in 2015, which led the *mcr-1* carried bacteria resistant to colistin. Since then, *mcr-1* or the *mcr* gene family-carrying bacteria have been reported in different species (such as *E. coli*, *klebsiella pneumoniae*, *Acinetobacter*, *Pseudomonas* and other gram-negative bacteria) isolated from food, animals, the environment and clinical samples worldwide.<sup>4-6</sup> To our knowledge, co-carriers of the colistin-resistant gene *mcr* and carbapenemase-resistant

genes (*bla*<sub>KPC</sub>, *bla*<sub>NDM</sub>, *bla*<sub>IMP</sub>, and *bla*<sub>VIM</sub>), microorganisms have potentially evolved into extensively drug-resistant or pan-drug-resistant isolates. Infections caused by these clinical isolates co-harboring the colistin-resistant gene (*mcr-1*) and carbapenem-resistant genes (*bla*<sub>KPC</sub>, *bla*<sub>NDM</sub>, *bla*<sub>IMP</sub>, and *bla*<sub>VIM</sub>) pose a serious threat because the antibiotic options would be much fewer.<sup>4,7,8</sup> *Aeromonas* species, one kind of the gram-negative bacteria, was identified in the 1980s as an enteric pathogen which can lead to severe diarrhea.<sup>9</sup> In addition, the carbapenem-resistant and/or colistin-resistant *Aeromonas* species strains have been reported increasingly in recent years to pose a serious threat in infection control.<sup>10–12</sup> In this study, we recovered a colistin and carbapenem-resistant *Aeromonas* strain from the diarrhea sample of a 45-year-old male patient with acute diarrhea in the affiliated hospital of Southwest Medical University, and the genomic information of this strain was characterized to gain insight into further infection control.

## Materials and Methods

### Isolation and Identification of *Aeromonas caviae* SCAC2001

*A. caviae* SCAC2001 was recovered from the diarrhea sample of a 45-year-old male patient with acute diarrhea in a hospital in Sichuan, China, in May, 2019. It was identified as *Aeromonas caviae* using the Vitek-2 compact system (bioMérieux, Marcy-l'Étoile, France). The presence of the acquired carbapenemase genes (*bla*<sub>KPC</sub>, *bla*<sub>NDM</sub>, *bla*<sub>GES</sub>, *bla*<sub>IMP</sub> and *bla*<sub>VIM</sub>) and *mcr* genes in SCAC2001 was determined by PCR amplification as described previously.<sup>13–15</sup>

### Antimicrobial Susceptibility Testing

In vitro susceptibility tests of *A. caviae* SCAC2001 against 17 antimicrobial agents (Solarbio, China) including meropenem, imipenem, cefepime, cefotaxime, ceftazidime, piperacillin-tazobactam, amoxicillin-clavulanic acid, gentamicin, amikacin, aztreonam, erythromycin, chloramphenicol, colistin, tigecycline, fosfomicin and ciprofloxacin and trimethoprim-sulfamethoxazole determined by broth microdilution method according to the recommendations of the Clinical and Laboratory Standards Institute (CLSI 2013, M100-S23), and the breakpoints of colistin were interpreted according to the European Committee on Antimicrobial Susceptibility

Testing (EUCAST) (<http://www.eucast.org/>); *E. coli* J53 was used as quality control.

## Genome Sequencing and Analysis

The genomic DNA of *A. caviae* SCAC2001 was extracted using the Axygen<sup>®</sup> DNA Gel Extraction Kits (Axygen, People's Republic of China) according to the manufacturer's protocol. Purified DNA was subjected to whole genomic sequencing on the Illumina system with the 150-bp paired-end approach and >180× coverage (Novogene, People's Republic of China). The reads were assembled using the software SOAP denovo (version 2.04).<sup>16</sup> Gene prediction was performed with GeneMarkS (version 4.17).<sup>17</sup> Gene annotation was achieved using the NCBI Prokaryotic Genome Annotation Pipeline. The pairwise alignment was performed by blastn search (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). The resistome was identified using ResFinder (<https://cge.cbs.dtu.dk/services/ResFinder/>)<sup>18</sup> (minimum threshold for identity, 80%; minimum coverage, 60%) and Comprehensive Antibiotic Resistance Database (CARD). The virulence factors were identified by the VFAnalyzer (<http://www.mgc.ac.cn/VFs/main.htm>). The pathogen–host interactions (PHI) genes were identified by comparison with the pathogen–host interactions database (minimum threshold for identity, 80%).<sup>19</sup> To determine the phylogenetic groups of the *A. caviae* SCAC2001 strain, the phylogenetic tree was constructed by aligning the core genome of *A. caviae* SCAC2001 strain with other representative *A. caviae* strains available in the genbank (Table 1). All the sequences were aligned using Mugsy and thereafter a maximum-likelihood phylogeny tree was generated using RAxML version 8 and MEGA7.0.<sup>20–22</sup> The genomic island sequences were predicted based on three different genomic islands (GIs) prediction softwares (IslandPATH-DIMOB, IslandPick, and SIGI-HMM)<sup>23–25</sup> and the prophages were predicted by using phiSpy.<sup>26</sup> The secondary metabolism products coding clusters were identified by the antiSMASH.<sup>27</sup>

## Results and Discussion

### Characteristics of the Isolate SCAC2001

A carbapenem-resistant gene *bla*<sub>KPC</sub> and a colistin-resistant gene *mcr-3.3* co-carried by *A. caviae* strain SCAC2001 was isolated and identified by the Vitek-2 compact system and resistance genes PCR detection from the diarrhea sample. The results of antimicrobial susceptibility testing showed that *A. caviae* SCAC2001 strain was resistant to meropenem, imipenem, cefepime, cefotaxime, ceftazidime, piperacillin-tazobactam, amoxicillin-clavulanic acid, gentamicin,

**Table 1** The Details of the Representative *A. Caviae* Strains Downloaded from Genbank

Isolate	Access Number	Country	Samples	Resistance Genes <sup>a</sup>
GSH8M-1 chromosomes and plasmids genome	AP019195.1	Japan	Wastewater	<i>bla</i> <sub>MOX-12</sub> , <i>bla</i> <sub>OXA-780</sub> , <i>mcr-3.18</i> , <i>bla</i> <sub>OXA-669</sub> , <i>aac(6)-Ia</i> , <i>aadA2</i> , <i>sul1</i> , <i>bla</i> <sub>KPC-2</sub> , <i>mph(A)</i>
WCWI-2 chromosomes genome	CP039832.1	People's Republic of China	Sewage	<i>bla</i> <sub>MOX-5</sub> , <i>bla</i> <sub>OXA-427</sub> , <i>catB3</i> , <i>cmlA1</i> , <i>floR</i> , <i>aac(6)-Ib-cr</i> , <i>qnrVC4</i> , <i>sul1</i> , <i>aac(6)-Ib3</i> , <i>aadA1</i> , <i>aph(3'')-Ib</i> , <i>aph(3')-Ia</i> , <i>aph(6)-Ic</i> , <i>dfrA14</i> , <i>dfrB4</i> , <i>bla</i> <sub>OXA-10</sub> , <i>tet(X4)</i>
T25-39 chromosomes genome	CP025706	People's Republic of China	Wastewater	<i>bla</i> <sub>MOX-6</sub> , <i>bla</i> <sub>OXA-427</sub>
R25-2 chromosomes genome	CP025777.1	People's Republic of China	Wastewater	<i>bla</i> <sub>MOX-6</sub> , <i>bla</i> <sub>OXA-427</sub> , <i>aph(3'')-Ib</i> , <i>aph(6)-Ic</i> , <i>tet(31)</i> , <i>sul2</i> , <i>floR</i>
NCTC12244 Chromosomes genome	LS483441.1	UK	N	<i>bla</i> <sub>MOX-6</sub> , <i>bla</i> <sub>OXA-427</sub>
Draft genome ZJ17-2	NXBR00000000.1	People's Republic of China	River water	<i>bla</i> <sub>MOX-6</sub> , <i>bla</i> <sub>OXA-427</sub> , <i>mcr-3</i> , <i>cat</i> , <i>tet(A)</i> , <i>aadA1</i> , <i>sul1</i>
Draft genome A23	LFXO00000000.1	People's Republic of China	Chicken sample	<i>bla</i> <sub>MOX-6</sub> , <i>bla</i> <sub>OXA-427</sub> , <i>catB3</i> , <i>mph(A)</i> , <i>bla</i> <sub>OXA-2</sub> , <i>bla</i> <sub>PER-3</sub> , <i>aadA1b</i> , <i>aadA1c</i> , <i>aac(6)-Ib-Hangzhou</i> , <i>aac(6)-Ib-cr</i> , <i>sul1</i>
Draft genome ZJ33-3	NXBW00000000.1	People's Republic of China	Human rectal swab	<i>bla</i> <sub>MOX-6</sub> , <i>bla</i> <sub>OXA-427</sub> , <i>tet(A)</i> , <i>mcr-3</i> , <i>cat</i> , <i>aadA1</i> , <i>sul1</i>
Draft genome AK245	JAAALU000000000.1	USA	Lake water	<i>bla</i> <sub>MOX-6</sub>
Draft genome Sch29	CAAKNG000000000.1	UK	Gastroenteritis samples	<i>bla</i> <sub>MOX-4</sub> , <i>bla</i> <sub>OXA-427</sub>
Draft genome LI2	JWJP00000000.1	Malaysia	Lake water	<i>bla</i> <sub>MOX-5</sub> , <i>mcr-3.15</i>
Draft genome TCO22	NMSG00000000.1	USA	Gut samples	<i>bla</i> <sub>MOX-6</sub> , <i>bla</i> <sub>OXA-427</sub> , <i>tetE</i> , <i>mcr3.12</i> , <i>mcr-3.15</i> , <i>aph(3')-Ia</i> , <i>mph(A)</i>
Draft genome strain D	VZQB00000000.1	South Africa	Seawater	<i>bla</i> <sub>MOX-6</sub> , <i>bla</i> <sub>OXA-427</sub>
Draft genome strain CHI29	MDSD00000000.1	Brazi	Seawater	<i>bla</i> <sub>MOX-6</sub>
Draft genome strain CHZ306	MDSC00000000.1	Brazi	Seawater	<i>bla</i> <sub>MOX-6</sub> , <i>bla</i> <sub>OXA-427</sub> , <i>tetE</i>
Draft genome strain ScAc2001	WUTZ00000000.1 (In this study)	People's Republic of China	Human sample	<i>bla</i> <sub>MOX-6</sub> , <i>bla</i> <sub>OXA-427</sub> , <i>aac(6)-Ib3</i> , <i>aadA1</i> , <i>bla</i> <sub>OXA-10</sub> , <i>catB3</i> , <i>sul1</i> , <i>mcr-3</i> , <i>dfrA5</i> , <i>bla</i> <sub>CTX-M-14</sub> , <i>bla</i> <sub>KPC-2</sub> , <i>bla</i> <sub>TEM-150</sub> , <i>bla</i> <sub>VEB-3</sub>

**Notes:** <sup>a</sup>The resistance genes were identified by the ResFinder (<https://cge.cbs.dtu.dk/services/ResFinder/>) (minimum threshold for identity, 80%; minimum coverage, 60%).  
**Abbreviations:** N, Not shown.

amikacin, aztreonam, erythromycin, chloramphenicol, colistin, ciprofloxacin and trimethoprim-sulfamethoxazole and sensitive to the tigecycline and fosfomycin. While the negative control *E.coli* J53 was sensitive to all the test antibiotics. To the best of our knowledge, the *Aeromonas species* isolated from environmental water samples received widespread

attention several years ago. However, there have been more reports of the *Aeromonas species* infection in humans worldwide in recent years,<sup>28,29</sup> because the *Aeromonas species* have always been identified as enteric pathogens which can lead to severe diarrhoea.<sup>9</sup> Unfortunately, what's more serious is that the carbapenem-

resistant and/or colistin-resistant strains produce emergencies in some countries and pose a serious threat to infectious diseases control and clinical treatment.<sup>30,31</sup>


## Draft Genome Characterization of SCAC2001 and Phylogenetic Analysis

A total of 1 Gigabases pairs (Gbp) of raw genome data was obtained from the Illumina system. Thereafter, we got about 800 million bases (Mbp) of clean data from the 1Gbp raw genome data by using the readfq (version 10). The 800Mbp clean data was assembled into a 340 scaffolds draft genome sequence of *Aeromonas caviae* strain SCAC2001 by the SOAP denovo, with a G+C content of 61.09%, for a total of 4487370bp. We predicted 4265 protein-coding sequences (CDS) in the draft genome. The genome encodes 103 tRNAs and 14 rRNAs, which contains five copies of the 5S rRNA gene, five copies of the 16S rRNA gene, and four copies of the 23S rRNA

genes. The core genome-based phylogenetic analysis showed that the *A. caviae* SCAC2001 strain was highly similar to the *A. caviae* strain R25-2 (Genbank accession number: CP025777.1) and T25-39 (Genbank accession number: CP025706), but distant from the clade grouped by the *mcr-3* carrying strain ZJ33-3 (NXBW00000000.1) and ZJ17-2 (NXBR00000000.1), and the *mcr-3* and *bla<sub>KPC-2</sub>* co-carried strain GSH8M-1 (AP019195.1)<sup>32</sup> (Figure 1).

## Identification of the Resistance Genes, Virulence Factors and PHI Genes

A total of 13 antibiotic drug resistance genes including the  $\beta$ -lactams (*bla<sub>KPC</sub>*, *bla<sub>CTX-M-14</sub>*, *bla<sub>TEM-1</sub>*, *bla<sub>OXA-10</sub>*, *bla<sub>OXA-427</sub>*, *bla<sub>VEB-3</sub>* and *bla<sub>MOX-6</sub>*), aminoglycosides (*aadA1*), fluoroquinolones (*aac(6)-Ib-cr*), phenicol resistance (*catB3*), sulfonamide (*sulI*) and trimethoprim (*dfrA5*) and colistin resistance gene (*mcr-3.3*) were detected in the genome of *Aeromonas*



**Figure 1** The phylogenetic tree was constructed by aligning the core genome of *Aeromonas caviae* SCAC2001 strain with 15 other representative *A. caviae* strains.

**Table 2** Distribution of the Resistance Genes in *Aeromonas caviae* SCAc2001

Resistance Gene	Identity (%)	Query/Length	Scaffold	Position in Scaffold	Predicted Phenotype	Accession Number
aac(6)-Ib3	100	555/555	Scaffold231	2433-2987	Fluoroquinolone and aminoglycoside resistance	X60321
aadA1	100	792/792	Scaffold231	29-820	Aminoglycoside resistance	JQ414041
bla <sub>OXA-10</sub>	100	801/801	Scaffold231	837-1637	Beta-lactam resistance Alternate name; PSE-2	J03427
catB3	100	633/633	Scaffold231	1706-2338	Phenicol resistance	U13880
sulI	100	840/840	Scaffold310	25-864	Sulphonamide resistance	U12338
mcr-3.3	100	1623/1623	Scaffold216	1393-3015	Polymyxin resistance	MF495680
dfpA5	100	474/474	Scaffold258	1177-1650	Trimethoprim resistance	X12868
bla <sub>CTX-M-14</sub>	100	876/876	Scaffold180	3068-3943	Beta-lactam resistance	AF252622
bla <sub>KPC-2</sub>	100	882/882	Scaffold72	15,640-6521	Beta-lactam resistance	AY034847
bla <sub>MOX-6</sub>	97.83	1153/1152	Scaffold128	10,749-11,900	Beta-lactam resistance AmpC-type	GQ152601
bla <sub>OXA-427</sub>	86.43	796/795	Scaffold165	5530-6324	Beta-lactam resistance	KX827604
bla <sub>TEM-150</sub>	99.82	570/861	Scaffold72	14,849-15,418	Beta-lactam resistance	AM183304
bla <sub>VEB-3</sub>	100	900/900	Scaffold240	750-1649	Beta-lactam resistance	AY536519

*caviae* strain ScAc2001c (Table 2). To the best of our knowledge, many reports are showing that the multi-drug-resistant *Aeromonas* species have been isolated from clinical, animal, food, and environmental water samples (Table 1). These isolates carried many more types of antibiotic resistance genes, especially the co-carried carbapenem- and colistin- resistance gene may be a huge risk for infectious disease control. Also, as

shown in Table 3, *Aeromonas caviae* strain ScAc2001 carries 54 putative virulence factors including type IV pilus, fimbria, flagellarthe and hemolysin A. What's more, we also identified 12 PHI genes including *csrA*, *lrp*, *crp*, *iscU*, *arcA*, *metJ*, *pykF*, *lon*, *dksA*, *fur*, *greB* and *hfq* from the genome (Table 4). It proved that these PHI genes are associated with diseases such as diarrheal, meningitis and urinary tract infections.<sup>33</sup> These

**Table 3** Distribution of the Virulence Factors in *Aeromonas caviae* Strain SCAc2001<sup>a</sup>

Virulence Factor	Scaffold	Position in Scaffold	Identity(%)	Characteristic
<i>tppE</i>	Scaffold14	20,853-21,356	99.4	Type IV pilus pseudopilin
<i>tppB</i>	Scaffold14	20,443-20,856	99.3	Type IV pilus modification protein PilV
<i>tppA</i>	Scaffold14	15,090-15,491	99.2	Type IV pilin
<i>tapY1</i>	Scaffold14	15,500-18,853	99.1	Type IV pilus biogenesis protein
<i>cheW-2</i>	Scaffold65	270-758	98.1	Chemotaxis protein CheW
<i>pomA2</i>	Scaffold18	20,514-21,245	97.9	Chemotaxis protein PomA
<i>hlyA</i>	Scaffold96	11,532-12,851	97.5	Hemolysin A
<i>exeG</i>	Scaffold3	14,115-14,546	97.2	General secretion pathway protein G
<i>exeE</i>	Scaffold3	15,920-17,425	96.2	General secretory pathway protein E
<i>tapT</i>	Scaffold84	9088-10,044	95.9	Twitching ATPase
<i>hutZ</i>	Scaffold20	28,013-28,570	95.7	Heme iron utilization protein
<i>exeF</i>	Scaffold3	14,698-15,918	95.6	General secretion pathway protein F
<i>tapB</i>	Scaffold45	118-1098	95.3	Type IV-A pilus assembly ATPase PilB
<i>exel</i>	Scaffold3	13,085-13,429	94.7	General secretion pathway protein I
<i>hutC</i>	Scaffold20	30,092-31,120	94.4	ABC-type heme transporter, permease protein
<i>tapB</i>	Scaffold288	85-870	94.4	Type IV pilus assembly protein TapB
<i>fleR/fliC</i>	Scaffold148	2516-3850	93.5	Transcriptional activator
<i>amoA</i>	Scaffold8	60,005-61,177	92.9	Isochorismate synthases
<i>tapW</i>	Scaffold25	28,637-29,758	92.8	Tfp pilus assembly protein, ATPase PilU
<i>tapC</i>	Scaffold45	1211-2467	92.5	Type 4 fimbrial assembly protein PilC
<i>fliB</i>	Scaffold148	1411-2445	92.2	Two-component system flagellar sensor histidine kinase FliB

**Notes:** <sup>a</sup>The resistance genes were identified by the virulence factors which were identified by the VFAnalyzer (<http://www.mgc.ac.cn/VFs/main.htm>).

**Table 4** Distribution of the PHI Genes in *Aeromonas caviae* Strain SCAc2001<sup>a</sup>

Gene Name	Scaffold	Position in Scaffold	Identity (%)	PHI-Base Accession	Disease Name	Gene Function
<i>csrA</i>	Scaffold 19	1661–1849	96.6	PHI:5061	Diarrheal diseases	Small RNA-binding protein involved in the regulation of a wide range of cellular processes.
<i>lrp</i>	Scaffold 106	12,874–13,143	92.1	PHI:6497	No data found	Global transcription factor
<i>crp</i>	Scaffold 20	14,048–14,686	89.6	PHI:2684	Food poisoning	Regulators of systemic infection
<i>iscU</i>	Scaffold 22	19,716–20,099	85.6	PHI:6892	Salmonellosis	Fe-S cluster sensor
<i>arcA</i>	Scaffold 24	11,882–12,601	84.8	PHI:6532	Meningitis	Aerobic respiration control protein
<i>metJ</i>	Scaffold 36	1313–1636	83.2	PHI:2695	Blackleg disease	Repressor of the methionine biosynthesis regulon
<i>pykF</i>	Scaffold 51	7554–8978	82.8	PHI:3134	Gut-associated diseases; diarrhoea; enteritis; colitis	Part of the pyruvate - tricarboxylic acid cycle node
<i>lon</i>	Scaffold 68	14,376–16,730	82.6	PHI:7160	Fire blight	Protease
<i>dkcA</i>	Scaffold 40	12,676–13,059	81.7	PHI:6508	Salmonellosis	Response Regulator
<i>fur</i>	Scaffold 49	144–572	81.7	PHI:4887	Urinary tract infections	global ferric uptake regulator
<i>greB</i>	Scaffold 83	192–665	81.5	PHI:7131	Salmonellosis	Transcription elongation factor
<i>hfq</i>	Scaffold 27	23,382–23,570	81.1	PHI:4064	Pneumonic plague	RNA-binding protein

results indicate the emergence of the co-location of a large number of resistance genes, a variety of virulence factors, and the PHI genes carrying *Aeromonas caviae* ScAc2001-like strain is a serious issue for public health.

## The Genetic Context of the Resistance Genes

Silicon analysis showed that the resistance gene *bla*<sub>KPC-2</sub> was located in scaffold72 and that it also carried the ESBL gene *bla*<sub>TEM-1</sub>. Sequence analysis showed that scaffold72 had 68%, 59% and 59% query cover and 99.95%, 99.98% and 99.98% sequence similarities with plasmid pGSH8M-1-2 (AP019197),<sup>32</sup> plasmid p1713-KPC (MH624132) and plasmid p198-KPC (MH624131) at nucleotide level, respectively. The linear structure of this genetic context is repA-orf-*klcA*-*korC*-*ISKpn6*-*bla*<sub>KPC</sub>-*bla*<sub>TEM</sub>-*ISKpn27*. However, the other sequence of scaffold72 was unique to *Aeromonas caviae* strain ScAc2001. The colistin resistance gene *mcr-3.3* was located in the scaffold216. Scaffold216 had 81%, 81% and 78% query cover and 99.89%, 99.06%, and 98.29% sequence similarities with the plasmid pGSH8M-1-2 (AP019197), *Aeromonas* ASNIH7 chromosome genome (CP026226), *Aeromonas veronii* 17ISAe chromosome genome (CP028133) at nucleotide level, respectively. The result showed the *mcr-3.3* carrying scaffold maybe derive from the plasmid and *Aeromonas* chromosome genome. In any case, the plasmid or chromosome borne colistin resistance gene *mcr* carried by the carbapenem-resistant -microorganisms is a risk factor in clinical control.

The ESBLs gene *bla*<sub>CTX-M-14</sub> carrying scaffold180 had 100%, 98% and 98% query cover and 100%, 99.96% and 99.95% sequence similarities with the *Vibrio* Vb0624 chromosome genome (CP041202), plasmid pKP96 (EU195449)<sup>34</sup> and plasmid pEC224\_4 (CP018944) at nucleotide level, respectively. Scaffold231 carried four resistance genes (*catB3*, *bla*<sub>OXA-10</sub>, *aadA1* and *aac(6)-Ib3*). Sequence analysis showed that scaffold231 had 99%, 99% and 99% query cover and 99.97%, 99.97% and 99.97% sequence similarities with the plasmid pC45\_002 (CP042553), plasmid pE33\_002 (CP042519) and plasmid pEC224\_4 (CP042480) at nucleotide level, respectively. It speculated that scaffold231 was derived from the plasmid.

The ESBLs gene *bla*<sub>MOX-6</sub> was carried by the scaffold128. Sequence analysis showed that scaffold128 had 99%, 98% and 98% query cover and 98.3%, 98.17% and 98.15% sequence similarities with the *Aeromonas caviae* GSH8M-1 complete genome (AP019195.1),<sup>32</sup> *Aeromonas caviae* strain T25-39 chromosome genome (CP025706.1) and *Aeromonas* strain R25-2 chromosome genome (CP025777.1) at nucleotide level, respectively.

Scaffold165 carried one class D β-lactamases (CHDLs) resistance gene *bla*<sub>OXA-427</sub>. Sequence analysis showed that scaffold165 had 98%, 98%, and 95% query cover and 95.83%, 95.83% and 95.67% sequence similarities with the *Aeromonas caviae* R25-2 genome (CP025777.1), *Aeromonas caviae* strain T25-39 genome (CP025706) and *Aeromonas caviae* GSH8M-1 genome (AP019195.1) at nucleotide level, respectively. This result showed that the resistance gene

**Table 5** Overall Features of the *Aeromonas caviae* SCAC2001 Genomic Islands and Prophages

Genomic Islands and Prophages		Location (Start-End)	Length (bp)	G+C%	Closest Match in Genbank (Query Cover and Identity)
Genomic Islands	GI_SCAC2001-1	Scaffold26 (22,433–29,614)	7182	54.58	<i>Aeromonas</i> sp. ASNIH7 chromosome genome (74%, 91.04%)
	GI_SCAC2001-2	Scaffold42 (14,540–21,489)	6950	63.42	<i>Aeromonas</i> sp. ASNIH3 chromosome genome (100, 99.07%)
	GI_SCAC2001-3	Scaffold57 (7135–15,483)	8349	62.27	<i>Aeromonas caviae</i> FDAARGOS_72 genome (98%, 99.19%)
	GI_SCAC2001-4	Scaffold5 (42,875–50,049)	7175	62.45	<i>Aeromonas caviae</i> WCWI-2 genome (100%, 98.27%)
Prophages	Pp_SCAC2001-1	Scaffold159 4884–7973	3090	58.93	plasmid pMCR5_045096 (100%, 100%)
	Pp_SCAC2001-2	Scaffold158 413–7151	6739	63.6	<i>Aeromonas</i> sp. ASNIH2 chromosome genome (100%, 99.6%)
	Pp_SCAC2001-3	Scaffold209 146–2968	2823	61.18	<i>Aeromonas caviae</i> GSHB1-1 genome (100%, 98.87%)
	Pp_SCAC2001-4	Scaffold38 8991–27,083	18,093	65.37	<i>Aeromonas</i> sp. ASNIH2 chromosome genome (98%, 97.62%)
	Pp_SCAC2001-5	Scaffold202 221–4124	3904	61.94	<i>Aeromonas caviae</i> strain R25-6 genome (100%, 99%)
	Pp_SCAC2001-6	Scaffold201 95–3149	3055	46.94	<i>Aeromonas media</i> VVS genome (98%, 99.87%)
	Pp_SCAC2001-7	Scaffold128 236–10,450	10,215	63.55	<i>Aeromonas caviae</i> FDAARGOS_72 genome (100%, 98.47%)
	Pp_SCAC2001-8	Scaffold129 479–4423	3945	58.93	<i>Aeromonas hydrophila</i> strain WCHAH045096 chromosome genome (80%, 94.6%)

*bla*<sub>OXA-427</sub> is chromosome borne. Another resistance gene *bla*<sub>VEB-3</sub> was carried by the scaffold240. Sequence analysis showed that scaffold240 had 90% and 90% query cover and 99.96% and 99.95% sequence similarities with *Aeromonas hydrophila* strain MX16A genome (CP018201) and JM45 plasmid p1 (CP006657.1) at nucleotide level, respectively. The context genetic of this resistance gene is the *Int1-bla*<sub>VEB-3-IS600-IS26</sub>. These results indicated co-carriage of a large number of resistance genes in genome making *Aeromonas caviae* strain highly resistant to almost all kinds of commonly used antibiotics, and brings a serious challenge for resistance control and clinical treatment.

## Characterization of the Genomic Islands and Prophages

As shown in Table 5, four genomic islands, named GI\_SCAC2001-1 to GI\_SCAC2001-4, were identified by the software IslandPATH-DIMOB, IslandPick, and SIGI-HMM. Silicon analysis showed that the length of the four genomic islands were 7,182, 6,950, 8,349 and 7,175bp with the G+C context of 54.585%, 63.425%, 62.275% and 62.45%, respectively. The sequences of four genomic islands are all the closest match to the *Aeromonas* sp. chromosome genome sequence in genbank. A total of eight prophages (length >2kbp), named Pp\_SCAC2001-1 to Pp\_SCAC2001-8, were identified by phiSpy. The size of the eight prophages ranged from 2823bp to 18,093bp with the average G+C content of 46.94%–65.37%, respectively. Among them, one of the prophage sequences' closest match was the corresponding region of plasmid pMCR5\_045096 and seven of the prophages were the closest match to the *Aeromonas* sp. chromosome genome in genbank. This indicated that the mobile genetic elements (genomic islands and prophages) can be excised and integrated from the chromosome and mobile genetic elements into each other. However, no resistance genes or virulence genes were found in the genomic islands and prophages. To the best of our knowledge, the mobile genetic elements (including the genomic islands and prophages) are effective integrative elements in bacterial evolution including the resistance, virulence and some function genes.<sup>35–37</sup>

## Characterization of Secondary Metabolism Products Coding Clusters

In this study, three secondary metabolism products coding clusters (nonribosomal peptide synthetases (nrps), hserlactone and bacteriocin), which are responsible for the biosynthesis of

**Table 6** Characterization of Secondary Metabolism Products' Coding Clusters

Cluster Name	Location (Start-End)	Length (bp)	G+C%	ORF Number	Predicted Protein
nrps	Scaffold8 (25,698–64,211)	38,514	65.27	27	TerC family protein, AroG, AroA, SerC, FixB, FixA, FAD-binding protein, CorA, DUF2919 family protein, hypothetical protein, amonabactin ABC transporter permease subunit 2, Amonabactin ABC transporter permease subunit 1, Amonabactin ABC transporter ATP-binding protein, 4'-phosphopantetheinyl transferase superfamily protein, ABC transporter substrate-binding protein, AmoH, AmoG, DhbA, Amof, Isochorismatase family protein, EntE isochorismate synthase, Lipoprotein, hypothetical protein
hserlactone	Scaffold70 (246–17,682)	17,438	63.52	13	HldE, LpxL, bifunctional 2',3'-cyclic-nucleotide 2'-phosphodiesterase/3'-nucleotidase, LysE family transporter, LuxR family transcriptional regulator, GNAT family N-acetyltransferase, ArgP/LysG family DNA-binding transcriptional, exoribonuclease II, PTS mannitol transporter subunit IICBA, mannitol-1-phosphate 5-dehydrogenase, MltR family transcriptional regulator, sodium:proton antiporter, hypothetical protein
bacteriocin	Scaffold139 (89–9975)	9887	64.36	12	Hypothetical protein, BCCT family transporter, DUF2282 domain-containing protein, DUF692 family protein, DUF2063 domain-containing protein, DoxX family protein, GNAT family N-acetyltransferase, PAS domain S-box protein, DUF3332 domain-containing protein, HD domain-containing protein

secondary metabolic products, were predicted using the search tool antiSMASH. The silicon analysis showed that the length of the three secondary metabolism coding clusters were 38,514, 17,438, and 9887 bp with the G+C context of 65.27%, 63.52% and 64.36%, respectively. Sequence analysis showed that the three putative gene clusters carrying 27, 13 and 12 ORFs, respectively (Table 6). It's proved that a large number of pharmaceutical agents, microbial natural products including the sterigmatocystin (carcinogen), penicillin vancomycin and (antibiotic), lovastatin (antihypercholesterolemic agent), and cyclosporin A (anti-inflammatories and immunosuppressants) are synthesized by the diverse array of the secondary metabolism products coding clusters. Researching the characterization of secondary metabolism products coding clusters can be seen as one of the potential ways to research the new drugs.<sup>38</sup>

## Conclusion

Isolates of *Aeromonas caviae* ScAc2001 harboring a lot of resistance genes including the carbapenem-resistant gene *bla<sub>KPC</sub>*, colistin resistance gene *mcr-3.3* and other  $\beta$ -lactams, aminoglycosides, fluoroquinolones, phenicol resistance (*catB3*), sulfonamide and a variety of virulence factors, PHI genes, four genomic islands and eight prophages may result in a possible risk to public health.

## Nucleotide Sequence Accession Numbers

This Whole Genome Shotgun project has been deposited at DDBJ/ENA/GenBank under the accession WUTZ000000001.

## Ethical Statement

This study was approved by the Experimentation Ethics Committee of Southwest Medical University.

## Funding

This work was supported by the National Natural Science Foundation of China (31500114) and United Funds of Luzhou and Southwest Medical University [2018LZ XNYD-ZK51].

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Shen Y, Xu C, Sun Q, et al. Prevalence and genetic analysis of *mcr-3*-positive *Aeromonas* species from humans, retail meat, and environmental water samples. *Antimicrob Agents Chemother.* 2018;62(9):e00404–e00418.
- Chen F, Deng X, Wang Z, Wang L, Wang K, Gao L. Treatment of severe ventriculitis caused by extensively drug-resistant *Acinetobacter baumannii* by intraventricular lavage and administration of colistin. *Infect Drug Resist.* 2019;12:241–247. doi:10.2147/IDR.S186646



3. Liu YY, Wang Y, Walsh TR, et al. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis*. 2016;16(2):161–168. doi:10.1016/S1473-3099(15)00424-7
4. Wang Y, Zhang R, Li J, et al. Comprehensive resistome analysis reveals the prevalence of NDM and MCR-1 in Chinese poultry production. *Nat Microbiol*. 2017;2(4):16260. doi:10.1038/nmicrobiol.2016.260
5. Snesrud E, Maybank R, Kwak YI, Jones AR, Hinkle MK, McGann P. Chromosomally encoded mcr-5 in colistin-nonsusceptible *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother*. 2018;62(8):e00679–e00618. doi:10.1128/AAC.00679-18
6. Nukui Y, Ayibieke A, Taniguchi M, et al. Whole-genome analysis of EC129, an NDM-5-, CTX-M-14-, OXA-10- and MCR-1-co-producing *Escherichia coli* ST167 strain isolated from Japan. *J Glob Antimicrob Resist*. 2019;18:148–150. doi:10.1016/j.jgar.2019.07.001
7. Wang Q, Zhang P, Zhao D, et al. Emergence of tigecycline resistance in *Escherichia coli* co-producing MCR-1 and NDM-5 during tigecycline salvage treatment. *Infect Drug Resist*. 2018;11:2241–2248. doi:10.2147/IDR.S179618
8. Lin YC, Kuroda M, Suzuki S, Mu JJ. Emergence of an *Escherichia coli* strain co-harboring mcr-1 and blaNDM-9 from a urinary tract infection in Taiwan. *J Glob Antimicrob Resist*. 2019;16:286–290. doi:10.1016/j.jgar.2018.10.003
9. Altwegg M. *Aeromonas caviae*: an enteric pathogen? *Infection*. 1985;13(5):228–230. doi:10.1007/BF01667217
10. Yang S, He T, Sun J, Sun S. Distinct antimicrobial resistance profiling of clinically important *Aeromonas* spp. in southwest China: a seven-year surveillance study. *Infect Drug Resist*. 2019;12:2971–2978. doi:10.2147/IDR.S216926
11. Goncalves Pessoa RB, de Oliveira WF, Marques DSC, Dos Santos Correia MT, de Carvalho E, Coelho L. The genus *Aeromonas*: a general approach. *Microb Pathog*. 2019;130:81–94. doi:10.1016/j.micpath.2019.02.036
12. Figueras MJ, Latif-Eugenin F, Ballester F, et al. ‘*Aeromonas intestinalis*’ and ‘*Aeromonas enterica*’ isolated from human faeces, ‘*Aeromonas crassostreae*’ from oyster and ‘*Aeromonas aquatilis*’ isolated from lake water represent novel species. *New Microbes New Infect*. 2017;15:74–76. doi:10.1016/j.nmni.2016.11.019
13. Fu L, Wang S, Zhang Z, et al. Co-carrying of KPC-2, NDM-5, CTX-M-3 and CTX-M-65 in three plasmids with serotype O89:H10 *Escherichia coli* strain belonging to the ST2 clone in China. *Microb Pathog*. 2019;128:1–6. doi:10.1016/j.micpath.2018.12.033
14. Liu X, Zhang J, Li Y, et al. Diversity and frequency of resistance and virulence genes in bla KPC and bla NDM co-producing *Klebsiella pneumoniae* strains from China. *Infect Drug Resist*. 2019;12:2819–2826. doi:10.2147/IDR.S214960
15. Rebelo AR, Bortolaia V, Kjeldgaard JS, et al. Multiplex PCR for detection of plasmid-mediated colistin resistance determinants, mcr-1, mcr-2, mcr-3, mcr-4 and mcr-5 for surveillance purposes. *Eurosurveillance*. 2018;23(6).
16. Li R, Zhu H, Ruan J, et al. De novo assembly of human genomes with massively parallel short read sequencing. *Genome Res*. 2010;20(2):265–272. doi:10.1101/gr.097261.109
17. John B, Alexandre L, Mark B. GeneMarkS: a self-training method for prediction of gene starts in microbial genomes. Implications for finding sequence motifs in regulatory regions. *Nucleic Acids Res*. 2001;29:2607.
18. Ea Z, Henrik H, Salvatore C, et al. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother*. 2012;67(11):2640–2644. doi:10.1093/jac/dks261
19. Winnenburg R, Baldwin TK, Urban M, Rawlings C, Köhler J, Hammond-Kosack KE. PHI-base: a new database for pathogen host interactions. *Nucleic Acids Res*. 2006;34(Database issue):D459. doi:10.1093/nar/gkj047
20. Alexandros S. RAXML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*. 2014;30(9):9.
21. Sudhir K, Glen S, Koichiro T. MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol*. 2016;33(7):7.
22. Angiuoli SV, Salzberg SL. Mugsy: fast multiple alignment of closely related whole genomes. *Bioinformatics*. 2011;27(3):334–342. doi:10.1093/bioinformatics/btq665
23. Bertelli C, Brinkman F. Improved genomic island predictions with IslandPath-DIMOB. *Bioinformatics*. 2018;34(13):2161–2167. doi:10.1093/bioinformatics/bty095
24. Langille MG, Hsiao WW, Brinkman FS. Evaluation of genomic island predictors using a comparative genomics approach. *BMC Bioinformatics*. 2008;9(1):329. doi:10.1186/1471-2105-9-329
25. Waack S, Keller O, Asper R, et al. Score-based prediction of genomic islands in prokaryotic genomes using hidden Markov models. *BMC Bioinformatics*. 2006;7(1):1–12. doi:10.1186/1471-2105-7-142
26. Langille MGI, Brinkman FSL. IslandViewer: an integrated interface for computational identification and visualization of genomic islands. *Bioinformatics*. 2009;25(5):664–665. doi:10.1093/bioinformatics/btp030
27. Kai B, Thomas W, Chevrette MG, et al. antiSMASH 4.0—improvements in chemistry prediction and gene cluster boundary identification. *Nucleic Acids Res*. 2017;45(W1):W36–W41. doi:10.1093/nar/gkx319
28. Khajanchi BK, Fadl AA, Borchardt MA, et al. Distribution of virulence factors and molecular fingerprinting of *Aeromonas* species isolates from water and clinical samples: suggestive evidence of water-to-human transmission. *Appl Environ Microbiol*. 2010;76(7):2313–2325. doi:10.1128/AEM.02535-09
29. Mendes-Marques CL, Nascimento LMD, Theophilo GND, Hofer E, Melo Neto OPD, Leal NC. Molecular characterization of *Aeromonas* spp. and *Vibrio cholerae* O1 isolated during a diarrhea outbreak. *Rev Do Inst De Med Trop De São Paulo*. 2012;54(6):299–304. doi:10.1590/S0036-46652012000600001
30. Adler A, Assous MV, Paikin S, et al. Emergence of VIM-producing *Aeromonas caviae* in Israeli hospitals. *J Antimicrob Chemother*. 2014;69(5):1211–1214. doi:10.1093/jac/dkt505
31. Anandan S, Gopi R, Ragupathi NKD, Sethuvel DPM, Veeraraghavan B. First report of bla OXA-181 mediated carbapenem resistance in *Aeromonas caviae* in association with pKP3-A: threat for rapid dissemination. *J Glob Antimicrob Resist*. 2017;10:310–314.
32. Sekizuka T, Inamine Y, Segawa T, Hashino M, Kuroda M, Kuroda M. Potential KPC-2 carbapenemase reservoir of environmental *Aeromonas hydrophila* and *Aeromonas caviae* isolates from the effluent of an urban wastewater treatment plant in Japan. *Environ Microbiol Rep*. 2019;11(4):589–597. doi:10.1111/1758-2229.12772
33. Ye F, Yang F, Yu R, et al. Molecular basis of binding between the global post-transcriptional regulator CsrA and the T3SS chaperone CesT. *Nat Commun*. 2018;9(1):1196. doi:10.1038/s41467-018-03625-x
34. Shen P, Jiang Y, Zhou Z, Zhang J, Yu Y, Li L. Complete nucleotide sequence of pKP96, a 67 850 bp multiresistance plasmid encoding qnrA1, aac(6'')-Ib-cr and blaCTX-M-24 from *Klebsiella pneumoniae*. *J Antimicrob Chemother*. 2008;62(6):1252–1256. doi:10.1093/jac/dkn397
35. Boyd EF, Almagro-Moreno S, Parent MA. Genomic islands are dynamic, ancient integrative elements in bacterial evolution. *Trends Microbiol*. 2009;17(2):0–53.

36. Penn K, Jenkins C, Nett M, et al. Genomic islands link secondary metabolism to functional adaptation in marine Actinobacteria. *Isme J.* 2009;3(10):1193–1203. doi:10.1038/ismej.2009.58
37. Bellanger X, Payot S, Leblond-Bourget N, Guédon G. Conjugative and mobilizable genomic islands in bacteria: evolution and diversity. *FEMS Microbiol Rev.* 2013;38(4):720–760.
38. Wang B, Guo F, Dong SH, Zhao H. Activation of silent biosynthetic gene clusters using transcription factor decoys. *Nat Chem Biol.* 2019;15(2):111–114. doi:10.1038/s41589-018-0187-0

## Infection and Drug Resistance

Dovepress

### Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>