

Nomogram Models for Predicting Delirium of Patients in Emergency Intensive Care Unit: A Retrospective Cohort Study

Yu Shi¹, Hai Wang², Li Zhang¹, Ming Zhang¹, Xiaoyan Shi¹, Honghong Pei¹, Zhenghai Bai¹

¹Emergency Department & EICU, The Second Affiliated Hospital of Xi'an Jiaotong University, Xi'an, Shaan Xi, 710004, Peoples' Republic of China;

²Department of Hepatobiliary Surgery, The First Affiliated Hospital of Xi'an Jiaotong University, Xi'an, Shaan Xi, 710061, Peoples' Republic of China

Correspondence: Honghong Pei; Zhenghai Bai, The Emergency Department & EICU, The Second Affiliated Hospital of Xi'an Jiaotong University, No. 157, Xiwu Road, Xincheng District, Xi'an, Shaanxi, Peoples' Republic of China, Email peihhjz@163.com; baizhenghaijz2021@163.com

Background: Intensive care unit (ICU) delirium is one of the most common clinical syndromes that results in many adverse events that affect patients, families, and hospitals. To date, there has been no tool for effectively predicting the occurrence of delirium in emergency intensive care unit (EICU) patients.

Methods: We conducted a retrospective cohort study and constructed a prediction model for 319 patients in EICU, who met our inclusion criteria. We analyzed the relationship between patients' clinical data within 24 hours of admission and delirium, applied univariate and multivariate logistic regression analyses to select the most relevant variables for construction of nomogram models, then applied bootstrapping for internal validation.

Results: A total of five variables, namely stomach and urinary tubes, as well as sedative, mechanical ventilation and APACHE-II scores, were selected for model construction. We generated a total of five sets of models (three sets of construction models and two sets of internal verification models), with similar predictive value. The optimal model was selected, and together with the 5 variables used to construct a nomogram. The AUC of the MFP model in all patients was 0.76 (0.70, 0.82), whereas that in non-elderly patients (<60 years old) for the full model was 0.83 (0.74, 0.91). In elderly patients (≥ 60 years old), the AUC of the MFP model was 0.82 (0.73, 0.91).

Conclusion: Overall, the five-marker-based prognostic tool, established herein, can effectively predict the occurrence of delirium in EICU patients.

Keywords: area under curve, delirium, emergency intensive care unit, model, prediction

Introduction

Although advancements in the field of scientific medicine have greatly improved treatment of critical diseases, ICU delirium still occurs in ICU.¹ Delirium refers to an acute and volatile change in mental state, which is characterized by disturbance of consciousness, and is often accompanied by sleep-wake cycle disorders, varying degrees of attention deficiency, as well as cognitive and affective disorders.² Delirium that occurs in ICU patients is known as ICU-delirium.

Previous studies have shown that the incidence of ICU-delirium is high, with a gradual growth trend.^{3,4} This incidence also varies across different patients. General ICU patients have an ICU-delirium incidence of 15–80%,^{5,6} whereas those under mechanical ventilation (MV) have a 60–80%.^{7–10}

Generally, the occurrence of ICU-delirium is associated with various adverse events.^{11,12} For example, patients experience short-term disease recovery time and long-term mental health problems,¹³ families encounter a high burden of disease and psychological pressure,¹⁴ while medical resources are subjected to increased physical and mental pressure for workers as well as high medical costs.¹⁵ Therefore, evaluating potential risk factors of ICU-delirium, as well as constructing prediction models, and development of prevention approaches are imperative to management of this condition and alleviation of suffering.

Although the risk factors and predictive models of ICU-delirium have been previously reported, little was known regarding the occurrence of ICU-delirium in EICU patients owing to the fact that most focus has been on ICU and coronary care (CCU) wards. EICU is a new organization and management model, an extremely important part of emergency medicine, and a brand-new sign of the modernization of the emergency department.^{16,17} In addition, several diseases have been reported in EICU patients, while composition of systemic diseases is different from that in ICU and CCU.¹⁶ Since the existing research evidences may not be suitable for EICU patients, we sought to identify risk factors of ICU-delirium within 24 hours of admission, and construct a nomogram model for predicting the risk of EICU patients, with the aim of providing a reference for early identification and intervention.

Methods

Study Design and Data Sources

This was a retrospective cohort study, comprising data collected from medical records of patients in the EICU at the Second Affiliated Hospital of Xi'an Jiaotong University between July 1st 2020 and December 1st 2020. The evaluation method used herein was one of the daily tasks performed in the ICU. All study procedures were performed in accordance with the guideline of the ethics committee of the Second Affiliated Hospital of Xi'an Jiaotong University, approval number 2021006. The research complied with the Declaration of Helsinki. In addition, the detailed information of participants was not collected in our research database to ensure anonymous usage data.

Inclusion and Exclusion Criteria

Participants who met the following criteria were enrolled in the study: 1) aged ≥ 18 years old; and 2) had EICU stay time ≥ 24 h. Conversely, those who met the following criteria were excluded: 1) had a history of mental illness or dementia; 2) exhibited severe hearing or visual impairment; 3) had severe mental retardation or aphasia; 4) those who were continuously in a coma or deep sedation, with a Glasgow Coma Scale (GSC) score less than 8 points; and 5) those who already had delirium when entering the EICU.

Participants

A total of 450 patients were screened, between July 1st to December 1st, 2020, of which 131 were excluded because of the following reasons: they were aged < 18 years old ($N = 5$); had EICU stay times < 24 h ($N = 14$); were severely mentally handicapped or had aphasia ($N = 4$); were in deep coma ($N = 30$); exhibited delirium prior to admission in the EICU ($N = 64$); and were missing important data ($N = 14$). Finally, a total of 319 cases were included in our study (Figure 1).

Calculation of Sample Size

Before the formal study, we conducted a pre-experiment with 50 samples and predicted that 5 factors would be required to establish the ICU delirium prediction model. Each factor was validated with a minimum of 5–10 cases. Based on previous data, the proposed incidence rate is 20%, and assuming that the follow-up loss rate was 10%, the minimum sample size required for this study was 278 ($(5 * 10 / 0.2) / 0.9 = 278$). A total of 319 samples were included in this study, which met the requirements.

Data Collection

Each patient's relevant medical records were collected, within 24 h of admission, and the following information recorded: (1) Demographic and clinical data, such as age, sex, diagnosis type, educational level, body mass index (BMI), smoking, drinking, diabetes, hypertension, sepsis, heart failure, and electrolyte disorders; (2) previous clinical treatment measures, including stomach and urinary tubes, central venous catheter (CVC), sedative, MV, and vasoactive drugs; and (3) disease severity index: Acute Physiology and Chronic Health Evaluation (APACHE-II) score and Sequential Organ Failure Assessment (SOFA) score.




Figure 1 Flow chart of patient recruitment.

Assessment of Delirium

Delirium was assessed using the Richmond Agitation-Sedation Scale (RASS) score and Confusion Assessment Method for ICU (CAM-ICU). Particularly, we assessed delirium by CAM-ICU if a patient's RASS score was -3 to $+4$, but not in those with a score of -4 to -5 . Delirium assessment based on CAM-ICU was performed based on four aspects, namely acute onset and fluctuating causes of symptoms, inattention, disorganized thinking and altered level of consciousness. A patient that met the first and second aspects, and either the third or fourth, was considered positive for delirium. This method was used to evaluate patients every 24 h, until their discharge or death.

Quality Control

RASS score and CAM-ICU are commonly used as evaluation tools in the world.^{18,19} Every patient was independently assessed by doctors and nurses, and in case of a disagreement, they would consult a third person in our team. All research in the patient study underwent specialized training and passed the assessment prior to evaluation.

Statistical Analysis

Missing value were less than 5% in all variables in the present study. These values had minimal influence on research conclusions, hence we did not special consideration during statistical analysis. Continuous variables were expressed as means \pm standard deviations ($\bar{x} \pm s$), whereas counts were expressed as numerical values and percentages. Variables with $p < 0.05$, after univariate analysis, were selected for multivariate logistic regression. Similarly, those with $p < 0.05$ after multivariate logistic regression analyses were selected for model construction. We adopted a bootstrapping approach for internal validation (resampling 500 times), due to the relatively small sample size of our study. All statistical analyses

Table 1 The Clinical Characteristics of Patients

Variables	Non-Delirium	Delirium	Statistics
Age (years)	57.65 ± 16.36	66.98 ± 14.87	60.46 ± 16.47
<60	113 (50.67%)	25 (26.04%)	138 (43.26%)
≥60	110 (49.33%)	71 (73.96%)	181 (56.74%)
Sex			
Male	144 (64.57%)	60 (62.50%)	204 (63.95%)
Female	79 (35.43%)	36 (37.50%)	115 (36.05%)
Diagnosis type			
Medicine group	191 (85.65%)	72 (75.00%)	263 (82.45%)
Surgical group	14 (6.28%)	12 (12.50%)	26 (8.15%)
Neurologic group	18 (8.07%)	12 (12.50%)	30 (9.40%)
Educational			
Illiteracy	17 (7.62%)	7 (7.29%)	24 (7.52%)
Lower education	152 (68.16%)	70 (72.92%)	222 (69.59%)
Higher education	54 (24.22%)	19 (19.79%)	73 (22.88%)
BMI (Kg/m²)	23.31 ± 4.48	23.91 ± 4.63	23.49 ± 4.52
Smoking	80 (36.04%)	29 (30.21%)	109 (34.28%)
Drinking	50 (22.52%)	21 (21.88%)	71 (22.33%)
Diabetes	44 (19.73%)	22 (22.92%)	66 (20.69%)
Hypertension	75 (33.63%)	45 (46.88%)	120 (37.62%)
Sepsis	59 (26.94%)	24 (25.26%)	83 (26.43%)
Heart failure	41 (18.39%)	25 (26.04%)	66 (20.69%)
Electrolyte disorder	45 (20.18%)	25 (26.04%)	70 (21.94%)
Stomach tube	16 (7.17%)	25 (26.04%)	41 (12.85%)
Urinary tube	78 (34.98%)	69 (71.88%)	147 (46.08%)
CVC	123 (55.16%)	62 (64.58%)	185 (57.99%)
Sedative	37 (16.59%)	41 (42.71%)	78 (24.45%)
MV	26 (11.66%)	25 (26.04%)	51 (15.99%)
Vasoactive drugs	39 (17.49%)	26 (27.08%)	65 (20.38%)
APACHE-II score	10.16 ± 5.05	12.36 ± 4.59	10.82 ± 5.01
SOFA score	5.00 (3.00, 6.00)	5.00 (3.00, 7.00)	5.00 (3.00, 6.75)
Delirium	223 (69.91%)	96 (30.09%)	319 (100%)

Abbreviations: BMI, body mass index; CVC, central venous catheter; MV, mechanical ventilation; APACHE, Acute Physiology and Chronic Health Evaluation; SOFA, Sequential Organ Failure Assessment.

were performed using EmpowerStats (<http://www.empowerstats.com>, X&Y Solutions, Inc, Boston, MA) and packages implemented in R software (<http://www.R-project.org>, The R Foundation).

Results

Clinical Characteristics of Patients

A summary of patients' clinical characteristics is provided in [Table 1](#). Briefly, a total of 319 patients met our inclusion and exclusion criteria, and were therefore included in the study. Before leaving the EICU, 96 (30.09%) patients developed delirium, while 223 (69.91%) did not. Their average age was 60.46 ± 16.47 years, while treatment measures included stomach tube (n = 41), urinary tube (n = 147), CVC (n = 185), sedative (n = 78), MV (n = 51), and vasoactive drugs (n = 65). APACHE-II and SOFA scores were 10.82 ± 5.01 and 4.98 ± 2.55 points, respectively ([Table 1](#)).

Univariate Analysis Results

Univariate analysis revealed that age, hypertension, stomach tube, urinary tube, sedative, MV, APACHE-II score and SOFA score were significantly associated with delirium (p < 0.05), while the other variables had no significant relationship with delirium (p > 0.05) ([Table 2](#)).

Table 2 The Results of Univariate Analysis

Variables	Non-Adjusted HR, P-value
Age (years)	1.04 (1.02, 1.06) <0.001
<60	1.0 (Reference)
≥60	2.92 (1.72, 4.94) <0.001
Sex	
Male	1.0 (Reference)
Female	1.09 (0.67, 1.80) 0.724
Diagnosis type	
Medicine group	1.0 (Reference)
Surgical group	2.27 (1.00, 5.15) 0.049
Neurologic group	1.77 (0.81, 3.85) 0.152
Educational	
Illiteracy	1.0 (Reference)
Lower education	1.12 (0.44, 2.82) 0.813
Higher education	0.85 (0.31, 2.38) 0.763
BMI (Kg/m²)	1.03 (0.98, 1.08) 0.281
Smoking	0.77 (0.46, 1.29) 0.315
Drinking	0.96 (0.54, 1.72) 0.899
Diabetes	1.21 (0.68, 2.16) 0.520
Hypertension	1.74 (1.07, 2.84) 0.026
Sepsis	0.92 (0.53, 1.59) 0.757
Heart failure	1.56 (0.89, 2.76) 0.123
Electrolyte disorder	1.39 (0.79, 2.44) 0.247
Stomach tube	4.56 (2.30, 9.02) <0.001
Urinary tube	4.75 (2.82, 8.02) <0.001
CVC	1.48 (0.90, 2.43) 0.119
Sedative	3.75 (2.19, 6.41) <0.001
MV	2.67 (1.45, 4.92) 0.002
Vasoactive drugs	1.75 (0.99, 3.09) 0.053
APACHE-II score	1.09 (1.04, 1.15) <0.001
SOFA score	1.10 (1.00, 1.21) 0.042

Abbreviations: BMI, body mass index; CVC, central venous catheter; MV, mechanical ventilation; APACHE, Acute Physiology and Chronic Health Evaluation; SOFA, Sequential Organ Failure Assessment.

Results of Multivariate Logistic Regression Analyses

Statistically significant variables ($p < 0.05$) from univariate analysis were selected for multivariate logistic regression analyses. Since the APACHE-II score included assessment of age and blood pressure, we only included stomach and urinary tubes, as well as sedative, MV, APACHE-II and SOFA scores for multivariate analysis in order to avoid over-fitting of the model. The principle of covariate screening was that after introducing the variable into the basic model, or excluding it from the complete model, the variable whose influence on the regression coefficient of the independent variable exceeds 10%. Adjusting covariates, including vasoactive drugs, heart failure, diagnosis and type, revealed that stomach and urinary tubes, sedative, mechanical ventilation and APACHE-II scores were associated with delirium of patients (Table 3).

Construction of a Prediction Model

The aforementioned five variables, with $P < 0.05$, from the multivariate logistic regression analyses above were selected and used for model construction. The areas under curve (AUC) of the receiver operating characteristic (ROC) were 0.76 (0.70, 0.82), 0.75 (0.69, 0.81) and 0.75 (0.69, 0.81) for Multiple Fractional Polynomial (MFP), full and stepwise models, respectively. Non-elderly patients, aged <60 years old, recorded AUCs of 0.82 (0.73, 0.91), 0.83 (0.74, 0.91) and 0.82 (0.73, 0.91) for MFP, full and stepwise models, respectively. On the other hand, elderly patients aged ≥60 years old had AUCs of 0.82 (0.73, 0.91), 0.71 (0.62, 0.79) and 0.71 (0.63, 0.78) for MFP, full and stepwise models, respectively (Table 4 and Figures 2–7).

Table 3 Multivariate Logistic Regression Analyses for Delirium

Exposure	Non-Adjusted HR, P-value	Adjust* HR, P-value
Stomach tube	4.56 (2.30, 9.02) <0.001	4.26 (2.11, 8.61) <0.001
Urinary tube	4.75 (2.82, 8.02) <0.001	4.23 (2.46, 7.26) <0.001
Sedative	3.75 (2.19, 6.41) <0.001	3.60 (2.00, 6.48) <0.001
MV	2.67 (1.45, 4.92) 0.002	2.62 (1.36, 5.03) 0.004
APACHE-II score	1.09 (1.04, 1.15) <0.001	1.10 (1.04, 1.15) <0.001
SOFA score	1.10 (1.00, 1.21) 0.042	1.10 (1.00, 1.22) 0.060

Note: *The model was adjusted for vasoactive drugs, heart failure, and diagnosis type.

Abbreviations: MV, mechanical ventilation; APACHE, Acute Physiology and Chronic Health Evaluation; SOFA, Sequential Organ Failure Assessment.

Validation of the Prediction Model

We adopted a bootstrapping approach for internal validation of our models, owing to the relatively small size of our study. The full bootstrap model (BS full) and stepwise bootstrap model (BS stepwise) were used to verify the accuracy and value of the full model and stepwise model. The AUCs of 0.75 (0.69, 0.81) and 0.75 (0.69, 0.81) were recorded for BS full and stepwise models, respectively. In patients aged <60 years old, the model recorded AUCs of 0.83 (0.74, 0.91) and 0.82 (0.73, 0.91) for BS full and stepwise, respectively, while in elderly patients (≥ 60 years old), AUC values were 0.71 (0.62, 0.79) and 0.71 (0.63, 0.78) for BS full and stepwise, respectively (Table 4 and Figures 2–Figures 7).

Decision Curve Analysis of Prediction Models

We performed decision curve analysis (DCA) of constructed models. The results showed that, whether it was a model constructed by all patients or models constructed by age stratification, there was a certain distance between the nomogram and the standard line, which also confirmed the good clinical practicability of models (Figures 8–10).

Table 4 Results of the Constructed Predictive Models

Model	MFP	Full	Stepwise	Bootstrap Full	Bootstrap Stepwise
Total					
AUC	0.76	0.75	0.75	0.75	0.75
95% CI	(0.70, 0.82)	(0.69, 0.81)	(0.69, 0.81)	(0.69, 0.81)	(0.69, 0.81)
Specificity	0.70	0.65	0.65	0.63	0.65
Sensitivity	0.73	0.75	0.76	0.76	0.76
Accuracy	0.70	0.68	0.68	0.67	0.68
<60 years					
AUC	0.82	0.83	0.82	0.83	0.82
95% CI	(0.73, 0.91)	(0.74, 0.91)	(0.73, 0.91)	(0.74, 0.91)	(0.73, 0.91)
Specificity	0.85	0.85	0.85	0.83	0.90
Sensitivity	0.68	0.68	0.68	0.68	0.60
Accuracy	0.81	0.81	0.81	0.80	0.84
≥ 60 years					
AUC	0.82	0.71	0.71	0.71	0.71
95% CI	(0.73, 0.91)	(0.62, 0.79)	(0.63, 0.78)	(0.62, 0.79)	(0.63, 0.78)
Specificity	0.85	0.65	0.60	0.65	0.60
Sensitivity	0.68	0.70	0.73	0.70	0.73
Accuracy	0.81	0.67	0.65	0.67	0.65

Abbreviations: AUC, area under curve; CI, confidence interval; MFP, multiple fractional polynomial.




Figure 2 ROC curves of the models in all patients.




Figure 3 A MFP Nomogram model for all patients.




Figure 4 ROC curves of the models in non-elderly patients.




Figure 5 A MFP nomogram model for non-elderly patients.




Figure 6 ROC curves of models in elderly patients.




Figure 7 A MFP nomogram model for elderly patients.




Figure 8 DCA of the model in all patients.

Discussion

Results of the present study revealed that the incidence of delirium was 30.09%, which was consistent with results from previous epidemiological surveys.^{5,6} However, incidence from our study was slightly higher than that reported by Tsuruta,²⁰ possibly due to the high use of sedatives in the present study. To date, numerous studies have demonstrated that sedatives can be used to increase incidence of delirium in patients.²¹ Our results further revealed a slightly lower delirium incidence relative to that reported by Yang,²² which might be due to differences in the average age of patients and APACHE scores between the study. Generally, older patients are more likely to develop ICU delirium.²³

Our results revealed five variables, namely stomach and urinary tubes, as well as sedative, mechanical ventilation and APACHE-II scores, that were significantly associated with delirium. We used these variables to construct 3 predictive models in the entire patient cohort, comprising elderly (≥ 60 years old), and non-elderly (< 60 years old) patients. We internally validated these models using a bootstrapping approach. Results revealed a simple and highly accurate model for predicting delirium. The model could accurately predict delirium in both young and elderly patients, as evidenced by high AUC values across these groups.

APACHE – II, a tool used to analyze disease severity and prognosis of critically ill patients, has been widely used in ICUs worldwide.^{24,25} Results of the present study showed that patients with high APACHE-II scores had a higher risk of delirium, consistent with previous, studies that have shown that this variable is an independent risk factor for patients with delirium.^{26,27} For example, results of a previous meta-analysis,²¹ with a sample size of 2440, showed that the APACHE-II score was significantly higher in the delirium than non-delirium group.




Figure 9 DCA of the model in non-elderly patients.

In addition, Margaret study²⁸ suggested that use of sedative was a controllable risk factor for delirium. In 2013, the guidelines for the management of pain, agitation/sedation, and delirium issued by the Society of Critical Care Medicine (SCCM) also emphasized the influence of sedatives on delirium.²⁹ Results of the present study showed that use of sedative was an independent risk factor for delirium, consistent with the findings of Huang³⁰ and Pan,²¹ who concluded this in ICU patients.

Although MV is a common treatment strategy with promising therapeutic effects in EICU patients, it is often associated with occurrence of physical and psychological discomfort. Results of the present study showed that MV was an independent risk factor for delirium, consistent with a meta-analysis by Zhang³¹ who found that MV could cause sleep disorders in patients and increase the risk of delirium in this group of patients. Particularly, patients with mechanical ventilation were more likely to develop lung infections and hypoxemia, which exacerbated the risk of delirium.³²

Insertion stomach and urinary tubes are common invasive treatment operations in EICU. Long-term indwelling of foreign bodies in the patient's body often predisposes a patient's body to a state of stress and causes bad moods, which increase occurrence of delirium. Our results showed that stomach and urinary tubes were independent risk factors for patients with delirium, hence their inclusion in construction of predictive models.

All five variables identified herein were closely associated with delirium, consistent with previous studies. These factors, coupled with application of multiple methods for controlling the influence of collinearity between variables, allowed us to construct an accurate and reliable model for predicting delirium.

We successfully constructed 3 optimal predictive models for all patients, as well as those below and above 60 years. The MFP model for all patients had an AUC of 0.76 (0.70, 0.82), and comprised 4 variables, namely stomach and urinary




Figure 10 DCA of the model in elderly patients.

tubes, as well as sedative, and APACHE-II scores. The full model for non-elderly patients (<60 years old) had an AUC of 0.83 (0.74, 0.91), and comprised all 5 aforementioned variables. In addition, the MFP model for elderly patients (≥ 60 years old) had an AUC value of 0.82 (0.73, 0.91), and comprised 4 variables, namely stomach and urinary tubes, as well as sedative, and APACHE-II scores. All these models were highly accurate in predicting delirium in patients.

Nomogram Application

Each variable in the figure is marked with a scale on the line segment, representing their value range, with the length of the line segment reflecting its contribution to the outcome event. The point in the figure represents a single score, and their single scores are added to obtain total points.^{33,34}

Limitations of the Study

Firstly, our prediction model requires additional validation, since it was constructed and validated from a single-center, and the verification was also internal, which may cause its prediction value to decline outside this hospital. Secondly, we used a small sample size.

Conclusion

We developed a five-marker-based prognostic tool, for effective predicting occurrence of delirium in EICU patients. This tool, if prospectively validated, could provide individualized risk estimation of delirium patients in EICU.

Ethics Statement

The agency's ethics committee approved the study, with an ethical batch number of 2021006.

Acknowledgments

This work was supported by fund from Science and Technology Planning Project of Xi'an (no. 2019115713YX012SF049) and academy fund from the Second Affiliated Hospital of Xi'an Jiaotong University. The authors are indebted to all individuals who participated in or helped with this research project.

Disclosure

The authors report no potential conflicts of interest in this work.

References

1. Motta E, Luglio M, Delgado AF, et al. Importance of the use of protocols for the management of analgesia and sedation in pediatric intensive care unit. *Rev Assoc Med Bras*. 2016;62(6):602–609. doi:10.1590/1806-9282.62.06.602
2. Delaney A, Hammond N, Litton E. Preventing delirium in the intensive care unit. *JAMA*. 2018;319(7):659–660. doi:10.1001/jama.2018.0159
3. Mao L. Research of morbidity and relation factors of delirium in intensive care unit. North Sichuan Medical College; 2015.
4. Estrup S, Kjer C, Poulsen LM, et al. Delirium and effect of circadian light in the intensive care unit: a retrospective cohort study. *Acta Anaesthesiol Scand*. 2018;62(3):367–375. doi:10.1111/aas.13037
5. Collinsworth AW, Priest EL, Campbell CR, et al. A review of multifaceted care approaches for the prevention and mitigation of delirium in intensive care units. *J Intensive Care Med*. 2016;31(2):127–141. doi:10.1177/0885066614553925
6. Moon KJ, Lee SM. The effects of a tailored intensive care unit delirium prevention protocol: a randomized controlled trial. *Int J Nurs Stud*. 2015;52(9):1423–1432. doi:10.1016/j.ijnurstu.2015.04.021
7. Yang J, Zhou Y, Kang Y, et al. Risk factors of delirium in sequential sedation patients in intensive care units. *Biomed Res Int*. 2017;2017:3539872. doi:10.1155/2017/3539872
8. Kanova M, Sklienka P, Roman K, et al. Incidence and risk factors for delirium development in ICU patients - a prospective observational study. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub*. 2017;161(2):187–196. doi:10.5507/bp.2017.004
9. Chen J, Yu J, Zhao S, et al. Research progress on the risk prediction model of delirium in ICU. *J Nurs*. 2019;26(5):15–19.
10. Ely EW, Shintani A, Truman B, et al. Delirium as a predictor of mortality in mechanically ventilated patients in the intensive care unit. *JAMA*. 2004;291(14):1753–1762. doi:10.1001/jama.291.14.1753
11. Yunlan J, Xujuan X, Jun S, et al. Research progress on early recognition of and preventive nursing strategies for delirium in trauma patients. *J Nurs Sci*. 2020;35(14):104–109.
12. Rongpeng X, Chun Y, Bin Z. Research progress of delirium in intensive care unit. *Chin Crit Care Med*. 2020;32(5):636–640. doi:10.3760/cma.j.cn121430-20200204-00164
13. Salluh JJ, Wang H, Schneider EB, et al. Outcome of delirium in critically ill patients: systematic review and meta-analysis. *BMJ*. 2015;350:h2538. doi:10.1136/bmj.h2538
14. Theuerkauf N, Guenther U. Delirium on the ICU: clinical impact, diagnostic workup, and therapy. *Med Klin Intensivmed Notfmed*. 2014;109(2):129–136. doi:10.1007/s00063-014-0354-3
15. Inouye SK, Westendorp RG, Saczynski JS. Delirium in elderly people. *Lancet*. 2014;383(9920):911–922. doi:10.1016/S0140-6736(13)60688-1
16. Pan Y. An analysis of status and effect factors of ICU syndrome of patients in emergency intensive care unit[D]. Dalian Medical University; 2016.
17. Yuan Y, Qin W, Lu Y, et al. Patients treated by EICU with ICD. *Chin J Crit Care Med*. 2009;29(4):356–357.
18. Wong CL, Holroyd-Leduc J, Simel DL, et al. Does this patient have delirium?: value of bedside instruments. *JAMA*. 2010;304(7):779–786. doi:10.1001/jama.2010.1182
19. Schieveld J, Strik J. Delirium in developmentally disabled PICU children: the Richmond agitation sedation scale and delirium fluctuation issue. *Pediatr Crit Care Med*. 2020;21(5):494–495. doi:10.1097/PCC.0000000000002250
20. Tsuruta R, Nakahara T, Miyauchi T, et al. Prevalence and associated factors for delirium in critically ill patients at a Japanese intensive care unit. *Gen Hosp Psychiatry*. 2010;32(6):607–611. doi:10.1016/j.genhosppsych.2010.09.001
21. Pan Y, Jiang Z, Zhang J, et al. Risk factors of ICU delirium in adult patients: a meta-analysis. *J Nurs Manag*. 2018;18(04):465–475.
22. Yang K, Wang Y. Risk factors and nursing strategies of delirium in critically ill patients in emergency department. *J Nurs Training*. 2016;31(16):1493–1495.
23. Wenjuan T, Zhenxian S, Caiyun Z, et al. Bibliometric analysis of risk factors of delirium in ICU of China from 2008 to 2018. *Med Higher Vocational Educ Mod Nurs*. 2019;2(4):294–297.
24. Hu M, Meng J, Qin X. Application and effect analysis of APACHE-II scoring during the emergency nursing for critically ill patients. *J Nurs Training*. 2010;25(20):1833–1835.
25. Udy AA, Roberts JA, Shorr AF, et al. Augmented renal clearance in septic and traumatized patients with normal plasma creatinine concentrations: identifying at-risk patients. *Crit Care*. 2013;17(1):R35. doi:10.1186/cc12544
26. Sosa FA, Roberti J, Franco MT, et al. Assessment of delirium using the PRE-DELIRIC model in an intensive care unit in Argentina. *Rev Bras Ter Intensiva*. 2018;30(1):50–56. doi:10.5935/0103-507X.20180010
27. Wu C, Zhu Y, Li G. Incidence and risk factors of delirium in ICU patients. *J Third Military Medl Univ*. 2018;40(11):1038–1043.
28. Pisani MA, Murphy TE, Araujo KLB, et al. Benzodiazepine and opioid use and the duration of intensive care unit delirium in an older population. *Crit Care Med*. 2009;37(1):177–183. doi:10.1097/CCM.0b013e318192fcf9

29. Barr J, Fraser GL, Puntillo K, et al. Clinical practice guidelines for the management of pain, agitation, and delirium in adult patients in the Intensive Care Unit: executive summary. *Am J Health Syst Pharm.* 2013;70(1):53–58. doi:10.1093/ajhp/70.1.53
30. Huang J, Xiao Q, Wu Y, et al. A meta-analysis of the risk factors of delirium in ICU. *Chin J Nurs.* 2010;45(01):6–9.
31. Zhang Z, Pan L, Ni H. Impact of delirium on clinical outcome in critically ill patients: a meta-analysis. *Gen Hosp Psychiatry.* 2013;35(2):105–111. doi:10.1016/j.genhosppsych.2012.11.003
32. Elpern EH, Scott MG, Petro L, et al. Pulmonary aspiration in mechanically ventilated patients with tracheostomies. *Chest.* 1994;105(2):563–566. doi:10.1378/chest.105.2.563
33. Wang H, Tang L, Zhang L, et al. Development a clinical prediction model of the neurological outcome for patients with coma and survived 24 hours after cardiopulmonary resuscitation. *Clin Cardiol.* 2020;43(9):1024–1031. doi:10.1002/clc.23403
34. Wang H, Zheng X, Bai ZH, et al. A retrospective population study to develop a predictive model of prediabetes and incident type 2 diabetes mellitus from a hospital database in Japan between 2004 and 2015. *Med Sci Monit.* 2020;26:e920880. doi:10.12659/MSM.920880

International Journal of General Medicine

Dovepress

Publish your work in this journal

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies across all disease areas. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-general-medicine-journal>