


Construction of Internet +home Care Quality Supervision Indicators in China Based on the Delphi Method

Guiyuan Ma¹, Jianmei Hou¹, Sha Peng¹, Yanhui Liu¹, Zhengkun Shi¹, Yuhua Fan¹, Jinghui Zhang¹⁻³ 

¹Teaching and Research Section of Clinical Nursing, Xiangya Hospital of Central South University, Changsha, 410008, People's Republic of China; ²National Clinical Research Center for Geriatric Disorders, Xiangya Hospital of Central South University, Changsha, 410008, People's Republic of China; ³Mobile Health-Ministry of Education Laboratory, Changsha, 410008, People's Republic of China

Correspondence: Jinghui Zhang, Teaching and Research Section of Clinical Nursing, Xiangya Hospital of Central South University, No. 87 Xiangya Road, Kaifu District, Changsha, People's Republic of China, 410008, Tel +86 15874032339, Email zhangjh-1206@163.com

Background: With the aging of China's population, the incidence rate of chronic diseases is rising. At the same time, residents' health awareness is also increasing. Implementing Internet +home care is an inevitable trend in adapting social development. Currently, Internet +home care is in the pilot stage in China, relevant institutional measures are neither standardized nor perfect, and there is no set of quality supervision indicators (QSIs).

Purpose: The construction of Internet +home care QSIs in China will provide a theoretical basis for Internet +home care quality supervision.

Materials and Methods: The Service-Quality model was used as the theoretical framework, and a literature review, semi-structured interviews, focus group discussions, and Delphi consultations determined the contents of the QSIs and the weight of each indicator.

Results: Internet +home care QSIs were constructed and included 5 first-level indicators, 17 secondary indicators, and 69 tertiary indicators. The effective recovery rates of the two rounds of Delphi expert consultation were 100% and 85%, the expert authority coefficients were 0.810 and 0.833, and the Kendall harmony coefficients were 0.189 and 0.125 ($P < 0.01$). The final set of Internet +home care QSIs was as follows: tangibility (4 secondary and 16 tertiary indicators), reliability (4 secondary and 19 tertiary indicators), guarantee (5 secondary and 20 tertiary indicators), responsiveness (2 secondary and 8 tertiary indicators), and empathy (2 secondary and 6 tertiary indicators).

Conclusion: The construction of Internet +home care QSIs based on the SERVQUAL model is scientifically valid, and the indicators are reliable. They provide guidance and reference values for the continuous improvement and promotion of Internet +home care. This work also provides a theoretical basis for researching and developing an Internet +home care quality supervision platform.

Keywords: SERVQUAL model, Internet +home care, delphi survey, quality supervision indicator

Introduction

With the development of the social economy and medical care in China, demands upon the health service are becoming increasingly diversified, and the provision and conceptualization of nursing are deepening and extending.¹ On February 12, 2019, the National Health Commission issued the Notice on Launching the "Internet +home care" Service Pilot Work in China.¹ Internet +home care refers to medical institutions using institution-registered nurses to provide door-to-door nursing services for discharged patients/patients suffering from diseases with limited mobility through the "online application, offline service" mode. In the same year, Internet +home care was piloted in some cities in China.² Initial results are now available, although the implementation is still in the exploratory stage. Relevant institutional measures are not yet standardized or perfect, and issues include high pricing and poor information connectivity, an unclear payment guarantee system, and an absence of a standardized nursing staff performance appraisal system.³ As such, constructing

scientifically based Internet +home care quality supervision indicators (QSIs) is essential for safeguarding the benefits for patients and medical staff.

In the United States, Japan, and other countries, Internet +home care was implemented earlier. These countries have established practical approaches to evaluating the quality of home care, including the Outcome and Assessment Information Set,⁴ the Resident Assessment Instrument-Home Care,⁵ the Quick Investigation of Quality, and other home care assessment tools.⁶ There are differences in the operation mode of Internet +home care in different countries. Moreover, differences in domestic and foreign national conditions, cultures, and customs existed. Therefore, quality supervision indicators as used abroad cannot be directly applied in China.

According to the main body of the service, the model of Internet +home care can be divided into two categories. One is that medical institution nurses provide a door-to-door service, which is the service model adopted by countries such as China and Japan. The other is when company nurses provide services, such as Uber in the United States and Portea Medical in India. There are differences in the operation mode of Internet +home care in different countries.⁷ In this study, the constructed Internet +home care QSIs are applicable to the service operation model in China.

In evaluating nursing service quality, the commonly used three-dimensional quality structure theory, Quality Control Circle, and Plan-Do-Check-Act cycle theory constitute active evaluation by service providers, while feelings about recipients may not be sufficiently considered. The SERVQUAL (Service-Quality) evaluation method proposed by Parasuraman is a method by which to evaluate service quality from consumers' perspectives.⁸ The service quality is evaluated according to the score differences of consumers' "feelings" and "expectations" of service from the five perspectives of tangibility, reliability, guarantee, responsiveness, and empathy.⁹ Therefore, based on the SERVQUAL model, we constructed a series of Internet +home care QSIs centered on patient experience, suitable for Chinese national conditions. Importantly, these QSIs focus on the connotation of nursing services and provide an empirical basis for standardizing and guiding the healthy development of Internet +home care.

Materials and Methods

Participants

The semi-structured interview participants were medical managers and nursing managers at tertiary hospitals, patients, and family members. The focus discussion group was composed of multidisciplinary experts from the Ministry of Nursing, Medical Care, Education of China-Mobile Laboratory, and other relevant departments.

The inclusion criteria for the Delphi experts were: 1) appointed at deputy high school level or above; 2) holding a bachelor's degree or above; 3) 10 years or above of hospital nursing management or clinical nursing work experience; 4) Familiar with SERVQUAL theory. Exclusion criteria included refusal to participate in the research or providing invalid questionnaire responses.

Methods

Literature Review

We searched PubMed, MEDLINE, Web of Science, China National Knowledge Infrastructure (CNKI), and Chinese Wanfang databases for eligible studies published between 2011 and 2021. The following keywords or medical terms were used: "Internet +home care, home care, quality supervision, SERVQUAL, nursing quality indicators, Delphi method."

The inclusion criteria for articles were: (a) published in English or Chinese; (b) content related to this current study with the title, abstract, keywords, or text containing the search terms. The exclusion criteria were: (a) duplicate publication; (b) defects in the study design and/or poor quality. The Johns Hopkins body of evidence grade and quality assessment standards were used to evaluate the quality of the included literature, and the quality indicators related to Internet +home care involved in the literature were extracted and classified.

Semi-Structured Interviews

The research team developed an interview outline based on the literature review and clinical experience (see Figure 1). Meaningful items and phrases were marked in the interview, the same or similar materials were integrated, and the interview contents were refined and classified. The face-to-face semi-structured interviews were conducted with

Outline of interviews with medical and nursing managers:	<p>(a) Do you understand the concept of Internet +home care quality supervision indicators? What is your experience?</p> <p>(b) Do you think it is necessary to formulate Internet +home care quality supervision indicators?</p> <p>(c) What aspects do you think should be considered for each part of Internet +home care quality supervision indicators based on the Service-Quality model?</p> <p>(d) How do you think the weight allocation of Internet +home care quality supervision indicators based on the Service-Quality model should be applied?</p>
Outline of interviews with patients and their families:	<p>(a) Do you understand the concept of Internet +home care quality supervision indicators? What is your experience?</p> <p>(b) Do you think it is necessary to formulate Internet +home care quality supervision indicators?</p> <p>(c) What do you think of Internet +home care?</p> <p>(d) Do you have any suggestions for Internet +home care?</p>

Figure 1 Semi-structured interview outline.

attendees wearing protective masks at a distance of 1 meter apart. Before the interview, the interviewee was introduced to the research purpose and method. After obtaining informed consent, the interview outline was followed in which participants were guided in describing their thoughts on the Internet +home care QSIs. The interview time was set to 30–45 minutes per interview, in which the interview quantity was information saturation and no new information appeared.

Focus Group Discussions

Based on the literature review results, semi-structured interviews, and the professional knowledge of the focus group, experts initially determined the Internet +home care QSIs based on five aspects: tangibility, reliability, guarantee, responsiveness, and empathy.⁷ All primary indicators were discussed and the connotation definition and qualitative measurement of each indicator were repeatedly demonstrated. The indicators were gradually refined and clarified at all levels to ensure the final QISs were more logical and scientifically based.

Delphi Consultations

The Delphi questionnaire consisted of four parts: (a) basic questionnaire introduction; (b) questionnaire regarding basic information about the experts; (c) evaluation of the importance and relevance of each indicator and suggestions from experts; (d) each experts' familiarity with the content of the questionnaire and the basis for judgment. After completing the first round of consultation, the analysis, feedback, and revision were organized and the second round was formed and distributed. It was agreed that the consultation would be stopped if the expert opinions showed a relatively concentrated trend; otherwise, the third round of consultation would be carried out until the Internet +home care QSIs based on the SERVQUAL model was determined.

Experts completed the Delphi questionnaires by email. The principles for selecting and revising indicators were as follows: (a) if the importance of the indicator was less than 3.5 and the coefficient of variation was greater than 0.30, the indicator would be deleted; (b) if experts suggested adding or deleting some indicators, the focus group would discuss and review the literature to decide whether to adopt and modify the indicators; (c) for the revision suggestions or questions raised by experts on a specific indicator, the researchers would refer to the suggestions and give feedback to the experts. After the first round of consultation, if an expert noted that they were not sufficiently familiar with the field or if the returned questionnaire did not meet the filing requirements or was of low quality, the expert was not invited for the subsequent consultation.

Statistical Analysis

The general demographic data were described in frequency, percentage, mean and standard deviation (mean±SD). The expert positive coefficient refers to the recovery rate of the questionnaire, calculated as the ratio of experts number

participating in the consultation relative to the total number of experts. Two factors determined the degree of authority (Cr): (Ca) the educational level of the expert and the basis for judgment (experience, theoretical analysis, reference to data at home and abroad, and intuitive feelings), and (Cs) the degree of the expert's familiarity with the questions such that: $Cr=(Ca+Cs)/2$. The degree of coordination of expert opinions is represented by a coordination coefficient (Kendal's concordance coefficient, W) and coefficient of variation. Factor analysis was used for weight analysis. The common factor variance of each indicator was obtained by principal component analysis. Moreover, the specific weight of each indicator was obtained by calculating the percentage of each common factor variance to the sum of the common factor variances. All statistical tests were two-sided, and analyses were performed using SPSS 24.0. $P<0.05$ determined statistical significance.

Results

Literature Review

A total of 208 articles were retrieved on the initial search, of which 6 articles were presented for review by the expert panel.^{10–15} The flow chart of the literature search and retrieval is shown in Figure 2. We identified 5 first-level, 20 secondary, and 72 tertiary indicators.

Semi-Structured Interviews

After the literature review, 3 medical managers, 5 nursing managers, 5 patients, and 5 family members from tertiary hospitals were selected for semi-structured interviews. In total, 1 secondary and 2 tertiary indicators were eliminated, and 5 first-level, 19 secondary, and 70 tertiary indicators remained (Figure 3).




Figure 2 Literature quality and grade assessment results: Level III (A) 2 articles, Level III (B) 3 articles, Level IV (A) 1 article.




Figure 3 Selection of Internet +home care quality supervision indicators.

Focus Group Discussions

The multidisciplinary experts' panel comprised three nursing staff, two medical staff, and three Chinese Ministry of Education-mobile laboratory staff. The panel discussed the significance and applicability of these indicators and finally reached a preliminary consensus. This process eliminated 1 secondary and 4 tertiary indicators, and facilitated the formation of the first round of the expert survey (Figure 3).

Delphi Consultations

Participants' General Characteristics

Two rounds of expert consultations using the Delphi method were conducted with 20 and 17 expert members (in rounds 1 and 2, respectively). In both rounds, the majority of participants were female (90.0% and 88.2% in rounds 1 and 2, respectively), and most participants were aged between 41 and 50 years old (80.0% and 88.2% in rounds 1 and 2, respectively). Over 90.0% of the participants held senior professional titles (95.0% and 100.0% in rounds 1 and 2, respectively), and over half of the participants had master's degrees (60.0% and 70.6% in rounds 1 and 2, respectively). Table 1 displays the general characteristics of the participants.

Reliability of the Delphi Method

The recovery rates for the two rounds of consultation were 100% and 85%, respectively, indicating that the experts were motivated and attached great importance to this topic. The Cr for the two rounds of expert inquiry were 0.810 and 0.833, respectively, and the average Cr was 0.822, indicating that the authoritativeness of the experts was high.

Kendall's coefficient of concordance was calculated for each QSI. The coefficient of variation for the first round of consultation ranged from 0 to 0.128, and Kendall's W was 0.189 ($P < 0.01$). The second-round consultation coefficient of variation ranged from 0 to 0.080, and Kendall's W was 0.125 ($P < 0.01$), indicating high consistency in experts' opinions.

Table 1 Demographic Characteristics of the Participants [n(%)]

Demographic/Variable	Round 1	Round 2
Gender		
Male	2(10.0)	2(11.8)
Female	18(90.0)	15(88.2)
Age		
≤40	3(15.0)	1(5.9)
41—50	16(80.0)	15(88.2)
≥51	1(5.0)	1(5.9)
Academic career		
PhD student	3(15.0)	3(17.6)
Postgraduate student	12(60.0)	12(70.6)
Undergraduate student	5(25.0)	2(11.8)
Professional title		
Primary-grade title	0(0.0)	0(0.0)
Middle-grade title	1(5.0)	0(0.0)
Senior-grade title	19(95.0)	17(100.0)
Job category		
Health management	6(30.0)	4(23.5)
Clinical nursing	4(20.0)	2(11.8)
Nursing management	6(30.0)	6(35.3)
Nursing education	4(20.0)	4(23.5)

Indicator System Revision and Weight Determination

The statistical results (means and standard deviations) and weights for the Internet +home care QSIs are presented in [Table 2](#). Three participants suggested the naming and connotation of the secondary indicators. For example, they believed “timeliness” in responsiveness should be defined as “service response speed”. In addition, “establish a particular working group for Internet +home care” in tangibility and “carry out safe and reliable Internet +home care service” in reliability were suggested by four participants. Two participants suggested that some names and connotations for the QSIs should be modified. For tertiary indicators, participants recommended adding “there are clear job responsibilities and work procedures for Internet office managers” and “there are clear job responsibilities and work procedures for Internet platform staff” in tangibility, for example. Moreover, pairs of tertiary indicators were proposed to be combined separately, such as “standard attire, wearing a work badge” and “dignified appearance and decent behavior” were combined into “service staff should dress in standard attire, wear work badges, be dignified, and behave appropriately”. Based on discussions with the research team and after considering the experts’ comments, the following consensus was reached: add two secondary and eight tertiary indicators, delete three secondary and three tertiary indicators, combine three pairs of tertiary indicators separately, and adjust the connotations of some items.

In the second round of consultation, one expert proposed adding one secondary indicator, “Internet +home care staff professional guarantee” to the guarantee. One secondary indicator was deleted and one was modified. In terms of the naming, connotation, deletion, and addition of tertiary indicators, three participants submitted their opinions. One tertiary

Table 2 Internet +home Care Quality Supervision Indicators: Importance and Weight Determination

First-Level Indicator	Secondary Indicators	Tertiary Indicators	Importance (Mean±SD)	Coefficient of Variation	Weight (%)	
I Tangibility	1.1 There are institutions for Internet + home care		4.82±0.53	0.11	4.40	
		1.1.1 The institution is qualified to carry out Internet +home care	4.94±0.24	0.05	1.48	
		1.1.2 The institution has an Internet +home care office managed by dedicated staff	4.41±0.79	0.18	1.50	
		1.1.3 The institution provides matching equipment, consumables and medicines, which are managed by dedicated staff in the office	4.82±0.39	0.08	1.42	
	1.2 Establish a particular working group for Internet +home care			4.76±0.56	0.12	8.50
		1.2.1 The Internet +home care particular working group includes office management staff, information platform maintenance staff, and Internet nursing staff		4.88±0.33	0.07	1.49
		1.2.2 There are clear job responsibilities and work procedures for office managers		4.88±0.33	0.07	1.49
		1.2.3 There are clear job responsibilities and work procedures for Internet platform staff		4.82±0.39	0.08	1.27
		1.2.4 There are clear job responsibilities and work procedures for Internet nurses		4.76±0.44	0.09	1.37
		1.2.5 Service staff should dress in standard attire, wear work badges, be dignified, and behave appropriately		4.82±0.39	0.08	1.41
	1.2.6 Service staff should be uniformly equipped with nurse work recorders, positioning and tracking systems, and one-click alarm systems, and carry outpatient boxes and medical waste disposal containers.		4.82±0.39	0.08	1.47	

(Continued)

Table 2 (Continued).

First-Level Indicator	Secondary Indicators	Tertiary Indicators	Importance (Mean±SD)	Coefficient of Variation	Weight (%)	
	I.3 There is an information platform for Internet +home care		4.82±0.53	0.11	5.74	
		I.3.1 There is a dedicated information platform for Internet +home care, supporting mobile phone, iPad, and computer operations	4.82±0.39	0.08	1.42	
		I.3.2 Information platform has complete user ports: patient reservations, dispatching orders, nurses receiving orders, and background data monitoring	4.88±0.33	0.07	1.46	
		I.3.3 The appearance design of the information platform is simple, it supports text, voice, and video, and meets all needs of users	4.94±0.24	0.05	1.39	
		I.3.4 The information platform can realize data connection with the in-hospital electronic medical record system	4.82±0.39	0.08	1.47	
	I.4 Standardize Internet +home care operating environment			4.71±0.59	0.13	4.31
		I.4.1 There is a quiet, clean, safe home environment with suitable temperature and humidity		4.24±0.75	0.18	1.46
		I.4.2 Community medical institutions provide service modes such as a family ward, patrol services, Internet diagnosis and treatment		4.35±0.70	0.16	1.46
		I.4.3 There are nearby nursing homes, nursing stations, or pension institutions		4.29±0.69	0.16	1.39

2 Reliability	2.1 Data regarding the Internet +home care information platform is safe and reliable		4.94±0.24	0.05	7.28
		2.1.1 Internet +home care information platform has a complete security system	4.88±0.33	0.07	1.47
		2.1.2 User registration information is connected with the public security system and the Health Commission System to ensure its authenticity and reliability	4.76±0.44	0.09	1.45
		2.1.3 Nursing staff' privacy information is protected and private on Internet +home care	4.94±0.24	0.05	1.44
		2.1.4 Patient information is protected and private in Internet +home care	4.94±0.24	0.05	1.46
		2.1.5 Payment system provides Internet +home care security guarantee	4.94±0.24	0.05	1.46
	2.2 Carry out safe and reliable Internet +home care service		4.88±0.33	0.07	4.15
		2.2.1 Internet +home care contents meet national policy requirements, and there are no negative list items	4.65±0.49	0.11	1.32
		2.2.2 Internet +home care contents meet the needs of patients and their families	4.94±0.24	0.05	1.44
		2.2.3 There are clear service instructions and agreements, and professional services can be provided according to the service agreements	4.94±0.24	0.05	1.39
	2.3 Internet +home care system is effective and reliable		4.59±0.51	0.05	5.78
		2.3.1 Capable of remote multidisciplinary consultation for Internet +home care	4.94±0.24	0.05	1.47
		2.3.2 Able to realize referral services for patients whose condition changes during the service process	4.94±0.24	0.05	1.44
		2.3.3 Can provide multiple forms of health education such as the provision of documents and videos	4.84±0.33	0.07	1.38
		2.3.4 Can allow two-way feedback and evaluation of nurses and patients	4.88±0.33	0.07	1.49

(Continued)

Table 2 (Continued).

First-Level Indicator	Secondary Indicators	Tertiary Indicators	Importance (Mean±SD)	Coefficient of Variation	Weight (%)	
	2.4 Reliable management of matching consumables, medicines, and equipment for Internet +home care		4.94±0.24	0.05	4.23	
		2.4.1 Medicines are managed by specially-assigned staff, reviewed by specially-assigned staff, fixed-point access, supervised and traceable	4.88±0.33	0.07	1.44	
		2.4.2 Consumables are managed by specially-assigned staff, reviewed by specially-assigned staff, fixed-point access, supervised and traceable	4.94±0.24	0.05	1.34	
		2.4.3 Matching equipment is managed by specially-assigned staff, reviewed by specially-assigned staff, fixed-point access, supervised and traceable	4.94±0.24	0.05	1.45	
	2.5 Safety protection of users and nursing staff of Internet +home care			4.76±0.56	0.12	5.55
		2.5.1 Nursing staff purchase relevant liability insurance, personal accident insurance		4.76±0.56	0.12	1.31
		2.5.2 Patients purchase medical accident insurance in the process of service		4.82±0.39	0.08	1.36
		2.5.3 The information platform can realize a one-click alarm and synchronously connect to the public security system and the operation management backstage		4.88±0.33	0.07	1.50
		2.5.4 Travel recorder can record the service process in real-time and throughout the whole process		4.94±0.24	0.05	1.38

3 Guarantee	3.1 Internet +home care supervision system is complete		4.88±0.33	0.07	7.39
		3.1.1 Complies with the national policy requirements for Internet +home care	4.82±0.39	0.08	1.52
		3.1.2 Provides a complete Internet +home care quality management and evaluation system	4.88±0.33	0.07	1.46
		3.1.3 Provides an Internet +home care disputation and complaint handling system	4.94±0.24	0.05	1.43
		3.1.4 Provides Internet +home care nurses and patients with an information security management system	4.88±0.33	0.07	1.46
		3.1.5 Provides a complete infection control system and medical waste disposal standard, which can be supervised and traced	4.76±0.44	0.09	1.52
	3.2 Internet +home care risk plan is complete		4.94±0.24	0.05	5.90
		3.2.1 Includes a complete Internet +home care risk assessment and response mechanism	4.88±0.33	0.07	1.49
		3.2.2 Includes occupational safety precautions and coping mechanisms for nursing staff	4.88±0.33	0.07	1.49
		3.2.3 Includes high-risk user identification and response mechanisms	4.88±0.33	0.07	1.49
		3.2.4 Includes contingency plans for patient accidents in the process of Internet +home care	4.94±0.24	0.05	1.43
	3.3 Internet +home care process specification		4.94±0.24	0.05	5.91
		3.3.1 The operation process for Internet +home care is detailed	4.94±0.24	0.05	1.47
		3.3.2 Provides hygiene standards and procedures in the Internet nursing work environment	4.88±0.33	0.07	1.44
		3.3.3 Provides an Internet +home care medical waste disposal process	4.88±0.33	0.07	1.50
		3.3.4 Provides a solution process for orders without response	4.88±0.33	0.07	1.50

(Continued)

Table 2 (Continued).

First-Level Indicator	Secondary Indicators	Tertiary Indicators	Importance (Mean±SD)	Coefficient of Variation	Weight (%)
	3.4 Internet +home care staff professional guarantee		4.94±0.24	0.05	10.29
		3.4.1 Qualifications of staff providing Internet +home care conform to the national policy; they have worked for more than 5 years and have the title of nurse	4.88±0.33	0.07	1.44
		3.4.2 Staff providing Internet +home care have received professional pre-job training and passed the examination	4.82±0.39	0.08	1.43
		3.4.3 No records of adverse events or complaints	4.94±0.24	0.05	1.46
		3.4.4 Staff providing Internet +home care have received primary life supportability training and passed the examination	4.82±0.39	0.08	1.46
		3.4.5 Those who perform specialized nursing programs shall have corresponding specialized nurse certificates	4.71±0.59	0.13	1.52
		3.4.6 Matching is ensured between nursing staff disciplines and patient disease	4.82±0.53	0.11	1.49
		3.4.7 Nursing staff follow relevant industry standards and guidance consensus	4.77±0.56	0.12	1.49
4 Responsiveness	4.1 Platform running speed		4.82±0.39	0.08	5.76
		4.1.1 The platform service interface runs smoothly and the operation is simple	4.76±0.56	0.12	1.28
		4.1.2 Client information can be updated synchronously in a timely fashion	4.82±0.39	0.08	1.45
		4.1.3 Platform functions can be optimized in a timely fashion according to patients' needs and suggestions	4.76±0.44	0.09	1.52
		4.1.4 Payment is convenient and electronic invoices for service fees are issued for patients in a timely fashion	4.88±0.33	0.07	1.51
	4.2 Service Response Speed		4.82±0.39	0.08	5.95
		4.2.1 Internet +home care staff receive orders and contact patients in time	4.77±0.56	0.12	1.52
		4.2.2 Internet +home care staff educate patients on health in a timely manner	4.88±0.33	0.07	1.52
		4.2.3 Internet +home care information maintenance staff solve technical problems in a timely manner	4.77±0.56	0.12	1.45
		4.2.4 Internet +home care management staff handle complaints in a timely manner	4.94±0.24	0.05	1.46

5 Empathy	5.1 Provide humanized service		4.88±0.33	0.07	4.56
		5.1.1 Staff pay attention to communication skills and patients' feelings	4.82±0.39	0.08	1.52
		5.1.2 Staff are gentle in their approach and pay attention to privacy protection	4.76±0.44	0.09	1.52
		5.1.3 Staff pay attention to the patient health demands and provide humanistic care	4.82±0.39	0.08	1.52
	5.2 Provide personalized service		4.59±0.51	0.11	4.21
		5.2.1 Staff develop personalized service items based on patient demands	4.94±0.24	0.05	1.32
		5.2.2 Staff establish personalized electronic files for patients	4.94±0.24	0.05	1.43
		5.2.3 Staff provide personalized health education based on patient characteristics	4.88±0.33	0.07	1.46

indicator was added and two tertiary indicators were modified. Ultimately, the Internet +home care QSIs contained 5 first-level, 17 secondary, and 69 tertiary indicators.

Discussion

The Internet +home Care QSIs Based on the SERVQUAL Model is Highly Scientific and Reliable

Based on the SERVQUAL theoretical framework applied to the situation in China, the Internet +home care QSIs constructed by literature review, semi-structured interviews, focus group discussions, and Delphi expert consultations are scientifically based.¹⁶ In the construction of the initial indicator database, after several focus group discussions and expert interviews, all aspects of Internet +home care QSIs were comprehensively considered and set, ensuring the reliability of the indicators.

The experts' qualifications, positive attitudes, authority, and consistency of opinions confirm the scientific basis and reliability of the research results. We selected multidisciplinary experts with abundant experience in clinical nursing, nursing management, and healthcare. The authority coefficients of the experts in the two rounds were above 0.8. Moreover, the experts' professional titles and educational levels also showed that they had high authority and reliability. The effective recovery rates of the two rounds of Delphi consultation were more than 70%, reflecting the active participation of experts participating in this research. Kendall's concordance coefficient for the two rounds indicated that the experts reached a consensus on each indicator's importance, ensuring the indicators' reliability. From the comprehensive data analysis, the Internet +home care QSIs achieved the expected goals in terms of scientificity and reliability. Overall, the indicators obtained in this study are reasonable and thus, can be used for Internet +home care quality supervision.

Content Analysis of Internet +home Care QSIs Based on the SERVQUAL Model

We established the Internet +home care QSIs, including 5 first-level, 17 secondary, and 69 tertiary indicators. Among the first-level indicators, the top two weights of indicators are reliability (29.49%) and guarantee (26.99%), which are consistent with the characteristics of the Internet +home care model. As a new service model, Internet +home care's complete hardware, software, and serviceability guarantee is its necessary condition. Medical institution qualifications and hardware and software configurations are directly proportioned to the level of serviceability of providing door-to-door nursing staff. This reminds us that medical institutions that carry out Internet +home care should further refine the requirements and improve the qualifications and service capabilities, to provide better care for patients needing Internet +home care. Furthermore, tangibility, responsiveness, and empathy quality indicators cannot be ignored.

Tangibility refers to the elements directly perceived by patients, including software and hardware configurations, and human resources. Tangibility in this study includes 4 secondary and 16 tertiary indicators, whereby the top 3 weights of secondary indicators are "establish a particular working group for Internet +home care", "there is an information platform for Internet +home care" and "there are institutions for Internet + home care". The indicators reflect the importance of a fully staffed work team, a software information platform with supporting functions, and a medical institution with complete qualifications to implement nursing services. In cases the inadequate qualifications of staff and the resources are insufficient, the credibility of service level will be low, and the quality of service will be affected.¹⁷

Reliability refers to performing service safely, reliably, and accurately. Focusing on the service platform and service levels, the guarantee of door-to-door nursing safety is the core element of Internet +home care.¹⁸ The highest weight in secondary reliability indicators is "Data regarding the Internet +home care information platform is safe and reliable". The big health data in Internet +home care is vast and contains much private and medical information. It is necessary to avoid network attacks, and at the same time, patients have the right to privacy, and medical staff must protect patients' privacy.¹⁹ Among the tertiary indicators of reliability, the highest weight is "the information platform can realize a one-click alarm and synchronously connect to the public security system and the operation management backstage". Internet +home care is currently in the trial operation stage in China, and safety issues cannot be avoided. It should be noted that users have concerns about the qualifications of nurses and the safety of medical care in Internet +home care.²⁰ Nurses'

concerns are concentrated on the safety of nursing operations, nurse-patient conflicts, and medical disputes,²¹ and so it is essential to improve security protection.

Guarantee refers to the standardization of management and the assurance of the professional skills of relevant service staff. The highest weight indicators among the secondary guarantee indicators are “Internet +home care staff professional guarantee” and “Internet +home care supervision system is complete”. Qualified Internet +home care staff provide sufficient guarantee for patients and interconnected systems should be established to guarantee the working years, professional titles, training, and assessment of nursing staff. Moreover, establishing complete and reasonable QSIs is the foundation of Internet +home care and an important quality and safety guarantee. Therefore, it is recommended that government and relevant management departments establish relevant policies and systems to provide platform users with better Internet +home care.²²

Responsiveness represents service staff responding quickly to patients demands and providing timely assistance. The secondary indicator with the highest weight in responsiveness is “service response speed”. The platform’s proper order dispatching strategy, timely order response, and active order confirmation will significantly reduce time consumption and improve efficiency. Among the tertiary indicators, the indicators with the highest weight are “platform functions can be optimized in a timely fashion according to patients’ needs and suggestions”, “Internet +home care staff receive orders and contact patients in time”, and “Internet +home care staff educate patients on health in a timely manner”. It is necessary to respond to patients’ orders in a timely fashion, pay attention to the requirements, evaluations and suggestions of patients, and take targeted improvement measures. It is recommended that nursing staff comprehensively evaluate the physical and psychological conditions of patients and provide appropriate health care education when providing nursing services to patients. In addition, such instruction should be personalized, easy to understand, and effective.

Empathy refers to concern about and caring for patients, being able to put oneself in the patients’ shoes, and providing individualized services for patients. After a two-round expert Delphi consultation about empathy, the secondary indicators are “provide humanized service” and “provide personalized service”. In nursing, being “people-oriented” should be implemented throughout the nursing process to maximize the nursing service experience for patients. Nursing staff must have a humanitarian spirit, and focus on patients’ physical and mental health, provide emotional and spiritual support, and ensure patients feel humanized. It is equally essential to provide personalized services to patients. Everyone is a unique individual; nursing staff should respect the individuality of patients, focus on the interests and demands of patients when providing services, and try their best to meet the psychological and physiological needs of patients as diverse social people. Therefore, nurses engaged in Internet +home care need to be able to identify patients’ needs, formulate personalized service contents, establish personalized electronic files for patients, and deliver personalized health education knowledge according to patients’ characteristics.²³

Limitations and Conclusions

The reliable and scientifically based Internet +home care QSIs identified here meet the needs of society, and the development of the Internet medical system has a guiding role in the improvement of Internet +home care. Cheng et al²⁴ found that Internet +home care is in high demand among Chinese residents and is highly accepted by nursing staff.²⁵ National policy vigorously promotes the development of Internet +home care, indicating a promising prospect in China. However, there is very little research on the quality supervision of Internet +home care in China. Therefore, based on the SERVQUAL theoretical system, we constructed the Internet +home care QSIs to standardize and guide the healthy development of Internet +home care, provide a reference for nursing managers and nursing educators in pilot hospitals, and provide a new direction for the standardized implementation of Internet +home care. Due to time constraints, we constructed the Internet +home care QSIs theoretically and did not conduct empirical research on Internet +home care. In future research, we will build an Internet +home care quality supervision platform based on the established Internet +home care QSIs, and continue to supervise and improve the quality of Internet +home care.

Ethics Approval

The Medical Ethics Committee of Xiangya Hospital, Central South University, Changsha, China, approved this study with approval code 202102060.

Acknowledgments

First, we would like to thank Xiangya Hospital of Central South University for approving and supporting this study. Second, we express our gratitude to Liqing Yue and Weihong Huang and all the participants for contributing to the consensus of the QSIs on Internet +home care. Finally, we thank Xiangya Hospital of Central South University, Hunan Provincial Social Science Committee, and the National Clinical Research Center for Geriatric Disorders for financial and material support.

Funding

This study was funded by the Management in Xiangya Hospital of Central South University (No. 2021GL23) and the General Funding Project of Hunan Provincial Social Science Review Committee (No. XSP22YBZ131). It was also supported by the Appropriate Technology Promotion Project of the National Clinical Research Center for Geriatric Diseases of Xiangya Hospital of Central South University (No. XYYYJSTG-04).

Disclosure

The authors declare that there are no conflicts of interest regarding this work.

References

1. General Office of the National Health Commission. (2019) Notice of the General Office of the National Health and Health Commission on Launching the Pilot Work of “Internet +home care Services”; 2019. Available from: <http://www.nhc.gov.cn/yzygj/s7657g/201902/bf0b25379ddb48949e7e21edae2a02da>. Accessed February 12, 2022.
2. Li QP, Li J, Pan HY. Effects of Online Home Nursing Care Model Application on Patients with Traumatic Spinal Cord Injury. *Risk Manag Healthc Policy*. 2021;14:1703–1709.
3. Zhao B, Zhang X, Huang R, et al. Barriers to accessing internet-based home Care for Older Patients: a qualitative study. *BMC Geriatr*. 2021;21(1):565.
4. Bowles KH, McDonald M, Barrón Y, et al. Surviving COVID-19 After Hospital Discharge: symptom, Functional, and Adverse Outcomes of Home Health Recipients. *Ann Intern Med*. 2021;174(3):316–325.
5. Aylin W. Home care quality indicators based on the Resident Assessment Instrument-Home Care (RAI-HC): a systematic review. *BMC Health Serv Res*. 2020;20(1):366.
6. Close MA, Barden-O’Fallon J, Mejia C. Quality of family planning services in HIV integrated and non-integrated health facilities in Malawi and Tanzania. *Reprod Health*. 2019;16(Suppl 1):58.
7. Huang YS, Yuan CR, Song XP, et al. Development status of “Internet + nursing service”. *Chine Nursing Res*. 2018;32(22):3499–3502.
8. Fan LH, Gao L, Liu X, et al. Patients’ perceptions of service quality in China: an investigation using the SERVQUAL mode. *PLoS One*. 2017;12(12):e0190123.
9. Fatima I, Humayun A, Iqbal U, et al. Dimensions of service quality in healthcare: a systematic review of literature. *Int J Qual Health Care*. 2019;31(1):11–29.
10. Wagner A, Zúñiga F, Rüesch P, et al. Selecting home care quality indicators based on the Resident Assessment Instrument-Home Care (RAI-HC) for Switzerland: a public health and healthcare providers’ perspective. *PLoS One*. 2020;15(12):e0244577.
11. Murphy F, Doody O, Lyons R, et al. The development of Nursing Quality Care Process Metrics and Indicators for use in Older Persons Care Settings: a Delphi-Consensus Study. *J Adv Nurs*. 2019;75(12):3471–3484.
12. Tan QH, Shi ZY, Li X, et al. Construction of quality evaluation index system of Internet plus nursing service based on three-dimensional quality theory. *Chin J Mod Nur*. 2021;27(33):4536–4540.
13. Srithumsuk W, Prachusilpa G, Thummapol O, et al. A Study of Home Healthcare Nursing Outcomes Quality Indicators for Older People in Thailand Using Delphi Technique. *Home Healthc*. 2021;39(3):154–159.
14. Zhuang HR, Zhang WY, Ma LL, et al. Construction of the “Internet + Nursing Service” quality evaluation index system based on the three-dimensional quality structure model. *Chin J Mod Nur*. 2021;27(25):3367–3373.
15. Wiig S, Ree E, Johannessen T, et al. Improving quality and safety in nursing homes and home care: the study protocol of a mixed-methods research design to implement a leadership intervention. *BMJ Open*. 2018;8(3):e020933.
16. McPherson S, Reese C, Wendler MC. Methodology Update: delphi Studies. *Nurs Res*. 2018;67(5):404–410.
17. Aiken LH, Sloane DM, Bruyneel L, et al. RN4CAST consortium. Nurse staffing and education and hospital mortality in nine European countries: a retrospective observational study. *Lancet*. 2014;383(9931):1824–1830.
18. Wiig S, Ree E, Johannessen T, et al. Improving quality and safety in nursing homes and home care: the study protocol of a mixed-methods research design to implement a leadership intervention. *BMJ Open*. 2018;8(3):e020933.
19. Kruse CS, Smith B, Vanderlinden H, et al. Security Techniques for the Electronic Health Records. *J Med Syst*. 2017;41(8):127.

20. Huang RD, Han J, Cui NX. Attitude and concerns of the public about “sharing nursing”: a text mining study based on Weibo comments. *Chine Nursing Res.* 2021;35(17):3021–3025.
21. Berland A, Bentsen SB. Medication errors in home care: a qualitative focus group study. *J Clin Nurs.* 2017;26(21–22):3734–3741.
22. Lin M, Ma L, Ying C. Matching daily home health-care demands with supply in service- sharing platforms[J]. *Transp Res E Logist Transp Rev.* 2021;145:102177.
23. Turjamaa R, Hartikainen S, Kangasniemi M, et al. Living longer at home: a qualitative study of older clients’ and practical nurses’ perceptions of home care. *J Clin Nurs.* 2014;23(21–22):3206–3217.
24. Cheng LN, Zhao Xie XF. Care willingness and demand of residents under 60 years of age in western China: a cross-sectional study[J]. *BMJ Open.* 2021;11(8):e046515.
25. Huang R, Xu M, Li X, et al. Internet-Based Sharing Nurse Program and Nurses’ Perceptions in China: cross-Sectional Survey. *J Med Internet Res.* 2020;22(7):e16644.

Risk Management and Healthcare Policy

Dovepress

Publish your work in this journal

Risk Management and Healthcare Policy is an international, peer-reviewed, open access journal focusing on all aspects of public health, policy, and preventative measures to promote good health and improve morbidity and mortality in the population. The journal welcomes submitted papers covering original research, basic science, clinical & epidemiological studies, reviews and evaluations, guidelines, expert opinion and commentary, case reports and extended reports. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/risk-management-and-healthcare-policy-journal>