

#### ORIGINAL RESEARCH

# Antidepressant-Like Activity of Solvent Fractions of the Root Bark of Carissa spinarum Linn. (Apocynaceae) in Rodents Involves Multiple Signaling Pathways

Hana Saif Ali (1), Ephrem Engidawork (1)

Department of Pharmacology and Clinical Pharmacy, School of Pharmacy, College of Health Sciences, Addis Ababa University, Addis Ababa, Ethiopia

Correspondence: Ephrem Engidawork, Department of Pharmacology and Clinical Pharmacy, School of Pharmacy, College of Health Sciences, Addis Ababa University, P.O. Box 9086, Addis Ababa, Ethiopia, Email ephrem.engidawork@aau.edu.et

Background: The root bark of Carissa spinarum Linn. (Apocynaceae) is claimed to be used for the management of depression in Ethiopian folkloric medicine, and the crude extract has been reported to possess antidepressant-like activity in rodents.

Objective: This study aimed to evaluate the antidepressant-like effect of different fractions of the root bark in rodents and the possible underlying mechanisms in rats.

Methods: A 70% ethanol extract of the root bark was successively fractionated with n-butanol, ethyl acetate, and water. Animals of both sexes received 2% Tween 80, imipramine (30 mg/kg), or various doses (50, 100, 200 mg/kg) of the fractions. Duration of immobility was determined using the tail suspension test and the forced swim test. Locomotor activity was evaluated in the open field test. Serum corticosterone levels, total phenols, flavonoids, and alkaloids were determined. Preliminary mechanistic studies were also performed to explore possible mechanisms of action of the active fraction.

**Results:** All fractions but the aqueous fraction significantly (p<0.001) decreased the duration of immobility in both tests, with the ethyl acetate fraction being the most active. The locomotor test revealed that the activity was not due to non-specific psycho-stimulant effects. Serum corticosterone levels were reduced by both fractions, with the ethyl acetate fraction again being the most effective. Mechanistic studies showed the involvement of multiple neurotransmission systems, including adrenergic, dopaminergic and cholinergic as well as L-Arginine-NO-cGMP pathway. Higher contents of phenols (42.42 vs 29.8 mgGAE/g), flavonoids (12.43 vs 2.07 mgQE/g), and alkaloids (0.17 vs 0.07 mgATE/g) were found in the ethyl acetate than in the n-butanol fraction.

**Conclusion:** The present findings collectively indicate that the ethyl acetate and n-butanol fractions are endowed with antidepressantlike activity due to the presence of phenols, flavonoids, and alkaloids, which are medium polar in nature.

**Keywords:** depression, Carissa spinarum Linn, antidepressant-like activity, solvent fractions, possible mechanisms

#### Introduction

Depression is a chronic and recurrent psychiatric disorder that affects mental and physical health, and has a significant impact on healthcare resources and costs.1 It is a heterogeneous mental disorder with psychological, behavioral, and physiological symptoms that affects one in five people worldwide. Moreover, it is a multifactorial disorder with different causes.<sup>2,3</sup> It is one of the most burdensome diseases in the middle years of life in both developing and developed countries. The highest prevalence is estimated in sub-Saharan Africa, North Africa, and the Middle East. 5

Despite the availability of several highly effective antidepressants that improve clinical and low work productivity, only 50-60% of patients receive adequate treatment, and those treated may develop resistance during treatment.<sup>4</sup> Almost all of the antidepressants are associated with various side effects<sup>6–9</sup> and are contraindicated in some diseases.<sup>6,10</sup> They also showed potential interactions with food and other medications, and have been associated with liver dysfunction and

increased risk of diabetes.<sup>11</sup> To overcome these side effects, low efficacy, and inaccessibility,<sup>12</sup> many scientists are seeking alternatives to conventional antidepressants from plant sources.

Plant-derived natural medicines have a crucial role in the development of natural drugs, which have antidepressant activity, as they are known to be affordable, safe and efficacious. Many plants that are used in the treatment of depression act by normalizing the disturbed neurotransmitters in the brain or related mechanisms, including the 5-HT system (hypericum species), the noradrenergic system (*Valeriana wallichii*), the dopamine system (*Rosmarinus officinalis*), the GABAergic system (*Asparagus racemosus*), the Glutamate system (*Siphocampylus verticillatus*), neurotrophins (*Polygala tenuifolia*), the HPA axis (*Ptychopetalum olacoides*) and the nitric oxide system (*Aloysia gratissima*).<sup>13,14</sup>

Carissa spinarum Linn. (Apocynaceae) (Supplementary Figure 1) is a medicinal plant known as a magic herb in most African countries and used in the treatment of various diseases, including depression. The plant harbors high content of bioactive secondary metabolites such as alkaloids, cardiac glycosides, saponins, flavonoids, tannins, anthraquinones, total phenols, and terpenoids, which are implicated in producing anti-depressant activity. C. spinarum has different vernacular names in Ethiopia, Hagmsa (in Affan Oromo) and Agam (in Amharic). The roots are extensively used in Ethiopian folk medicine to treat various ailments, either alone or in the form of mixtures. 15,19–23

Moreover, pharmacological studies have confirmed most of the claimed uses, including diuretic, <sup>24,25</sup> antinociceptive, <sup>26,27</sup> anticonvulsant, <sup>28,29</sup> anti-inflammatory and antioxidant, <sup>30</sup> and antimicrobial <sup>31–33</sup> activity. The crude root bark extract of *C. spinarum* is also reported to possess antidepressant activity. <sup>34</sup> Therefore, the aim of this study was to evaluate the antidepressant-like activity of the solvent fractions using a series of paradigms and to explore the possible mechanisms.

### **Materials and Method**

## Drugs, Chemicals, and Reagents

Cyproheptadine HCl (Algorithm S.A.L - Lebanon); Diazepam BP (Gland Pharma Limited - Denmark); Ketamine Hydrochloride USP (Rotex Medica - Germany); L-Arginine HCl (Vita Pharmaceutical - Syria); Prazosin HCl (Mylan Inst., RKFD - USA); Standard Imipramine HCL BP (Remedica - Cyprus); Sulpiride (Sanofi-Aventis - France); Yohimbine (Sun Naturals - USA); Atropine (BDH Chemicals - England); Absolute Ethanol 99.9% (Iso Lab Chemicals - India); Dimethyl sulfoxide (DMSO) (Riedel.de Haen - Germany); Ethyl Acetate (Lobe Chemie - India); Methylene Blue (BDH Chemicals - England); N-Butanol (Loba Chemie - India); Normal saline Solution (Sansheng Pharmaceutics - Ethiopia); Petroleum Ether 40–60°C (Loba Chemie - India); Tween 80 (Uni-Chem - India), Aluminum Chloride Hexahydrate (Loba Chemie - India); BCG (Sisco research laboratories - India); Citric acid (Avonchem - UK); Disodium hydrogen orthophosphate Na<sub>2</sub>HPO<sub>4</sub> (BDH Chemicals - England); Follin Ciocalteu; Gallic acid (Merch - Germany); Potassium Acetate (Blulux laboratories - India); Quercetin dihydrate (Sigma Aldrich - Germany); Sodium hydroxide pellets (Loba Chemie - India); and distilled water. All drugs were purchased from pharmacies in Ethiopia, Yemen, and the USA, while the chemicals were purchased from their respective vendors in Ethiopia. All were of analytical grades.

# Plant Material Collection and Preparation

The roots of *C. spinarum* were collected in August 2020 from Ashewa Meda, Burayu city administration (near Addis Ababa, the capital), Oromia region, Ethiopia. Identification and authentication of the plant sample was performed by a taxonomist, and a voucher specimen (HS001) was deposited at the National Herbarium, College of Natural and Computational Sciences, Addis Ababa University for future reference. The root was cleaned and carefully washed with tap water to remove dirt and soil debris. The root bark was then separated from the root wood, cut into small pieces and air dried in the shade. The dried root bark was pulverized using mortar and pestle.

## **Experimental Animals**

Healthy Swiss albino mice (20–30 g, 6–8 weeks old) and Sprague-Dawley rats (200–250 g, 8–12 weeks old) of both sexes were bred and obtained from the animal house of the School of Pharmacy, Addis Ababa University, Addis Ababa, Ethiopia. All animals were housed in groups of 6 per cage and had access to standard laboratory pellets and water ad

libitum under controlled conditions (12 h light and dark cycle and temperature of 23–25 °C). Before performing the pharmacological studies, the animals were acclimatized to laboratory conditions for 5 days. In addition, all animals used were handled and cared for according to internationally accepted guidelines for the care and use of laboratory animals.<sup>35</sup> Ethical approval was obtained from an ethical review board of the School of Pharmacy, College of Health Sciences, Addis Ababa University (Reference number. ERB/SOP/176/12/2020).

#### Plant Extraction and Fractionation

#### Crude Extract

Five hundred and fifty grams of the ground root bark was defatted overnight with 1L of petroleum ether. It was then filtered and allowed to dry on a clean piece of aluminum. After complete evaporation of the petroleum ether, the plant material was macerated with 1.5 L of 70% ethanol for three days. The maceration process was repeated twice with the same volume. The extract was then filtered with a cotton gauze followed by Whatman filter paper No. 1, and filtration was accelerated with a vacuum pump. The filtrate was then concentrated under reduced pressure using a rotary evaporator at 40° C and stored in a refrigerator at 4° C. A sticky, dark brown colored crude extract was obtained with a percentage yield of 9.22% (w/w).

#### Solvent Fractionation

Thirty-three grams of the extract was mixed with acid-washed sand and placed in a small thimble in a Soxhlet apparatus for fractionation with various solvents. The sequence of solvents used for fractionation was based on their polarity index (n-butanol, ethyl acetate, and water).<sup>36</sup> The n-butanol (CSB) and ethyl acetate (CSE) fractions were then collected sequentially and dried under reduced pressure in a rotary evaporator at 40° C. Finally, the marc left was macerated with 1 L of distilled water and freeze-dried using a lyophilizer (Labconco, Console Freeze Dry system, USA) to obtain the aqueous fraction (CSA). The final samples obtained were weighed and the percentage yields were 21.00% for the n-butanol, 11.55% for the ethyl acetate, and 61.7% for the aqueous fraction. The n-butanol and ethyl acetate fractions were sticky and light brown in color, while the aqueous fraction was black. The fractions were stored in tightly closed containers and kept in a refrigerator at -20° C until use.

# Determination of Phytochemical Constituents and TLC Analysis Determination of Total Phenolic Content

Total phenolic content (TPC) of the fractions was determined using the Follin-Ciocalteu method as described elsewhere<sup>37</sup> with minor modifications. Follin-Ciocalteu reagent (2N) was diluted with distilled water (1:20). A series of concentrations (25, 12.5, 6.25, and 3.125  $\mu$ g/mL) were then prepared in methanol using a standard solution of gallic acid (50  $\mu$ g/mL). One mL of the standard solvent and each fraction (250  $\mu$ g/mL) were mixed separately with 0.5 mL of Follin-Ciocalteu reagent in a test tube and allowed to stand at room temperature for 8 min. Sodium carbonate (2 mL of a 7.5% solution) was added and the mixture was allowed to stand at room temperature for 30 min. The absorbance was then measured at 765 nm using a UV- spectrophotometer. The same procedure was used to prepare a blank solution. The test was performed in triplicate and the average value was recorded. A linear calibration curve was plotted (Figure 1A)), with an equation (Y = 0.009296\*X + 0.05072) and a correlation coefficient (R<sup>2</sup>) of 0.9997, and used to calculate the concentration of TPC, which was expressed as milligrams of gallic acid equivalent per gram (mgGAE/g).

#### Determination of Total Flavonoid Content

Total flavonoid content (TFC) of the solvent fractions was determined by the colorimetric Aluminum chloride method as described elsewhere with minor modifications. A series of concentrations (25, 12.5, 6.25, 3.125, and  $1.5625~\mu g/mL$ ) were prepared in methanol using quercetin (1mg/mL) as a standard solution. Ten mg of each solvent fractions was dissolved in 10 mL of methanol to obtain 1 mg/mL of a stock solution. Half a milliliter of each solvent fraction and one milliliter of the standard solution were mixed separately in a test tube with 100  $\mu$ L of 10% aluminum chloride, 100  $\mu$ L of 1M potassium acetate, and 2.8 mL of distilled water. The mixture was incubated at room temperature for 30 min. The absorbance was measured at 415 nm using a UV- spectrophotometer. The same




Figure I Calibration curve for Gallic acid (A), Quercetin (B) and Atropine (C) standard solutions.

procedure was used to prepare a blank solution. The assay was performed in triplicate and the average value was noted. A linear calibration curve was plotted (Figure 1B)), with an equation of (Y = 0.02413\*X - 0.01900) and a correlation coefficient  $(R^2)$  of 0.9986, and used to determine TFC, which was expressed as milligrams of quercetin equivalent per gram (mgQE/g).

#### Determination of Total Alkaloid Content

Total alkaloid content (TAC) of the solvent fractions was determined using bromocresol green solution (BCG) and phosphate buffer solution prepared as described elsewhere.<sup>39</sup> A series of concentrations (0.5, 0.25, 0.125, 0.0625, and 0.03125 mg/mL) were prepared using a standard pure atropine (1mg/mL) dissolved in methanol. Two mL of each solvent fraction (1 mg/mL) was dissolved in 2N HCl and then filtered using Whatman filter paper No.1. One mL of the filtered solution was then transferred to a separatory funnel and washed twice with 5 mL of chloroform. The pH was neutralized with 0.1 N NaOH. Five mL of the BCG solution and phosphate buffer were then added and shaken vigorously. Five mL of chloroform was added and the complex formed was transferred to a 10 mL volumetric flask and diluted to volume with chloroform. The absorbance of the complex formed was measured at 470 nm. A blank solution was prepared by the same procedure without the addition of atropine.<sup>40</sup> The assay was performed in triplicate and the average value was noted. A linear calibration curve was plotted (Figure 1C)), with an equation of Y = 0.002184\*X + 0.03608 and a correlation coefficient (R<sup>2</sup>) of 0.9935, and used to determine TAC, which was expressed as milligrams of Atropine equivalent per gram (mgATE/g).

# Thin Layer Chromatography Analysis

Thin layer chromatography (TLC) was performed on precoated silica gel plates using the conventional one-dimensional ascending method. The plate was cut using a glass cutter and marked with a soft pencil. The active fraction was dissolved

in methanol and spotted with a glass capillary at a distance of 1 cm. The plate was kept in a pre-saturated chamber with a solvent system containing a mixture of petroleum ether and ethyl acetate in a 1:2 ratio, covered and allowed to run 3/4th of the plate. The plate was removed and dried at room temperature for 30 min and observed under UV light (254 nm and 366 nm) (Figure 2). The movement of the active compound was calculated and expressed as retention factor (Rf) using the following equation:

Retention factor = 
$$\frac{\text{Distance travelled by the solute}}{\text{Distance travelled by the solvent}}$$

## Grouping and Dosing of Experimental Animals

For each model and solvent fraction, five groups of animals of both sexes were randomly assigned with 6 animals per group. Group I, served as a negative control and received vehicle (2% Tween 80) (TW80), group II served as a positive control and received standard drug, imipramine (30 mg/kg) (IP30), group III–V received increasing doses of the solvent fractions 50, 100, and 200 mg/kg, respectively. The solvent fractions and standard drug were dissolved in 2% Tween 80, prepared and administered orally one hour before the experimental sessions. The maximum volume administered was 10 mL/kg. Dose selection for the standard drug as well as the fractions was based on previous studies and a pilot experiment conducted before the main experiment.

## **Experimental Paradigms**

### Tail Suspension Test

The tail suspension test (TST) was performed as described elsewhere. Following 1 h after administration of vehicle/ standard/fractions, each mouse was suspended by its tail upside-down, using an adhesive tape placed 1 cm from the tip of the tail, on a countertop at a height of 35–50 cm above the floor. The test was recorded with a digital video camera. The duration of immobility was measured in seconds using a stopwatch software, the XNote Timer, for the entire duration of the test. The duration of immobility is defined as the absence of any movement of the head and body. def. 46,47

#### Forced Swim Test

The forced swim test (FST) or the Porsolt Despair test in rats was performed as described elsewhere <sup>48</sup> with some modifications. The FST consists of two sessions. The initial 15-min pretest session followed by a 5-min test session 24 h later. Rats were forced to swim in an inescapable situation in a transparent glass cylinder (40 cm height; 18 cm diameter) filled with water 20 cm deep,




Figure 2 TLC profile of Ethyl acetate fraction under UV light at a wavelength of 254 nm (A) and 366 nm (B).

with their paws not touching the bottom. The solvent fractions were administered 1, 4, and 24 h before the test. After swimming, the rat was taken out and dried with a clean towel, to maintain body temperature and prevent hypothermia, and returned to the cage. The water was also changed and replaced. Only the test sessions were recorded using a digital video camera, which was then used to evaluate the duration of immobility. The duration of immobility was measured using a stopwatch software, the XNote Timer. Floating and absence of any struggling behavior except those necessary to keep the head above the water for breathing were scored as immobile behavior, implying the development of an adaptive response to the stressful situation in the water.<sup>49</sup>

#### Open Field Test

The open field test (OFT) measures general locomotor activity and rule out any false-positive effect of the extract in mice. The open field apparatus is a rectangular inescapable box (68 cm x 68 cm x 45 cm) with the surface marked horizontally and vertically with lines forming a grid of 16 squares and a 60 W lamp placed above it. The solvent fractions were administered orally one hour before the test. The mouse was then placed in the center and crossing of the central and peripheral squares were tracked and counted for 5 min. The inner surface of the box was swapped and cleaned with alcohol and absorbent cotton to avoid any olfactory cues.<sup>50</sup>

## Serum Corticosterone Assay

After completion of the FST and TST, each animal was immediately anesthetized with ketamine (80 mg/kg, i.p) and diazepam (10 mg/kg, i.p), and a cardiac puncture was performed. Blood was collected in an SST tube and kept at room temperature for 60 min. It was then centrifuged at 1000 g force for 10 min at 4°C and the separated serum was stored at -80°C until assayed. Corticosterone is a glucocorticoid hormone in rodents that is secreted by the adrenal cortex under various stressful conditions such as the forced swim test and tail suspension test. Therefore, it was analyzed and measured using an electrochemiluminescence Immunoassay (ECLIA) according to the manufacturer's instructions (Cobas e 41-Roche Diagnostics GmbH, Mannheim, Germany).

## Evaluation of Possible Mechanism(s) of Action

According to the pathophysiology of depression and the mechanism of action of conventional antidepressants, the possible mechanism (s) of action of the most active solvent fraction of the root bark of *C. spinarum* was investigated. This was performed by assessing the involvement of noradrenergic, serotonergic, dopaminergic, nitric oxide, and cholinergic-muscarinic system using different drugs at doses that do not modify the locomotor behaviors of rats in FST.<sup>51,52</sup> Rats were randomly divided into nine different groups of 6 rats each. All drugs and the extract were dissolved in 0.5% v/v DMSO in normal saline. Prazosin (1 mg/kg, i.p., an α1-adrenoceptor antagonist) and yohimbine (1 mg/kg, i.p., an α2-adrenoceptor antagonist) were used for the noradrenergic system; cyproheptadine (3 mg/kg, i.p., a 5-HT<sub>2</sub> receptor antagonist) for the serotonergic system; sulpiride (50 mg/kg, i.p., a dopamine D2 receptor antagonist) for the dopamine system; atropine (1 mg/kg, i.p., a muscarinic cholinergic receptor antagonist) for the cholinergic system; and L-Arginine (750 mg/kg, i.p. a precursor of NO) and methylene blue (10 mg/kg, i.p., an inhibitor of soluble guanylate cyclase (sGC)) used for the NO system. All drugs were administered 15 min before CSE (50 mg/kg, P.O.) and rats were subjected to FST 60 min after administration of the fraction.

# Statistical Analysis

Data were expressed as mean ± standard error of the mean (SEM). Analyses were carried out using the Statistical Package for the Social Sciences (SPSS), version 25. One-way analysis of variance (ANOVA) followed by a Tukey's post hoc test was used for testing and significance level was set at a *P*-value of less than 0.05. Curves for the quantification of the phytochemical constituents were generated using Graph Pad Prism software version 8.00 for Windows (Graph Pad Software Inc., San Diego, California, USA).

## Results

## Quantification of Secondary Metabolites and TLC Analysis

Phytochemical analysis of CSB and CSE, as these were found to be active in the experimental paradigms used, revealed that TPC was 29.8 mgGAE/g and 42.42 mgGAE/g, respectively. Likewise, TFC was 2.07 mgQE/g and 12.43 mgQE/g, and TAC was 0.07 mgATE/g and 0.17 mgATE/g, respectively, in CSB and CSE. TFC, TPC and TAC were higher in CSE than CSB. In addition, TAC was lower in both fractions compared to TFC and TPC, the order being TPC > TFC > TAC.

TLC study of CSE (as it is the most active fraction) showed the presence of three different spots, one major spot with an  $R_f$  value 0.43 and two minor spots with  $R_f$  values of 0.69 and 0.77 respectively (Figure 2).

## Tail Suspension Test

Data obtained for the three fractions in the TST model are shown in Tables 1 and 2. In mice treated with the three doses of CSE, there was a significant reduction (p< 0.001) in the duration of immobility compared to TW80. The duration of immobility appeared to decrease with dose, with the middle (100 mg/kg, CSE100) and higher (200 mg/kg, CSE200) doses producing a significantly greater reduction (p < 0.05) than the lower dose (50 mg/kg, CSE50). CSB also produced the same pattern as CSE, except that CSB200 had a comparable effect to IP30. Indeed, the reduction in the duration of immobility produced by IP30 was significantly greater (p<0.05) than all doses of CSE as well as CSB50 and CSB100. By contrast, CSA did not significantly reduce the duration of immobility at the doses used in the present study. A comparison between CSE and CSB revealed no apparent difference between the different doses of the two fractions, except that CSB200 (p<0.001) produced a significantly greater reduction than CSE50.

**Table I** Antidepressant-Like Activity of Ethyl Acetate and n-Butanol Fractions of the Root Bark of *C. spinarum* in Mice Using Tail Suspension Test

Treatment	Dose (mg/kg)	Duration of Immobility (sec)	% Reduction
TW80	10 mL/kg	218.10 ± 4.36	-
IP	30	77.40 ± 10.25 <sup>a3</sup>	64.41%
CSE	50	$152.60 \pm 3.21^{a3b3d2e2}$	30.03%
	100	110.30 ± 2.84 <sup>a3b1c1</sup>	46.43%
	200	108.40 ± 8.16 <sup>a3b1c2</sup>	50.30%
CSB	50	157.30 ± 4.88 <sup>a3b3d1e3</sup>	27.88%
	100	122.60 ± 8.82 <sup>a3b1c1</sup>	43.79%
	200	96.40 ± 10.10 <sup>a3c3</sup>	55.80%

**Notes**: Values are presented as mean  $\pm$  SEM and analyzed using One-way ANOVA followed by Tukey's Post hoc test; n = 6; TW80, controls receiving 2%Tween 80;  $^1p < 0.05$ ;  $^2p < 0.01$ ;  $^3p < 0.001$ ;  $^a$ Compared to control;  $^b$ Compared to imipramine;  $^c$ Compared to 50 mg/kg;  $^d$ Compared to 100 mg/kg;  $^e$ Compared to 200 mg/kg.

**Abbreviations**: IP, imipramine; CSE, Carissa spinarum Linn. ethyl acetate fraction; CSB, Carissa spinarum Linn. n-butanol fraction.

**Table 2** Antidepressant-Like Activity of an Aqueous Fraction of the Root Bark of *C. spinarum* in Mice Using Tail Suspension Test

Treatment	Dose (mg/kg)	Duration of Immobility (Seconds)	% Reduction
TW80	10 mL/kg	188.70 ± 8.59	_
IP	30	III.90 ± 2.98 <sup>a2</sup>	40.70%
CSA	50	136.90 ± 10.57	27.45%
	100	192.30 ± 25.94 <sup>b2</sup>	-1.91%
	200	164.10 ± 13.01	13.04%

**Notes**: Values are presented as mean  $\pm$  SEM and analyzed using One-way ANOVA followed by Tukey's Post hoc test; n=6; TW80, controls receiving 2%Tween 80;  $^2p < 0.01$ ;  $^a$ Compared to control;  $^b$ Compared to imipramine. **Abbreviations**: IP, imipramine; CSA, *Carissa spinarum* Linn aqueous fraction.

## Forced Swim Test

Since the CSA had no effect in the TST, the FST was performed for the CSE and CSB fractions. Table 3 summarizes the data obtained from the FST using the two fractions at different doses used in the study. Accordingly, both fractions significantly reduced (p<0.001) the duration of immobility in a dose-dependent manner compared with TW80. In addition, the lower doses of both fractions had a significantly smaller reduction (p<0.01) than the middle and higher doses, although no detectable difference was observed between the latter two doses. CSE50 and CSB50 produced a significantly smaller reduction (p<0.001) in the duration of immobility than IP30. Although CSB100 and CSB200 also produced a significantly lower (p<0.001) duration of immobility than IP30, no apparent difference was observed between IP30 and the other two CSE doses (CSE100 & CSE200). In contrast to TST, the effect of CSE on FST was consistently higher than that of the corresponding doses of CSB for each dose used. For example, CSE200 showed a significantly greater reduction in immobility time than CSB100 (p<0.01) and CSB200 (p<0.05).

## Open Field Test

Table 4 shows a summary of data obtained from the OFT. Accordingly, neither the standard nor the fractions resulted in a significantly different number of crosses compared to TW80. Similarly, no apparent difference was found between the

Table 3 Antidepressant-Like Activity of Ethyl Acetate and n-Butanol Fractions of the Root Bark of C. spinarum in Rats Using Forced Swim Test

Treatment	Dose (mg/kg)	Duration of Immobility (Seconds)	% Reduction
TW80	10 mL/kg	190.46 ± 1.73	-
IP	30	$67.25 \pm 3.60^{a3}$	61.59%
CSE	50	$132.70 \pm 8.28^{a3b3d3e3}$	32.84%
	100	$85.10 \pm 3.85^{a3c3}$	56.93%
	200	$78.30 \pm 3.03^{a3c3}$	60.37%
CSB	50	$136.80 \pm 6.62^{a3b3d3e3}$	28.17%
	100	$103.80 \pm 2.66^{a3b3c2}$	45.50%
	200	$100.60 \pm 3.65^{a3b3c3}$	47.18%

Notes: Values are presented as mean ± SEM and analyzed using One-way ANOVA followed by Tukey's Post hoc test; n = 6; TW80, controls receiving 2%Tween 80;  $^2$ p < 0.01;  $^3$ p < 0.001;  $^3$ Compared to control; <sup>b</sup>Compared to imipramine; <sup>c</sup>Compared to 50mg/kg; <sup>d</sup>Compared to 100mg/kg; <sup>e</sup>Compared to 200mg/kg. Abbreviations: IP, imipramine; CSE, Carissa spinarum Linn. ethyl acetate fraction; CSB, Carissa spinarum Linn. n-butanol fraction.

Table 4 The Effect of Ethyl Acetate and n-Butanol Fractions of the Root Bark of C. spinarum on Locomotion in Mice Using Open Field Test

Treatment	Dose (mg/kg)	Number of Squares Crossed		
		Peripheral	Central	Total
TW80	10 mL/kg	80.00 ± 5.51	6.83 ± 2.10	86.83 ± 7.06
IP	30	77.50 ± 9.58	7.50 ± 1.36	85.00 ± 9.52
CSE	50	73.67 ± 17.30	7.33 ± 1.65	81.00 ± 18.02
	100	68.50 ± 11.29	7.67 ± 1.84	76.17 ± 12.48
	200	82.33 ± 12.81	10.67 ± 4.23	93.00 ± 15.78
CSB	50	69.00 ± 15.01	5.67 ± 2.59	74.67 ± 16.49
	100	65.67 ± 16.49	5.83 ± 1.14	71.50 ± 16.18
	200	62.00 ± 21.88	4.67 ± 2.35	66.67 ± 23.72

Notes: Values are presented as mean ± SEM and analyzed using One-way ANOVA followed by Tukey's Post hoc test; n = 6; TW80, controls receiving 2%Tween 80.

Abbreviations: IP, imipramine; CSE, Carissa spinarum Linn. ethyl acetate fraction; CSB, Carissa spinarum Linn. n-butanol fraction.

different doses of the fractions and the standard, ruling out the possibility that the antidepressant-like activity was due to nonspecific psychostimulation.

## Serum Corticosterone Assay

The effect of CSE and CSB on serum corticosterone levels in TST is shown in Table 5. CSE significantly (p<0.01) decreased serum corticosterone levels at all doses compared with TW80 in a dose-dependent manner. Although IP30 significantly decreased the level compared to CSE50 (p<0.05), no obvious difference was observed between IP30 and the other two doses. All CSB doses had no significant effect on corticosterone levels, as no detectable changes were observed compared to TW80. In addition, with the exception of CSB50 (p<0.05), no significant difference was observed for the other doses compared to IP30.

Serum corticosterone levels were also measured in rats subjected to FST. As shown in Table 6, CSE50, CSB50, and CSB100 had no effect on corticosterone levels. In contrast, CSE100 (p < 0.05) and CSE200 (p < 0.05) significantly reduced levels compared to TW80. On the other hand, a significantly greater reduction was seen with IP30 compared to all doses of CSE (p < 0.05). However, larger individual differences precluded statistically significance between IP30 and CSB.

**Table 5** The Effect of Ethyl Acetate and n-Butanol Fractions of the Root Bark of *C. spinarum* on Serum Corticosterone Level in Mice Using Tail Suspension Test

Treatment	Dose (mg/kg)	Serum Corticosterone Level (nmol/l)	% Reduction
TW80	10 mL/kg	63.72 ± 7.80	-
IP	30	19.57 ± 2.65 <sup>a3</sup>	69.29%
CSE	50	39.18 ± 3.91 <sup>a2b1</sup>	38.51%
	100	$31.11 \pm 2.62^{a2}$	51.27%
	200	$27.09 \pm 3.96^{a3}$	57.49%
CSB	50	50.59 ± 8.48 <sup>b1</sup>	20.61%
	100	44.96 ± 11.57	29.44%
	200	34.35 ± 7.26	46.09%

**Notes**: Values are presented as mean  $\pm$  SEM and analyzed using One-way ANOVA followed by Tukey's Post hoc test; n = 5; TW80, controls receiving 2%Tween 80;  $^{1}p < 0.05$ ;  $^{2}p < 0.01$ ;  $^{3}p < 0.001$ ;  $^{a}$ Compared to control;  $^{b}$ Compared to imipramine.

**Abbreviations**: IP, imipramine; CSE, *Carissa spinarum* Linn. ethyl acetate fraction; CSB, *Carissa spinarum* Linn. n-butanol fraction.

**Table 6** The Effect of Ethyl Acetate and n-Butanol Fractions of the Root Bark of *C. spinarum* on Serum Corticosterone Level in Rat Using Forced Swim Test

Treatment	Dose (mg/kg)	Serum Corticosterone Level (nmol/l)	% Reduction
TW80	10 mL/kg	72.66 ± 5.14	_
IP	30	$28.13 \pm 3.86^{a3}$	61.29%
CSE	50	59.92 ± 4.38 <sup>b3</sup>	17.53%
	100	$53.08 \pm 1.53^{a1b2}$	26.95%
	200	48.61 ± 3.53 <sup>a2b1</sup>	33.10%
CSB	50	51.91 ± 6.02	28.56%
	100	48.59 ± 5.59	33.13%
	200	45.65 ± 8.90 <sup>a1</sup>	37.17%

**Notes**: Values are presented as mean  $\pm$  SEM and analyzed using One-way ANOVA followed by Tukey's Post hoc test; n = 5; TW80, controls receiving 2%Tween 80;  $^{1}p < 0.05$ ;  $^{2}p < 0.01$ ;  $^{3}p < 0.001$ ;  $^{a}$ Compared to control;  $^{b}$ Compared to imipramine.

**Abbreviations**: IP, imipramine; CSE, *Carissa spinarum* Linn. ethyl acetate fraction; CSB, *Carissa spinarum* Linn. n-butanol fraction.




Figure 3 The effects of various pharmacological drugs on the antidepressant-like activity of ethyl acetate fraction of the root bark of C. spinarum using forced swim test in rats; Data are presented as mean  $\pm$  SEM and analyzed using One-way ANOVA followed by Tukey's post hoc test; n=6; Control received normal saline (10 mL/kg);  $^{1}p < 0.05$ ;  $^{2}p < 0.01$ ;  $^{3}P < 0.001$ ;  $^{3}Compared$  to control;  $^{5}Compared$  to CSE50.

Abbreviations: CSE50, Carissa spinarum Linn. ethyl acetate fraction 50 mg/kg; PZN, Prazosin; YOH, Yohimbine; CYP, Cyproheptadine; SUL, Sulpiride; ATR, Atropine; L-Arg, L-Arginine; MB, Methylene blue.

## Evaluation of the Possible Mechanism(s) of Action

According to the results of the antidepressant-like activity in the current study, CSE proved to be the most active fraction. Therefore, this fraction was used to investigate the possible mechanism(s) of action involved using different pharmacological agents.

The effect of different pharmacological agents (drugs) on the antidepressant-like activity of CSE using the FST model is shown in Figure 3. Pretreatment with yohimbine, sulpiride, atropine, and L-arginine significantly reversed the antidepressant-like activity of CSE50 (p < 0.05). In contrast, pretreatment with prazosin, cyproheptadine, and methylene blue had no effect on the antidepressant-like activity of CSE50. Among the agents that reversed the activity, only sulpiride and L-arginine were able to return the duration of immobility to the level of TW80.

## **Discussion**

In the present study, an attempt was made to evaluate the antidepressant-like activity and explore the possible mechanism (s) of action of solvent fractions of a root bark of a plant used in Ethiopia to treat depression.

Several experimental models are essentially used to assess and evaluate the efficacy and potency of agents with antidepressant-like activity. In this study, the FST (Despair Behavior test) and TST were used due to their ease of use, high reliability, and specificity, as well as for their sensitivity to all classes of antidepressants.<sup>53</sup>

Oral administration of the solvent fractions of the root bark of *C. spinarum* at different doses indicated that CSE and CSB exhibited antidepressant-like activity in the rodent models of depression used in the present study. CSA on the other hand, did not show significant antidepressant-like activity, as reported elsewhere. <sup>34</sup> Moreover, a nonsignificant increase in the total number of squares crossed in OFT with CSE could be an indicator of some anxiolytic effect. However, there was no apparent effect of CSE and CSB on the locomotor activity, indicating that the effect was not due to a nonspecific psychostimulant effect of the fractions.

Depression as a byproduct of stress or as a stress-related disorder triggers a neuronal, endocrine, and behavioral response. In FST and TST, rodents are exposed to an inescapable situation that leads to endocrine disruption and a change in corticosterone levels as a result of activating the major neuroendocrine stress system, the HPA axis. 54,55 Therefore, corticosterone level can be used as a possible marker of depression. CSE, especially at the middle and higher doses, consistently decreased corticosterone levels in both TST and FST, although such a consistent decrease was not observed in CSB. This effect may be used as additional evidence for the antidepressant-like activity of the fractions, with CSE showing better activity than CSB.

The antidepressant-like activity observed in the present study could be attributed to the presence of phenols, flavonoids, and alkaloids in the fractions. Many scientific studies have reported that plants with antidepressant-like activity are rich in these secondary metabolites<sup>52,56</sup> and produce effect at doses used in the present study. 13,57 Quantitative analysis showed that CSE contained more secondary metabolites than CSB, which, at least, in part explains the difference in activity between the two fractions. The difference in polarity of the solvents used for fractionation could be the basis for the selective distribution of secondary metabolites in CSE compared to CSB. These secondary metabolites are thought to exert their antidepressant-like effect individually or in synergy by influencing the various neurotransmitters involved in the neurobiology of depression. Phenols, flavonoids (polyphenol) and alkaloids, which were shown to be constituents of *C. spinarum*, are among the best studied secondary metabolites whose mechanisms may involve stimulation of 5-HT, noradrenaline, dopamine, and GABAergic neurotransmission systems. 56,58,59 Indeed, alkaloids have been found to decrease immobility behavior in FST and several preclinical studies have demonstrated their antidepressant-like activity, although their concentration in *C. spinarum* was very low. TLC study of CSE indicates the presence of different spots with different R<sub>f</sub> values, which reinforces the notion that the root bark is endowed with different secondary metabolites, which contribute to its antidepressant-like activity.

Since CSE was found to be the most active fraction in the variety of depression paradigms used in the present study, further investigation of the possible mechanism(s) using this fraction was conducted based on several hypotheses proposed to explain the pathophysiology of depression.

 $\alpha$ 2-Adrenoceptors are inhibitory autoreceptors that autoregulate the noradrenergic system. The affinity and density of these receptors are greatly increased in depressed patients,  $^{62}$  so that activation of the receptors has a crucial influence on behavioral activation and target neuron function. Therefore, dysregulation of  $\alpha$ 2-adrenoceptors is associated with depression. The fact that yohimbine, but not prazosin, was able to reverse the antidepressant-like activity of CSE suggests that the fraction mediates its effect via  $\alpha$ 2- adrenoceptors but not  $\alpha$ 1-adrenoceptors. This finding is consistent with reports indicating yohimbine that plant extracts preferentially interact with  $\alpha$ 2-adrenoceptors to produce antidepressant-like effects. Indeed, studies suggest that some flavonoids interact with  $\alpha$ 2-adrenoceptors to produce depression-like symptoms in animals, while some flavonoids exhibit antidepressant-like effects.

Pretreatment with cyproheptadine did not reverse the antidepressant-like activity of CSE, suggesting that the effect was not mediated via 5-HT<sub>2</sub> receptor neurotransmission. Similar results have been reported elsewhere.<sup>70</sup> The dopaminergic system is an extremely important target involved in the regulation of depression through its important role in reward stimulus processing, cognition, mood, attention, and learning.<sup>71,72</sup> Pretreatment with sulpiride significantly reversed the antidepressant-like activity of CSE, suggesting that the effect may also be related to the dopaminergic

system. Indeed, several studies have suggested the role of sulpiride in altering the antidepressant-like activity of various plant extracts in rodents<sup>66,73</sup> as well as the influence of some flavonoids on the dopaminergic system. <sup>74,75</sup>

The cholinergic system plays an important role in the regulation of central functions such as arousal, attention, cognition, and memory. The cholinergic neurons are connected to the hippocampus and the VTA, and are involved in function of the reward system and mood regulation. Impairment of this system is responsible for development of the cognitive symptoms observed in patients with depression. Pretreatment with atropine significantly reversed the anti-depressant-like activity of CSE, suggesting that the muscarinic cholinergic system is involved in the antidepressant-like activity of the fraction. Similar observations have been made in other studies.<sup>73,76</sup> Moreover, reports are available showing that the antidepressant-like activity of some isolated flavonoids is abolished by atropine, suggesting the involvement of the cholinergic system in their action.<sup>77</sup>

The involvement of nitric oxide in physiological neuronal functions such as synaptic plasticity, including depression and neurological disorders is well recognized. <sup>78,79</sup> NO is formed from L-arginine by the enzyme NO synthase and inhibited by L-arginine analogs. Pretreatment with L-arginine reversed the antidepressant-like activity of CSE, indicating the involvement of NO in the activity of the fraction. NO probably plays a dual role in the modulation of depression in TST and FST, as either increased or decreased synthesis can produce an antidepressant-like effect. <sup>80,81</sup> Although it is generally believed that L-arginine enhances the antidepressant-like activity of various agents, there are cases in which other effects have been reported. For example, moderate doses of L-arginine have antidepressant-like activity, whereas higher doses have no effect. <sup>80</sup> In addition, co-administration of L-arginine with an effective dose of modafinil has been shown to attenuate the antidepressant-like activity of modafinil. <sup>82</sup> Thus, attenuation of the activity of CSE by L-arginine could be related to either the use of high dose (750 mg/kg) or co-administration with the effective dose of CSE. In a parallel experiment, pretreatment with methylene blue (an inhibitor of nitric oxide synthase and an inhibitor of soluble guanylate cyclase (sGC)) did not reverse the antidepressant-like activity of CSE. By contrast, reports in the literature show the antidepressant-like activity of methylene blue in FST, <sup>83,84</sup> which could not be replicated in the present study. Such discrepancies may be related to the multifaceted role of NO in depression and the achievement of an optimal concentration of nitrites in the brain during treatment with agents acting through this pathway to produce an antidepressant-like effect. <sup>81</sup>

Although this study found that adrenergic, dopaminergic, cholinergic and NO may be involved in the mechanism of action of the antidepressant-activity of CSE, only L-arginine and sulpiride were able to return the duration of immobility to control levels, suggesting that dopaminergic and NO are the most likely mechanisms through which CSE mediates its effect.

Individual variation and sample size could be factors that contribute to variations observed in the measurement of corticosterone level. Measuring brain corticosterone than serum level could have given a better picture. Nevertheless, the use of several models for the study could offset these limitations.

#### Conclusions

The results of this study shows that CSE and CSB from the root bark of *C. spinarum* have antidepressant-like activities in TST and FST, the former appearing to be more active than the latter. This was further enhanced by a decrease in serum corticosterone levels. The study suggests that the antidepressant-like activity of CSE might be mediated via α2-adrenergic, D2-dopaminergic receptors, the muscarinic cholinergic system, and the L-arginine-NO pathway. The quantification study suggests that flavonoids, phenols, and alkaloids present in the solvent fractions could be responsible for the observed activity. Further studies are needed to isolate compounds from the CSE fraction that could be used to develop new compounds with anti-depressant activity.

#### **Abbreviations**

ECLIA, Electro-chemiluminescence Immunoassay; FST, Forced swim test; OFT, Open field test; TLC, Thin layer Chromatography; TST, Tail suspension test; TW80, Tween 80.

## **Authors' Details**

Department of Pharmacology and Clinical Pharmacy, School of Pharmacy, College of Health Sciences, Addis Ababa University, P.O. Box 9086, Addis Ababa, Ethiopia.

## **Data Sharing Statement**

The data sets used and/or analyzed during the current work are available and included in this article.

## **Ethics Approval**

The protocol was approved by institutional review board of the School of Pharmacy with Reference no. (ERB/SOP/176/12/2020).

## **Acknowledgments**

The authors are grateful to the Ethiopian Public Health Institute (EPHI) for sponsoring the biochemical analysis (Serum corticosterone). HS would like to acknowledge Addis Ababa University for providing laboratory facilities.

#### **Author Contributions**

Both authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

## **Funding**

No funding was received.

#### **Disclosure**

The authors declare that they have no conflicts of interest in relation to this work nor regarding the publication of this paper. This paper was uploaded to the Addis Ababa University repository as a thesis in March 2022 (<a href="http://etd.aau.edu.et/handle/123456789/31685">http://etd.aau.edu.et/handle/123456789/31685</a>). 85

## References

- 1. Arnaud A, Suthoff E, Tavares RM, Zhang X, Ravindranath AJ. The increasing economic burden with additional steps of pharmacotherapy in major depressive disorder. *Pharmacoeconomics*. 2021;39(6):691–706. doi:10.1007/s40273-021-01021-w
- Filatova EV, Shadrina MI, Slominsky PA. Major depression: one brain, one disease, one set of intertwined processes. Cells. 2021;10(6):1283. doi:10.3390/cells10061283
- 3. Surana AR, Wagh RD. Phytochemical analysis and antidepressant activity of ixora coccinea extracts in experimental models of depression in mice. *Turk J Pharm Sci.* 2018;15:130–135. doi:10.4274/tjps.14622
- 4. Wang PS, Simon G, Kessler RC. The economic burden of depression and the cost-effectiveness of treatment. *Int J Methods Psychiatr Res.* 2003;12 (1):22–33. doi:10.1002/mpr.139
- Santomauro DF, Mantilla Herrera AM, Shadid J, et al. Global prevalence and burden of depressive and anxiety disorders in 204 countries and territories in 2020 due to the COVID-19 pandemic. *Lancet*. 2021;398:1700–1712. doi:10.1016/S0140-6736(21)02143-7
- 6. Friedman ES, Anderson IM. Handbook of Depression. Springer Healthcare Limited. 2nd ed. 2014.
- 7. Haller E, Geier M, Finley P. Antidepressants, pharmacology. In: Encyclopedia of the Neurological Sciences. Elsevier. 2014:219–223.
- 8. Laban TS, Saadabadi A, Monoamine Oxidase Inhibitors (MAOI), Treasure Island (FL); StatPearls; 2021.
- 9. Yildiz A, Gönül AS, Tamam L. Mechanism of actions of antidepressants: beyond the receptors; 2002.
- 10. Petersen TJ, Hopkins G, Kunwar A, Schwartz TL. Drug development, psychotherapy development, and clinical use. In: *Depression: Treatment Strategies and Management*. Philadelphia, PA, US: Taylor & Francis; 2006:41–83.
- 11. Ashraf H, Salehi A, Sousani M, Sharifi MH. Use of complementary alternative medicine and the associated factors among patients with depression. Evid Based Complement Altern Med. 2021;27:2021.
- 12. Lee G, Bae H. Therapeutic effects of phytochemicals and medicinal herbs on depression. Biomed Res Int. 2017;2017. doi:10.1155/2017/6596241
- 13. Farahani MS, Bahramsoltani R, Farzaei MH, Abdollahi M, Rahimi R. Plant-derived natural medicines for the management of depression: an overview of mechanisms of action. *Rev Neurosci.* 2015;26(3):305–321. doi:10.1515/revneuro-2014-0058
- 14. Zeni AL, Zomkowski AD, Dal-Cim T, Maraschin M, Rodrigues AL, Tasca CI. Antidepressant-like and neuroprotective effects of Aloysia gratissima: investigation of involvement of L-arginine-nitric oxide-cyclic guanosine monophosphate pathway. *J Ethnopharmacol*. 2011;137 (1):864–874. doi:10.1016/j.jep.2011.07.009
- 15. Moravec I, Fernandez E, Vlkova M, Milella L. Ethnobotany of medicinal plants of northern Ethiopia. *Boletín Latinoamericano y del Caribe de Plantas Medicinales y Aromáticas*. 2014;13(2):126–134.
- 16. Ansari I, Patil DT, Brief A. Review on phytochemical and pharmacological profile of Carissa spinarum L. Asian J Pharm Clin Res. 2018;11 (9):12–18. doi:10.22159/ajpcr.2018.v11i9.26316
- 17. Nantongo JS, Odoi JB, Abigaba G, Gwali S. Variability of phenolic and alkaloid content in different plant parts of Carissa edulis Vahl and Zanthoxylum chalybeum Engl. *BMC Res Notes*. 2018;11(1):1–5. doi:10.1186/s13104-018-3238-4

18. Teke GN, Kuete V. Acute and subacute toxicities of african medicinal plants. In: *Toxicological Survey of African Medicinal Plants*. Elsevier; 2014:63–98.

- 19. Esubalew ST, Belete A, Lulekal E, Gabriel T, Engidawor E, Asres K. Review of ethnobotanical and ethnopharmacological evidences of some Ethiopian medicinal plants traditionally used for the treatment of cancer. *Ethiop J Health Dev.* 2017;31:161.
- 20. Gidey M, Asfaw Z. Review on ethnobotanical studies on traditional medicinal plants used to treat livestock and human ailments in Tigray Region, Ethiopia. *Adv J Biol Sci Res.* 2015;3:8–36.
- 21. Moges A, Moges Y. Ethiopian common medicinal plants: their parts and uses in traditional medicine ecology and quality control. In: *Plant Science Structure, Anatomy and Physiology in Plants Cultured in vivo and in Vitro*. IntechOpen; 2020:89.
- 22. Tefera BN, Kim Y-D. Ethnobotanical study of medicinal plants in the Hawassa Zuria District, Sidama Zone, Southern Ethiopia. *J Ethnobiol Ethnomed*. 2019;15(1):25. doi:10.1186/s13002-019-0302-7
- 23. Teklehaymanot T. Ethnobotanical study of knowledge and medicinal plants use by the people in Dek Island in Ethiopia. *J Ethnopharmacol*. 2009;124(1):69–78. doi:10.1016/j.jep.2009.04.005
- 24. Kebamo S, Makonnen E, Debela A, Geleta B. Evaluation of diuretic activity of different solvent fractions of methanol extract of Carissa edulis root bark in rats. *Med Chem (Los Angeles)*. 2015;5:472–478. doi:10.4172/2161-0444.1000304
- 25. Nedi T, Mekonnen N, Urga K. Diuretic effect of the crude extracts of Carissa edulis in rats. *J Ethnopharmacol*. 2004;95(1):57–61. doi:10.1016/j.jep.2004.06.017
- 26. Maina GS, Kelvin JK, Maina MB, et al. Antinociceptive properties of dichloromethane: methanolic leaf and root bark extracts of Carissa edulis in rats. J Phytopharmacol. 2015;4(2):106–112.
- 27. Ngulde SI, Sandabe UK, Barkindo AA, Tijjani MB, Hussaini IM. Antinociceptive and Anti-inflammatory activities of the ethanol extract of Carissa edulis vahl. root bark in rats and mice. *Int J Modern Biol Med.* 2013;4(2):85–95.
- 28. Ya'u J, Yaro AH, Malami S, et al. Anticonvulsant activity of aqueous fraction of Carissa edulis root bark. *Pharm Biol.* 2015;53:1329–1338. doi:10.3109/13880209.2014.981280
- 29. Ya'u J, Yaro AH, Abubakar MS, Anuka JA, Hussaini IM. Anticonvulsant activity of Carissa edulis (Vahl) (Apocynaceae) root bark extract. *J Ethnopharmacol*. 2008;120(2):255–258. doi:10.1016/j.jep.2008.08.029
- 30. Woode E, Ansah C, Ainooson GK, Abotsi WM, Mensah AY, Duweijua M. Anti-inflammatory and antioxidant properties of the root extract of Carissa edulis (forsk.) Vahl (Apocynaceae). *J Sci Tech*. 2007;27:5–15.
- 31. Berhanu G, Atalel D, Kandi V. A review of the medicinal and antimicrobial properties of Carissa spinarum L. Am J Biomed Res. 2020;8:54-58.
- 32. Jenson BN Pharmacognostic and antimicrobial studies on the leaf extracts of Carissa Edulis Vahl. (Apocynaceae) and Senna Alata L. (Fabaceae) used in the management of skin infections; 2018:165.
- 33. Ngulde SI, Sandabe UK, Tijjani MB, Barkindo AA, Hussaini IM. Phytochemical constituents, antimicrobial screening and acute toxicity studies of the ethanol extract of Carissa edulis Vahl. root bark in rats and mice. *Am J Res Commun*. 2013;1(9):99–110.
- 34. Ya U, Malami S, Ngura HE, Ngura H. Antidepressant activity of ethanol extract and residual aqueous fraction of Carissa edulis (Apocynaceae) root bark in mice. *Trop J Nat Prod Res.* 2017;1(3):129–132. doi:10.26538/tjnpr/v1i3.9
- 35. National Research Council. Guide for the Care and Use of Laboratory Animals. 8th ed. National Research Council; 2010.
- 36. Sarker SD, Latif Z, Gray AI. Natural product isolation. In: Sarker SD, Latif Z, Gray AI, editors. *Natural Products Isolation*. Totowa, NJ: Humana Press; 2006:1–25.
- 37. Maria R, Shirley M, Xavier C, et al. Preliminary phytochemical screening, total phenolic content and antibacterial activity of thirteen native species from Guayas province Ecuador. *J King Saud Univ Sci.* 2018;30(4):500–505. doi:10.1016/j.jksus.2017.03.009
- 38. Gandagule U, Duraiswamy B, Bhurat M, Nagdev S, Zalke A, Gupta L. Estimation of total phenolic content and total flavonol content in leaves and stem extracts of ventilago maderspatana Gaertn. and leaves, stem and stem bark extracts of ZiZiphus xylopyrus (Retz) Willd. *Inventi J.* 2018;2019:5.
- 39. Van Tan P. The determination of total alkaloid, polyphenol, flavonoid and saponin contents of Pogang gan (Curcuma sp.). *Int J Biol.* 2018;10:4.
- 40. Tabasum S, Khare S, Jain K. Spectrophotometric quantification of total phenolic, flavonoid, and alkaloid contents of abrus precatorius L. Seeds. *Asian J Pharm Clin Res.* 2016;9(2):371–374.
- 41. Dawa I, Mamza U, Sodipo O, Abdulrahman FI, Balami V, Yakubu J. Phytochemical evaluation and gas chromatography-mass spectrometric analysis of column fractions of Carissa edulis leaf extract. Chem Sci. 2021;10:59–68.
- 42. Oduor LP. Investigation of in vitro antiplasmodial activities of Carissa edulis, Azadirachta indica, Cassia siamea and Harrisonia abyssinica on Plasmodium falciparum Doctoral dissertation. Egerton University; 2016:98.
- 43. Diehl K-H, Hull R, Morton D, et al. A good practice guide to the administration of substances and removal of blood, including routes and volumes. *J Appli Toxicol*. 2001;21(1):15–23. doi:10.1002/jat.727
- 44. Porsolt RD, Le Pichon M, Jalfre M. Depression: a new animal model sensitive to antidepressant treatments. *Nature*. 1977;266:730–732. doi:10.1038/266730a0
- 45. Steru L, Chermat R, Thierry B, Simon P. The tail suspension test: a new method for screening antidepressants in mice. *Psychopharmacology*. 1985;85(3):367–370. doi:10.1007/BF00428203
- 46. Can A, Dao DT, Terrillion CE, Piantadosi SC, Bhat S, Gould TD. The tail suspension test. J Vis Exp. 2012;3769. doi:10.3791/3769
- 47. Costa-Nunes JP, Cline BH, Araújo-Correia M, et al. Animal models of depression and drug delivery with food as an effective dosing method: evidences from studies with celecoxib and dicholine succinate. *Biomed Res Int.* 2015;2015. doi:10.1155/2015/596126
- 48. Porsolt RD, Anton G, Blavet N, Jalfre M. Behavioural despair in rats: a new model sensitive to antidepressant treatments. *Eur J Pharmacol*. 1978;47:379–391. doi:10.1016/0014-2999(78)90118-8
- 49. Krishnan V, Nestler EJ. Animal models of depression: molecular perspectives. Curr Top Behav Neurosci. 2011;7:121-147.
- 50. Demissie S, Shibeshi W, Engidawork E. Evaluation of the antidepressant-like activity of solvent fractions of the aerial parts of Hypericum revolutum (Hypericaceae) in rodent models. *Ethiop Pharm J.* 2017;33(2):95–106. doi:10.4314/epj.v33i2.3
- 51. Abiola AL, Ayofe AM, Adedoyin A. Neurobehavioural mechanism of antidepressant effect of methanol stem bark extract of Adansonia digitata (Linn) in tail suspension test in mice. *Adv Pharmacol Pharm*. 2019;7(1):5–13. doi:10.13189/app.2019.070102
- 52. Akinpelu L, Adegbuyi A, Agboola S, et al. Antidepressant activity and mechanism of aqueous extract of Vigna Unguiculata ssp. Dekindtiana (L.) walp dried aerial part in mice. *Int J Neurosci Behavioral Sci.* 2017;5(1):7–18. doi:10.13189/ijnbs.2017.050102

53. Cryan JF, Markou A, Lucki I. Assessing antidepressant activity in rodents: recent developments and future needs. *Trends Pharmacol Sci.* 2002;23 (5):238–245. doi:10.1016/S0165-6147(02)02017-5

- Bangasser DA, Valentino RJ. Sex differences in stress-related psychiatric disorders: neurobiological perspectives. Front Neuroendocrinol. 2014;35:303–319.
- 55. Brigitta B. Pathophysiology of depression and mechanisms of treatment. *Dialogues Clin Neurosci*. 2002;4(1):7–20. doi:10.31887/DCNS.2002.4.1/bbondy
- Malik H, Javaid S, Fawad Rasool M, et al. Amelioration of scopolamine-induced amnesic, anxiolytic and antidepressant effects of ficus benghalensis in behavioral experimental models. *Medicina*. 2020;56(3):144. doi:10.3390/medicina56030144
- 57. Kumar A, Lakshman K, Velmurugan C, Sridhar SM, Saran G. Antidepressant activity of methanolic extract of amaranthus spinosus. *Basic Clin Neurosci*. 2014;5(1):11–17.
- 58. García-Ríos RI, Mora-Pérez A, Ramos-Molina AR, Soria-Fregozo C. Neuropharmacology of secondary metabolites from plants with anxiolytic and antidepressant properties. In: *In Behavioral Pharmacology from Basic to Clinical Research*. IntechOpen; 2020.
- 59. Kennedy DO, Wightman EL. Herbal extracts and phytochemicals: plant secondary metabolites and the enhancement of human brain function. *Adv Nutr.* 2011;2(1):32–50. doi:10.3945/an.110.000117
- 60. Al-Youssef HM, Hassan WHB. Phytochemical and pharmacological aspects of Carissa edulis vahl: a review. *Int J Curr Res Chem Pharm Sci.* 2014;1(9):12–24.
- 61. Umar UA, Hassan LG, Maradun KL. TLC analysis and antioxidant activity of garden egg leaves. *Bayero J Pure Appl Sci.* 2019;12:443–448. doi:10.4314/bajopas.v12i1.67S
- 62. Maletic V, Eramo A, Gwin K, Offord SJ, Duffy RA. The role of norepinephrine and its α-adrenergic receptors in the pathophysiology and treatment of major depressive disorder and schizophrenia: a systematic review. *Front Psychiatry*. 2017;8:42. doi:10.3389/fpsyt.2017.00042
- 63. Stone EA, Lin Y, Rosengarten H, Kramer HK, Quartermain D. Emerging evidence for a central epinephrine-innervated α1-adrenergic system that regulates behavioral activation and is impaired in depression. *Neuropsychopharmacol*. 2003;28(8):1387–1399. doi:10.1038/sj.npp.1300222
- 64. Cottingham C, Wang Q. α2 adrenergic receptor dysregulation in depressive disorders: implications for the neurobiology of depression and antidepressant therapy. *Neurosci Biobehav Rev.* 2012;36(10):2214–2225. doi:10.1016/j.neubiorev.2012.07.011
- 65. Ishola I, Akinyede A, Sholarin A. Antidepressant and anxiolytic properties of the methanolic extract of momordica charantia Linn (Cucurbitaceae) and its mechanism of action. *Drug Res.* 2013;64(7):368–376. doi:10.1055/s-0033-1358712
- Bettio LEB, Machado DG, Cunha MP, et al. Antidepressant-like effect of extract from Polygala paniculata: involvement of the monoaminergic systems. *Pharm Biol.* 2011;49(12):1277–1285. doi:10.3109/13880209.2011.621958
- 67. Mensah JA Antidepressant effect and the possible mechanism (S) of action of secondary metabolites from trichilia monadelpha in murine models Doctoral dissertation. University of Ghana; 2016.
- 68. Kaur R, Chopra K, Singh D. Role of α2 receptors in quercetin-induced behavioral despair in mice. *J Med Food*. 2007;10(1):165–168. doi:10.1089/imf.2005.063
- 69. Butterweck V, Jürgenliemk G, Nahrstedt A, Winterhoff H. Flavonoids from hypericum perforatum show antidepressant activity in the forced swimming test. *Planta Med.* 2000;66(1):3–6. doi:10.1055/s-2000-11119
- 70. Ishola IO, Chatterjee M, Tota S, et al. Antidepressant and anxiolytic effects of amentoflavone isolated from Cnestis ferruginea in mice. *Pharmacol Biochem Behav.* 2012;103(2):322–331. doi:10.1016/j.pbb.2012.08.017
- 71. Dunlop BW, Nemeroff CB. The role of dopamine in the pathophysiology of depression. Arch Gen Psychiatry. 2007;64:327. doi:10.1001/archpsyc.64.3.327
- 72. Ploski JE, Vaidya VA. The neurocircuitry of PTSD and major depression: insights into overlapping and distinct circuit dysfunction a tribute to Ron Duman. *Biol Psychiatry*. 2021;90:109–117. doi:10.1016/j.biopsych.2021.04.009
- 73. Ishola IO, Olayemi SO, Yemitan OK, Umeh EA. Antidepressant and anxiolytic effects of the methanol root extract of Capparis thonningii: involvement of monoaminergic, cholinergic and GABAergic systems. *Drug Res.* 2015;65(4):205–213. doi:10.1055/s-0034-1376963
- 74. Pannu A, Sharma PC, Thakur VK, Goyal RK. Emerging role of flavonoids as the treatment of depression. *Biomolecules*. 2021;11(12):1825. doi:10.3390/biom11121825
- 75. Wang X, Zhang L, Hua L, Xing D, Du L. Effect of flavonoids in scutellariae radix on depression-like behavior and Brain Rewards: possible in dopamine system. *Tinshhua Sci Technol*. 2010;15:460–466. doi:10.1016/S1007-0214(10)70088-2
- 76. Owope T, Ishola I, Akinleye M, Oyebade R, Adeyemi O. Antidepressant effect of cnestis ferruginea Vahl ex DC (Connaraceae): involvement of cholinergic, monoaminergic and L-arginine-nitric oxide pathways. *Drug Res.* 2016;66:235–245. doi:10.1055/s-0035-1565174
- 77. Onasanwo A, Ilenre K, Faborode O. The impact of kolaviron (A bioflavonoid of garcinia kola seed) on depression status in laboratory rodents: roles of monoaminergic systems. *Annals Depress Anxiety*. 2015;2(1):1042.
- 78. Dhir A, Kulkarni SK. Nitric oxide and major depression. Nitric Oxide. 2011;24:125-131. doi:10.1016/j.niox.2011.02.002
- Zhou Q-G, Zhu X-H, Nemes AD, Zhu D-Y. Neuronal nitric oxide synthase and affective disorders. IBRO Report. 2018;5:116–132. doi:10.1016/j. ibror.2018.11.004
- 80. da Silva G, Matteussi AS, Dos Santos ARS, Calixto JB, Rodrigues ALS. Evidence for dual effects of nitric oxide in the forced swimming test and in the tail suspension test in mice. *NeuroReport*. 2000;11(17):3699–3702. doi:10.1097/00001756-200011270-00022
- 81. Inan SY, Yalcin I, Aksu F. Dual effects of nitric oxide in the mouse forced swimming test: possible contribution of nitric oxide-mediated serotonin release and potassium channel modulation. *Pharmacol Biochem Behav.* 2004;77(3):457–464. doi:10.1016/j.pbb.2003.12.024
- 82. Omidi-Ardali H, Badi AG, Saghaei E, Amini-Khoei H. Nitric oxide mediates the antidepressant-like effect of modafinil in mouse forced swimming and tail suspension tests. *J Basic Clin Physiol Pharmacol*. 2021;32(2):25–31. doi:10.1515/jbcpp-2020-0021
- 83. Delport A, Harvey BH, Petzer A, Petzer JP. Methylene blue and its analogues as antidepressant compounds. *Metab Brain Dis.* 2017;32 (5):1357–1382. doi:10.1007/s11011-017-0081-6
- 84. Heiberg IL, Wegener G, Rosenberg R. Reduction of cGMP and nitric oxide has antidepressant-like effects in the forced swimming test in rats. Behav Brain Res. 2002;134:479–484. doi:10.1016/S0166-4328(02)00084-0
- 85. Saif H. Antidepressant-like activity of solvent fractions of the root bark of Carissa spinarum Linn. (Apocynaceae) in rodents. Thesis. Addis Ababa University; 2022.

### Journal of Experimental Pharmacology

# **Dove**press

## Publish your work in this journal

The Journal of Experimental Pharmacology is an international, peer-reviewed, open access journal publishing original research, reports, reviews and commentaries on all areas of laboratory and experimental pharmacology. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

 $\textbf{Submit your manuscript here:} \ \texttt{https://www.dovepress.com/journal-of-experimental-pharmacology-journal-of-experiment$ 

