

# Analysis of the Outcomes of the Screen-Time Reduction in Computer Vision Syndrome: A Cohort Comparative Study [Letter]

Khaled Abdelaziz<sup>1</sup>, Mahrous Shaheen<sup>2</sup>

<sup>1</sup>Department of Ophthalmology, Faculty of Medicine, Beni Suef University, Beni Suef, Egypt; <sup>2</sup>Department of Ophthalmology, Faculty of Medicine, Fayoum University, Fayoum, Egypt

Correspondence: Khaled Abdelaziz, Email [Khaled.ophthalmologist@yahoo.com](mailto:Khaled.ophthalmologist@yahoo.com)

## Dear editor

We have read with great interest the study by Iqbal et al<sup>1</sup> titled “Analysis of the Outcomes of the Screen-Time Reduction in Computer Vision Syndrome: A Cohort Comparative Study”. We would like to congratulate Iqbal et al<sup>1</sup> for their novel study that documented the existence of the screen-induced foveal dysfunction in computer vision syndrome (CVS) using the multifocal electroretinogram (mfERG) examination as they concluded that this screen-induced foveal dysfunction is reversible with strict screen-time reduction, thus improving visual performance.

To a great extent, we really agree with their unique outcomes and conclusions. Meanwhile, despite the fact that Iqbal et al<sup>1</sup> investigated a very important clinical question in the clinical and ophthalmic fields nowadays, and pointed to one of the great drawbacks of the mandated computer system use program; however, we have some concerns that need further explanation and clarification. In addition, some untouched and warranted issues need further discussion.

First, in their conclusion, Iqbal et al<sup>1</sup> stated that the improvements in the mfERG foveal responses were associated with corresponding improvements in the visual performances. We think that this conclusion is missing statistical evidence to document such a relationship between the foveal responses and visual acuity.

Second, we understand that the authors applied extraordinary methods to ensure that the students’ screen-time was limited to only 1 screen-hour daily for 4 weeks. However, no mention has been made about how the authors guaranteed their prescribed methods and we really wonder how they managed to control and assess the students’ use of their digital screens, ie, did the authors observe these students, for example using surveillance cameras? Moreover, the authors mentioned that they allowed the students to watch the TV screens from a proper distance, what was this proper distance in particular?

Another third issue, in the Methods section; Iqbal et al<sup>1</sup> mentioned that they recruited the participants based on a CVS-F3 questionnaire<sup>2,3</sup> and Iqbal’s four major diagnostic criteria for accurate CVS diagnosis.<sup>2-4</sup> They anticipated a large number of students to participate but a limited number actually agreed to be recruited in their trial as the authors faced difficulties in convincing the students to participate in the trial. How did the authors choose these actually recruited participants from the total mentioned anticipated large number of participants?

Furthermore, the authors described certain interesting instructions to the recruited students to change their screen-time and screen-style. Are these instructions applicable for all cases suffering from CVS complaints in general?

We are very grateful to the Editorial Board of the *Clinical Ophthalmology Journal* for publishing such remarkable novel studies and congratulate Iqbal et al<sup>1</sup> for their recent unique publication.

## Disclosure

The authors report no conflicts of interest in this communication.

## References

1. Iqbal M, Soliman A, Ibrahim O, Gad A. Analysis of the outcomes of the screen-time reduction in computer vision syndrome: a cohort comparative study. *Clin Ophthalmol.* 2023;17:123–134. doi:10.2147/OPHT.S399044
2. Iqbal M, Said O, Ibrahim O, Soliman A. Visual sequelae of computer vision syndrome: a cross-sectional case-control study. *J Ophthalmol.* 2021;2021:6630286. doi:10.1155/2021/6630286
3. Iqbal M, Elzembely H, El-Massry A. Computer vision syndrome prevalence and ocular sequelae among medical students: a university-wide study on a marginalized visual security issue. *Open Ophthalmol J.* 2021;15:156–170. doi:10.2174/1874364102115010156
4. Iqbal M, Ibrahim Elzembely H, Said OM. Letter to the editor: “self-reported student awareness and prevalence of computer vision syndrome during COVID-19 pandemic at Al-Baha University” [Letter]. *Clin Optom.* 2022;14:193–194. doi:10.2147/OPTO.S391171

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Clinical Ophthalmology ‘letters to the editor’ section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Clinical Ophthalmology editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

### Clinical Ophthalmology

Dovepress

### Publish your work in this journal

Clinical Ophthalmology is an international, peer-reviewed journal covering all subspecialties within ophthalmology. Key topics include: Optometry; Visual science; Pharmacology and drug therapy in eye diseases; Basic Sciences; Primary and Secondary eye care; Patient Safety and Quality of Care Improvements. This journal is indexed on PubMed Central and CAS, and is the official journal of The Society of Clinical Ophthalmology (SCO). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-ophthalmology-journal>

<https://doi.org/10.2147/OPHT.S405004>