

# Predicting Short-Term Mortality in Older Patients Discharged from Acute Hospitalizations Lasting Less Than 24 Hours

Amalia Lærke Kjær Heltø <sup>1,2</sup>, Emilie Vangsgaard Rosager <sup>1,2</sup>, Martin Aasbrenn <sup>3</sup>, Cathrine Fox Maule <sup>4</sup>, Janne Petersen <sup>4,5</sup>, Finn Erland Nielsen <sup>1</sup>, Charlotte Suetta <sup>3</sup>, Rasmus Gregersen <sup>1,4,5</sup>

<sup>1</sup>Department of Emergency Medicine, Bispebjerg and Frederiksberg Hospital, Copenhagen, Denmark; <sup>2</sup>Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark; <sup>3</sup>Department of Geriatrics and Palliative Medicine, Bispebjerg and Frederiksberg Hospital, Copenhagen, Denmark; <sup>4</sup>Center of Clinical Research and Prevention, Bispebjerg and Frederiksberg Hospital, Copenhagen, Denmark; <sup>5</sup>Section of Biostatistics, Department of Public Health, University of Copenhagen, Copenhagen, Denmark

Correspondence: Amalia Lærke Kjær Heltø, Bispebjerg and Frederiksberg Hospital, Department of Emergency Medicine, Ebba Lunds Vej 40A, Building 67, 2. floor, Copenhagen, NV, 2400, Denmark, Tel +45 60654575, Email [amalia.laerke.kjaer.heltoe.01@regionh.dk](mailto:amalia.laerke.kjaer.heltoe.01@regionh.dk)

**Purpose:** Over coming decades, a rise in the number of short, acute hospitalizations of older people is to be expected. To help physicians identify high-risk patients prior to discharge, we aimed to develop a model capable of predicting the risk of 30-day mortality for older patients discharged from short, acute hospitalizations and to examine how model performance changed with an increasing amount of information.

**Methods:** This registry-based study included acute hospitalizations in Denmark for 2016–2018 lasting  $\leq 24$  hours where patients were permanent residents,  $\geq 65$  years old, and discharged alive. Utilizing many different predictor variables, we developed random forest models with an increasing amount of information, compared their performance, and examined important variables.

**Results:** We included 107,132 patients with a median age of 75 years. Of these, 3.3% ( $n=3575$ ) died within 30 days of discharge. Model performance improved especially with the addition of laboratory results and information on prior acute admissions (AUROC 0.835), and again with comorbidities and number of prescription drugs (AUROC 0.860). Model performance did not improve with the addition of sociodemographic variables (AUROC 0.861), apart from age and sex. Important variables included age, dementia, number of prescription drugs, C-reactive protein, and eGFR.

**Conclusion:** The best model accurately estimated the risk of short-term mortality for older patients following short, acute hospitalizations. Trained on a large and heterogeneous dataset, the model is applicable to most acute clinical settings and could be a useful tool for physicians prior to discharge.

**Keywords:** machine learning, prediction model, register-based, geriatric, emergency medicine, early discharge

## Introduction

The older population is increasing more quickly than any other age-group and by 2050 one in four persons in Europe and North America could be aged  $\geq 65$  years.<sup>1</sup> Currently, older patients already comprise  $\geq 40\%$  of acute emergency department (ED) contacts.<sup>2–4</sup> Thereby, an associated rise in the need for urgent care of older patients over the next few decades is to be expected and will require optimization and development of our current health-care system, eg, through improved preventive care along with fewer and possibly shorter hospitalizations. Shorter length of stay not only leads to lower hospitalization costs but is also associated with a lower risk of many adverse events, such as nosocomial infections, delirium, and complications related to immobility like deep vein thrombosis and loss of muscle mass.<sup>5–8</sup> The average length of stay in hospitals has decreased over the last decade, particularly for older age groups.<sup>9,10</sup> However, although shorter hospitalizations are in many ways to be preferred, concern has been raised that premature discharges among older patients could lead to preventable readmissions and even deaths.<sup>2,9–13</sup> Previous studies have identified

several factors associated with short-term mortality following hospital discharge, such as age, frailty, comorbidity burden, number of medications, and laboratory results.<sup>13–16</sup> To our knowledge, no prediction model has been developed that focuses on the short-term mortality of older patients following early discharge. Valid prediction models may assist clinicians in safely planning early discharge for the right group of older patients. Therefore, the aim of this study was to develop a reliable prediction algorithm using random forest to predict the risk of 30-day mortality for acutely hospitalized older patients following early discharge from hospital ( $\leq 24$  hours). Further, we examined how predictive performance changed with an increasing number of variables.

## Methods

### Design and Study Population

This was a registry-based cohort study. We included contacts where the patients were permanent residents  $\geq 65$  years of age who were discharged alive from acute hospitalizations lasting  $\leq 24$  hours between January 1, 2016 and December 1, 2018 in Denmark. To standardize the cohort across, different organizational hospital structures and mimic patients who could be handled in shared-entrance emergency departments, we included only hospitalizations with discharges from emergency, medical, or surgical departments. To consider the entire length of the patients' hospital stay, we combined sequential Danish National Patient Registry (DNPR) contacts within  $\leq 4$  hours, considering contacts with both somatic and psychiatric departments and all types of inpatient/outpatient and acute/elective contacts.<sup>4</sup> We only included contacts with at least one acute and one inpatient contact. Thereby, patients with a contact with the ED without outright hospitalization were not included. Hospitalizations from the Central Denmark Region were excluded, since we did not have access to their laboratory results. Hospitalizations from the remaining regions were excluded if no laboratory tests were registered during the hospitalization. Hospitalizations with registered laboratory results were excluded if they were still missing one or more laboratory test that was otherwise present in  $\geq 90\%$  of the study population. Hospitalizations where the patient was discharged from psychiatric, ear–nose–throat, dermatology, or ophthalmology departments were excluded, after an initial overview revealed a discrepancy in discharge diagnoses from these departments compared to the other medical, emergency, and surgical departments. Hospitalizations without information on the sociodemographic variables of patients were excluded. Hospitalizations where the patient had registered encounters for palliative care within 5 years prior to and including their index hospitalization were excluded. Finally, all but the first eligible hospitalization for each patient were excluded.

### Setting

Denmark has a tax-funded universal health-care system that provides access for every citizen to primary, hospital, and home-care services.<sup>17</sup> Citizens in need of acute hospital care are almost exclusively referred, either by their general practitioners or emergency medical services, and are generally first assessed in the ED, after which they are either discharged or hospitalized.<sup>17</sup> Each citizen has a unique identification number (CPR number), which allows for the continuous tracking of Danish citizens over time as well as linkage among different national registers.<sup>18,19</sup>

### Data Sources and Predictor Variables

Information on the time and date of admission and discharge, the primary discharge diagnosis, and comorbidities at the time of hospitalization were available through the DNPR. Laboratory results from the index hospitalization were available through the Register of Laboratory Results for Research. Information on prescriptive medication was collected via the Danish National Prescription Registry. Sociodemographic and economic factors were collected via the DNPR, Central Person Register, Income Statistics Register, Building and Housing Register, and Danish Education Register from the end of the calendar year prior to the date of admission.

Diseases from the Charlson Comorbidity Index (CCI) were used individually, resulting in a total of 17 binary categorical comorbidity variables instead of a single CCI score. For descriptive purposes, the CCI score was calculated using the Quan modified CCI score.<sup>20</sup> Patients were considered as having a comorbidity if the relevant ICD-10 diagnostic code was registered as a primary or secondary diagnosis up to 5 years prior to their index hospitalization. Living situation

was derived from combining the marital status of the patient and the number of people registered living at the same address as the patient. Education was assessed using the International Standard Classification of Education (ISCED) 2011. For descriptive purposes, disposable income was compared to quintiles of the general population stratified by age. Number of prescription drugs was examined using the Anatomical Therapeutic Chemical Classification System (ATC) level 4, collected up to 6 months prior to hospitalization. Since four-digit ICD-10 grouping has been shown to be an inadequate way of categorizing diseases, we conducted our own clinically meaningful grouping of discharge diagnoses for our population ([Supplementary Materials; Table S1](#)).<sup>4,21,22</sup> We selected 14 laboratory blood tests to represent a standard ED package, and reference values were drawn directly from the respective laboratories. Initial screening of the frequency of these tests was done: tests missing for >10% of the population were handled as categorical variables with up to four levels: “missing”, “below normal”, “normal”, and “above normal.” Otherwise, the tests were handled as continuous. Continuous laboratory tests were hemoglobin, thrombocytes, leukocytes, C-reactive protein, creatinine, eGFR, sodium ion, and potassium ion. INR, albumin, bilirubin, alanine transaminase, alkaline phosphatase, and glucose were categorical variables. For hospitalizations with repeated laboratory results, the last result prior to discharge was chosen.

## Statistics

Data preparation and descriptive analyses were performed using SAS 9.4 (SAS Institute, Cary, NC, US) while R version 4.1.3 (R Core Team, Vienna, Austria), was used for model development and visualizations.<sup>23,24</sup> For descriptive purposes, continuous variables are displayed as medians and quartiles (Q1–Q3), while categorical variables are presented as percentages. Risk differences between proportions are reported using exact differences between proportions with 95% CIs. For continuous variables, the Hodges–Lehmann estimator of location of shift was used to display the median of differences and appertaining 95% CIs. Due to the size of the cohort and complexity of the calculation, this was done on 50,000 patients randomly selected from the cohort.

Random forest was chosen as the classification algorithm.<sup>25,26</sup> Initially, each patient was randomly assigned to either the training (70%), validation (15%), or test dataset (15%) while ensuring an equal distribution of survivors and nonsurvivors between the datasets. The dataset suffered from class imbalance, a common problem when utilizing random forest for classification purposes.<sup>27</sup> Therefore, we utilized the random forest quantile classifier during model training and hyperparameter tuning on the training set.<sup>28</sup> Hyperparameter optimization was explored in a stepwise fashion through narrowing grid searches for each model level until no further improvement could be achieved based on out-of-bag Brier score. Hyperparameter-optimization was performed on number of trees, tree depth, node size, and number of variables available to choose from at each split. Final testing for each model was done on the assigned test set with no weighing or sampling of survivors/nonsurvivors. The validation set was set aside as a secondary dataset on which to retest the model in the case that the results from testing on the test set showed the need for further model alterations. A cutoff point was selected for each model using the Youden index, and discriminative model performance was measured by ROC (receiver-operating characteristic) curves, AUROCs (area under receiver-operating characteristic) curves, Brier scores, sensitivity, and specificity on the test set. To gain insight into the model’s decision-making process, we utilized variable importance and partial dependence plots.

Variables were selected based on availability at discharge through electronic health records and expected correlation with mortality, as suggested by previous studies.<sup>13–16</sup> To investigate how model performance changed with additional information, variables were gradually added groupwise in an estimated order of most to least easily available to the physician. The order in which variables were added was: level 0 (age, sex), level 1 (level 0 + discharge diagnosis, specialty), level 2 (level 1 + time of day at discharge, day of the week at time of discharge, length of stay), level 3 (level 2 + laboratory results from the index hospitalization, previous hospitalization during the last 30 days prior to index hospitalization), level 4 (level 3 + diseases of Charlson Comorbidity Index, number of different prescription drugs redeemed within 6 months prior to the index hospitalization), and level 5 (level 4 + living situation, marital status, education, yearly disposable income). Finally, we developed a one-step backward-selection model using only variables from the level 5 model with positive importance.

## Results


### Study Population Baseline Characteristics

We considered 253,534 hospitalizations, of which 108,322 were excluded due to either missing laboratory results from the index hospitalization (including all hospitalizations from Central Denmark Region), discharge from psychiatric, ear-nose-throat, dermatology, or ophthalmology departments, lacking information on socioeconomic variables, or encounters for palliative care registered within 5 years prior to/including index hospitalization. Finally, we excluded all but the first eligible hospitalization for each patient, and the final population was comprised of 107,132 patients. Of these, 3,575 (3.3%) died within 30 days of discharge (Figure 1).

Descriptive analyses revealed notable differences between nonsurvivors and survivors (Table 1). Nonsurvivors were older than survivors and used more prescription medications. Nonsurvivors were more likely to have been acutely admitted to hospital within 30 days prior to the index hospitalization and were more often single and living alone compared to survivors. Discharge diagnoses also differed between the two groups (Supplementary Materials; Table S1). For example, “Dehydration. Other disorders of fluid, electrolyte, and acid–base balance” was more common among the nonsurvivors, while “Encounters for medical observation for suspected diseases and conditions ruled out” was more common among the survivors. Nonsurvivors had a greater occurrence of all comorbidities than survivors, but notable differences between the two groups included dementia, congestive heart failure, and cancer (Supplementary Materials; Table S2). Notable laboratory tests include higher C-reactive protein and creatinine and lower hemoglobin, albumin, and eGFR for nonsurvivors than survivors (Table 2). Comparison between those included and excluded from the study revealed a slightly lower mortality rate in those excluded from the study compared to those included (Supplementary Materials; Table S3). The excluded group had a larger proportion of men and was also younger than those who were included. There was no difference in Charlson Comorbidity Index scores.

### Predictive Performance

The level 0, 1, and 2 models had AUROCs of 0.723–0.755, similar ROC curves, and Brier scores of 0.0311–0.0313 (Figure 2 and Table 3). The level 3 model achieved an AUROC of 0.835 and a Brier score of 0.0286. The level 4, 5, and backward-selection



**Figure 1** Flowchart of cohort selection process, with inclusion and exclusion criteria.

**Table 1** Population characteristics of the total cohort and according to survival status

	Subcategories	Survivors, n= 103,557 (96.7%)	Nonsurvivors, n=3575 (3.3%)	Risk difference (95% CI)*
Sex	Female	52.7%	51.6%	1.1% (-0.6, 2.8)
Age, years	Median (Q1–Q3)	75.0 (70.0–82.0)	83.0 (76.0–89.0)	-6.5 (-7.0, -6.0)
Acute hospitalization within last 30 days	Yes	9.6%	26.0%	-16.4% (-17.8, -14.9)
Prescription drugs, n	Median (Q1–Q3)	7.0 (4.0–11.0)	10.0 (7.0–14.0)	-3.0 (-3.0, -3.0)
Charlson Comorbidity Index score	Median (Q1–Q3)	0.0 (0.0–2.0)	2.0 (0.0–3.0)	-1.0 (-1.0, -1.0)
Marital status	Divorced	15.8%	14.7%	1.1% (-0.1, 2.3)
	Married	52.0%	36.3%	15.7% (14.1, 17.3)
	Unmarried	5.4%	6.9%	-1.5% (-2.3, -0.7)
	Widow/widower	26.8%	42.1%	-15.3% (-17.0, -13.7)
Living situation	Couple, living with at least one other person	53.0%	33.0%	20.1% (18.5, 21.7)
	Single, living alone	43.8%	64.0%	-10.1% (-21.7, -18.5)
	Single, living with at least one other person	3.3%	3.3%	0.0 (-0.6, 0.6)
Education	Missing or no primary education	3.0%	7.7%	-4.7% (-5.6, -3.8)
	Primary or lower secondary (ISCED 0–2)	38.8%	45.6%	-6.8% (-8.5, -5.2)
	Upper secondary or postsecondary, non-tertiary (ISCED 3–4)	38.5%	32.4%	6.1% (4.5, 7.6)
	Short-cycle tertiary or above (ISCED 5–8)	19.8%	14.3%	5.5% (4.3, 6.6)
Disposable income, DKK	Median (Q1–Q3)	170,728 (134,587–220,142)	172,242 (142,728–207,262)	-312 (-3154.9, 2531.3)
Time of day of discharge	08:00–15:59	53.7%	52.7%	1.1% (-0.6, 2.7)
	16:00–23:59	40.7%	42.8%	-2.1% (-3.8, -0.4)
	24:00–7:59	5.6%	4.5%	1.0% (0.4, 1.7)
Day of week of discharge	Monday–Friday	77.0%	75.7%	1.3% (-0.2, 2.7)
	Saturday–Sunday	23.0%	24.3%	-1.3% (-2.7, 0.2)
Length of stay, hours	Median (Q1–Q3)	11.0 (5.0–19.0)	13.0 (6.0–20.0)	-1.0 (-1.0, -1.0)

**Notes:** \*For categorical variables, risk difference is defined as the absolute difference between proportions (survivors – nonsurvivors). For continuous variables, the median of difference is displayed using the Hodges–Lehmann estimator of location of shift (survivors – nonsurvivors).

**Table 2** Laboratory results from index hospitalization of the total cohort and according to survival status

	Subcategories	Survivors, n=103,557 (96.7%)	Nonsurvivors, n=3575 (3.3%)	Risk Difference (95% CI)*
Hemoglobin, B (mmol/L)	Median (Q1–Q3)	8.2 (7.4–8.8)	7.4 (6.4–8.3)	0.7 (0.7, 0.8)
	Above normal	2.5%	1.8%	0.7% (0.2, 1.1)
	Normal range	64.3%	39.4%	24.9% (23.3, 26.5)
Thrombocytes, B ( $10^9/L$ )	Below normal	33.2%	58.7%	-25.5% (-27.2, -23.9)
	Median (Q1–Q3)	238.0 (194.0–292.0)	256.0 (195.0–332.0)	-18.5 (-23.0, -14.0)
	Above normal	6.7%	15.5%	-8.7% (-9.9, -7.6)
Leukocytes, B ( $10^9/L$ )	Normal range	86.5%	74.0%	12.6% (11.1, 14.0)
	Below normal	6.7%	10.6%	-3.8% (-4.8, -2.8)
	Median (Q1–Q3)	7.8 (6.3–9.7)	8.9 (6.8–11.7)	-1.0 (-1.2, -0.9)
C-reactive protein, P (mg/L)	Above normal	32.8%	49.5%	-16.7% (-18.4, -15.0)
	Normal range	65.7%	48.6%	17.1% (15.4, 18.8)
	Below normal	1.4%	1.8%	-0.4% (-0.9, 0.1)
Creatinine, P ( $\mu\text{mol/L}$ )	Median (Q1–Q3)	5.0 (2.9–18.0)	21.0 (6.5–52.0)	-10.4 (-11.6, -9.1)
	Above normal	38.1%	69.8%	-31.8% (-33.3, -20.2)
	Normal range	61.9%	30.2%	31.8% (30.2, 33.3)
Creatinine, P ( $\mu\text{mol/L}$ )	Median (Q1–Q3)	79.0 (66.0–97.0)	87.0 (66.0–120.0)	-7.5 (-9.0, -6.0)
	Above normal	23.2%	39.0%	-15.7% (-17.4, -14.1)
	Normal range	71.8%	52.3%	19.5% (17.8, 21.2)
	Below normal	5.0%	8.7%	-3.7% (-4.7, 2.8)


(Continued)

**Table 2** (Continued).

	Subcategories	Survivors, n=103,557 (96.7%)	Nonsurvivors, n=3575 (3.3%)	Risk Difference (95% CI)*
eGFR (mL/min)	Median (Q1–Q3)	72.0 (56.0–85.0)	62.0 (41.0–81.0)	8.0 (7.0, 9.0)
	Normal range	71.0%	52.8%	18.2% (16.5, 19.8)
	Below normal	29.0%	47.2%	–18.2% (–19.8, –16.5)
Sodium ion, P (mmol/L)	Median (Q1–Q3)	139.0 (137.0–141.0)	139.0 (136.0–142.0)	0.0 (0.0, 0.0)
	Above normal	1.9%	8.6%	–6.7% (–7.6, –5.6)
	Normal range	76.6%	62.0%	14.5% (12.9, 16.1)
Potassium ion, P (mmol/L)	Median (Q1–Q3)	3.9 (3.7–4.2)	4.0 (3.7–4.3)	–0.1 (–0.1, 0.0)
	Above normal	6.4%	13.5%	–7.1% (–8.2, 6.0)
	Normal range	83.2%	72.9%	10.3% (8.8, 11.8)
INR	Median (Q1–Q3)	1.0 (1.0–1.1)	1.1 (1.0–1.2)	0.0 (–0.1, 0.0)
	Above normal	13.1%	19.6%	–6.5% (–7.8, –5.1)
	Normal range	80.9%	72.4%	8.5% (7.0, 10.0)
Albumin, P (g/L)	Median (Q1–Q3)	36.0 (33.0–40.0)	32.0 (28.0–36.0)	4.0 (4.0, 4.0)
	Above normal	2.3%	0.6%	1.7% (1.4, 1.9)
	Normal range	64.3%	39.1%	25.2% (23.6, 26.8)
Bilirubin, P (µmol/L)	Median (Q1–Q3)	8.0 (6.0–12.0)	9.0 (6.0–14.0)	–1.0 (–1.0, –1.0)
	Above normal	1.9%	5.4%	–3.5% (–4.3, –2.7)
	Normal range	73.5%	74.5%	–1.0% (–2.5, 0.5)
Alanine transaminase, P (U/L)	Median (Q1–Q3)	22.0 (16.0–30.0)	20.0 (13.0–30.0)	2.5 (2.0–3.0)
	Above normal	5.1%	8.1%	–2.9% (–3.8, –2.0)
	Normal range	77.9%	71.8%	6.1% (4.6, 7.6)
Alkaline phosphatase, P (U/L)	Median (Q1–Q3)	74.0 (61.0–93.0)	87.0 (68.0–123.0)	–13.5 (–15.0, –12.0)
	Above normal	13.2%	30.4%	–17.2% (–18.7, –15.6)
	Normal range	71.7%	58.3%	13.5% (11.8, 15.1)
Glucose, P (mmol/L)	Median (Q1–Q3)	6.4 (5.7–7.5)	6.6 (5.8–8.0)	–0.2 (–0.2, –0.1)
	Above normal	28.6%	34.5%	–5.9% (–7.4, –4.3)
	Normal range	59.7%	51.4%	8.3% (6.6, 10.0)
	Below normal	0.2%	0.5%	–0.3% (–0.5, 0.0)
	Missing	11.4%	13.5%	–2.2% (–3.3, –1.0)

**Notes:** \*Median difference is displayed using the Hodges–Lehmann estimator of location of shift (survivors – nonsurvivors). For subcategories, risk difference is defined as the absolute difference between proportions (survivors – nonsurvivors).

model (excluding 54 variables with no or negative importance) achieved AUROCs of 0.859–0.861, Brier scores of 0.0284–0.0286, and shared indistinguishable ROC curves. The specificities for the level 0, 1, and 2 models were 0.681–0.710, and their sensitivities were 0.669–0.680. The level 3 model achieved a specificity of 0.771 and a sensitivity of 0.737, while the level 4, 5, and backward-selection model had specificities of 0.726–0.755 and sensitivities of 0.803–0.836. Model performances on the validation set were similar to performances on the test set ([Supplementary Materials; Table S4](#)). For the level 0, 1, and 2 models,




**Figure 2** Receiver-operating characteristic curves and appertaining area under curve for the seven models when evaluated on the test set.  
**Abbreviations:** AUROC, area under receiver-operating characteristic curve; BSM, backward-selection model.

median risk predictions were 2.2% for survivors and 4.7%–5.3% for nonsurvivors ([Supplementary Materials; Table S5](#)). Median risk predictions from the level 3 model were 1.5% for survivors and 8.7% for nonsurvivors. For the level 4, 5, and backward-selection model, median predictions for survivors ranged between 1.3%–1.4% and 10.0%–10.7% for nonsurvivors. The models were well calibrated below predicted probabilities <25% - the range in which most of the predictions were located ([Supplementary Materials; Figure S1](#)).

**Table 3** Model performance on test set: area under receiver-operating characteristic curve (AUROC), Brier score, specificity, and sensitivity for each model

	AUROC	Brier score	Specificity*	Sensitivity*
Level 0 (age, sex)	0.723	0.0313	0.694	0.669
Level 1 (level 0 + discharge diagnosis, specialty)	0.755	0.0311	0.681	0.710
Level 2 (level 1 + time of day at discharge, day of the week at time of discharge, length of stay)	0.754	0.0311	0.710	0.680
Level 3 (level 2 + laboratory results from the index hospitalization, previous hospitalization during the last 30 days prior to index hospitalization)	0.835	0.0286	0.771	0.737
Level 4 (level 3 + diseases of Charlson Comorbidity Index, number of different prescription drugs redeemed within 6 months prior to index hospitalization)	0.860	0.0284	0.726	0.836
Level 5 (level 4 + living situation, marital status, education, disposable yearly income)	0.861	0.0284	0.751	0.818
Backward-selection model (variables from level 5 model with positive variable importance)	0.859	0.0285	0.755	0.803

**Notes:** \*The Youden index was used to select the cutoff point from which the displayed specificities and sensitivities are derived.



**Figure 3** Variable importance relative to the most important variable—age. Top 20 ranking variables.

**Abbreviation:** LRTI, lower respiratory tract infections.

## Important Variables

Variable importance revealed that from the level 5 model, nine of the 20 most important features were laboratory results, three were related to acute health status, four were related to chronic health status, four were sociodemographic, and none were administrative (Figure 3; [Supplementary Materials; Table S6](#)). The partial dependence plots, which show the marginal effect of a variable on predicted outcome, revealed that a comorbidity of dementia resulted in higher mortality predictions, along with older age, higher levels of C-reactive protein, and previous acute admissions 30 days prior, to mention a few ([Supplementary Materials, Figure S2](#)).<sup>26</sup>

## Discussion

Using nationally representative data from more than 100,000 patients, we developed random forest models to predict the risk of short-term mortality of older patients after early discharge from acute hospitalizations. Predictive ability improved especially with the addition of laboratory results from hospitalization and information on previous acute admissions up to 30 days prior (level 3), and again with the addition of comorbidities and number of prescription drugs (level 4). Model performance did not improve with the addition of sociodemographic variables, other than age and sex (level 5). The model with all variables (level 5) found the most important variables to be age, a comorbidity of dementia, number of different prescription drugs, C-reactive protein, and eGFR. The observed mortality rate was 3.3%.

Predictions for survivors and nonsurvivors became more distinct with the addition of variables, but when predicted risk of mortality increased, calibration also declined. However, considering that the vast majority of patients were predicted to have a 30-day mortality risk of  $\leq 25\%$ , overall model calibration must be considered satisfactory for the level



3, 4, 5, and backward-selection model. This difficulty in gauging short-term mortality in patients at higher risk of mortality is a known problem for statisticians and clinicians, and is yet to be solved.<sup>29</sup> Considering that patients who are discharged soon after acute admissions have been assessed and deemed fit for discharge by a clinician, it seems reasonable that we did not achieve near-perfect discriminative abilities and that median predicted risk of nonsurvivors was 10.5% (level 5 model), especially when considering the heterogeneity of the cohort and lack of clinical information from the hospitalization. The level 0, 2, and 3 models achieved higher specificity than sensitivity, while the level 1, 4, 5, and backward-selection models achieved higher sensitivity than specificity. The level 4 model achieved good discrimination (AUROC 0.860) and was comparable to other models that predict mortality in older patients, though we have not been able to find other models that focus on early discharge.<sup>30–33</sup> However, besides improving predictive performance, introducing comorbidities and number of prescription medications (level 4) makes the model more labor-intensive for the physician, as this information would often require the manual tallying of the patient's comorbidities and medications. The level 5 model performed comparably to the level 4 model, but with slightly higher specificity and lower sensitivity. Considering the additional work required to attain the sociodemographic information and the minimal difference in performance between the two models, the level 4 model must be considered superior. The backward-selection model also performed comparably to the level 4 model while only requiring half as many variables, and could possibly be more easily implementable in a clinical setting.

Rather than applying imputation, we excluded patients with no laboratory results because of the assumption that a majority of these tests were missing not at random: in Denmark, most patients have a standardized laboratory package taken upon admission to hospital, and it can be assumed that hospitalizations without any tests were of a less severe character. This assumption is supported by the fact that those excluded from the study were younger and had a slightly lower mortality rate than those included. Many older patients have multiple comorbidities and underlying health conditions that can present in many different ways in an acute hospital setting.<sup>34</sup> Therefore, rather than only including patients with a specific diagnosis, we included patients from emergency, medical, and surgical departments with many different discharge diagnoses and comorbidities. This approach is concordant with the future treatment of acute patients in Denmark, in which all patients will be seen in the joint emergency department for initial diagnosis and treatment.<sup>35</sup> Random forest was chosen for model development as it has been shown to potentially achieve higher precision for large, complex data structures containing many variables with complicated interactions, such as ours, compared to other algorithms.<sup>36–38</sup> As computational capacity and data availability increases, the predictive ability of machine-learning models such as random forest will continue to improve, and these are likely to become more commonly used in the field of medical research.<sup>39</sup>

A comorbidity of dementia was very high-ranking in terms of variable importance, and is—along with cognitive decline—a known predictor of poor outcomes in older patients.<sup>30,40–42</sup> This high ranking likely partly contributed to the correlation between dementia and age and possibly frailty, which analyses of variable importance do not account for.<sup>43–46</sup> We had no information on where patients were discharged to, which could hold significant value. However, in our population, 3.3% were registered as single, but living with at least one other person, which is concordant with the proportion of people above the age of 65 years living in nursing homes in Denmark.<sup>47</sup> Though patients were excluded if they were receiving palliative care, it is plausible that some were willingly discharged from hospital despite a poor prognosis, with the intent of dying at home. We did not exclude patients with a comorbidity of cancer, since they comprise a considerable part of older patients. However, cancer and metastatic disease were high-ranking in terms of variable importance, and short-term mortality for patients with a history of cancer and/or metastatic disease is known to differ from patients without.<sup>30,48</sup> Recent acute admission and number of prescription drugs were also important variables, and are both possible indicators of patient morbidity and disease burden. Previous studies found associations between number of medications and mortality, but we could not find studies that examined previous recent admissions as a prognostic factor of mortality.<sup>14,31</sup> A discharge diagnosis of pneumonia and a self-reported feeling of physical fatigue have also previously been shown to be independent prognostic factors of death in older patients.<sup>13,33,49</sup> Previous studies also found a correlation between mortality and routine laboratory tests from hospitalization, such as C-reactive protein, creatinine, sodium, hemoglobin, and glucose.<sup>15,50–52</sup> We have not been able to find studies that have examined eGFR, bilirubin, or alkaline phosphatase as prognostic factors of mortality, though these were all high-ranking variables in our

model. Adding administrative information from the hospitalization did not improve predictive ability. Adding socio-demographic information (apart from age and sex) did not improve overall model performance, although marital status and living situation were high-ranking variables. This discrepancy between high variable importance and minimal improvement of model performance can possibly be partly attributed to variable interactions between marital status and living situation and other variables, such as age, though previous studies have shown marital status and living situation to be independent risk factors of mortality.<sup>13,53,54</sup> Although initial testing showed no significant difference in mortality between males and females, sex did have a high importance ranking and male sex was associated with slightly higher mortality predictions than female sex, which is supported by previous findings.<sup>13,31,55</sup> Our mortality rate is concordant with previous studies that reported 30-day mortality rates of 3%–5%, though none of these focused on early discharge.<sup>13,50,56</sup>

## Strengths and Limitations

This study had good statistical power with the inclusion of more than 100,000 patients through a 3-year period, covering almost the entirety of Denmark. Registry-based studies in Denmark are known to have almost complete follow-up.<sup>18</sup> Furthermore, the models are trained on a highly heterogeneous cohort consisting of hospitalizations from a wide range of medical and surgical specialties. This, coupled with the fact that EDs in Denmark diagnose and treat patients from most medical and surgical specialties, makes the model applicable to most acute clinical settings as a useful tool for physicians prior to discharge.<sup>35</sup> We lacked possibly important factors, such as the patient's vital signs from the index hospitalization (eg, blood pressure, pulse and respiratory rate, temperature, oxygen saturation, Glasgow Coma Scale), presenting symptoms, or information on frailty and functional impairment, which have previously been shown to be strongly correlated with short-term mortality.<sup>16,32,40,57</sup> Future models could include these parameters, possibly improving predictive performance.<sup>58</sup> Though we only included hospitalizations with a duration of  $\leq 24$  hours, we cannot account for randomness in the length of stay, eg, due to inconvenient timing of discharge (nights) or a delay in the discharge process caused by differences in daily workload and patient flow. The department from which patients were discharged is also liable to some randomness due to overcrowding or other administrative difficulties. We excluded hospitalizations from the Central Denmark Region since we did not have access to their laboratory results. However, a previous study of sociodemographic and health-related homogeneity in Denmark found minimal discrepancy among Danish regions, and the cohort should thereby be representative of the entirety of Denmark.<sup>59</sup> We excluded patients who were missing a specific laboratory test that was present in >90% of the study population, to avoid imputation. Consequently, the model is only applicable to patients with a laboratory package similar to the one in our model. Few of the 66 discharge diagnoses were found to be important in the level 5 model, while none of the broader diagnostic groups, eg, "Other diseases of the respiratory system", had positive variable importance, indicating that this was an insufficient way of categorizing disease. Due to the sheer number and variety of diagnostic codes in our data, we could not include each patient's specific discharge diagnosis in our model. Consequently, the new diagnostic groupings may have led to selection bias and affected performance accordingly. Another contributing factor could be misclassification of diagnoses in a busy hospital environment, but the validity of diagnostic registration in Danish medical departments has previously been examined and deemed satisfactory.<sup>60</sup>

## Implications

We have developed a model with good predictive ability that can detect patients at risk of short-term mortality following early discharge. The patients in the cohort were assessed by physicians and deemed well enough to be discharged early, and should thus be expected to have a good short-term prognosis. Despite this, the observed 30-day mortality rate was 3.3%. Thus, integration of the model into the Danish electronic health record systems would make this model a useful additional tool for physicians to aid in the safe planning of early discharge for the right group of older patients. Patients at high estimated risk could qualify for interventions such as continued hospitalization, postdischarge follow-up appointments at the hospital or at home, or discharge to rehabilitation facilities instead of their own homes. With limited hospital capacity and an increasing number of acute hospitalizations of older patients, physicians will need to carefully consider which patients will benefit from continued hospitalization. However, even with the best possible treatment, care, and

follow-up, certain deaths will remain unavoidable, as it is ultimately impossible to change the natural life course of some patients. This study also shows that random forest prediction modeling is feasible and meaningful for a heterogeneous cohort of elderly ED patients. Before clinical testing and possible implementation, we recommend redeveloping the model with the inclusion of relevant clinical information, such as vital signs, frailty, symptoms, and where the patient is discharged to.

## Conclusion

In this registry-based cohort study, we developed random forest models capable of predicting the risk of short-term mortality of older patients following hospitalizations  $\leq 24$  hours and examined how an increasing amount of information changed the predictive ability. Model performance improved substantially with the addition of laboratory results and information on previous acute hospitalization, and further with the addition of comorbidities and number of prescription drugs. The addition of sociodemographic variables, apart from age and sex, did not improve predictive ability. Important variables included age, a comorbidity of dementia, number of different prescription drugs, C-reactive protein, and eGFR. Trained on a large and highly heterogeneous dataset, the best model achieved good discriminatory ability (AUROC 0.860) and could be a useful tool for physicians prior to early discharge in most acute clinical settings.

## Data Sharing

Register-based data is protected by Danish legislations and cannot be shared neither publicly, nor privately. Danish research institutions can apply for equivalent data through the Danish Health Data Authority and Statistics Denmark.

## Ethics Approval

Ethics approval is not required for registry-based studies in Denmark. The project was approved by Statistics Denmark (project 707838), Danish Health Data Authority (FSEID-00004732), and Data Protection Agency (P-2019-616).

## Funding

The project was financed by the Bispebjerg and Frederiksberg Hospital Research Committee and the Department of Emergency Medicine. The financial support was used for salary support. The funding sources did not play any part in the design process, conduction, or analysis of the project.

## Disclosure

The authors have no conflicts of interest to declare for this work.

## References

1. United Nations Department of Economic and Social Affairs Population Division. World population ageing 2019 (ST/ESA/SER.A/444); 2020.
2. Andersen AL, Houliind MB, Nielsen RL, et al. Optimization of Nutrition And Medication (OptiNAM) for acutely admitted older patients: protocol for a randomized single-blinded controlled trial. *Trials*. 2021;22(1):616. doi:10.1186/s13063-021-05456-6
3. West Health Institute. Geriatric emergency department factsheet (2013 HCUP-NEDS data). <https://www.westhealth.org/resource/geriatric-emergency-department-ged-fact-sheet/>. Accessed May 20, 2022.
4. Gregersen R, Maule CF, Bak-Jensen HH, et al. Profiling bispebjerg acute cohort: methodology, acute contact characteristics, and comparisons to urban and rural hospitals in Denmark. *Clin Epidemiol*. 2022;14:1713–1717. doi:10.2147/CLEP.S338149
5. Bail K, Draper B, Berry H, Karmel R, Goss J. Predicting excess cost for older inpatients with clinical complexity: a retrospective cohort study examining cognition, comorbidities and complications. *PLoS One*. 2018;13(2):e0193319. doi:10.1371/journal.pone.0193319
6. Kaboli PJ, Go JT, Hockenberry J, et al. Associations between reduced hospital length of stay and 30-day readmission rate and mortality: 14-year experience in 129 veterans affairs hospitals. *Ann Intern Med*. 2012;157(12):837. doi:10.7326/0003-4819-157-12-201212180-00003
7. Strom C, Mollerup TK, Kromberg LS, Rasmussen LS, Schmidt TA. Hospitalisation in an emergency department short-stay unit compared to an internal medicine department is associated with fewer complications in older patients - an observational study. *Scand J Trauma Resusc Emerg Med*. 2017;25(1):80. doi:10.1186/s13049-017-0422-9
8. Strom C, Rasmussen LS, Lowe AS, et al. Short-stay unit hospitalisation vs. standard care outcomes in older internal medicine patients-a randomised clinical trial. *Age Ageing*. 2018;47(6):810–817. doi:10.1093/ageing/afy090
9. Wittenberg R, Sharpin L, McCormick B, Hurst J. Understanding emergency hospital admission of older people; 2014.
10. Jakobsen GJ. De ældste patienter er indlagt i markant kortere tid [The oldest patients are admitted for markedly shorter]. Kommunernes Landsforening [Local Government Denmark]; 2020. Danish. Available from: <https://www.kl.dk/nyheder/momentum/2020/2020-16/de-aeldste-patienter-er-indlagt-i-markant-kortere-tid/>. Accessed May 15, 2023.

11. Bryan K. Policies for reducing delayed discharge from hospital. *Br Med Bull.* 2010;95:33–46. doi:10.1093/bmb/ldq020
12. Klinge M, Aasbrenn M, Ozturk B, et al. Readmission of older acutely admitted medical patients after short-term admissions in Denmark: a nationwide cohort study. *BMC Geriatr.* 2020;20(1):203. doi:10.1186/s12877-020-01599-4
13. Aasbrenn M, Christiansen CF, Esen BO, Suetta C, Nielsen FE. Mortality of older acutely admitted medical patients after early discharge from emergency departments: a nationwide cohort study. *BMC Geriatr.* 2021;21(1):410. doi:10.1186/s12877-021-02355-y
14. Brockhattingen KK, Anru PL, Masud T, Petrovic M, Ryg J. Association between number of medications and mortality in geriatric inpatients: a Danish nationwide register-based cohort study. *Eur Geriatr Med.* 2020;11(6):1063–1071. doi:10.1007/s41999-020-00390-3
15. Klausen HH, Petersen J, Bandholm T, et al. Association between routine laboratory tests and long-term mortality among acutely admitted older medical patients: a cohort study. *BMC Geriatr.* 2017;17(1):62. doi:10.1186/s12877-017-0434-3
16. Hao Q, Zhou L, Dong B, Yang M, Dong B, Weil Y. The role of frailty in predicting mortality and readmission in older adults in acute care wards: a prospective study. *Sci Rep.* 2019;9(1):1207. doi:10.1038/s41598-018-38072-7
17. Schmidt M, Schmidt SAJ, Adelborg K, et al. The Danish health care system and epidemiological research: from health care contacts to database records. *Clin Epidemiol.* 2019;11:563–591. doi:10.2147/CLEP.S179083
18. Schmidt M, Pedersen L, Sorensen HT. The Danish civil registration system as a tool in epidemiology. *Clin Epidemiol.* 2014;29(8):541–549. doi:10.1007/s10654-014-9930-3
19. Schmidt M, Schmidt SAJ, Sandegaard JL, Ehrenstein V, Pedersen L, Sorensen HT. The Danish national patient registry: a review of content, data quality, and research potential. *Clin Epidemiol.* 2015;7:449–490. doi:10.2147/CLEP.S91125
20. Quan H, Li B, Couris CM, et al. Updating and validating the Charlson comorbidity index and score for risk adjustment in hospital discharge abstracts using data from 6 countries. *Am J Epidemiol.* 2011;173(6):676–682. doi:10.1093/aje/kwq433
21. Wockenfuss R, Frese T, Herrmann K, Claussnitzer M, Sandholzer H. Three- and four-digit ICD-10 is not a reliable classification system in primary care. *Scand J Prim Health Care.* 2009;27(3):131–136. doi:10.1080/02813430903072215
22. Vest-Hansen B, Riis AH, Sorensen HT, Christiansen CF. Acute admissions to medical departments in Denmark: diagnoses and patient characteristics. *Eur J Intern Med.* 2014;25(7):639–645. doi:10.1016/j.ejim.2014.06.017
23. Ishwaran H, Kogalur UB. Fast Unified Random Forests for Survival, Regression, and Classification (RF-SRC); 2022.
24. Villanueva RA, Chen ZJ. *ggplot2: Elegant Graphics for Data Analysis.* Springer-Verlag New York; 2016.
25. Breiman L. Random Forests. *Mach Learn.* 2001;45:5–32. doi:10.1023/A:1010933404324
26. Hastie T, Tibshirani R, Friedman J. *The Elements of Statistical Learning: Data Mining, Inference, and Prediction.* 2nd ed. New York: Springer; 2009.
27. Chen C, Liaw A, Breiman L. *Using Random Forest to Learn Imbalanced Data.* Berkeley Statistics; 2004.
28. O'Brien R, Ishwaran H, Random Forests A. Quantile classifier for class imbalanced data. *Pattern Recognit.* 2019;90:232–249. doi:10.1016/j.patcog.2019.01.036
29. Henderson R, Keiding N. Individual survival time prediction using statistical models. *J Med Ethics.* 2005;31(12):703–706. doi:10.1136/jme.2005.012427
30. Smolin B, Levy Y, Sabbach-Cohen E, Levi L, Mashiah T. Predicting mortality of elderly patients acutely admitted to the department of internal medicine. *Int J Clin Pract.* 2015;69(4):501–508. doi:10.1111/ijcp.12564
31. de Gelder J, Lucke JA, Heim N, et al. Predicting mortality in acutely hospitalized older patients: a retrospective cohort study. *Intern Emerg Med.* 2016;11(4):587–594. doi:10.1007/s11739-015-1381-7
32. Edmans J, Bradshaw L, Gladman JR, et al. The Identification of Seniors at Risk (ISAR) score to predict clinical outcomes and health service costs in older people discharged from UK acute medical units. *Age Ageing.* 2013;42(6):747–753. doi:10.1093/ageing/afh054
33. Steinmeyer Z, Piau A, Thomazeau J, Kai SHY, Nourhashemi F. Mortality in hospitalised older patients: the WHALES short-term predictive score. *BMJ Support Palliat Care.* 2021. doi:10.1136/bmjspcare-2021-003258
34. Juul-Larsen HG, Christensen LD, Bandholm T, et al. Patterns of multimorbidity and differences in healthcare utilization and complexity among acutely hospitalized medical patients (>=65 years) - A latent class approach. *Clin Epidemiol.* 2020;12:245–259. doi:10.2147/CLEP.S226586
35. Sundhedsstyrelsen [Danish Health Authority]. Styrket akutberedskab - planlægningsgrundlag for det regionale sundhedsvæsen [Improved emergency response – suggestions for the regional health services]. Danish. Available from: <https://www.sst.dk/-/media/Udgivelser/2007/Publ2007/PLAN/Akutberedskab/StyrketAkutberedskab,-d,.pdf.ashx>. Accessed May 15, 2023.
36. Makar M, Ghassemi M, Cutler DM, Obermeyer Z. Short-term mortality prediction for elderly patients using medicare claims data. *Int J Mach Learn Comput.* 2015;5(3):192–197. doi:10.7763/IJMLC.2015.V5.506
37. Couronne R, Probst P, Boulesteix AL. Random forest versus logistic regression: a large-scale benchmark experiment. *BMC Bioinform.* 2018;19(1):270. doi:10.1186/s12859-018-2264-5
38. Parikh RB, Manz C, Chivers C, et al. Machine learning approaches to predict 6-month mortality among patients with cancer. *JAMA Netw Open.* 2019;2(10):e1915997. doi:10.1001/jamanetworkopen.2019.15997
39. Editorial. Ascent of machine learning in medicine. *Nat Mater.* 2019;18(5):407. doi:10.1038/s41563-019-0360-1
40. Anpalahan M, Gibson SJ. Geriatric syndromes as predictors of adverse outcomes of hospitalization. *Intern Med J.* 2008;38(1):16–23. doi:10.1111/j.1445-5994.2007.01398.x
41. Campbell SE, Seymour DG, Primrose WR. A systematic literature review of factors affecting outcome in older medical patients admitted to hospital. *Age Ageing.* 2004;33(2):110–115. doi:10.1093/ageing/afh036
42. Graversen SB, Pedersen HS, Sandbaek A, Foss CH, Palmer VJ, Ribe AR. Dementia and the risk of short-term readmission and mortality after a pneumonia admission. *PLoS One.* 2021;16(1):e0246153. doi:10.1371/journal.pone.0246153
43. Petermann-Rocha F, Lyall DM, Gray SR, et al. Associations between physical frailty and dementia incidence: a prospective study from UK Biobank. *Lancet Health Longev.* 2020;1(2):e58–e68.
44. Bai G, Wang Y, Kuja-Halkola R, et al. Frailty and the risk of dementia: is the association explained by shared environmental and genetic factors? *BMC Med.* 2021;19(1):248. doi:10.1186/s12916-021-02104-3
45. Rogers NT, Steptoe A, Cadar D. Frailty is an independent predictor of incident dementia: evidence from the English longitudinal study of ageing. *Sci Rep.* 2017;7(1):15746. doi:10.1038/s41598-017-16104-y
46. Koria LG, Sawan MJ, Redston RM, Gnjudic D. The prevalence of frailty among older adults living with dementia: a systematic review. *J Am Med Dir Assoc.* 2022;23:1807–1814. doi:10.1016/j.jamda.2022.01.084

47. Sundhedsdatastyrelsen [Danish Health Data Authority]. Mange bor længe på plejehjem [many live in nursing homes for a long time]; 2022. Danish. Available from: [https://sundhedsdatastyrelsen.dk/da/nyheder/2020/plejehjemsbeboere\\_161220](https://sundhedsdatastyrelsen.dk/da/nyheder/2020/plejehjemsbeboere_161220). Accessed May 15, 2023.
48. Buurman BM, van Munster BC, Korevaar JC, Abu-Hanna A, Levi M, de Rooij SE. Prognostication in acutely admitted older patients by nurses and physicians. *J Gen Intern Med.* 2008;23(11):1883–1889. doi:10.1007/s11606-008-0741-7
49. Glynn NW, Gmelin T, Renner SW, et al. Perceived physical fatigability predicts all-cause mortality in older adults. *J Gerontol.* 2022;77(4):837–841. doi:10.1093/gerona/glab374
50. Kristensen M, Iversen AKS, Gerds TA, et al. Routine blood tests are associated with short term mortality and can improve emergency department triage: a cohort study of >12,000 patients. *Scand J Trauma Resusc Emerg Med.* 2017;25(1):115. doi:10.1186/s13049-017-0458-x
51. Sahni N, Simon G, Arora R. Development and validation of machine learning models for prediction of 1-year mortality utilizing electronic medical record data available at the end of hospitalization in multicondition patients: a proof-of-concept study. *J Gen Intern Med.* 2018;33(6):921–928. doi:10.1007/s11606-018-4316-y
52. Faisal M, Howes R, Steyerberg EW, Richardson D, Mohammed MA. Using routine blood test results to predict the risk of death for emergency medical admissions to hospital: an external model validation study. *QJM.* 2017;110(1):27–31. doi:10.1093/qjmed/hcw110
53. Ng N, Santosa A, Weinehall L, Malmberg G. Living alone and mortality among older people in Vasterbotten county in Sweden: a survey and register-based longitudinal study. *BMC Geriatr.* 2020;20(1):7. doi:10.1186/s12877-019-1330-9
54. Robards J, Evandrou M, Falkingham J, Vlachantoni A. Marital status, health and mortality. *Maturitas.* 2012;73(4):295–299. doi:10.1016/j.maturitas.2012.08.007
55. Rosella LC, Calzavara A, Frank JW, Fitzpatrick T, Donnelly PD, Henry D. Narrowing mortality gap between men and women over two decades: a registry-based study in Ontario, Canada. *BMJ Open.* 2016;6(11):e012564. doi:10.1136/bmjopen-2016-012564
56. Kellett J, Deane B. The simple clinical score predicts mortality for 30 days after admission to an acute medical unit. *Q J Med.* 2006;99(11):771–781. doi:10.1093/qjmed/hcl112
57. Blomaard LC, Speksnijder C, Lucke JA, et al. Geriatric screening, triage urgency, and 30-day mortality in older emergency department patients. *J Am Geriatr Soc.* 2020;68(8):1755–1762. doi:10.1111/jgs.16427
58. Kamper RS, Schultz M, Hansen SK, et al. Biomarkers for length of hospital stay, changes in muscle mass, strength and physical function in older medical patients: protocol for the Copenhagen PROTECT study—a prospective cohort study. *BMJ Open.* 2020;10(12):e042786. doi:10.1136/bmjopen-2020-042786
59. Henriksen DP, Rasmussen L, Hansen MR, Hallas J, Pottegard A. Comparison of the five Danish regions regarding demographic characteristics, healthcare utilization, and medication use—a descriptive cross-sectional study. *PLoS One.* 2015;10(10):e0140197. doi:10.1371/journal.pone.0140197
60. Vest-Hansen B, Riis AH, Christiansen CF. Registration of acute medical hospital admissions in the Danish national patient registry: a validation study. *Clin Epidemiol.* 2013;5:129–133. doi:10.2147/CLEP.S41905

## Clinical Epidemiology

Dovepress

### Publish your work in this journal

Clinical Epidemiology is an international, peer-reviewed, open access, online journal focusing on disease and drug epidemiology, identification of risk factors and screening procedures to develop optimal preventative initiatives and programs. Specific topics include: diagnosis, prognosis, treatment, screening, prevention, risk factor modification, systematic reviews, risk & safety of medical interventions, epidemiology & biostatistical methods, and evaluation of guidelines, translational medicine, health policies & economic evaluations. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use.

Submit your manuscript here: <https://www.dovepress.com/clinical-epidemiology-journal>