

# A Response to Article “Effectiveness of Secretome from Human Umbilical Cord Mesenchymal Stem Cells in Gel (10% SM-hUCMSC Gel) for Chronic Wounds (Diabetic and Trophic Ulcer) – Phase 2 Clinical Trial” [Letter]

Annisa Dwicantika<sup>1</sup>, Rilianawati<sup>2</sup>, Ratih Rinendyaputri<sup>3</sup>

<sup>1</sup>Faculty of Pharmacy, Pancasila University, Jakarta, Indonesia; <sup>2</sup>Vaccine and Drug Research Center, National Research and Innovation Agency, Serpong, Indonesia; <sup>3</sup>Center for Biomedical Research, Research Organization for Health, National Research and Innovation Agency, Bogor, Indonesia

Correspondence: Annisa Dwicantika, Email [dwicantikaa@gmail.com](mailto:dwicantikaa@gmail.com)

## Dear editor

We have read the published article and we appreciate the authors who have reported the article about “Effectiveness of Secretome from Human Umbilical Cord Mesenchymal Stem Cells in Gel (10% SM-hUCMSC Gel) for Chronic Wounds (Diabetic and Trophic Ulcer) – Phase 2 Clinical Trial”, published in the Journal of Multidisciplinary Healthcare 2023;16:1763–1777.

Mesenchymal stem cell secretomes have promising prospects to be produced as drugs for regenerative medicine. Many studies suggest that mesenchymal stem cell secretomes can be used as therapy for alopecia, acute hind limb ischemia, ischemic stroke, chronic kidney disease, and acute and chronic wound healing.<sup>1</sup> The research carried out by Tan ST et al reports that secretome from human umbilical stem cells can be made into a topical gel for chronic wounds (diabetic and trophic ulcers) and has been proven effective in accelerating the process for wound healing. This is can be a new innovation for secretome-based treatment for chronic wound healing.<sup>2</sup>

Previous research has shown that three-dimensional (3D) bioprinting has also been used to reconstruct skin through layer-by-layer deposition of cells with scaffolding materials over the wound areas.<sup>3</sup> Guo X et al used porcine small intestinal submucosa (SIS) as a scaffold that can mimic the extracellular matrix of skin, facilitating cell growth, organization, and functional differentiation tissues.<sup>4</sup> Other study has investigated the effects of MSC-CM-T in topical gel preparation in accelerating optimal wound healing through analyzing PDGF levels, wound closure rate percentages, and fibroblast density appearances.<sup>5</sup> This can be used as a reference for Tan ST et al<sup>2</sup> research in addition to using gel as a scaffold material for cell growth in topical gel preparations.

This study directly determines the level of secretome used for topical gel preparation without providing some clear reasons. Meanwhile, in our opinion, it is also necessary to mention the characteristics of UCMSCs as a source of SH-hUCMSCs according to the International Society of Stem Cell Therapy/ISCT, and compare several levels of SM-hUCMSC used in this gel formulation so that it can be known which level can produce the best treatment in terms of quality, speed or time, and accuracy for healing diabetic chronic wounds and trophic ulcers in the future.

## Disclosure

The authors report no conflicts of interest in this communication.

## References

1. Pawitan JA. Prospect of stem cell conditioned medium in regenerative medicine. *Biomed Res Int.* 2014;2014:965849. doi:10.1155/2014/965849
2. Tan ST, Aisyah PB, Firmansyah Y, Nathasia N, Budi E, Hendrawan S. Effectiveness of secretome from human umbilical cord mesenchymal stem cells in gel (10% SM-hUCMSC gel) for chronic wounds (diabetic and trophic ulcer) - phase 2 clinical trial. *J Multidiscip Healthc.* 2023;16:1763–1777. doi:10.2147/JMDH.S408162
3. Abdul Kareem N, Aijaz A, Jeschke MG. Stem cell therapy for burns: story so far. *Biologics.* 2021;15:379–397. doi:10.2147/BTT.S259124
4. Guo X, Xia B, Lu XB, et al. Grafting of mesenchymal stem cell-seeded small intestinal submucosa to repair the deep partial-thickness burns. *Connect Tissue Res.* 2016;57(5):388–397. doi:10.1080/03008207.2016.1193173
5. Putra A, Ibrahim S, Muhar AM, et al. Topical gel of mesenchymal stem cells-conditioned medium under TNF- $\alpha$  precondition accelerates wound closure healing in full-thickness skin defect animal model. *J Med Life.* 2022;15(2):214–221. doi:10.25122/jml-2019-0103

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Journal of Multidisciplinary Healthcare 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Journal of Multidisciplinary Healthcare editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Journal of Multidisciplinary Healthcare

Dovepress

### Publish your work in this journal

The Journal of Multidisciplinary Healthcare is an international, peer-reviewed open-access journal that aims to represent and publish research in healthcare areas delivered by practitioners of different disciplines. This includes studies and reviews conducted by multidisciplinary teams as well as research which evaluates the results or conduct of such teams or healthcare processes in general. The journal covers a very wide range of areas and welcomes submissions from practitioners at all levels, from all over the world. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-inflammation-research-journal>

<https://doi.org/10.2147/JMDH.S446608>