



Neutrophil/Lymphocyte Ratio as a Predictive Biomarker of NIV Failure in AECOPD: The Unveiled Facts [Letter]

Dipasri Bhattacharya ¹, Antonio M Esquinas ², Mohanchandra Mandal ³

¹Department of Anaesthesiology, Pain Medicine and Critical Care, R.G.Kar Medical College & Hospital, Kolkata, West Bengal, India; ²Critical Care Specialist and Staff Physician, Intensive Care Unit, Hospital Morales, Murcia, Spain; ³Department of Anaesthesiology, Institute of Post Graduate Medical Education and Research, Kolkata, West Bengal, India

Correspondence: Mohanchandra Mandal, Department of Anaesthesiology, Institute of Post Graduate Medical Education and Research, Kolkata, West Bengal, India, Email drmcmandal@gmail.com

Dear editor

We appreciate the work of Sun et al,¹ who observed that Neutrophil Lymphocyte Ratio (NLR) can be an effective marker for predicting Non Invasive Ventilation (NIV) failure in Acute Exacerbation of Chronic Obstructive Pulmonary Disease (AECOPD). They also observed that NLR >8.9 had higher risk of NIV failure requiring intubation. However, we raise some points for further clarification.

NLR gets elevated in AECOPD compared with stable state. Serial monitoring of NLR can predict NIV failure better than single measurement.² NLR is an independent risk factor for 28-day mortality in patients with AECOPD,² and the mortality rate increases with NLR ≥ 4 .³ We are curious to know whether continual monitoring of NLR was done and compared with that of stable state in the same patient. In addition, it would be interesting to know whether they assessed mortality outcome, given that the length of hospital stay was observed already.

NLR encompasses both neutrophilia and lymphopenia, the indicators of inflammation and decreased immune response, respectively. Thus, it can reflect the severity of COPD better than when neutrophilia or lymphopenia are monitored alone. Another novel inflammatory marker, PLR (serum absolute count ratio of platelet to lymphocyte) can help predict the severity of clinical condition and mortality in AECOPD patients.³

The authors observed respiratory rate and arterial blood gas analysis. We are curious to know whether they used simple bedside parameters or scores predictive of NIV failure such as initial high respiratory rate, low PaO₂/FiO₂ ratio, HACOR score >5 at the end of 1 hour of NIV, and initial hs-CRP.⁴

Individuals with comorbidities and decreased functional capabilities are at higher risk for NIV failure. Patients with poor baseline performance status and having higher DECAF (dyspnea, eosinopenia, consolidation, acidemia, atrial fibrillation) score on admission can predict severity of acute clinical deterioration.⁵ We wonder whether they had a scope to observe those in their study.

Funding

No external funding supported the preparation of this manuscript. The authors have no financial and/or personal relationships with other people or organizations that might inappropriately influence our work.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Sun W, Luo Z, Jin J, Cao Z, Ma Y. The neutrophil/lymphocyte ratio could predict noninvasive mechanical ventilation failure in patients with acute exacerbation of chronic obstructive pulmonary disease: a retrospective observational study. *Int J Chron Obstruct Pulmon Dis*. 2021;16:2267–2277. Erratum in: *Int J Chron Obstruct Pulmon Dis*. 2023;18:3047–3048. doi:10.2147/COPD.S320529
2. Yang L, Gao C, He Y, et al. The neutrophil-to-lymphocyte ratio is associated with the requirement and the duration of invasive mechanical ventilation in acute respiratory distress syndrome patients: a retrospective study. *Can Respir J*. 2022;2022:1581038. doi:10.1155/2022/1581038
3. Allena N, Khanal S, Jog A, et al. Decoding the Chronic Obstructive Pulmonary Disease (COPD) Puzzle: investigating the significance of exacerbation scores in triage decision-making. *Cureus*. 2023;15(7):e41975. doi:10.7759/cureus.41975
4. Mathen PG, Kumar KG, Mohan N, et al. Prediction of noninvasive ventilation failure in a mixed population visiting the emergency department in a Tertiary Care Center in India. *Indian J Crit Care Med*. 2022;26(10):1115–1119. doi:10.5005/jp-journals-10071-24338
5. Echevarria C, Steer J, Bourke SC. Comparison of early warning scores in patients with COPD exacerbation: DECAF and NEWS score. *Thorax*. 2019;74(10):941–946. doi:10.1136/thoraxjnl-2019-213470

Dove Medical Press encourages responsible, free and frank academic debate. The content of the International Journal of Chronic Obstructive Pulmonary Disease 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the International Journal of Chronic Obstructive Pulmonary Disease editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

International Journal of Chronic Obstructive Pulmonary Disease

Dovepress

Publish your work in this journal

The International Journal of COPD is an international, peer-reviewed journal of therapeutics and pharmacology focusing on concise rapid reporting of clinical studies and reviews in COPD. Special focus is given to the pathophysiological processes underlying the disease, intervention programs, patient focused education, and self management protocols. This journal is indexed on PubMed Central, MedLine and CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-chronic-obstructive-pulmonary-disease-journal>

<https://doi.org/10.2147/COPD.S465448>