

Involuntary Falls in Patients with Chronic Kidney Diseases on Nephrology Wards: Research Advances and Future Perspectives

Yumei Liao^{1,*}, Aihong Wang^{1,*}, Wenjuan Sui², Abbas Khan³, Zibo Xiong^{1,4,5}, Guang Yang^{1,4,5}

¹Division of Renal Medicine, Peking University Shenzhen Hospital, Peking University, Shenzhen, Guangdong, 518036, People's Republic of China;

²Institute of City Strategy Studies, Guangdong University of Foreign Studies, Guangzhou, Guangdong, 510006, People's Republic of China;

³Department of Nutrition and Health Promotion, University of Home Economics Lahore Pakistan, Lahore, 54000, Pakistan; ⁴Institute of Nephrology, Shenzhen Peking University-Hong Kong University of Science and Technology (PKU-HKUST) Medical Center, Shenzhen, Guangdong, 518036, People's Republic of China; ⁵Shenzhen Clinical Research Centre for Urology and Nephrology, Shenzhen, Guangdong, 518036, People's Republic of China

*These authors contributed equally to this work

Correspondence: Zibo Xiong; Guang Yang, Email xiong0301@163.com; guang.yang@pkusz.com

Background: Nephrology nursing faces an inevitable challenge when it comes to falls. This study aims to review existing literature on falls in chronic kidney disease (CKD) patients and to discuss potential solutions.

Methods: This review explores the characteristics and causes of falls in CKD patients, elucidate the underlying mechanisms, shares better nursing practices, and discusses both current prevention strategies as well as potential future interventions.

Results: CKD patients often experience consciousness impairment and motor dysfunction due to factors such as muscle atrophy, nerve damage, malnutrition, anemia, fluid-electrolyte imbalances, hormonal disorders, and dialysis. Additionally, this study not only shares insights from nursing experience with these patients, but also reviews current evidence-based fall prevention strategies and proposes potential new interventions that could be tested in future research.

Conclusion: CKD may increase the risk of involuntary falls in patients due to consciousness impairment and motor dysfunction. Advocating for a standardized expert consensus on fall risk assessment and comprehensive care for CKD patients would enhance the nursing framework in nephrology wards and help patients decrease their fall risk and elevate their quality of life.

Keywords: falls, chronic kidney diseases, dialysis, nephrology nursing, anemia

Introduction

Falls among hospitalized patients represent a prevalent and consequential adverse event, capable of causing severe physical harm such as fractures, head trauma, and soft tissue injuries.^{1,2} Additionally, they can significantly impact a patient's self-esteem and confidence, potentially leading to long-lasting psychological trauma.^{3,4} Approximately one-third of individuals over the age of 65 experience a fall each year.⁵ Within this age bracket, chronic kidney disease (CKD) increases the rate of falls by 1.81 times.⁶ Moreover, middle-aged adults with CKD had a higher likelihood of falling (2.30, 95% CI 2.07–2.55) and sustaining fall-related injuries (1.54, 95% CI 1.32–1.80) compared to those without CKD.⁷ Several factors contribute to the increased fall risk in CKD patients, including osteoporosis, anemia, water-electrolyte imbalances, blood pressure fluctuations, toxin accumulation, muscle atrophy, and nerve damage.^{8–13} For this reason, they commonly exhibit motor dysfunction and impaired subjective consciousness, rendering them unable to effectively maintain balance.^{10,14} Furthermore, hospital-specific factors such as physical deterioration and medication-related side effects can further heighten the risk of falls.^{15,16} Consequently, preventing falls among patients with CKD is a crucial responsibility in nephrology nursing.^{17,18}

To optimize patient care, a series of studies on this issue were conducted, proposing several preventive measures that have effectively reduced the rate of fall injuries.¹⁴ Yet, sporadic falls among CKD patients persist, underscoring the need

for improved fall prevention strategies in nephrology care. This study aims to analyze the dominant characteristics and causes of fall injuries in CKD patients, understand their underlying mechanisms, share nursing expertise, and investigate potential advancements in prevention methods and interventions. This knowledge will serve as a foundation for the continued development of nephrology nursing strategies.

Characteristics of Falls in CKD Patients

High-Risk Groups

CKD affects a wide demographic, but the risk of falls is not uniformly distributed across all patients. The prevailing consensus suggests that individuals with advanced CKD face a higher risk of falling.^{19–21} The exact reasons will be detailed in subsequent sections. Also, the elderly group is at high risk, which may be related to their weaker physical functioning.^{4,7,22} Moreover, there have been several studies that have identified risk factors associated with falls in CKD patients,^{2,23} but all have been related to the course of CKD and age. Although CKD encompasses a range of primary conditions and complications, there's a lack of comprehensive research exploring the relationship between these conditions and fall risk. This leaves a gap in our understanding of which CKD subgroups are most prone to falls. In clinical practice, assessments are routinely conducted to identify patients with elevated fall risks. Common evaluation methods involve measuring muscle strength using a grip strength device and evaluating the patient's mental state. Patients presenting with diminished muscle strength, depression, anxiety, and recurrent episodes of subjective confusion are deemed to be at a higher risk of falls.^{24–26}

Spatio-Temporal Characteristics of Falls

Hospitalized CKD patients exhibit specific areas prone to falls, such as toilets and bathrooms.^{14,26} The act of standing and walking to restrooms can pose a fall risk, often due to unstable legs or a sudden drop in blood pressure. Slippery surfaces in bathrooms, especially during showers, further compound the risk. Additionally, walking, whether within the ward or the hospital corridor, is a common scenario for falls. Factors such as unsteady gait, visual impairment, and misuse of walking aids intensify this risk. Falls are also prevalent during actions like turning, standing up, transferring, or shifting, often stemming from physical instability and uncoordinated movements.²⁷ These findings emphasize the importance of prioritizing fall prevention in restrooms, hallways, and bedside areas, advising patients against abrupt postural changes, and encouraging them to request assistance when needed.

In terms of timing, falls in hospitalized CKD patients tend to be concentrated during nighttime and early morning hours. There's a heightened frequency of falls at night, possibly because patients tend to be less alert.¹⁴ Moreover, patients might experience dizziness upon rising, which may be closely linked to postural hypotension.²⁷ Given these trends, enhancing monitoring during the night and early morning hours can help promptly detect and respond to falls. As such, healthcare institutions and professionals should adopt preventive measures tailored to each patient's unique situation. Possible strategies include offering appropriate mobility aids, intensifying nursing rounds, and installing bedside alarms. These actions aim to effectively mitigate the risk of falls among hospitalized patients.

Summary

From the insights presented (Figure 1), it is evident that preventive measures are paramount for CKD patients, especially those in advanced stages or with nephrotic syndrome. Ensuring patient safety by improving the living environment is vital, both at home and in the hospital. This entails keeping pathways clear, minimizing clutter, ensuring dry floors, and using non-slip mats. Furthermore, to further mitigate fall risks, patients and their families should also be educated about home safety precautions.




Figure 1 Who, when, where, and what activities are at high-risk for falls in CKD patients.

Reasons and Mechanisms of Falls in CKD Patients (Figure 2)

Muscle Atrophy

A critical facet of motor impairment is muscle atrophy, which can be attributed to several intertwined factors. First, CKD results in the accumulation of metabolites, which results in increased protein breakdown and reduced synthesis, culminating in muscle tissue breakdown and atrophy.^{28,29} Second, CKD disrupts glucose, fat, and protein metabolism, leading to insufficient energy supply and subsequent muscle tissue atrophy.^{30,31} Third, CKD induces chronic inflammatory and oxidative stress, triggering the release of inflammatory factors that inhibit protein synthesis and promote protein catabolism, ultimately resulting in muscle atrophy.^{28,31} Lastly, CKD patients often experience fatigue, weakness, and reduced exercise endurance, which restricts physical activity and exacerbates muscle atrophy. Therefore, it is imperative for CKD patients to ensure optimal nutritional uptake, embrace moderate physical activity, and lay emphasis on muscle fortification. Tailored supplementation and nutritional interventions are equally pivotal to safeguarding muscle vitality.




Figure 2 Schematic illustration of the reasons and mechanisms of falls in CKD patients. These images are our originals, and the icons in them are freely available icons, originating from WPS office software.

Nerve Damage

Nerve damage and functional abnormalities play a pivotal role in motor dysfunction and impaired subjective consciousness in CKD patients.^{6,25} Several principal factors underlie this phenomenon. Firstly, the accumulation of toxins resulting from uremia directly impacts the nervous system.³² Secondly, the accumulation of toxins incites an inflammatory response, releasing inflammatory mediators like cytokines and oxygen-free radicals, which can compromise nerve cell structure and function.³³ Thirdly, CKD-associated malnutrition hampers the adequate supply of nutrients to nerve cells, thereby affecting their function.³⁴ Lastly, CKD patients often undergo long-term treatment with various medications, some of which may have adverse effects on the nervous system.³⁵ Overall, these factors collectively contribute to the structural and functional damage of nerve cells.

Malnutrition

Malnutrition is a common concern among CKD patients, manifesting in symptoms such as weakness and fatigue.²⁴ This emerges from an interplay of various factors. First, CKD patients often experience a loss of appetite, which can be attributed to abnormalities in taste and smell senses caused by uremia, as well as the impact of metabolite accumulation on the central nervous system.^{36,37} Second, vomiting and nausea are prevalent among CKD patients, often caused by metabolite buildup and medication side effects.^{38–40} Third, digestive complications, including diarrhea and constipation, frequently afflict CKD patients. Arising from electrolyte imbalances, medication side effects, or disrupted bowel function, these complications obstruct the effective absorption and digestion of vital nutrients.⁴¹ Fourth, CKD patients may experience proteinuria, characterized by the excessive loss of protein through urine.⁴² Protein is a vital nutrient for maintaining normal physiological function, and its restriction contributes to a decline in nutritional status.⁴³ The task of restricting protein intake and enhancing absorption has been particularly challenging. Addressing this nutritional challenge requires a careful approach to nutritional management. Despite extensive research on nutrition for individuals with CKD, there is limited practical application. There is a prevailing preference for pharmacological interventions, as they are perceived as more effective in improving a single nutritional indicator.

Anemia

Anemia, a frequent complication in CKD patients, which can induce fatigue, weakness, and breathlessness, thereby compromising physical mobility and equilibrium and elevating fall susceptibility. First, Impaired kidney function leads to reduced erythropoietin (EPO) secretion, which is an essential hormone governing erythropoiesis, resulting in decreased erythropoiesis and subsequent anemia.^{39,44} Second, CKD patients often exhibit iron metabolism disorders, including impaired iron absorption, abnormal transportation, and reduced storage, leading to inadequate iron supply and affecting erythropoiesis.^{44,45} Third, CKD patients frequently experience chronic inflammatory responses, which can encumber erythropoiesis and truncate the lifespan of red blood cells.^{46–49} Fourth, the accumulation of metabolites, such as urea and creatinine, adversely affects bone marrow hematopoiesis and inhibits erythropoiesis.^{50,51} Fifth, malnutrition can impact the production and function of red blood cells.⁵² Therefore, recognizing and addressing anemia with alacrity is paramount for CKD patients. Treatment options for anemia include EPO replacement therapy, iron supplementation, and red blood cell transfusion.

Fluid and Electrolyte Balance

CKD impairs the kidney's aptitude for water and electrolyte regulation, resulting in imbalances that can disrupt nerve and muscle regulation. For instance, sodium retention can induce hypertension and edema, while hyperkalemia can have detrimental effects on cardiac function, posing potential life-threatening risks.^{11,53–56} Hyperphosphatemia and hypocalcemia can lead to abnormal mineral metabolism and osteoporosis.⁹ Abnormal blood potassium, hyponatremia, and hypocalcemia can give rise to symptoms like muscle twitching, limb weakness, and numbness. Additionally, CKD patients may develop acidosis or alkalosis, further disrupting this equilibrium.⁵⁷ Therefore, meticulous monitoring and management of these imbalances in CKD care is indispensable. Developing an individualized treatment plan tailored to

the patient's specific condition is essential, which may involve limiting fluid intake, controlling dietary sodium, potassium, and phosphorus intake, and utilizing medications as necessary.

Hormonal Disorder

CKD can induce various hormonal dysfunctions, ultimately impacting the body's mobility. For instance, CKD stimulates the release of parathyroid hormone (PTH) from the parathyroid glands.⁵⁸ PTH primarily enhances intestinal calcium absorption, inhibits renal tubular reabsorption of phosphate, and promotes calcium release from bones. This cycle perpetuates and culminates in hyperparathyroidism. Parathyroid excision can significantly improve negative physical and psychological symptoms and enhance the quality of life.⁵⁹ Despite the high effectiveness of this procedure in treating the aforementioned conditions, it is essential to acknowledge the existence of potential risks and hazards. One notable risk is the development of hypocalcemia following parathyroidectomy,⁶⁰ which can manifest as muscle spasms, neurological complications, cardiac arrhythmias, and in severe instances, life-threatening situations. Consequently, the implementation of postoperative calcium and vitamin D supplementation therapy becomes imperative.

Moreover, CKD is associated with the synthesis and metabolism of thyroid hormones, such as hypothyroidism or hyperthyroidism.^{12,61,62} Hypothyroidism is characterized by reduced levels of thyroxine (T3 and T4) and elevated levels of thyroid-stimulating hormone (TSH), manifesting symptoms such as fatigue, weight gain, and constipation. Conversely, hyperthyroidism is characterized by increased thyroxine levels and decreased TSH levels, resulting in symptoms like palpitations, weight loss, and excessive perspiration.

Additionally, CKD may impact the synthesis and secretion of growth hormones, leading to either deficiencies or excesses.¹² Growth hormone deficiency affects the growth and development of children, while excessive growth hormone supplementation can result in conditions like gigantism or acromegaly.⁶³ It is essential to highlight that CKD-induced hormone dysfunction is a complex process potentially involving abnormalities in various hormones. The specific circumstances and implications of this dysfunction should be determined based on the individual patient's condition and hormone levels.

Dialysis

Patients are susceptible to falls following dialysis due to several potential reasons. First, dialysis can result in hypotension or hypertension manifesting as dizziness and vertigo.¹⁰ Second, dialysis can further exacerbate CKD-induced muscle atrophy and weakness.⁶⁴ Third, dialysis can accelerate the loss of calcium.^{9,64} Fourth, there is a high probability that dialysis will cause hypoglycemia.⁶⁵ Lastly, patients may require medications like anti-hypertensives and diuretics during dialysis, which can induce side effects such as hypotension and dizziness. Consequently, patients are typically advised to rest and gradually adjust their posture before leaving the bed after dialysis to minimize the risk of falls.

Aging

Apart from the aforementioned factors, aging is another factor contributing to an increased susceptibility to falls among the elderly.⁶⁶ As age advances, there is a gradual decline in physical function and balance, weakened muscle strength, slower reaction time, and impaired vision and hearing, all of which collectively elevate the risk of falling. Moreover, the elderly often experience conditions such as muscle atrophy and osteoporosis, rendering their muscles and bones fragile and more susceptible to injury. Additionally, conditions like arthritis can result in reduced joint mobility, further augmenting the likelihood of falls.⁶⁷ However, it is essential to note that the relationship between age-associated falls and CKD-associated falls is not absolute, albeit there may be a correlation. This is primarily because aging inherently leads to a decline in kidney function and other organ functions. Moreover, elderly individuals with CKD may have a history of long-term CKD accompanied by multiple complications. Irrespective of the underlying cause, it is imperative to acknowledge that elderly CKD patients constitute a high-risk group for falls and therefore require specialized attention and care.




Figure 3 A set of assessment and prevention strategies to reduce fall risk in CKD patients. These images are our originals, and the icons in them are freely available icons, originating from WPS office software.

Methods of Fall Prevention and Management in Wards

Fall prevention is a vital concern in nephrology nursing, and hospitals employ various strategies to ensure patient safety and well-being. These strategies encompass holistic patient assessment, monitoring, furnishing of apt amenities, medication management, and a robust emphasis on patient education and training. In this section, we present our management experience spanning over two decades, spanning domains such as risk assessment, science popularization, and nursing management (Figure 3).

Risk Assessment

For efficacious fall prevention, a comprehensive fall risk assessment was performed on each patient before admission (Table 1). This assessment will encompass factors such as age, health condition, medication usage, mobility, cognitive function, and other relevant considerations to determine the patient's susceptibility to falls. Additionally, the patient's vital signs, including blood pressure and heart rate, were continuously monitored to detect any abnormalities, such as

Table 1 Criteria for Assessing Fall Risk in CKD Patients

Criteria for Assessing Fall Risk in CKD Patients	
Property	Standard
Age	≥ 65 years
History of falls	Six months prior to admission
Subjective consciousness disorders	Depression, irritability, hallucinations, Alzheimer's disease
Motor dysfunction	Limb weakness, deformity, paralysis, limited mobility
Cardiovascular function abnormalities	Hypoglycemia, abnormal blood pressure, arrhythmia
Sensory impairment	Reduced visual or auditory abilities
Toileting frequency	≤ 2 hours
Treatment and comorbidities	Surgery, seizures, flipping over fences, taking sedatives

Notes: According to the aforementioned eight scoring criteria, meeting two-point suggests a potential risk of falling, while meeting three-point is indicative of a high risk of falling. However, it is important to consider that patients with CKD, particularly those in CKD stage five, undergoing dialysis, or experiencing nephrotic syndrome, may not experience a pronounced sensation of limb weakness due to the long-term nature of their chronic condition or the presence of edema. Consequently, relying solely on the risk assessment score may not accurately reflect the true risk of falling for these patients.

hypotension, in a timely manner. In cases where patients exhibit higher risk scores, persistent reminders to both the patients and their caregivers about the latent fall risks are crucial. This also involves equipping them with foundational knowledge to amplify awareness and bolster future preventive tactics.

Science Popularization

In fall prevention, educating patients and their families about fall-related knowledge is a vital component.⁶⁸ This education includes teaching correct mobility skills, safety precautions, measures, and protocols for handling emergencies such as falls. Given the age and potential cognitive challenges of many CKD patients, educating family members often proves more effective than directly instructing the patients. We actively encourage the involvement of the patient's family members in fall prevention efforts, such as assisting the patient with movement and reminding them to prioritize safety. A variety of methods were used to widen the outreach of fall prevention knowledge, including conducting educational discussions with patients, displaying informative videos in patient rooms periodically, and placing posters in corridors. Using these methods to prevent falls is both straightforward and effective.

Nursing Management

The provision of assistive devices is fundamental to fall prevention. For example, the installation of various safety facilities, such as handrails, non-slip flooring, and guardrails to offer support and protection for patient mobility. For patients with mobility challenges, supportive aids such as crutches and walkers are also available to enhance their stability and mobility. Additionally, the use of exoskeletons to facilitate rehabilitation training for patients with muscle weakness is also an option. Effective patient medication management is another pivotal facet of fall prevention, ensuring to sidestep drugs that may precipitate adverse effects, such as hypotension or dizziness. Furthermore, some places have established a comprehensive fall monitoring and reporting system to promptly record and analyze fall incidents, enabling nurses to identify areas for improvement and implement preventive measures. This innovative approach lays the groundwork for future endeavors in precision medicine and bespoke fall prevention methodologies.

Hot Topics in Fall Prevention Research and Future Outlook

While these measures have proven to be highly effective in reducing fall incidences among nephrology inpatients, falls still occur. To address this persisting challenge, researchers have developed additional modalities that have demonstrated effectiveness in preventing patient falls (Figure 4).




Figure 4 Hot topics in fall prevention research and future outlook. These images are our originals, and the icons in them are freely available icons, originating from WPS office software.

Convalescent Training

Engaging in appropriate exercise can effectively enhance a patient's balance, muscle strength, and flexibility, consequently reducing the risk of falls.^{69,70} For example, incorporating balance training exercises can improve patients' sense of balance and control. Examples of such exercises include standing on one foot, walking along a tracing line, and performing exercises with closed eyes. Muscle-strengthening exercises can enhance patients' muscle strength, thereby improving stability while walking and standing. Examples of these exercises include weightlifting, squatting, and wall-resistance exercises. Likewise, flexibility training aids in maintaining optimal joint mobility and body movement. Stretching exercises, yoga, and Tai Chi exemplify flexibility-boosting activities. Gait training can assist patients in improving their walking posture and gait, ultimately reducing the risk of falls. This can be achieved through practicing proper walking techniques, improving walking speed, and utilizing assistive devices when necessary. Additionally, exercise should be conducted safely, avoiding high-risk activities in unstable environments.

Nutrition

By enhancing nutritional status and providing adequate nutritional support, the risk of falls among hospitalized patients can be mitigated.⁷¹ However, dietary considerations specific to CKD patients must be accounted for. Certain nutrients such as sodium, potassium, and phosphorus may require restriction based on the patient's specific condition.⁷¹ Patients experiencing malnutrition or muscle weakness may necessitate additional protein supplementation. However, it is crucial to supplement protein intake with high-quality protein to prevent proteinuria.^{72,73} Patients with osteoporosis may require additional calcium and vitamin D supplementation.^{13,74} Meanwhile, maintaining proper hydration is vital to avoid dehydration and fluid overload.⁷⁵ For patients unable to meet their nutritional needs orally, nutritional support through intravenous or gastrointestinal routes may be necessary. Therefore, regular nutritional assessments should be conducted to monitor patients' nutritional status and allow for timely adjustments to nutritional interventions.

Equipment

In addition to enhancing the patient's bodily health to prevent falls in hospitalized patients, there exists a range of assistive devices that can be employed for fall prevention. The most common are bedside rails, mobility aids, non-slip mats, non-slip shoes, and non-slip socks. Bed rails play a crucial role in preventing falls, especially during transitions like getting up or changing positions. Presently, most medical beds lack sufficient bed rails or feature inadequately long ones. Future advancements should prioritize user-friendliness and smart functionalities. For instance, incorporating features like voice prompts or lighting cues that activate when the system detects the patient rising can serve as reminders for the patient to prioritize safety.

Moreover, the development of relevant and user-friendly assistive devices is a burgeoning research area. For instance, a fall warning device is an electronic tool that monitors a patient's activity and automatically alarms whenever a patient falls.⁷⁶ Furthermore, safety mats with softer surfaces and wearable airbags are utilized to mitigate the impact of a patient's fall and provide additional support.⁷⁷⁻⁷⁹ These assistive devices and measures should be carefully tailored to the individual patient's needs and circumstances.

Others

The medical and healthcare industries have witnessed significant advancements with the development of data science. Increasingly, big data and artificial intelligence technologies are being utilized in nursing, and these technologies can also play a crucial role in the prevention of patient falls. For example, through the collection and analysis of extensive fall-related data, it becomes possible to construct fall prediction models that aid in assessing a patient's susceptibility to falls.⁷⁶ In addition, the integration of virtual reality and augmented reality technologies in the development of training and intervention resources for fall prevention can significantly enhance patients' balance and movement control, ultimately reducing the risk of falling.⁸⁰⁻⁸² Moreover, in addition to prioritizing fall prevention within healthcare organizations, future research endeavors could concentrate on extending fall prevention initiatives to the community level.^{83,84} Implementing measures such as public education and the establishment of community support networks can effectively enhance fall awareness and prevention among older adults residing in the community. Furthermore, aggressive treatment of CKD-related complications, such as arthritis and neurological disorders, can help reduce falls.^{6,7}

Limitations

This study presents several limitations. Firstly, the discussion is confined to the common causes of involuntary falls, management strategies, and research trends within the Nephrology Ward. The applicability of these measures to other healthcare departments or home settings warrants further investigation. Secondly, due to the sporadic nature of falls, there is a dearth of pharmacological research. Specifically, quantitative studies focusing on the use of medications to enhance patients' physical and mental well-being for fall prevention are lacking. Consequently, the emphasis in current research lies on fall prevention through nursing interventions and the utilization of equipment to mitigate risks post-fall. Thirdly, CKD patients often present with multiple comorbidities and are on multiple medications, rendering the identification of high-risk factors for falls challenging. Addressing these complexities represents a crucial avenue for future research.

Conclusions

In conclusion, prolonged CKD results in diminished motor function and impaired subjective consciousness by disturbing homeostasis. As a result, patients are prone to involuntary falls under the influence of external and unexpected factors. Presently, the primary approach to prevention revolves around status assessment, science popularization, and nursing management. Patients are also advocated for physical rehabilitation through physical training and nutritional interventions, as well as to utilize assistive devices to support walking. Future research should focus on the development of early warning systems for falls, intelligent assistive tools, personalized prevention programs, and community outreach activities. To summarize, an integrated, multifaceted approach across the healthcare continuum is needed to proactively assess risks, optimize nursing care, encourage patient engagement in recovery, and ultimately create safer hospital environments that minimize preventable harm.

Abbreviations

CRRT, continuous renal replacement therapy; CKD, chronic kidney disease; EPO, erythropoietin; PTH, parathyroid hormone; TSH, thyroid-stimulating hormone.

Acknowledgments

We are grateful to all our colleagues in the clinical departments and laboratories.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

This work was supported by the Shenzhen Science and Technology Innovation Commission (JCYJ20220530150412026), Guangdong Basic and Applied Basic Research Foundation (2023A1515111112), Shenzhen San-Ming Project of Medicine (SZSM201812097), and Shenzhen Clinical Research Center for Urology and Nephrology.

Disclosure

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

1. Cook WL, Tomlinson G, Donaldson M, et al. Falls and fall-related injuries in older dialysis patients. *Clin J Am Soc Nephrol*. 2006;1(6):1197–1204. doi:10.2215/cjn.01650506
2. Kutner NG, Zhang R, Huang Y, Wasse H. Falls among hemodialysis patients: potential opportunities for prevention? *Clin Kidney J*. 2014;7(3):257–263. doi:10.1093/ckj/sfu034
3. Ren J, Zhang X, Lin H, Zeng JH. Influencing factors of fear of falling among glaucoma patients in west China: a cross-sectional study. *Int J Ophthalmol*. 2023;16(4):563–570. doi:10.18240/ijo.2023.04.10

4. Sakthivadivel V, Geetha J, Gaur A, Kaliappan A. Performance-Oriented mobility assessment test and timed up and go test as predictors of falls in the elderly - A cross-sectional study. *J Family Med Prim Care*. 2022;11(11):7294–7298. doi:10.4103/jfmpc.jfmpc_1294_22
5. O'Loughlin JL, Robitaille Y, Boivin JF, Suissa S. Incidence of and risk factors for falls and injurious falls among the community-dwelling elderly. *Am J Epidemiol*. 1993;137(3):342–354. doi:10.1093/oxfordjournals.aje.a116681
6. Kistler BM, Khubchandani J, Jakubowicz G, Wilund K, Sosnoff J. Falls and fall-related injuries among US adults aged 65 or older with chronic kidney disease. *Prev Chronic Dis*. 2018;15:E82. doi:10.5888/pcd15.170518
7. Kistler BM, Khubchandani J, Wiblishauser M, Wilund KR, Sosnoff JJ. Epidemiology of falls and fall-related injuries among middle-aged adults with kidney disease. *Int Urol Nephrol*. 2019;51(9):1613–1621. doi:10.1007/s11255-019-02148-8
8. Bortolussi-Courval É, Podymow T, Trinh E, et al. Electronic decision support for deprescribing in patients on hemodialysis: clinical research protocol for a prospective, controlled, quality improvement study. *Can J Kidney Health Dis*. 2023;10:20543581231165712. doi:10.1177/20543581231165712
9. Aguilar A, Gifre L, Ureña-Torres P, et al. Pathophysiology of bone disease in chronic kidney disease: from basics to renal osteodystrophy and osteoporosis. *Front Physiol*. 2023;14:1177829. doi:10.3389/fphys.2023.1177829
10. Hall RK, Landerman LR, O'Hare AM, Anderson RA, Colón-Emeric CS. Chronic kidney disease and recurrent falls in nursing home residents: a retrospective cohort study. *Geriatr Nurs*. 2015;36(2):136–141. doi:10.1016/j.gerinurse.2014.12.012
11. Falcão L, Raimundo M, Fernandes S, et al. At least one hyperkalemia episode is associated with mortality and disease progression in stage 3 CKD patients. *J Nephrol*. 2023;36(9):2655–2656. doi:10.1007/s40620-023-01647-3
12. Gungor O, Ulu S, Hasbal NB, Anker SD, Kalantar-Zadeh K. Effects of hormonal changes on sarcopenia in chronic kidney disease: where are we now and what can we do? *J Cachexia Sarcopenia Muscle*. 2021;12(6):1380–1392. doi:10.1002/jcsm.12839
13. Shen Y. Role of nutritional vitamin D in chronic kidney disease-mineral and bone disorder: a narrative review. *Medicine*. 2023;102(14):e33477. doi:10.1097/md.00000000000033477
14. Liao Y, Zhang L, Peng Y, et al. Characteristics of accidental falls in the patients with chronic kidney diseases: a 14-year retrospective study and review. *medRxiv*. 2023;3:1.
15. van der Velde N, Seppala LJ, Hartikainen S, et al. European position paper on polypharmacy and fall-risk-increasing drugs recommendations in the world guidelines for falls prevention and management: implications and implementation. *Eur Geriatr Med*. 2023. doi:10.1007/s41999-023-00824-8
16. Novotna K, Zeiselova J, Stanickova B, Kovari M, Uher T, Janatova M. Interest in telerehabilitation among patients with mild to severe multiple sclerosis: results of the Czech Republic. *Telemed J E Health*. 2023. doi:10.1089/tmj.2023.0065
17. Bowling CB, Hall RK, Khakharia A, Franch HA, Plantinga LC. Serious fall injury history and adverse health outcomes after initiating hemodialysis among older U.S. ADULTS. *J Gerontol a Biol Sci Med Sci*. 2018;73(9):1216–1221. doi:10.1093/gerona/glx260
18. Kear T, Ulrich B. Patient Safety and patient safety culture in nephrology nurse practice settings: issues, solutions, and best practices. *Nephrol Nurs J*. 2015;42(2):113–22; quiz 123.
19. Shiomi K, Saito C, Nagai K, et al. Ratio of serum creatinine to cystatin C is related to leg strength in predialysis CKD patients. *Clin Exp Nephrol*. 2021;25(10):1079–1086. doi:10.1007/s10157-021-02050-7
20. Rampersad C, Brar R, Connelly K, et al. Association of physical activity and poor health outcomes in patients with advanced CKD. *Am J Kidney Dis*. 2021;78(3):391–398. doi:10.1053/j.ajkd.2020.12.018
21. Assimom MM, Flythe JE. Zolpidem versus trazodone initiation and the risk of fall-related fractures among individuals receiving maintenance hemodialysis. *Clin J Am Soc Nephrol*. 2020;16(1):88–97. doi:10.2215/cjn.10070620
22. Chao YT, Kuo FH, Lee YS, et al. Characteristics and outcome determinants of hospitalized older patients with cognitive dysfunction. *Int J Environ Res Public Health*. 2022;19:1.
23. Takase R, Nakata T, Aoki K, et al. The relationship between edema and body functions in patients with chronic kidney disease: a preliminary study. *Cureus*. 2022;14(7):e27118. doi:10.7759/cureus.27118
24. Dávalos-Yerovi V, Pérez-Sáez MJ, Faura-Vendrell A, et al. Association between GLIM-defined malnutrition and hospitalizations in kidney transplant candidates: a post hoc analysis of a cohort study. *JPEN J Parenter Enteral Nutr*. 2023;14. doi:10.1002/jpen.2532
25. Cogley C, Carswell C, Bramham J, et al. Improving kidney care for people with severe mental health difficulties: a thematic analysis of twenty-two healthcare providers' perspectives. *Front Public Health*. 2023;11:1225102. doi:10.3389/fpubh.2023.1225102
26. Krauss MJ, Nguyen SL, Dunagan WC, et al. Circumstances of patient falls and injuries in 9 hospitals in a Midwestern healthcare system. *Infect Control Hosp Epidemiol*. 2007;28(5):544–550. doi:10.1086/513725
27. Roberts RG, Kenny RA, Brierley EJ. Are elderly haemodialysis patients at risk of falls and postural hypotension? *Int Urol Nephrol*. 2003;35(3):415–421. doi:10.1023/b:urol.0000022866.07751.4a
28. Gamboa JL, Roshanravan B, Towse T, et al. Skeletal muscle mitochondrial dysfunction is present in patients with CKD before initiation of maintenance hemodialysis. *Clin J Am Soc Nephrol*. 2020;15(7):926–936. doi:10.2215/cjn.10320819
29. Thome T, Kumar RA, Burke SK, et al. Impaired muscle mitochondrial energetics is associated with uremic metabolite accumulation in chronic kidney disease. *JCI Insight*. 2020;6:1.
30. Wang XH, Mitch WE, Price SR. Pathophysiological mechanisms leading to muscle loss in chronic kidney disease. *Nat Rev Nephrol*. 2022;18(3):138–152. doi:10.1038/s41581-021-00498-0
31. Karava V, Dotis J, Kondou A, Printza N. Malnutrition patterns in children with chronic kidney disease. *Life*. 2023;13:3.
32. Zhao Y, Yi Y, Gu B, Wang H, Ma J, Guo Z. Echinacoside protects adenine-induced uremic rats from sciatic nerve damage by up-regulating α -Klotho. *J Musculoskelet Neuronal Interact*. 2021;21(3):413–421.
33. Tan RZ, Li JC, Zhu BW, et al. Neuropeptide Y protects kidney from acute kidney injury by inactivating M1 macrophages via the Y1R-NF- κ B-Mincle-dependent mechanism. *Int J Biol Sci*. 2023;19(2):521–536. doi:10.7150/ijbs.80200
34. Korucu B, Erten YT, Yeter HH, et al. Hypothalamic energy regulatory peptides in chronic kidney disease. *Ther Apher Dial*. 2019;23(5):437–443. doi:10.1111/1744-9987.12798
35. Del Colle S, Morello F, Rabbia F, et al. Antihypertensive drugs and the sympathetic nervous system. *J Cardiovasc Pharmacol*. 2007;50(5):487–496. doi:10.1097/FJC.0b013e318135446c
36. Yildiz S, Heybeli C, Smith L, Soysal P, Kazancioglu R. The prevalence and clinical significance of loss of appetite in older patients with chronic kidney disease. *Int Urol Nephrol*. 2023;3. doi:10.1007/s11255-023-03540-1

37. Ofori EK, Adekena CN, Boima V, et al. Serum leptin levels in patients with chronic kidney disease and hypertensive heart disease: an observational cross-sectional study. *Health Sci Rep.* 2023;6(1):e1053. doi:10.1002/hsr2.1053
38. Collister D, Herrington G, Delgado L, et al. Patient views regarding cannabis use in chronic kidney disease and kidney failure: a survey study. *Nephrol Dial Transplant.* 2023;38(4):922–931. doi:10.1093/ndt/gfac226
39. Moussa O, Feng C, Wang JX, Li XS, Zhang FX, Tang XH. Safety and effectiveness of roxadustat in dialysis-dependent patients with renal anemia: a hospital-based cohort study. *Cureus.* 2022;14(4):e24427. doi:10.7759/cureus.24427
40. Silva MZC, Cederholm T, Gonzalez MC, Lindholm B, Avesani CM. GLIM in chronic kidney disease: what do we need to know? *Clin Nutr.* 2023;42(6):937–943. doi:10.1016/j.clnu.2023.04.019
41. Mirmohammadali SN, Gallant KMH, Biruete A. Oh, My Gut! New insights on the role of the gastrointestinal tract and the gut microbiome in chronic kidney disease-mineral and bone disorder. *Curr Opin Nephrol Hypertens.* 2024;33(2):226–230. doi:10.1097/mnh.0000000000000961
42. Jiang S, Fang J, Li W. Protein restriction for diabetic kidney disease. *Cochrane Database Syst Rev.* 2023;1(1):Cd014906. doi:10.1002/14651858.CD014906.pub2
43. Spahia N, Rroji M, Barbullushi M, Spasovski G. The role of protein restriction in the progression of chronic kidney disease. *Pril.* 2023;44(2):53–63. doi:10.2478/prilozi-2023-0025
44. Olivari V, Di Modica SM, Lidonnici MR, et al. A single approach to targeting transferrin receptor 2 corrects iron and erythropoietic defects in murine models of anemia of inflammation and chronic kidney disease. *Kidney Int.* 2023;104(1):61–73. doi:10.1016/j.kint.2023.03.012
45. Hain D, Bednarski D, Cahill M, et al. Iron-deficiency anemia in CKD: a narrative review for the kidney care team. *Kidney Med.* 2023;5(8):100677. doi:10.1016/j.xkme.2023.100677
46. Alvarenga L, Saldanha JF, Stockler-Pinto MB, Fouque D, Soulage CO, Mafra D. Effects of resveratrol on inflammation and oxidative stress induced by the uremic toxin indoxyl sulfate in Murine macrophage-like RAW 264.7. *Biochimie.* 2023;213:22–29. doi:10.1016/j.biochi.2023.05.001
47. Yang G, Tan L, Yao H, Xiong Z, Wu J, Huang X. Long-term effects of severe burns on the kidneys: research advances and potential therapeutic approaches. *J Inflamm Res.* 2023;16:1905–1921. doi:10.2147/JIR.S404983
48. Yang G, Waheed S, Wang C, Shekh M, Li Z, Wu J. Exosomes and Their bioengineering strategies in the cutaneous wound healing and related complications: current knowledge and future perspectives. review. *Int J Bio Sci.* 2023;19(5):1430–1454. doi:10.7150/ijbs.80430
49. Zoccali C, Mallamaci F. Innate Immunity system in patients with cardiovascular and kidney disease. *Circ Res.* 2023;132(8):915–932. doi:10.1161/circresaha.122.321749
50. Bowling CB, Bromfield SG, Colantonio LD, et al. Association of reduced eGFR and albuminuria with serious fall injuries among older adults. *Clin J Am Soc Nephrol.* 2016;11(7):1236–1243. doi:10.2215/cjn.11111015
51. Koury MJ, Haase VH. Deregulating iron-erythropoiesis regulation: transferrin receptor 2 as potential target for treating anemia in CKD. *Kidney Int.* 2023;104(1):25–28. doi:10.1016/j.kint.2023.04.017
52. Mishina S, Waratani M, Onozawa S, Okumura H, Ito Y, Yasuda Y. A retrospective database analysis of erythropoiesis-stimulating agent treatment patterns and associated healthcare resource use in patients with non-dialysis-dependent chronic kidney disease-related anaemia in Japan. *Nephrology.* 2023;28(8):446–455. doi:10.1111/nep.14168
53. Hassanein M, Arragain S, Schold JD, et al. Dysnatremias, mortality, and kidney failure in CKD: findings From the Chronic Renal Insufficiency Cohort (CRIC) study. *Kidney Med.* 2022;4(12):100554. doi:10.1016/j.xkme.2022.100554
54. Shin J, Lee CH. The roles of sodium and volume overload on hypertension in chronic kidney disease. *Kidney Res Clin Pract.* 2021;40(4):542–554. doi:10.23876/j.krcp.21.800
55. Jo SM. Understanding and treatment strategies of hypertension and hyperkalemia in chronic kidney disease. *Electrolyte Blood Press.* 2023;21(1):24–33. doi:10.5049/ebp.2023.21.1.24
56. Chang HH, Chiang JH, Tsai CC, Chiu PF. Predicting hyperkalemia in patients with advanced chronic kidney disease using the XGBoost model. *BMC Nephrol.* 2023;24(1):169. doi:10.1186/s12882-023-03227-w
57. Pietribiasi M, Waniewski J, Leypoldt JK. Mathematical modelling of bicarbonate supplementation and acid-base chemistry in kidney failure patients on hemodialysis. *PLoS One.* 2023;18(2):e0282104. doi:10.1371/journal.pone.0282104
58. Tartaglione L, Rotondi S, Aucella F, et al. Parathyroidectomy and survival in a cohort of Italian dialysis patients: results of a multicenter, observational, prospective study. *J Nephrol.* 2023;23. doi:10.1007/s40620-023-01658-0
59. Ionova TI, Buzanakov DM, Chernikov RA, et al. Quality of life in patients with primary hyperparathyroidism before and after parathyroidectomy: long term single center experience. *BMC Endocr Disord.* 2023;23(1):87. doi:10.1186/s12902-023-01344-z
60. Carsote M, Nistor C. Forestalling hungry bone syndrome after parathyroidectomy in patients with primary and renal hyperparathyroidism. *Diagnostics.* 2023;13:11.
61. Echterdiek F, Ranke MB, Schwenger V, Heemann U, Latus J. Kidney disease and thyroid dysfunction: the chicken or egg problem. *Pediatr Nephrol.* 2022;37(12):3031–3042. doi:10.1007/s00467-022-05640-z
62. Rhee CM. Thyroid disease in end-stage renal disease. *Curr Opin Nephrol Hypertens.* 2019;28(6):621–630. doi:10.1097/mnh.0000000000000542
63. Drube J, Wan M, Bonthuis M, et al. Clinical practice recommendations for growth hormone treatment in children with chronic kidney disease. *Nat Rev Nephrol.* 2019;15(9):577–589. doi:10.1038/s41581-019-0161-4
64. Wang Y, Ma W, Pu J, Chen F. Interrelationships between sarcopenia, bone turnover markers and low bone mineral density in patients on hemodialysis. *Ren Fail.* 2023;45(1):2200846. doi:10.1080/0886022x.2023.2200846
65. Ling J, Ng JKC, Chan JCN, Chow E. Use of continuous glucose monitoring in the assessment and management of patients with diabetes and chronic kidney disease. *Front Endocrinol.* 2022;13:869899. doi:10.3389/fendo.2022.869899
66. Long S, Hu L, Luo Y, Li Y, Ding F. Incidence and risk factors of falls in older adults after discharge: a prospective study. *Int J Nurs Sci.* 2023;10(1):23–29. doi:10.1016/j.ijnss.2022.12.010
67. Tanaka Y. Rheumatoid arthritis. *Inflamm Regen.* 2020;40(1):20. doi:10.1186/s41232-020-00133-8
68. Sullivan R, Hemsley B, Skinner I, Harding K. Hospital policies on falls in relation to patients with communication disability: a scoping review and content analysis. *Aust Health Rev.* 2023. doi:10.1071/ah22289
69. Bae S, Lee S, Park H, et al. Position statement: exercise guidelines for osteoporosis management and fall prevention in osteoporosis patients. *J Bone Metab.* 2023;30(2):149–165. doi:10.11005/jbm.2023.30.2.149

70. Bennett PN, Bohm C, Yee-Moon Wang A, et al. An International survey of peritoneal dialysis exercise practices and perceptions. *Kidney Int Rep.* 2023;8(7):1389–1398. doi:10.1016/j.ekir.2023.04.024
71. Sharma B, Yadav DK. L-Carnitine and Chronic kidney disease: a comprehensive review on nutrition and health perspectives. *J Pers Med.* 2023;13(2).
72. Chang L, Tian R, Guo Z, et al. La dieta baja en proteínas suplementada con inulina reduce los niveles de toxinas unidas a proteínas en pacientes con enfermedad renal crónica en estadio 3b-5: un estudio controlado aleatorio [Low-protein diet supplemented with inulin lowers protein-bound toxin levels in patients with stage 3b-5 chronic kidney disease: a randomized controlled study]. *Nutr Hosp.* 2023. doi:10.20960/nh.04643
73. Rhee CM, Wang AY, Biruete A, et al. Nutritional and dietary management of chronic kidney disease under conservative and preservative kidney care without dialysis. *J Ren Nutr.* 2023. doi:10.1053/j.jrn.2023.06.010
74. Torregrosa JV, Bover J, Rodríguez Portillo M, et al. Recommendations of the Spanish Society of Nephrology for the management of mineral and bone metabolism disorders in patients with chronic kidney disease: 2021 (SEN-MM). *Nefrologia.* 2023;43(Suppl 1):1–36. doi:10.1016/j.nefro.2023.03.003
75. Hao G, Lu W, Huang J, et al. Predialysis fluid overload linked with quality of sleep in patients undergoing hemodialysis. *Sleep Med.* 2018;51:140–147. doi:10.1016/j.sleep.2018.07.011
76. Lin HC, Chen MJ, Lee CH, Kung LC, Huang JT. Fall recognition based on an imu wearable device and fall verification through a smart speaker and the IoT. *Sensors.* 2023;23:12.
77. Fukaya K, Uchida M. Protection against impact with the ground using wearable airbags. *Ind Health.* 2008;46(1):59–65. doi:10.2486/indhealth.46.59
78. Tarbert RJ, Singhatat W. Real world evidence of wearable smartbelt for mitigation of fall impact in older adult care. *IEEE J Transl Eng Health Med.* 2023;11:247–251. doi:10.1109/jtehm.2023.3256893
79. Botonis OK, Harari Y, Embry KR, et al. Wearable airbag technology and machine learned models to mitigate falls after stroke. *J Neuroeng Rehabil.* 2022;19(1):60. doi:10.1186/s12984-022-01040-4
80. Percy D, Phillips T, Torres F, Chaleunphonh M, Sung P. Effectiveness of virtual reality-based balance and gait in older adults with fear of movement: a systematic review and meta-analysis. *Physiother Res Int.* 2023;e2037. doi:10.1002/pri.2037
81. Martínez Montilla LA, López Cruces K, Calderón Erazo HS, et al. Effectiveness of virtual reality in balance training for fall prevention in older adults: systematic review. *Sports Med Arthrosc Rev.* 2023;31(2):41–48. doi:10.1097/jsa.0000000000000367
82. Zhao R, Zhao X, Guan J, Zhang C, Zhu K. The effect of virtual reality technology on anti-fall ability and bone mineral density of the elderly with osteoporosis in an elderly care institution. *Eur J Med Res.* 2023;28(1):204. doi:10.1186/s40001-023-01165-9
83. Bally ELS, Ye L, van Grieken A, et al. Factors associated with falls among hospitalized and community-dwelling older adults: the APPCARE study. *Front Public Health.* 2023;11:1180914. doi:10.3389/fpubh.2023.1180914
84. Leal JC, Belo VS, Santos IM, Ferreira RV, de Melo SN, da Silva ES. Exergames in older adult community centers and nursing homes to improve balance and minimize the risk of falls in older adults: a systematic review and meta-analysis. *Healthcare.* 2023;11:13.

Nursing: Research and Reviews

Dovepress

Publish your work in this journal

Nursing: Research and Reviews is an international, peer-reviewed, open access journal publishing original research, reports, reviews and commentaries on all aspects of nursing and patient care. These include patient education and counseling, ethics, management and organizational issues, diagnostics and prescribing, health outcomes, economics and resource management, improving patient safety in all settings. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/nursing-research-and-reviews-journal>