



Effect of SophorOx[®] on Oxidative Stress and Body Composition in Individuals with High BMI: A Randomized Controlled Trial

James M Roza ¹, Shalini Srivastava ²

¹Research & Development, Layn Natural Ingredients, Irvine, CA, 92602, USA; ²Clinical Development Department, Vedic Lifesciences Pvt. Ltd, Mumbai, Maharashtra, India

Correspondence: Shalini Srivastava, Vedic Lifesciences Pvt. Ltd, Morya House, 118-B, off Link Road, Andheri West, Mumbai, Maharashtra, 400053, India, Tel +91 9619477885, Email shalini.s@vediclifesciences.com

Purpose: The study aimed to investigate the efficacy and safety of SophorOx[®] (LN-OS-22) on oxidative stress and body composition in adults with excessive body weight and obesity.

Participants and Methods: The 56-days randomized, double-blind, placebo-controlled, parallel-group, multi-centric clinical trial had individuals aged 30–60 years with body mass index (BMI) ≥ 25 to ≤ 34.9 kg/m². 68 participants were randomly allocated to LN-OS-22 or placebo groups. The primary outcome was improvement in the oxidative stress. Secondary outcomes were changes in plasma lipopolysaccharide (LPS) and serum malondialdehyde (MDA) levels, weight and waist circumference, inflammatory markers, and quality of life.

Results: At day 56, a statistically significant change in the 8-Isoprostane levels between LN-OS-22 vs placebo was observed ($p = 0.0222$). As compared to placebo, at the end of study, statistically significant reductions were demonstrated in body weight, waist circumference and BMI in the LN-OS-22 group ($p < 0.0001$). Also, a statistically significant change when compared to placebo for the energy/stamina domain ($p = 0.0300$) of the Impact of Weight on Quality of Life-Lite-Clinical Trials Version (IWQOL-Lite-CT) questionnaire was depicted in LN-OS-22 group.

Conclusion: The study demonstrates that LN-OS-22 was effective in reducing the oxidative stress, anthropometrics and improving the quality of life in individuals with overweight and obesity.

Keywords: obesity, oxidative stress, inflammation, weight loss, body composition

Introduction

In a healthy individual, the balance between pro-oxidant generation such as Reactive Oxygen Species (ROS) and Reactive Nitrogen Species (RNS) and antioxidant defense is autonomic in nature. Exposure to toxic xenobiotics coupled with the sedentary lifestyle, results in an imbalance between the pro-oxidants and antioxidants, leading to oxidative stress. Pro-oxidant antioxidant balance in the cell is shifted towards the pro-oxidants if the production of oxygen species is increased greatly or when levels of antioxidants are diminished. This state is called oxidative stress.¹ This oxidative stress is known to be one of the primary mechanism through which these lifestyle changes adversely affect individual health leading to a disruption of redox signaling and control and/or molecular damage.^{2–5} As demonstrated by several studies, overproduction of ROS can lead to damage of cellular molecules gradually resulting into white adipose tissue accumulation and release of several pro-inflammatory cytokines.^{6,7} Oxidative stress is known to trigger the deposition of white adipose tissue by increasing preadipocyte proliferation, adipocyte differentiation, and the size of mature adipocytes.⁸ Other contributing factors of oxidative stress are low antioxidant defense,⁹ chronic inflammation,¹⁰ and postprandial ROS generation.¹¹ Therefore, obesity served as an ideal condition to investigate the antioxidant activity of LN-OS-22 extract.

Several dietary practices have been shown to alleviate oxidative stress thereby minimizing the health associated hazards.¹² *Sophora japonica* L (*Styphnolobium japonicum*) has a rich history of use in traditional Chinese medicine. A wide range of biological activities of *Sophora japonica* L in vitro and in vivo have been reported such as anti-inflammatory, antibacterial, antiviral, anti-osteoporotic, antioxidant, radical scavenging, anti-hyperglycemic, anti-obesity, antitumor, and hemostatic effects.^{13,14} *Sophora Japonica* L is one of the richest sources of rutin and quercetin flavonoids. Both of these flavonoids have been studied extensively for exploiting their use in improving human health.^{15–20} Layn Natural Ingredients adopted *Sophora japonica* L under the “Adopt-an-Herb” initiative of the American Botanical Council and derived LN-OS-22. It was extracted from the flower bud of the plant, underwent ethanol/water extraction, and has been standardized to contain $\geq 90\%$ flavonoids, primarily, quercetin, a rutin and other minor flavonoids. This extract has been tested in vitro in LPS-stimulated RAW 264.7 macrophages for its anti-inflammatory and antioxidant activity. The extract showed significant inhibition of pathways of inflammation and oxidative stress.²¹ The biological activities were then confirmed by an in vivo study. Oral administration of this extract in exercised animals resulted in a significant decrease in oxidative stress and inflammatory biomarkers associated with the high-intensity exercise in a rodent model (Sprague-Dawley rat model).²² The validation in humans of these in vitro and in vivo antioxidant activities was studied in the current randomized clinical trial.

A meta-analysis²³ of 50 clinical studies showed that individuals with overweight and obesity have a comparatively increased levels of 8-isoprostane, a reliable and stable biomarker of oxidative stress,²⁴ than healthy population. A reduction in the body weight is known to decrease oxidation markers in the body and thereby reduce obesity-related pathological risk factors.⁸ Obesity is also known to be associated with chronic low-grade inflammation.²⁵ A significant impact on the health-related quality of life is also seen in individuals with overweight and obesity.²⁶ Thus, the primary objective of the study was to assess the effect of the investigational product (IP), LN-OS-22 on oxidative stress. Body composition (BMI, body weight and waist circumference) were included as secondary objectives as clinical indicators of reduction in oxidative stress and inflammation. Furthermore, lipid peroxidation, metabolic endotoxemia as well as quality of life were also included as secondary objectives. The safety of 2-month administration of the Investigational Product (IP), LN-OS-22, was evaluated by monitoring the vitals as well as liver and kidney function biomarkers [serum glutamic oxaloacetic transaminase (SGOT), serum glutamic pyruvic transaminase (SGPT), alkaline phosphatase (ALP), and creatinine] throughout the study.

Materials and Methods

Study Approval and Registration

This study was conducted in compliance with the Declaration of Helsinki, International Conference on Harmonization (ICH) recommendation on Good Clinical Practice (GCP)-E6 (R2), 2016, and National Ethical Guidelines for Biomedical and Health Research involving Human Participants, 2017, India. The ethical approval for conduct of the study at three sites was provided by the independent ethics committee - Harmony Ethical Research Committee, Mumbai, India (Reg. No. ECR/1411/Inst/MH/2020). The eligible participants were enrolled in the study only after providing written consent for participation. The study was registered with the NIH ClinicalTrials.gov (Identifier: NCT05404217) and Clinical Trials Registry, India (Registration Number: CTRI/2022/06/043394). Vedic Lifesciences, Mumbai, India, monitored and audited the study to ensure the study protocol and ICH-GCP compliance. The study report conformed to the Consolidated Standard Reporting of Trials (CONSORT) guidelines.

Study Design

The current study is a randomized, double-blind, placebo-controlled, parallel-group multi-centric study intended for evaluating the effect of 56-days administration of LN-OS-22 on oxidative stress and body composition in individuals with overweight and obesity. The study was conducted at three clinical sites in Mumbai, Maharashtra, India, under the supervision of qualified physicians from June 2022 to March 2023.

Participants

The present study was conducted among adults aged ≥ 30 to ≤ 60 years with a BMI range of ≥ 25 to ≤ 34.9 kg/m². Individuals with at least three of the following five metabolic risk factors were included in the study: 1) waist circumference: male: ≥ 102 cm;

female ≥ 88 cm, 2) triglycerides > 150 mg/dL, 3) systolic blood pressure ≥ 130 mm Hg and/or diastolic blood pressure ≥ 85 mm Hg, 4) fasting blood glucose ≥ 100 mg/dL, 5) high-density lipoproteins: male: < 40 mg/dL; female: < 50 mg/dL. Also, individuals who were physically inactive for 1/3rd of their wake time as per the Longitudinal Aging Study Amsterdam (LASA) sedentary behavior questionnaire were included. Individuals meeting the following criteria were excluded from the study: known cases of lactose intolerance, thyroid function test (TSH) level < 0.4 to > 4.2 mIU/L, known cases of type I diabetes mellitus and presence of uncontrolled type II diabetes mellitus [indicated by glycosylated hemoglobin (HbA1c) ≥ 6.5]. The other eligibility criteria are described in [Table S1 \(Supplementary Material\)](#).

The prospective participants fulfilling inclusion and exclusion criteria were randomized in the ratio of 1:1 for LN-OS-22 or placebo by the statistician who was not directly involved in the study. Based on the computer generated randomization chart, the participant identification numbers were generated which were printed on the product labels using a stratified block randomization method with a block of four. The randomization chart was generated by an independent statistician using StatsDirect software (Ver. 3.1.17). The participant, research staff, and Investigator were blinded to the study product allocation.

Intervention

The investigational product (IP), LN-OS-22, is a polyphenolic supplementation comprising of rutin, quercetin, and other minor flavonoids. Previous studies have shown that quercetin up to a dose of 2000 mg and Rutin in a dose of 500 mg when administered for 6–8 weeks is safe in humans.^{27–30} Thus, in the present study, 500 mg capsule of LN-OS-22 was administered to participants in one of the arms. The detailed percentage composition of the study products is given in [Table 1](#). The placebo capsule contained 500 mg of Maltodextrin. All the participants were instructed to take two capsules a day orally with a glass of milk after breakfast daily. At each follow-up visit, the compliance to the intervention was ensured by counting the remaining number of capsules in the bottles. Any concomitant medication taken was recorded in the source document and Electronic-Case Record Form (e-CRF). The participants having at least 90% compliance during the intervention period were considered “completed intervention”.

To preserve the blinding, the study products and the placebo capsules were matched for size, shape, color, and texture and packed in identical packaging. Layn Natural Ingredients, USA supplied the products for the study. The products were manufactured in compliance with good manufacturing practices and applicable regulations.

Study Procedures

During the screening visit, the study procedures were thoroughly explained to the research participants in a language well understood by them, following which he/ she duly signed and dated informed consent. The participants were screened for the eligibility criteria, and those who qualified were instructed to report on day 0 for randomization and baseline assessments. The follow-up assessments were conducted on days 28 and 56, as depicted in ([Figure 1](#)). Demographic and baseline characteristics such as anthropometric parameters (height, weight and BMI), vital signs (pulse rate and blood pressure), clinical examination

Table 1 Investigational Product Composition

Study Product	Active Ingredients	Percentage	Total Composition
LN-OS-22	Quercetin	51.7%	94.81%
	Rutin	40.75%	
	Kaempferol	0.88%	
	Isorhamnetin	1.33%	
	Genistin	0.14%	
	Genistein	0.005%	
Placebo	Maltodextrin	100%	100%




Figure 1 Study flow chart.
Abbreviation: n; number of participants.

and medical history with prior and concomitant medication were recorded. Participants were asked to self-administer the LASA questionnaire. Participants were required to have a minimum of 90% Investigational Product (IP) compliance during the study. The detailed method of assessments is explained in the [Supplementary Material](#).

Study Outcomes

Primary Outcome

The primary objective of the study was to assess the effect of 56 days of IP intake on the oxidative stress as compared to placebo. This was assessed by the change in serum levels of 8-isoprostane (8-iso-PGF₂α) which is considered a reliable and pivotal biomarker of generalized as well as cellular oxidative stress.³¹ The 8-iso-PGF₂α levels were analyzed using sandwich ELISA technique.

Secondary Outcomes

The secondary objective was to assess the effect of IP on body weight, waist circumference along with the inflammatory (CRP and IL-6) and lipid peroxidation biomarkers (MDA) along with the serum lipopolysaccharide (LPS) levels, a marker for the bacterial pollution. Lastly, effect of IP was assessed on the quality of life using Impact of Weight on Quality of Life-Lite-Clinical Trials Version (IWQOL-Lite-CT). The detailed study flow chart is shown in [Figure 1](#). The participants were instructed to maintain their routine diet and physical activity levels throughout the study.

Safety Outcomes

The safety of the IP was evaluated by measuring the liver and kidney function panel [serum glutamic-oxaloacetic transaminase (SGOT), serum glutamic-pyruvic transaminase (SGPT), serum alkaline phosphatase (ALP) and serum creatinine]. Blood pressure, pulse rate along with adverse events were monitored throughout the study.

Statistical Analysis

The sample size calculation was based on previous studies conducted to evaluate the efficacy of other products with similar objectives.^{32–34} The null hypothesis was considered to be rejected in case LN-OS-22 was able to significantly reduce oxidative stress and body composition in a population with overweight and obesity as compared to the placebo. Data was checked for normality using Shapiro Wilk/Kolmogorov–Smirnov test. Mean change of serum levels of 8-isoprostane at 56 days from baseline was compared between the IP and placebo groups using ANCOVA. Mean change in the serum MDA, plasma LPS levels at 56 days from baseline were assessed on similar line of primary endpoints. Change in serum CRP, serum IL-6, weight and waist circumference, and IWQOL-Lite-CT from baseline to Day 28 and Day 56, in the IP and placebo groups was analyzed using multivariate ANCOVA. Data analyses were performed using R/R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/version4.0.5> and XLSTAT statistical and data analysis solution, New York, USA. <https://www.xlstat.com/version2021.3.1>.

Quality Assurance

Vedic Lifesciences monitored and audited the study to ensure compliance with the study protocol and ICH-GCP E6 (R2) guidelines.

Results

Out of 153 screened individuals, 68 met the predefined inclusion-exclusion criteria and were randomized into the study. Of these, 50 participants completed the study. The detailed participant disposition is presented in [Figure 2](#).

Demographic and Baseline Characteristics

Eligible individuals were randomized in a 1:1 ratio to receive either LN-OS-22 (n = 33) or placebo (n = 35). The enrolled participants had a mean age of 42 years and a mean BMI of 30.25 kg/m². The demographic and baseline characteristics of all the randomized participants are summarized in [Table 2](#).

Oxidative Stress

At the baseline, 8-Isoprostane levels matched between the groups (p = 0.1016). At day 28, the serum levels increased in both the LN-OS-22 and the placebo (p = 0.7634) groups. At day 56, the increase in the levels of the 8-isoprostane was significantly higher in the placebo group as compared to the IP (p = 0.0222) and this increase was significantly higher as compared to the baseline (p = 0.0710). Thus, at the end of the study, the LN-OS-22 helped restore the balance between anti- and pro-oxidative levels ([Table 3](#)).

Body Composition

The LN-OS-22 study group demonstrated not only a statistically significant reduction in the body weight and BMI but also showed a clinically meaningful reduction in the waist circumference when compared with the placebo. Eight weeks' administration of the IP reduced the body weight by 2.07 (1.62) kgs as opposed to the placebo group which showed only a marginal reduction by 0.14 (1.53) kgs (p < 0.0001). The IP group, thus, also showed a statistically significant reduction in the BMI as compared to placebo (p < 0.0001). Similarly, waist circumference reduced by 2.58 (1.69) cm in the LN-OS-22 group as compared to the placebo which showed only a 0.31 (1.71) cm of reduction over 56 days of administration (p < 0.0001). This decrease in waist circumference resulted in the significant reduction in the waist-to-height ratio (p = 0.0005) ([Table 4](#)).

Inflammatory & Lipid Peroxidation Markers

At baseline, the groups seemed statistically similar for inflammatory and lipid peroxidation markers (p > 0.05). After 8 weeks of study period, the LN-OS-22 group showed lower levels of the CRP as compared to the placebo, however, the levels of CRP increased in both the groups from baseline. Similarly, the IL-6 levels were also lower in the intervention




Figure 2 Participant Disposition.
Abbreviation: n; number of participants,

group as compared to the placebo group however the results were not statistically significant (Table 5) By the end of the study, a reduction in the mean MDA levels was observed in both study groups. The LN-OS-22 showed a greater mean reduction of 5.52 (70.99) EU/mL of LPS levels in comparison to the placebo study group (Table 5).

Table 2 Demographics and Baseline Characteristics of Randomized Participants

Parameters	Categories	LN-OS-22	Placebo	Total	*p-value
		(N=33)	(N=35)	(N=68)	
Age (Years)	Mean	42.85 (7.42)	40.57 (5.91)	41.68 (6.73)	0.1652 (T)
	(Min, Max)	(31.00, 57.00)	(31.00, 52.00)	(31.00, 57.00)	
Gender	Male	10 (30.30%)	12 (34.29%)	22 (32.35%)	0.7257 (C)
	Female	23 (69.70%)	23 (65.71%)	46 (67.65%)	
Height (cm)	Mean	158.34 (7.96)	157.83 (8.98)	158.08 (8.44)	0.8065 (T)
	(Min, Max)	(143.06, 179.31)	(140.85, 175.00)	(140.85, 179.31)	
Weight (kg)	Mean	75.66 (8.89)	76.07 (10.47)	75.87 (9.66)	0.8644 (T)
	(Min, Max)	(57.80, 98.00)	(60.00, 102.30)	(57.80, 102.30)	
BMI (kg/m ²)	Mean	30.14 (2.84)	30.36 (2.64)	30.25 (2.72)	0.7390 (T)
	(Min, Max)	(25.68, 34.51)	(25.48, 34.82)	(25.48, 34.82)	

Notes: *p values were calculated using two sample t-test (T) for continuous variables and Chi Square (C) test for categorical variables. Percentages were calculated using respective column header count as denominator.

Abbreviations: SD; standard deviation, N; number of participants.

Table 3 Summary of 8-Isoprostane Levels of the mITT Population

Parameters	Visit	LN-OS-22 (N=23)		Placebo (N=28)		p value [§]
		Mean (SD)	95% CI	Mean (SD)	95% CI	
8-Isoprostane (pg/mL)	Day 0	102.41 (60)	(76.46, 128.35)	175.26 (218.87)	(90.39, 260.13)	0.1016 (T)
	Day 56	109.03 (73.72)	(77.16, 140.91)	216.33 (205.7)	(136.56, 296.09)	0.0147 (T)
	Change from baseline at day 56	6.63 (85.19)	(-30.21, 43.47)	41.07 (115.64)	(-3.77, 85.91)	
	p value [#]	0.0222 (A)				
	p value*	0.7126 (T)		0.0710 (T)		

Notes: [§]p-value was calculated using two sample t-test (T). *p-values were calculated using paired t-test (T). [#]p-values were calculated using ANCOVA with treatment as factor and baseline value, age and BMI as covariate vs placebo.

Abbreviations: CI; confidence interval, N; number of participants, mITT; modified intention to treat, SD; standard deviation.

Table 4 Summary of Anthropometric Measurements of the mITT Population

Parameters	Visit	LN-OS-22 (N=23)		Placebo (N=28)		p value [§]
		Mean (SD)	95% CI	Mean (SD)	95% CI	
Body Weight (kgs)	Day 0	77.25 (7.24)	(74.12, 80.38)	75.99 (10.74)	(71.82, 80.15)	0.6317 (T)
	Day 56	75.18 (7.1)	(72.11, 78.25)	76.12 (10.58)	(72.02, 80.23)	0.7169 (T)
	Change from baseline at day 56	-2.07 (1.62)	(-2.77, -1.37)	0.14 (1.53)	(-0.45, 0.73)	
	p value [#]	<0.0001 (A)				
	p value*	<0.0001 (T)		0.6339 (T)		

(Continued)

Table 4 (Continued).

Parameters	Visit	LN-OS-22 (N=23)		Placebo (N=28)		p value [§]
		Mean (SD)	95% CI	Mean (SD)	95% CI	
Waist circumference (cm)	Day 0	103.16 (7.3)	(100.00, 106.32)	102.53 (7.94)	(99.45, 105.61)	0.7706 (T)
	Day 56	100.58 (6.85)	(97.62, 103.54)	102.22 (7.55)	(99.29, 105.15)	0.4264 (T)
	Change from baseline at day 56	-2.58 (1.69)	(-3.31, -1.85)	-0.31 (1.71)	(-0.98, 0.35)	
	p value [#]	<0.0001 (A)				
	p value*	<0.0001 (T)		0.3457 (T)		
Waist to height ratio	Day 0	0.65 (0.06)	(0.62, 0.68)	0.65 (0.04)	(0.63, 0.67)	0.8763 (T)
	Day 56	0.63 (0.06)	(0.61, 0.66)	0.65 (0.04)	(0.63, 0.66)	0.2995 (T)
	Change from baseline at day 56	-0.02 (0.01)	(-0.02, -0.01)	0 (0.01)	(-0.01, 0.00)	
	p value [#]	0.0005 (A)				
	p value*	<0.0001 (T)		0.1018 (T)		
BMI (kg/m ²)	Day 0	30.45 (2.71)	(29.28, 31.62)	30.44 (2.69)	(29.39, 31.48)	0.9888 (T)
	Day 56	29.65 (2.88)	(28.40, 30.89)	30.5 (2.62)	(29.48, 31.51)	0.2759 (T)
	Change from baseline at day 56	-0.8 (0.61)	(-1.06, -0.54)	0.06 (0.6)	(-0.17, 0.29)	
	p value [#]	<0.0001 (A)				
	p value*	<0.0001 (T)		0.5984 (T)		

Notes: [§]p-value was calculated using two sample t-test (T). *p-values were calculated using paired t-test (T). [#]p-values were calculated using ANCOVA with treatment as factor and baseline value, age and BMI as covariate vs placebo.

Abbreviations: BMI, body mass index, CI; confidence interval, mITT; modified intention to treat, N; number of participants SD; standard deviation.

Table 5 Summary of Inflammatory and Lipid Peroxidation Biomarkers of the mITT Population

Parameters	Visit	LN-OS-22 (N=23)		Placebo (N=28)		p value [§]
		Mean (SD)	95% CI	Mean (SD)	95% CI	
CRP (mg/L)	Day 0	5.63 (6.17)	(2.96, 8.30)	5.75 (5.39)	(3.65, 7.84)	0.9433 (T)
	Day 56	6.3 (9.18)	(2.33, 10.27)	7.45 (7.03)	(4.72, 10.17)	0.6157 (T)
	Change from baseline at day 56	0.67 (9.92)	(-3.62, 4.96)	1.7 (6.17)	(-0.69, 4.10)	
	p value [#]	0.8116 (A)				
	p value*	0.7481 (T)		0.1557 (T)		
IL-6 (pg/mL)	Day 0	7.03 (9.49)	(2.93, 11.14)	77.45 (374.07)	(-67.60, 222.50)	0.3282 (T)
	Day 56	7.5 (5.74)	(5.02, 9.98)	13.03 (18.16)	(5.99, 20.07)	0.1374 (T)
	Change from baseline at day 56	0.46 (11.89)	(-4.68, 5.60)	-64.42 (376.76)	(-210.51, 81.67)	
	p value [#]	0.1786 (A)				
	p value*	0.8535 (T)		0.3736 (T)		

(Continued)

Table 5 (Continued).

Parameters	Visit	LN-OS-22 (N=23)		Placebo (N=28)		p value [§]
		Mean (SD)	95% CI	Mean (SD)	95% CI	
MDA (nmol/mL)	Day 0	0.68 (0.36)	(0.52, 0.83)	0.66 (0.39)	(0.51, 0.81)	0.9013 (T)
	Day 56	0.5 (0.37)	(0.34, 0.67)	0.38 (0.4)	(0.22, 0.53)	0.2545 (T)
	Change from baseline at day 56	-0.17 (0.64)	(-0.45, 0.10)	-0.29 (0.47)	(-0.47, -0.10)	
	p value [#]	0.3871 (A)				
	p value*	0.2102 (T)		0.0036 (T)		
LPS (EU/mL)	Day 0	72.61 (47.79)	(51.94, 93.27)	78.21 (55.45)	(56.71, 99.71)	0.7041 (T)
	Day 56	67.09 (87.45)	(29.27, 104.90)	74.75 (61.22)	(51.01, 98.49)	0.7150 (T)
	Change from baseline at day 56	-5.52 (70.99)	(-36.22, 25.18)	-3.46 (52.62)	(-23.87, 16.94)	
	p value [#]	0.9093 (A)				
	p value*	0.7127 (T)		0.7303 (T)		

Notes: [§]p-value was calculated using two sample t-test (T). *p-values were calculated using paired t-test (T). [#]p-values were calculated using ANCOVA with treatment as factor and baseline value, age and BMI as covariate vs placebo.

Abbreviations: CI, confidence interval; CRP, C-reactive protein; IL-6, interleukin-6; LPS, lipopolysaccharide; MDA, malondialdehyde; mITT, modified intention to treat; N, number of participants; SD, standard deviation.

Quality of Life

On days 28 and 56, the change in the quality of life in the IP group was statistically significant as indicated by $p = 0.0029$ and $p = 0.0004$, respectively, when compared to baseline. After 56 days of intervention, LN-OS-22 group demonstrated a statistically significant change when compared to placebo for the energy/stamina domain ($p = 0.0300$) of the IWQOL-Lite-CT questionnaire. Also, at day 56, when compared to baseline, statistically significant results were demonstrated in the IP group in the physical impact ($p = 0.0047$), emotions ($p = 0.0003$), self-confidence/self-esteem ($p = 0.0236$) domains (Table 6).

Table 6 Summary of IWQOL-Lite-CT of the mITT Population

Parameters	Visit	LN-OS-22 (N=23)		Placebo (N=28)		p value [§]
		Mean (SD)	95% CI	Mean (SD)	95% CI	
Physical Impact	Day 0	10.22 (4.84)	(8.12, 12.31)	9.61 (4.57)	(7.84, 11.38)	0.6460 (T)
	Day 56	7.00 (4.72)	(4.96, 9.04)	7.64 (4.6)	(5.86, 9.43)	0.6259 (T)
	Change from baseline at day 56	-3.22 (4.91)	(-5.34, -1.10)	-1.96 (5.55)	(-4.12, 0.19)	
	p value [#]	0.4721 (A)				
	p value*	0.0047 (T)		0.0718 (T)		
Energy/Stamina	Day 0	2.43 (1.38)	(1.84, 3.03)	1.96 (1.23)	(1.49, 2.44)	0.2039 (T)
	Day 56	1.57 (1.38)	(0.97, 2.16)	1.89 (1.29)	(1.39, 2.39)	0.3847 (T)
	Change from baseline at day 56	-0.87 (1.18)	(-1.38, -0.36)	-0.07 (1.33)	(-0.59, 0.44)	
	p value [#]	0.0300 (A)				
	p value*	0.0019 (T)		0.7787 (T)		

(Continued)

Table 6 (Continued).

Parameters	Visit	LN-OS-22 (N=23)		Placebo (N=28)		p value [§]
		Mean (SD)	95% CI	Mean (SD)	95% CI	
Emotions	Day 0	12.26 (4.87)	(10.15, 14.37)	11.32 (5.16)	(9.32, 13.32)	0.5105 (T)
	Day 56	8.39 (4.39)	(6.49, 10.29)	9.46 (5.14)	(7.47, 11.46)	0.4323 (T)
	Change from baseline at day 56	-3.87 (4.26)	(-5.71, -2.03)	-1.86 (5.68)	(-4.06, 0.35)	
	p value [#]	0.1929 (A)				
	p value*	0.0003 (T)		0.0951 (T)		
Self-confidence/Self-esteem	Day 0	11.17 (6.1)	(8.53, 13.81)	10.32 (5.62)	(8.14, 12.50)	0.6065 (T)
	Day 56	8.83 (4.5)	(6.88, 10.77)	8.54 (5.41)	(6.44, 10.63)	0.8380 (T)
	Change from baseline at day 56	-2.35 (4.63)	(-4.35, -0.35)	-1.79 (5.51)	(-3.92, 0.35)	
	p value [#]	0.7382 (A)				
	p value*	0.0236 (T)		0.0980 (T)		
Sexual life	Day 0	1.65 (1.19)	(1.14, 2.17)	1.36 (1.16)	(0.91, 1.81)	0.3766 (T)
	Day 56	1.3 (0.88)	(0.93, 1.68)	1.39 (1.03)	(0.99, 1.79)	0.7456 (T)
	Change from baseline at day 56	-0.35 (1.3)	(-0.91, 0.21)	0.04 (1.4)	(-0.51, 0.58)	
	p value [#]	0.4867 (A)				
	p value*	0.2130 (T)		0.8937 (T)		
Social life	Day 0	1.48 (1.24)	(0.94, 2.01)	1.46 (1.17)	(1.01, 1.92)	0.9672 (T)
	Day 56	1.52 (1.08)	(1.05, 1.99)	1.39 (1.23)	(0.92, 1.87)	0.6957 (T)
	Change from baseline at day 56	0.04 (0.93)	(-0.36, 0.44)	-0.07 (1.27)	(-0.57, 0.42)	
	p value [#]	0.7373 (A)				
	p value*	0.8243 (T)		0.7691 (T)		
Productivity	Day 0	1.91 (1.44)	(1.29, 2.54)	1.89 (1.4)	(1.35, 2.43)	0.9599 (T)
	Day 56	1.52 (1.34)	(0.94, 2.10)	1.61 (1.4)	(1.07, 2.15)	0.8260 (T)
	Change from baseline at day 56	-0.39 (1.27)	(-0.94, 0.16)	-0.29 (1.36)	(-0.81, 0.24)	
	p value [#]	0.5402 (A)				
	p value*	0.1536 (T)		0.2750 (T)		
Total score	Day 0	41.13 (18.28)	(33.23, 49.03)	37.93 (18.24)	(30.86, 45.00)	0.5360 (T)
	Day 56	30.13 (16.08)	(23.18, 37.08)	31.93 (18.34)	(24.82, 39.04)	0.7144 (T)
	Change from baseline at day 56	-11 (12.8)	(-16.54, -5.46)	-6 (19.16)	(-13.43, 1.43)	
	p value [#]	0.3045 (A)				
	p value*	0.0004 (T)		0.1091 (T)		

Notes: [§]p-value was calculated using two sample t-test (T). ^{*}p-values were calculated using paired t-test (T). [#]p-values were calculated using ANCOVA with treatment as factor and baseline value, age and BMI as covariate vs placebo.

Abbreviations: CI, confidence interval; IWQOL-Lite-CT, Impact of Weight on Quality of Life-Lite-Clinical Trials Version; N, number of participants; mITT, modified intention to treat; SD, standard deviation.

Safety Outcomes

No significant change in safety parameters was observed during the study in the IP group compared to baseline and placebo. Only five adverse events were reported during the study in the IP group – 1) high SGPT and ALP levels, 2) dry cough, 3) high IL-6 levels, 4) high CRP levels, 5) high SGPT and SGOT levels. None of the adverse events were found related to the study products. No serious adverse event was reported during the study ([Table S2](#)) ([Supplementary Material](#)).

Discussion

Obesity has been purported as a global pandemic in recent times with a well-established relationship with oxidative imbalance. Studies have shown that individuals with higher BMI have increased productions of 8-isoprostane.^{6,35–37} The levels of 8-isoprostane are known to increase in response to oxidative stress.³⁸ Hence, 8-Isoprostane, which is considered as the most stable biomarker of oxidative stress was assessed in the serum of the participants enrolled in the study.³⁹ In the present study, participants with increased body weight and at a risk of metabolic syndrome showed a trend of increasing oxidative stress over next eight weeks. Dietary polyphenols (or flavonoids) serve as effective scavengers for free radicals and ROS. They possess the capacity to neutralize ROS or inhibit cellular oxidative stress, thereby preventing damage to biomolecules such as lipids, proteins, and DNA. This phenomenon is commonly referred to as the antioxidant effects of dietary polyphenols.⁴⁰ The IP consistent with this known antioxidant effect of polyphenols, helped to cease the increase of 8-isoprostane, whereas in the placebo group the oxidative stress continued to increase till the end of the study. Oxidative imbalance arises due to a deficiency in the antioxidant defense system (ADS) resulting from impaired production or distribution of antioxidants (AO) or surplus of ROS.⁴¹ These results of the present study are similar to the effect of Vitamin E, wherein 6-month supplementation led to the reduction in the 8-Isoprostane levels thus ceasing oxidative stress.⁴² In another study conducted by Askari et al, eight weeks of quercetin supplementation proved to be efficacious in reducing oxidative stress.³²

Obesity is known to disrupt the antioxidant tissue defenses and this disruption has been found to be related to a low intake of dietary phytochemicals and antioxidants that possess antioxidant capacity. An inverse correlation of oxidative stress with high BMI, waist circumference, waist-to-hip ratio and plasma oxidative stress was demonstrated in a study conducted by Vincent.⁴³ Polyphenols exhibit potential anti-obesity properties, with proposed mechanisms for weight management including the promotion of thermogenesis and increased energy expenditure,⁴⁴ inhibition of adipocyte differentiation and growth,⁴⁵ augmentation of lipolysis and induction of β -oxidation,⁴⁶ and reduction of appetite.⁴⁷ The anti-obesity effect of our present study is in line with the known effects of polyphenols like rutin and quercetin. After 56 days of intervention, a mean reduction in the BMI of 0.8 (0.61) kg/m² was seen in the LN-OS-22 study group as opposed to placebo where the mean BMI showed an increase by 0.06 (0.6) kg/m²; and this result of the IP group was statistically significant when compared to placebo as well as baseline ($p < 0.0001$). Also, at the end of the study, when compared with placebo, a statistically significant reduction was observed in the waist circumference ($p < 0.0001$), waist-to height ratio ($p = 0.0005$) as well as body weight ($p < 0.0001$) in the intervention group. Thus, the anthropometric endpoints of the present study proved to be positive in the LN-OS-22 study group.

Oxidative stress and the increased body weight can have a significant impact on the quality of life.⁴⁸ In the present study, a statistically significant change in the total scores of the IWQOL-Lite-CT was depicted after 56 days of intervention in the IP group ($p = 0.0004$). When compared to baseline, the IP group showed significant improvement in the physical impact, energy/stamina, emotions and self-confidence/self-esteem domains of the IWQOL-Lite-CT questionnaire. Thus, it can be concluded that the study group LN-OS-22 positively influenced the quality of life of individuals. No statistically significant results were demonstrated in the intervention group on inflammatory (CRP and IL-6), lipid peroxidation (MDA) as well as metabolic endotoxemia (LPS) biomarkers. This can be attributed to the small sample size of the study. Nevertheless, the investigational product LN-OS-22 proved to be efficacious in reducing the oxidative stress, body weight, BMI and waist circumference of individuals with overweight and obesity as well as improved the quality of life of individuals with overweight and obesity.

Conclusion

The present randomized clinical trial concludes that the LN-OS-22 has antioxidant potential thereby helping to maintain healthy body composition. It also has shown promise of reducing the risk of metabolic dysfunction by maintaining healthy waist to height ratio. The study product LN-OS-22 was found to be well tolerated.

Abbreviations

BMI, Body Mass Index; BP, Blood Pressure; CONSORT, Consolidated Standards Of Reporting Trials; CRP, C-Reactive Protein; EC, Ethics Committee; e-CRF, Electronic-Case Record Form; GMP, Good Manufacturing Practice; IP, Investigational Product; ICH-GCP, International Conference On Harmonization-Good Clinical Practice; IL, Interleukin; LASA, Longitudinal Aging Study Amsterdam (LASA); LPS, Lipopolysaccharide; MDA, Malondialdehyde; PR, Pulse rate; ROS, Reactive Oxygen Species; RNS, Reactive Nitrogen Species; ALP, Serum Alkaline Phosphatase; SGOT, Serum Glutamic-Oxaloacetic Transaminase; SGPT, Serum Glutamic-Pyruvic Transaminase; IWQOL-Lite-CT, Weight on Quality of Life-Lite-Clinical Trials Version.

Data Sharing Statement

The data used in the study is available on reasonable request from the corresponding author with due permission from the sponsor.

Acknowledgments

The authors are grateful to all the participants who volunteered for the study. The authors thank Layn Natural Ingredients, USA, for providing the study products and Vedic Lifesciences, India, for facilitating the study.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

Layn Natural Ingredients, USA provided financial support and investigational products for the trial.

Disclosure

Mr James Roza is an employee for Layn, outside the submitted work; and Layn Corp. will use this research to market this ingredient (SophorOx[®]) as a dietary supplement worldwide. The authors report no other conflicts of interest in this work.

References

1. Videla LA. Oxidative stress signaling underlying liver diseases and hepatoprotective mechanism. *World J Hepatic.* 2009;31(1):72–78. doi:10.4254/wjh.v1.i1.72
2. Laufs U, Wassmann S, Czech T, et al. Physical inactivity increases oxidative stress, endothelial dysfunction, and atherosclerosis. *Arterioscler Thromb Vasc Biol.* 2005;25(4):809–814. doi:10.1161/01.ATV.0000158311.24443.af
3. Aleksandrova K, Koelman L, Rodrigues CE. Dietary patterns and biomarkers of oxidative stress and inflammation: a systematic review of observational and intervention studies. *Redox Biol.* 2021;42:101869. doi:10.1016/j.redox.2021.101869
4. Sies H, Jones D. Oxidative Stress. *Encyclop Stress.* 2007;45–48. doi:10.1016/B978-012373947-6.00285-3
5. Sies H. Oxidative stress: a concept in redox biology and medicine. *Redox.* 2015;Biol(4):180–183. doi:10.1016/j.redox.2015.01.002
6. Keaney JF, Larson MG, Vasan R, et al. Obesity and systemic oxidative stress: clinical correlates of oxidative stress in the Framingham Study. *Arterioscler Thromb Vasc Biol.* 2003;23:434–439. doi:10.1161/01.ATV.0000058402.34138.11
7. Biswas SK. Does the Interdependence between oxidative stress and inflammation explain the antioxidant paradox? *Oxid Med Cell Longev.* 2016;2016:5698931. doi:10.1155/2016/5698931
8. Manna P, Jain SK. Obesity, oxidative stress, adipose tissue dysfunction, and the associated health risks: causes and therapeutic strategies. *Metabolic Syndr Rel Disord.* 2015;13(10):423–444. doi:10.1089/met.2015.0095

9. Chrysohoou C, Panagiotakos DB, Pitsavos C, et al. The implication of obesity on total antioxidant capacity in apparently healthy men and women: the ATTICA study. *Nutr Metab Cardiovasc Dis*. 2007;17(8):590–597. doi:10.1016/j.numecd.2006.05
10. Fernández-Sánchez A, Madrigal-Santillán E, Bautista M, et al. Inflammation, oxidative stress, and obesity. *Int J Mol Sci*. 2011;12(5):3117–3132. doi:10.3390/ijms12053117
11. Patel C, Ghanim H, Ravishankar S, et al. Prolonged reactive oxygen species generation and nuclear factor-kappaB activation after a high-fat, high-carbohydrate meal in the obese. *J Clin Endocrinol Metab*. 2007;92(11):4476–4479. doi:10.1210/jc.2007-0778
12. Jiang S, Liu H, Li C. Dietary regulation of oxidative stress in chronic metabolic diseases. *Foods*. 2021;10(8):1854. doi:10.3390/foods10081854
13. He X, Bai Y, Zhao Z, et al. Local and traditional uses, phytochemistry, and pharmacology of *Sophora japonica* L.: a review. *J Ethnopharmacol*. 2016;187:160–182. doi:10.1016/J.JEP.2016.04.014
14. Abd-Alla HI, Souguir D, Radwan MO. Genus *Sophora*: a comprehensive review on secondary chemical metabolites and their biological aspects from past achievements to future perspectives. *Arch Pharm Res*. 2021;44(11):903. doi:10.1007/S12272-021-01354-2
15. Ha AT, Rahmawati L, You L, Hossain MA, Kim JH, Cho JY. Anti-inflammatory, antioxidant, moisturizing, and antimelanogenesis effects of quercetin 3-O- β -D-glucuronide in human keratinocytes and melanoma cells via activation of NF- κ B and AP-1 pathways. *Int J Mol Sci*. 2021;23. doi:10.3390/IJMS23010433
16. Alizadeh SR, Ebrahimzadeh MA. Quercetin derivatives: drug design, development, and biological activities, a review. *Eur J Med Chem*. 2021;229. doi:10.1016/J.EJMECH.2021.114068
17. Bernini R, Velotti F. Natural polyphenols as immunomodulators to rescue immune response homeostasis: quercetin as a research model against severe COVID-19. *Molecules*. 2021;26(19):5803. doi:10.3390/MOLECULES26195803
18. Di Petrillo A, Orrù G, Fais A, Fantini MC. Quercetin and its derivatives as antiviral potentials: a comprehensive review. *Phytother Res*. 2022;36(1):266–278. doi:10.1002/PTR.7309
19. Shen P, Lin W, Deng X, et al. Potential implications of quercetin in autoimmune diseases. *Front Immunol*. 2021;12. doi:10.3389/FIMMU.2021.689044
20. Yi H, Peng H, Wu X, et al. The therapeutic effects and mechanisms of quercetin on metabolic diseases: pharmacological data and clinical evidence. *Oxid Med Cell Longev*. 2021;2021:1–16. doi:10.1155/2021/6678662
21. Shanmugasundaram D, Roza JM. Assessment of anti-inflammatory and antioxidant activity of quercetin–rutin blend (SophorOx™) – an *in vitro* cell based assay. *J Complement Integr Med*. 2022;19:637–644. doi:10.1515/jcim-2021-0568
22. Shanmugasundaram D, Roza JM. Assessment of anti-inflammatory and antioxidant activities of a proprietary preparation of quercetin-rutin blend (SophorOx™) in exercised rats. *ScientificWorldJournal*. 2024;2024:9063936. doi:10.1155/2024/9063936 PMID: 38371227; PMCID: PMC10874291.
23. Van't Erve TJ, Kadiiska MB, London SJ, Mason RP. Classifying oxidative stress by F2-isoprostane levels across human diseases: a meta-analysis. *Redox Biol*. 2017;12(March):582–599. doi:10.1016/j.redox.2017.03.024
24. Van't Erve TJ, Lih FB, Jelsema C, et al. Reinterpreting the best biomarker of oxidative stress: the 8-iso-prostaglandin F2 α /prostaglandin F2 α ratio shows complex origins of lipid peroxidation biomarkers in animal models. *Free Radic Biol Med*. 2016;95:65–73. doi:10.1016/j.freeradbiomed.2016.03.001
25. Castro AM, Macedo-de la Concha LE, Pantoja-Meléndez CA. Low-grade inflammation and its relation to obesity and chronic degenerative diseases. *Revista Médica Del Hospital General de México*. 2017;80(2):101–105. doi:10.1016/J.HGMX.2016.06.011
26. Rozjabeck H, Fastenau J, Laprade A, Sternbach N. Adult obesity and health-related quality of life, patient activation, work productivity, and weight loss behaviors in the United States. *Diabet Metabo Syndr obesit*. 2020;13:2049–2055. doi:10.2147/DMSO.S245486
27. Dehghani F, Sezavar Seyedi Jandaghi SH, Janani L, Sarebanhassanabadi M, Emamat H, Vafa M. Effects of quercetin supplementation on inflammatory factors and quality of life in post-myocardial infarction patients: a double blind, placebo-controlled, randomized clinical trial. *Phytother Res PTR*. 2021;35(4):2085–2098. doi:10.1002/ptr.6955
28. Han MK, Barreto TA, Martinez FJ, Comstock AT, Sajjan US. Randomised clinical trial to determine the safety of quercetin supplementation in patients with chronic obstructive pulmonary disease. *BMJ Open Respir Res*. 2020;7(1):e000392. doi:10.1136/bmjresp-2018-000392
29. Javadi F, Ahmadzadeh A, Egtesadi S, et al. The effect of quercetin on inflammatory factors and clinical symptoms in women with rheumatoid arthritis: a double-blind, randomized controlled trial. *J Am Coll Nutr*. 2017;36(1):9–15. doi:10.1080/07315724.2016.1140093
30. Boyle SP, Dobson VL, Duthie SJ, Hinselwood DC, Kyle J, Collins AR. Bioavailability and efficiency of rutin as an antioxidant: a human supplementation study. *Eur J Clin Nutr*. 2020;54:774–782. doi:10.1038/sj.ejcn.1601090
31. Liu ZH, Cai Y, He J. High serum levels of 8-OHdG are an independent predictor of post-stroke depression in Chinese stroke survivors. *Neuropsychiatr Dis Treat*. 2018;14:587–596. doi:10.2147/NDT.S155144
32. Askari G, Ghiasvand R, Feizi A, Ghanadian SM, Karimian J. The effect of quercetin supplementation on selected markers of inflammation and oxidative stress. *J Res Med Sci*. 2021;17(7):637–641.
33. Wiswedel I, Hirsch D, Kropf S, et al. Flavanol-rich cocoa drink lowers plasma F(2)-isoprostane concentrations in humans. *Free Radic Biol Med*. 2004;37(3):411–421. doi:10.1016/j.freeradbiomed.2004.05.013
34. O'Reilly JD, Mallet AI, McAnlis GT, et al. Consumption of flavonoids in onions and black tea: lack of effect on F2-isoprostanes and autoantibodies to oxidized LDL in healthy humans. *Am J Clin Nutr*. 2001;73(6):1040–1044. doi:10.1093/AJCN/73.6.1040
35. Furukawa S, Fujita T, Shimabukuro M, et al. Increased oxidative stress in obesity and its impact on metabolic syndrome. *J Clin Invest*. 2004;114(12):1752–1761. doi:10.1172/JCI21625
36. Morrow JD. Quantification of isoprostanes as indices of oxidant stress and the risk of atherosclerosis in humans. *Arterioscler Thromb Vasc Biol*. 2005;25(2):279–286. doi:10.1161/01.ATV.0000152605.64964.c0
37. Urakawa H, Katsuki A, Sumida Y, et al. Oxidative stress is associated with adiposity and insulin resistance in men. *J Clin Endocrinol Metab*. 2003;88(10):4673–4676. doi:10.1210/jc.2003-030202
38. Herasymchuk NM. 8-isoprostane as the main marker of oxidative stress. *Zaporozhskij Medicinskij Zhurnal*. 2018b;1(6). doi:10.14739/2310-1210.2018.6.146780
39. Meera S, Sarangarajan R, Rajkumar K. 8-Isoprostane: a salivary oxidative stress biomarker for oral submucous fibrosis and oral squamous cell carcinoma. *J Oral Maxillofac Pathol*. 2020;24(2):279–284. doi:10.4103/jomfp.JOMFP_235_19 PMID: 33456237; PMCID: PMC7802855.
40. Rudrapal M, Khairnar SJ, Khan J, et al. Dietary polyphenols and their role in oxidative stress-induced human diseases: insights into protective effects, antioxidant potentials and mechanism(s) of action. *Front Pharmacol*. 2022;13. doi:10.3389/fphar.2022.806470
41. Montuschi P, Barnes PJ, Roberts LJ. Isoprostanes: markers and mediators of oxidative stress. *THE FASEB Journal*. 2004;18(15):1791–1800. doi:10.1096/fj.04-2330rev

42. Sutherland WH, Manning PJ, Walker RJ, de Jong SA, Ryalls AR, Berry EA. Vitamin E supplementation and plasma 8-isoprostane and adiponectin in overweight subjects. *Obesity*. 2007;15(2):386–391. doi:10.1038/oby.2007.546
43. Vincent HK, Bourguignon CM, Taylor AG. Relationship of the dietary phytochemical index to weight gain, oxidative stress and inflammation in overweight young adults. *J Hum Nutr Diet*. 2010;23(1):20–29. doi:10.1111/j.1365-277X.2009.00987.x
44. Dulloo AG, Duret C, Rohrer D, et al. Efficacy of a green tea extract rich in catechin polyphenols and caffeine in increasing 24-h energy expenditure and fat oxidation in humans. *The American Journal of Clinical Nutrition*. 1999;70(6):1040–1045. doi:10.1093/ajcn/70.6.1040
45. Sy M, Yang HS, Sg S, et al. Cocoa polyphenols suppress adipogenesis in vitro and obesity in vivo by targeting insulin receptor. *Int J Obesity*. 2012;37(4):584–592. doi:10.1038/ijo.2012.85
46. Rupasinghe HV, Sekhon-Loodu S, Mantso T, Panayiotidis MI. Phytochemicals in regulating fatty acid β -oxidation: potential underlying mechanisms and their involvement in obesity and weight loss. *Pharmacol Ther*. 2016;165:153–163. doi:10.1016/j.pharmthera.2016.06.005
47. Boix-Castejón M, Herranz-López M, Gago AP, et al. Correction: hibiscus and lemon verbena polyphenols modulate appetite-related biomarkers in overweight subjects: a randomized controlled trial. *Food Funct*. 2018;9(7):4037. doi:10.1039/c8fo90028k
48. Kolotkin RL, Meter K, Williams GR. Quality of life and obesity. *Obes Rev*. 2001;2(4):219–229. doi:10.1046/j.1467-789x.2001.00040.x

Diabetes, Metabolic Syndrome and Obesity

Dovepress

Publish your work in this journal

Diabetes, Metabolic Syndrome and Obesity is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert opinion and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-journal>